

Cambridge University Press

978-052-1-11389-2 - The Insects: Structure and Function: Fifth Edition

Edited by R. F. Chapman, Stephen J. Simpson and Angela E. Douglas

Excerpt

[More information](#)

Part I

The head, ingestion, utilization and distribution of food

Cambridge University Press
978-052-1-11389-2 - The Insects: Structure and Function: Fifth Edition
Edited by R. F. Chapman, Stephen J. Simpson and Angela E. Douglas
Excerpt
[More information](#)

1


Head

REVISED AND UPDATED BY **STEPHEN J. SIMPSON**

INTRODUCTION

Insects and other arthropods are built up on a segmental plan, and their characteristic feature is a hard, jointed exoskeleton. The cuticle, which forms the exoskeleton, is continuous over the whole of the outside of the body and consists of a series of hard plates, the sclerites, joined to each other by flexible membranes, which are also cuticular. Sometimes the sclerites are articulated together so as to give precise movement of one on the next. Each segment of the body primitively has a dorsal sclerite, the tergum, joined to a ventral sclerite, the sternum, by lateral membranous areas, the pleura. Arising from the sternopleural region on each side is a jointed appendage.

In insects, the segments are grouped into three units, the head, thorax and abdomen, in which the various basic parts of the segments may be lost or greatly modified. Typical walking legs are only retained on the three thoracic segments. In the head, the appendages are modified for sensory and feeding purposes and in the abdomen they are lost, except that some may be modified as the genitalia and in Apterygota some pregenital appendages are retained. This chapter introduces the structures of the head (Section 1.1), neck (Section 1.2) and antennae (Section 1.3). Chapter 2 concerns the mouthparts and feeding.



The Insects: Structure and Function (5th edition), ed. S. J. Simpson and A. E. Douglas.
Published by Cambridge University Press. © Cambridge University Press 2013.

1.1 Head

The insect head is a strongly sclerotized capsule joined to the thorax by a flexible, membranous neck. It bears the mouthparts, comprising the labrum, mandibles, maxillae and labium, and also the antennae, compound eyes and ocelli. On the outside it is marked by grooves, most of which indicate ridges on the inside; some of these inflexions extend deep into the head, fusing with each other to form an internal skeleton. These structures serve to strengthen the head and provide attachments for muscles, as well as supporting and protecting the brain and foregut.

The head is derived from the primitive pre-oral and post-oral segments. Molecular studies of *Drosophila* suggest that there are seven head segments: labral, ocular, antennal, intercalary, mandibular, maxillary and labial. The last three segments are post-oral and are innervated by the three neuromeres of the subesophageal ganglion (see Chapter 20). They are often called the gnathal segments because their appendages form the mouthparts of the insect. The pre-oral segments are innervated by the brain, but their nature and number remains contentious. The protocerebrum (forebrain) innervates the compound eyes, the deutocerebrum (midbrain) innervates the antennae, and the labrum receives its innervation from the tritocerebrum (hindbrain) (see Chapter 20).

1.1.1 Orientation

The orientation of the head with respect to the rest of the body varies (Fig. 1.1). The hypognathous condition, with the mouthparts in a continuous series with the legs, is probably primitive. This orientation occurs most commonly in phytophagous species living in open habitats. In the prognathous condition the mouthparts point forwards and this is found in predaceous species that actively pursue their prey, and in larvae, particularly of Coleoptera, which use their mandibles in burrowing. In Hemiptera, the




Figure 1.1 Orientation of the head. (a) Hypognathous – mouthparts ventral, in a continuous series with the legs (grasshopper). (b) Prognathous – mouthparts in an anterior position (beetle larva). (c) Opisthorhynchous – sucking mouthparts with the proboscis extending back between the front legs (aphid).

elongate proboscis slopes backwards between the forelegs. This is the opisthorhynchous condition.
The mouthparts (labrum with a basal segment called the clypeus, mandibles, labium and maxillae)




Figure 1.2 Pre-oral cavity and some musculature. Diagrammatic vertical section through the head of an insect with biting and chewing mouthparts. Sclerites associated with the hypopharynx are black with white spots. Muscles attached to these sclerites move the hypopharynx (after Snodgrass, 1947).

enclose a cavity, the pre-oral cavity, which is divided into two sections by the hypopharynx (Fig. 1.2). The larger of these sections, the cibarium, is enclosed between the hypopharynx and the labrum/clypeus, with the true mouth (the opening to the foregut) at its inner end. Between the hypopharynx and the labium is a smaller cavity known as the salivarium, into which the salivary duct opens.

1.1.2 Rigidity

The head is a continuously sclerotized capsule with no outward appearance of segmentation, but it is marked by a number of grooves. Most of these grooves are sulci (singular: sulcus), marking lines along which the cuticle is inflected to give increased rigidity. The term “suture” should be retained for grooves marking the line of fusion of two formerly distinct plates. The groove which ends between the points of attachment of maxillae and labium at the back of the head is generally believed to represent the

line of fusion of the maxillary and labial segments, and is therefore known as the postoccipital suture.

Since the sulci are functional mechanical developments to resist the various strains imposed on the head capsule, they are variable in position in different species and any one of them may be completely absent. However, the needs for strengthening the head wall are similar in the majority of insects, so some of the sulci are fairly constant in occurrence and position (Fig. 1.3). The most constant is the epistomal (frontoclypeal) sulcus, which acts as a brace between the anterior mandibular articulations. At each end of this sulcus is a pit, the anterior tentorial pit, which marks the position of a deep invagination to form the anterior arm of the tentorium. The lateral margins of the head above the mandibular articulations are strengthened by a horizontal inflexion indicated externally by the subgenal sulcus. This sulcus is generally a continuation of the epistomal sulcus to the postoccipital suture. The part of the subgenal sulcus above the mandible is called the pleurostomal sulcus; the part behind the mandible is the hypostomal sulcus. Another commonly occurring groove is the circumocular sulcus, which strengthens the rim of the eye and may develop into a deep flange protecting the inner side of the eye. Sometimes this sulcus is connected to the subgenal sulcus by a vertical subocular sulcus; the inflexions associated with these sulci act as a brace against the pull of the muscles associated with feeding. The circumantennal ridge, marked by a sulcus externally, strengthens the head at the point of insertion of the antenna, while running across the back of the head, behind the compound eyes, is the occipital sulcus.

The areas of the head defined by the sulci are given names for descriptive purposes, but they do not represent primitive sclerites. Since the sulci are variable in position, so too are the areas which they delimit. The front of the head, the frontoclypeal area, is divided by the epistomal sulcus into the frons above and the clypeus below (Fig. 1.3). It is common to regard the arms of the ecdysial cleavage line as delimiting




Figure 1.3 Common lines or grooves on the insect head and the areas which they define (*italicized*) (modified after Snodgrass, 1960).

the frons dorsally, but this is not necessarily so. From the frons, muscles run to the pharynx, the labrum and the hypopharynx; from the clypeus arise the dilators of the cibarium. The two groups of muscles are always separated by the frontal ganglion and its connectives to the brain (Fig. 1.2). Dorsally the frons continues into the vertex and posteriorly this is separated from the occiput by the occipital sulcus. The occiput is divided from the postocciput behind it by the postoccipital suture, while at the back of the head, where it joins the neck, is an opening, the occipital foramen, through which the alimentary canal, nerve cord and some muscles pass into the thorax.

The lateral area of the head beneath the eyes is called the gena, from which the subgena is cut off below by the subgenal sulcus, and the postgena behind by the occipital sulcus. The region of the subgena above the mandible is called the pleurostoma, and that part behind the mandible is the hypostoma.

In hypognathous insects with a thick neck, the posterior ventral part of the head capsule is

membranous. The postmentum of the labium is contiguous with this membrane, articulating with the subgena on either side. The hypostomal sulci bend upwards posteriorly and are continuous with the postoccipital suture (Fig. 1.4a). In insects with a narrow neck, permitting greater mobility of the head, and in prognathous insects, the cuticle of the head below the occipital foramen is sclerotized. This region has different origins. In Diptera, the hypostomata of the two sides meet in the midline below the occipital foramen to form a hypostomal bridge that is continuous with the postocciput (Fig. 1.4b). In other cases, such as Hymenoptera and the water bugs *Notonecta* and *Naucoris* (Hemiptera), a similar bridge is formed by the postgenae, but the bridge is separated from the postocciput by the postoccipital suture (Fig. 1.4c). Where the head is held in the prognathous position, the lower ends of the postocciput fuse and extend forwards to form a median ventral plate, the gula (Fig. 1.4d), which may be a continuous sclerotization with the labium. Often the gula is




Figure 1.4 Sclerotization at the back of the head. Notice the position of the bridge below the occipital foramen with reference to the posterior tentorial pit. Membranous areas stippled, compound eyes cross-hatched. The names of areas defined by sulci are italicized (after Snodgrass, 1960). (a) Generalized condition, no ventral sclerotization; (b) hypostomal bridge (*Deromyia*, Diptera); (c) postgenal bridge (*Vespula*, Hymenoptera); (d) gular bridge formed from the postoccipital sclerites (*Epicauta*, Coleoptera).




Figure 1.5 Tentorium. Cutaway of the head capsule to show the tentorium and its relationship with the grooves and ridges of the head (after Snodgrass, 1935).

reduced to a narrow strip by enlargement of the postgenae and sometimes the postgenae meet in the midline, so that the gula is obliterated. The median ventral suture which is thus formed at the point of contact of the postgenae is called the gular suture.

In all insects, the rigidity of the head is increased by four deep cuticular invaginations, known as apodemes, which usually meet internally to form a brace for the head and for the attachment of muscles. The structure formed by these invaginations is called the tentorium (Fig. 1.5). Its two anterior arms arise from the anterior tentorial pits, which in Apterygota and Ephemeroptera are ventral and medial to the mandibles. In Odonata, Plecoptera and Dermaptera the pits are lateral to the mandibles, while in most higher insects they are facial at either end of the epistomal sulcus. The posterior arms arise from pits at the ventral ends of the postoccipital suture and they unite to form a bridge running across the head from one side to the other. In Pterygota the anterior arms also join up with the bridge, but the development of the tentorium as a whole is very variable. Sometimes a pair of dorsal arms arise from the anterior arms and they may be attached

to the dorsal wall of the head by short muscles. In Machilidae (Archaeognatha) the posterior bridge is present, but the anterior arms do not reach it, while in Lepismatidae (Thysanura) the anterior arms unite to form a central plate near the bridge and are joined to it by very short muscles.

1.1.3 Molting

Immature insects nearly always have a line along the dorsal midline of the head dividing into two lines on the face to form an inverted Y (Fig. 1.3). There is no groove or ridge along this line, and it is simply a line of weakness, continuous with that on the thorax, along which the cuticle splits when the insect molts (see Fig. 16.11). It is therefore called the ecdysial cleavage line, but has commonly been termed the epicranial suture. The anterior arms of this line are very variable in their development and position and, in Apterygota, they are reduced or absent. The ecdysial cleavage line may persist in the adult insect, and sometimes the cranium is inflected along this line to form a true sulcus. Other ecdysial lines may be present on the ventral surface of the head of larval insects.

1.2 Neck

The neck or cervix is a membranous region which gives freedom of movement to the head. It extends from the postocciput at the back of the head to the prothorax, and possibly represents the posterior part of the labial segment together with the anterior part of the prothoracic segment. Laterally in the neck membrane are the cervical sclerites. Sometimes there is only one, as in Ephemeroptera, but there may be two or three. In *Schistocerca* (Orthoptera) the first lateral cervical sclerite, which articulates with the occipital condyle at the back of the head, is very small. The second sclerite articulates with it by a ball and socket joint, allowing movement in all planes. Posteriorly it meets the third (posterior) cervical sclerite; movement at this joint is restricted to the vertical plane. The third cervical sclerite connects with the prothoracic episternum, relative to which it can move in all planes. Muscles arising from the postocciput and the pronotum are inserted on the cervical sclerites

(Fig. 1.6a) and their contraction increases the angle between the sclerites so that the head is pushed forward (Fig. 1.6b). A muscle arising ventrally and inserted onto the second cervical sclerite may aid in retraction or lateral movements of the head. Running through the neck are longitudinal muscles, dorsal muscles from the antecostal ridge of the mesothorax to the postoccipital ridge, and ventral muscles from the sternal apophyses of the prothorax to the postoccipital ridge or the tentorium. These muscles serve to retract the head onto the prothorax, while their differential contraction will cause lateral movements of the head. *Schistocerca* has 16 muscles on each side of the neck, each of which is innervated by several axons, often including an inhibitory fiber. This polyneuronal innervation, together with the versatility of the cervical articulations and the complexity of the musculature, permit movement of the head in a highly versatile and accurately controlled manner.




Figure 1.6 Neck and cervical sclerites of a grasshopper. (a) Seen from the inside to show the muscles (after Imms, 1957). (b) Diagrams showing how a change in the angle between the second and third cervical sclerites retracts or protracts the head (the first cervical sclerite is small and is not shown). Arrows indicate points of articulation.

1.3 Antennae

All insects possess a pair of antennae, but they may be greatly reduced, especially in larval forms. Among the non-insect Hexapoda, Collembola and Diplura have antennae, but Protura do not.

1.3.1 Antennal structure

The antenna consists of a basal scape, a pedicel and a flagellum. The scape is inserted into a membranous region of the head wall and pivoted on a single marginal point, the antennifer (Fig. 1.7a), so it is free to move in all directions. Frequently the flagellum is divided into a number of similar annuli joined to each other by membranes so that the flagellum as a whole is flexible. The term “segmented” should be avoided with reference to the flagellum of insects since the annuli are not regarded as equivalent to leg segments.

The antennae of insects are moved by levator and depressor muscles arising on the anterior tentorial arms and inserted into the scape, and by flexor and extensor muscles arising in the scape and inserted into the pedicel (Fig. 1.8a). There are no muscles in the flagellum, and the nerve which traverses the flagellum is purely sensory. This is the annulated type of antenna. In Collembola and Diplura the musculature at the base of the antenna is similar to that in insects, but, in addition, there is an intrinsic musculature in each unit of the flagellum (Fig. 1.8b), and, consequently, these units are regarded as true segments.

The number of annuli is highly variable between species. Adult Odonata, for example, have five or fewer annuli, while adult *Periplaneta* (Blattodea) have over 150, increasing from about 48 in the first-stage larva.

The form of the antenna varies considerably depending on its precise function (Fig. 1.7). Sometimes the modification produces an increase




Figure 1.7 Antennae. Different forms occurring in different insects. Not all to the same scale.