

Index

- absorption of alcohol, 19
 acamprosate, 209–210
 accidents, alcohol as risk factor, 63
 acute alcohol intoxication, 54
 acute pancreatitis, 58
 acute pharmacological effects of alcohol, 21–22
 adolescents, polydrug use, 96
 adult respiratory distress syndrome (ARDS), 62
 aggression, 79
 and crime, 45–47
 dealing with, 107–109
 Al-Anon, 12-step fellowship, 225
 alcohol, 17
 alcohol amblyopia, 68
 alcohol dehydrogenase (ADH), 20
 alcohol dependence
 (syndrome), 8
 clinical genesis of concept, 8–10
 collating evidence for, 151–152
 diagnosis of, 8, 14
 elements of, 10–13
 importance of understanding, 14–15
 neurology, 22–24
 time element, 14
 tolerance increase, 11
 validity, 10
 withdrawal symptoms, 11–13
 alcohol-induced amnesic episodes, 79–80
 alcohol-induced pseudo-Cushing's syndrome, 59
 alcohol-induced psychotic disorder with delusions, 78–79
 alcohol-related arrhythmias, 54–55
 alcohol-related brain damage, 67–68
 alcohol use disorders, 6
 alcohol dependence
 (syndrome), 8–10
 harmful drinking, 7–8
 hazardous drinking, 6–7
 alcohol withdrawal seizures, 63–64, 173–174
 alcoholic cardiomyopathy, 56
 alcoholic cerebellar degeneration, 64
 alcoholic cirrhosis, 57–58
 alcoholic dementia, 90
 alcoholic hallucinosis, 78
 alcoholic hepatitis, 57
 alcoholic ketoacidosis, 60
 alcoholic liver disease, 56–58
 alcoholic pellagra encephalopathy, 67
 Alcoholics Anonymous (AA), 218
 alternative organizations, 225–226
 as a pluralist spirituality, 229
 change processes, 221–223
 history of, 229–230
 meetings, 219–221
 other 12-step fellowships, 225
 research on, 224–225
 role of therapist in maximizing benefits of, 223–224
 variety of groups, 223
 alcoholism as a disease, 189–190
 alcohols, chemical group, 170
 aldehyde dehydrogenase 2 (ALDH2), 20
 amblyopia, 68
 amnesia, 79–80
 anaemia, 62
 anxiety disorder, 85–87
 post-traumatic stress disorder (PTSD), 87
 appointments, spacing of, 193
 arrhythmias, 54–55
 assessment, 143
 15-and 5-minute, 154–157
 audit of alcohol-related disabilities, 152–153
 case formulation, 161–163
 evidence for dependence, collating 151–152
 help-seeking and motivation, 153–154
 personality, 147, 149–150
 physical examination and lab tests, 154
 setting the tone, 144
 typical drinking day, analysis of, 148–151
 with partner/significant other/spouse, 157, 161
 see also history-taking
 ‘at risk’ drinking, 5
 ‘at risk’ population, 130
 attention deficit hyperactivity disorder (ADHD), 89
 AUDIT (Alcohol Use Disorders Identification Test), 134–135
 availability of alcohol, 27–28
 aversion therapy, 200–201
 behaviour contracting, 200
 behavioural self-control training (BSCT), 199–200
 benzodiazepines
 for withdrawal symptoms, 168, 170–171
 misuse of, 99–100
 to avert withdrawal seizures, 173
 withdrawal from, 176–177
 binge drinking, 6–7, 53
 biomarkers of alcohol consumption, 132–134
 bipolar disorder, 85
 blackouts, amnesic episodes, 79–80
 blood alcohol concentration (BAC), 133–134
 and higher tissue damage in women, 19
 levels affecting the CNS, 22
 scientific notation, 17–18
 blood disorders, 62
 blood pressure, rise in, 55
 BMI *see* brief motivational intervention (BMI)

- body temperature, effects of
 alcohol on, 22
- bone density, effects of alcohol
 on, 61
- brain
 effect of alcohol on CNS, 22
 neurotransmitters, 22, 24
- brain damage, 67–68
 cause and consequence of
 drinking, 90–91
 patients with cognitive
 impairment, 111–112
 to the unborn child, 69
- breath alcohol concentration
 (BrAC), 19–20, 170
- brief motivational intervention
 (BMI), 138–140
- Buddhism, 230
- CAGE questionnaire, 135
- cancer, 54, 62
- cannabis use, 89, 96, 98–99
- carbohydrate-deficient
 (desialylated) transferrin
 (CDT) test, 133
- ‘cardio-protective’ effect of
 alcohol, 55
- cardiomyopathy, 56
- cardiovascular disease, 54–56
- case-finding and detection,
 129–131, 135–136
- case formulation, 161–163
- causes of drinking problems,
 27
 application of knowledge of,
 32–33
 availability of the drug
 disorder and chronic stress, 32
 economic factors, 29–30
 genetics, 30–32
 norms and values, 28–29
- CBT *see* cognitive behavioural
 therapy
- central nervous system
 (CNS), effect of alcohol
 on, 22
- central pontine myelinolysis, 68
- cerebrovascular disease, 55
- children
 at risk, safeguarding,
 116–117
 experiences of, 43
 help for, 192–193
 physical injury to, 44
 psychological problems, 43–44
 resilience of adult children,
 44
- Christianity, 230, 233
- chronic pancreatitis, 58, 97
- cirrhosis, 57–58
- classification systems, 152
- clinical presentations, 105
- cognitively impaired patient,
 111–112
- cultural differences,
 113–114
 hospital ward, 109–110
 polydrug use, 110–111
 the older patient, 112–113
 the child at risk, 116–117
 the pregnant drinker, 110
 the public figure, 115–116
 the violent patient
 the young drinker, 105, 107
 working with intermediaries,
 114–115
- clues for detection of drinking
 problems, 130–131
- co-morbid psychiatric
 disorders, 81
 anxiety disorder, 85
 attention deficit hyperactivity
 disorder (ADHD), 89
 bipolar disorder, 85
 depression, 81
 eating disorders, 89
 epidemiology, 73
 pathological jealousy, 88
 personality disorder, 88
 post-traumatic stress
 disorder (PTSD), 87
 psychosis and
 schizophrenia, 89
 suicide, 84
 see also brain damage
- cocaine use with alcohol, 94, 98
- cognitive behavioural therapy
 (CBT), 86–87, 187, 198
 aversion therapy
 behavioural self-control
 training (BSCT), 199–200
 broader spectrum
 approaches, 202
 cognitive behavioural
 marital therapy, 200
 Community Reinforcement
 Approach (CRA), 199
 coping and social skills
 training, 198–199
 cue exposure, 200
 social behaviour and net-
 work therapy (SBNT), 199
- cognitively impaired patients,
 111–112
- COMBINE study, 209–210
- community detoxification, 166,
 168
- community reinforcement
 approach (CRA), 199
- control, loss of, 13, 239
- controlled drinking *see* normal
 drinking
- coping styles
 of partners, 41
 of patients, response
 to poor, 248
- coronary heart disease
 (CHD), 55
- covert sensitisation, 201
- craving, 13, 154
 and exposure to risky
 situations, 240
 anti-craving medication, 207
- crime, 45, 127
- cue exposure, 200
- cultural differences, 113
- cultural norms and drinking
 patterns, 29
- defences of patients
 defence of sickness,
 249–250
 romantic defence, 250–251
- degree of dependence, 15, 236
- delirium tremens (DTs), 75
 aetiology and course, 77
 possibilities of diagnostic
 confusion, 77–78
 surgical ward case history
 symptoms of, 75, 77
 treatment of, 173, 175
- delusions, 76, 78–79
- dependence *see* alcohol depend-
 ence (syndrome)
- depression, 81
 depressive illness, 81–82
 practical importance of
 diagnosing, 83–84
 relationship to drinking
 problems, 83
- detection of drinking
 problems, 129, 135
- detoxification
 community
 in-patient, 170, 173
- diagnosis, 129
 and polydrug use, 102
 classification systems, 152
 in the absence of withdrawal
 symptoms, 14
 laboratory tests, 131

	Index	257
diagnosis (<i>cont.</i>)		
practical conclusions on		
case-finding, 135–136		
screening questionnaires,		
134		
Disability Adjusted Life Years (DALYs), 50		
distribution of alcohol in body,		
19		
disulfiram, 206–207		
dopamine system, effects of		
alcohol on, 23–24		
drink-driving offences, 47		
drinking guidelines, 4, 6		
for pregnant women, 70,		
110, 138		
drinking problems, 3		
categories of alcohol misuse,		
6, 10		
elements of alcohol		
dependence, 10–13		
‘messy’ treatment of, 123,		
125		
population perspective,		
4, 6		
range of, 15–16		
settings where prevalent,		
127, 129		
varied manifestations of,		
3–4		
drug abuse <i>see</i> polydrug use		
drug treatment <i>see</i>		
pharmacotherapy		
DSM-IV-TR, 150		
criteria for alcohol		
dependence, 8		
early intervention, importance		
of, 126–127		
eating disorders, 89		
economic factors causing		
drinking problems,		
29–30		
elderly people, 128		
electrolyte imbalance, 175		
employment problems, 45		
encephalopathy		
alcoholic pellagra		
encephalopathy, 67		
hepatic encephalopathy,		
68–69		
Wernicke’s encephalopathy,		
65–66, 172		
endocrine and metabolic		
disorders, 59–60		
endogenous opioid system,		
effects of alcohol on, 24		
engagement, coping strategy of		
partners, 42		
environmental factors,		
31–32		
epidemiological studies,		
co-morbidity, 73–74		
ethanol, chemical properties,		
17		
excretion of alcohol, 19–20		
faith <i>see</i> religion		
familial causes of drinking		
problems, 30–32		
family complications, 39		
children, 43–44		
partners, 39, 43		
family history-taking,		
145–146		
family interventions, 33,		
192–193, 202		
fatty liver, 57		
fetal alcohol spectrum		
disorders (FASD),		
69–70		
financial problems, 30, 47–48		
FRAMES mnemonic for BMI,		
139–140		
GABA (gamma-aminobutyric		
acid) receptor complex,		
23		
gamma-glutamyl transferase		
(GGT), 133		
gastritis, 58–59		
gastroenterological disorders,		
56, 59		
gastrointestinal tract, effect of		
alcohol on, 22		
glutamate system, effect of		
alcohol on, 23		
gout, 60		
group therapy, 193, 203–204		
haematological effects of		
alcohol, 85		
hallucinations, 74–76, 78		
harmful drinking, 7–8		
hazardous drinking, 6–7		
health education, 33		
heart disease, 55		
help-seeking, 50, 52		
history of, 151		
pathways, 121–122		
settings, 121		
hepatic alcohol metabolism,		
20		
hepatic encephalopathy, 68		
hepatitis, 57		
hereditary issues, 30–32		
heroin use, 99		
history-taking, 143–144		
background and family		
history, 145–146		
drinking history,		
evolution of problems,		
147–148		
essential framework, 154		
from partner/spouse,		
39–40		
initial interview, setting		
tone of, 142		
of past help-seeking		
of typical recent heavy		
drinking day, 148		
personal history, 146		
personality, 147		
previous illnesses, 147		
time spent on, 144–145		
hospital wards, clinical		
presentation, 109–110		
housing problems, 48		
hyperlipidaemia, 60		
hypertension, 55		
hypoglycaemia, 59–60		
hypomania, 85		
ICD-10, 152		
criteria for alcohol		
dependence syndrome,		
8		
immune system suppression,		
60		
individual behavioural analysis,		
240		
infections, respiratory, 62		
in-patient treatment of		
withdrawal, 169		
alcohol withdrawal		
seizures, 63–64		
delirium tremens, 173–175		
drug treatments, 170–173		
monitoring procedures,		
169–170		
insight-oriented psychotherapy,		
203		
intake levels leading to physical		
harm, 52–53		
intoxication, emergency		
treatment, 54		
iron deficiency, 62		
Islam, 230		
jealousy, pathological, 88		
journey syndrome, 80		

- ketoacidosis, 60
 kidney, diuretic effect of
 alcohol, 22
 Korsakoff's syndrome, 66–67
- laboratory tests, 131
 liver disease
 alcoholic, 56–58
 counselling patients with,
 50–52
 liver function tests, 133
- macrocytosis, 62
 male hypogonadism, 59
 Mallory–Weiss Syndrome,
 59
mania à potu, 79
 Marchiafava–Bignami disease,
 68
 marital problems, 40–41
 marital therapy
 MAST questionnaire, 135
 mean corpuscular volume
 (MCV), 132–133
 medical conditions *see* physical
 complications
 medical management (MM),
 140
 COMBINE study, 209
 medical negligence, 251–252
 medication *see*
 pharmacotherapy
 memory loss, 79–80
 mental health problems,
 dealing with, 187–188
 mental health services, 128–129
 mental illness
 and normal drinking,
 237–238
see also psychiatric disorders
 metabolism of alcohol,
 20–21
 methanol poisoning, 54
 microsomal ethanol-oxidizing
 system (MEOS), 20
 mood disturbance, 12, 191
see also depression
 motivation to change, 166, 182,
 184
 and relationship with
 therapist, 181
 motivational enhancement
 therapy (MET), 197–198
 motivational interviewing,
 186–187
 muscles, effect of alcohol on,
 61
- see also* Alcoholics
 Anonymous (AA)
 myopathy, 61
 cardiomyopathy, 56
- naltrexone, 207, 209
 COMBINE study, 209
 Native American religion,
 230
 neurological disorders
 alcohol amblyopia, 68
 alcoholic cerebellar
 degeneration, 64
 alcoholic pellagra
 encephalopathy, 67
 brain damage, 67–68
 central pontine
 myelinolysis, 68
 hepatic encephalopathy,
 68–69
 Korsakoff's syndrome,
 66–67
 Marchiafava–Bignami
 disease, 68
 peripheral neuropathy, 64
 seizures, 63–64
 Wernicke–Korsakoff
 syndrome, 65
 Wernicke's encephalopathy,
 65
- neurotransmitters, effect of
 alcohol, 22, 24
 neutropaenia, 62
 nicotine use, link to alcohol use
 disorders, 97–98
 non-medical use of
 prescription drugs
 (NMUPD), 101
 non-specialist settings,
 126–127
 brief motivational
 intervention, 138–140
 case-finding and
 detection, 129–131,
 135–136
 differences from specialist
 settings, 129
 information provision, 138
 in which drinking
 problems are prevalent,
 127, 129
 laboratory tests, 131
 linking to specialist care,
 140
 medical management, 140
 screening questionnaires,
 134
- triage and intervention, 136,
 138
see also specialist treatment
 normal drinking, 4–6
 and exposure to risky
 situations, 240–241
 involving the partner, 241
 limits on goal choice, 242
 monitoring of progress and
 treatment termination,
 241
 self-control strategies, 238,
 240
 sobriety as first step, 238
 versus abstinence, 242
 working towards, 236,
 238
- norms and values, causal
 factors, 28–29
- opiates
 withdrawal from, 177
 opioid system, effect of alcohol
 on, 24
 over-the-counter (OTC)
 medicines, 96–97
 overdose risks, 61
- parental alcoholism, effects
 on children, 43–44,
 116–117
 partners of drinkers, 39
 account of patient's drinking
 problems, 159, 161
 both partners with drinking
 problem, 41
 coping styles of, 160
 initial assessment with,
 157–158
 personal history, 158–159
 relationship problems,
 40–41
 therapeutic work with,
 192
 'pathological intoxication', 79
 pathological jealousy, 88
 peptic ulceration, 58–59
 percentage alcohol by volume
 (%ABV), 19
 peripheral neuropathy, 64
 personality, 147, 149–150
 personality disorder, 88
 pharmacology, 17, 19
 absorption and distribution,
 19
 acute effects of alcohol,
 21–22

	Index	259
pharmacology (<i>cont.</i>)		
brain sites involved, 22–24		
excretion and metabolism, 19, 21		
of alcohol (ethanol), 17		
reinstatement after abstinence, 152		
pharmacotherapy		
anti-craving medications, 207–210		
for anxiety disorder, 86		
for out-patient detoxification, 168		
for psychiatric co-morbidity, 210		
for relapse prevention, 205, 207		
physical complications, 50, 52–53		
accidents and trauma, 63		
acute alcohol intoxication and coma, 154		
alcoholic muscle, skin and bone disease, 61		
and help-seeking, 50–52		
and the immune system, 60		
cancers, 54		
cardiovascular disease, 54–56		
endocrine and metabolic disorders, 59–60		
fetal alcohol spectrum disorders (FASD), 69–70		
gastroenterological disorders, 56–59		
haematological effects, 62		
neurological disorders, 63, 69		
respiratory disease, 62		
surgery, 63		
policy making, 33		
polydrug use, 94, 97		
practical implications, 101–103		
specific combinations, 97, 101		
post-traumatic stress disorder (PTSD), 87		
practical application of knowledge about causes, 32–33		
pregnancy and drinking, 110, 128		
and fetal damage, 69–70		
medically supervised withdrawal, 171		
safe drinking guidelines, 70		
screening instruments, 135		
primary care setting, 128		
Project MATCH, treatment study, 204–205		
pseudo-Cushing's syndrome, 59		
psoriasis, 61		
psychiatric disorders, 73		
alcohol-induced disorders, 74, 80		
co-morbid disorders, 81, 91		
epidemiological studies, 73–74		
general implications, 91		
psychiatric services, 128–129		
psychodynamic psychotherapy, 203		
psychosis, 89		
public figures, 115–116		
questionnaires		
alcohol dependence, 152		
measuring intensity of problem experience and craving, 153		
screening tools for non-medical settings, 134		
questions for detection of drinking problems, 130		
readiness of patient to change, 183–184		
reinstatement after abstinence, 13, 152		
relapse		
dealing positively with, 190–191		
prevention therapy, 201–202		
reinstatement after abstinence, 13, 152		
relapse prevention therapy, 201		
relationships		
effect of drinking on, 40–41		
marital therapy, 198		
<i>see also</i> therapeutic relationship		
religion		
and individual therapy, 233–234		
attitudes to addiction, 230–231		
categories of Faith-Based Organisations, 233		
definitions, 228–229		
reducing alcohol consumption, 29		
treatment approaches, 233		
respiratory disease, 62		
rewards of sobriety, 188–189		
salience of drinking, 10–11, 151		
schizophrenia, 89		
and alcoholic hallucinosis, 78, 245		
scientific notation, blood alcohol concentrations, 17–18		
screening, 134		
laboratory tests, 131		
questionnaires, 134		
time-limited assessments		
utility of different procedures, 135–136		
seizures, 64		
alcohol withdrawal, 63–64, 175–176		
treatment to avert, 173		
self-control strategies, 238–240		
sensible drinking guidelines <i>see</i> drinking guidelines		
sensitivity of laboratory tests, 131–132		
skeletal muscle, effects of alcohol on, 61		
skin disorders, 61		
smoking and alcohol use, 97–98		
social behaviour and network therapy (SBNT), 199		
social complications, 36		
difficulty of pinpointing, 36–37		
family and children, 39, 44		
financial and housing stability, 47–48		
individual differences, 37–38		
moderated by person's resources, 38–39		
social contexts, selection of, 38		
violence and other crime, 45–47		
work and education, 127		
social skills training, 198–199		
specialist treatment, 197		
anti-craving medication, 207, 209		
cognitive behavioural therapy (CBT), 198, 202		
COMBINE study, 209		
differences from non-specialist, 127		
drug therapy for relapse		

- specialist treatment (*cont.*)
 prevention, 203–205
 effectiveness of, 210
 implications for real world, 211
 motivational enhancement therapy, 197–198
 other psychosocial treatment modalities, 202
 Project MATCH (USA), 204–205
 referral to, 140–141
 research evidence base for, 197
 resistance to referral, 137
 United Kingdom Alcohol Treatment Trial (UKATT), 205
see also non-specialist settings
- spirituality, 227
 and individual therapy, 233–234
 definition of, 227–228
 nature of drinking problems, 231–232
 origins of AA, 229–230
 philosophy and practice of AA, 232
- spouse *see* partners of drinkers
- standard drinks, 5, 19
- stepped care model, 122
 resistance to, 136–138
- stress
 cause of drinking problems, 30, 32
 post-traumatic stress disorder (PTSD), 87
- stroke, 55
- subjective awareness of
 compulsion to drink, 13
- suicide risk, 131
- support, importance of, 167–168, 170, 238
- surgical complications, 63
- surgical ward, clinical presentation, 109–110
- systematic reviews, 211
- tests, laboratory, 131, 134
- therapeutic difficulties, 244
 absolution expected by patient, 250–251
 losing the balances, 244–248
- medical negligence, 251–252
- poor coping strategies, 248, 250
- positive response to, 252–253
- therapeutic relationship, 181–182
- therapeutic work, 185, 187
 rewards of sobriety, 188–189
see also normal drinking; specialist treatment; withdrawal states and treatment
- thiamine (vitamin B₁)
 and Korsakoff's syndrome, 66–67
 for Wernicke's encephalopathy, 65, 172
 thrombocytopaenia, 62
- tolerance, coping strategy of partners, 42
- tolerance to alcohol, 151
- topiramate, 210
- trajectories of help-seeking and treatments, 121–125
- transient hallucinatory experience, 74–76
- trauma, post-traumatic stress disorder (PTSD), 87
- treatment, 'basic work', 180
 allied problems, 187–188
 appointments, spacing of, 193
 check-list, 194–195
 duration and termination of, 193–194
 family work, 39
 group therapy, 193
 motivation of patient, 182, 184
 patient work, guiding principles, 184–185
 relapse, dealing positively with, 190–191
 relationship with therapist, 181–182
- tremor, withdrawal symptom, 11–12, 76
- triage process, 136, 138
- 'Twelve Steps', AA, 220–221, 232
- Type II diabetes, 59
- typical drinking day, 148
- 'unit of alcohol', UK, 20
- United Kingdom Alcohol Treatment Trial (UKATT), 205
- values and norms, causal factors, 28–29
- 'very important' patients, 115–116
- violence
 heavy drinking and crime, 45, 47
 individuals with schizophrenia, 89
 in-patients who are drunk, 107–109
- vitamin B deficiency
 neurological conditions caused by, 64, 67
 thiamine treatment for, 175
- Wernicke–Korsakoff syndrome, 65
- Wernicke's encephalopathy, 65–66
 vitamin treatment, 175
- withdrawal, coping strategy of partners, 42
- withdrawal states and treatment, 165–166
 alcohol withdrawal seizures, 63–64, 175–176
 case history, 176
 community management, 166
 delirium tremens, 173–175
 drugs other than alcohol, 176–177
 in-patient setting, 169, 173
 withdrawal symptoms, 11–12, 151–152, 177
 absence of, impact on diagnosis, 14
 avoidance of by further drinking, 12–13, 152
 out-patient management, 167
 workplace problems, 127
- young people, 105
 'at risk' drinking, 5
 intervention with, 107
 polydrug use, 110–111
 possibilities for growth, 105–106
 special difficulties, 106–107
- Zieve's syndrome, 58