

Cambridge University Press

978-0-521-17229-5 - Fundamentals of Mid-Tertiary Stratigraphical Correlation

F. E. Eames, F. T. Banner, W. H. Blow and W. J. Clarke

[Table of Contents](#)[More information](#)

## CONTENTS

### PART 1: REVIEW AND REVISION

BY F. E. EAMES, F. T. BANNER, W. H. BLOW & W. J. CLARKE

I	Introduction . . . . .	<i>page</i> 2
II	Acknowledgments . . . . .	2
III	Type regions for stages of the Tertiary . . . . .	2
IV	Assessment of fossil faunas for dating and correlation . . . . .	3
V	Dating of beds within the limits Oligocene and Aquitanian in the Old World . . . . .	4
	A General Remarks . . . . .	4
	B The Oligocene and Aquitanian marine molluscan faunas of parts of Europe and their bearing on correlation (by Dr L. R. Cox) . . . . .	5
	C Correlation of the type stages with their equivalents in the Mediterranean Region . . . . .	10
VI	Correlation of Oligocene and Lower Miocene stages of Europe and the Mediterranean Region with successions in the Middle East, East Africa, and the Indo-Pacific Region . . . . .	13
VII	Consequent commentary on and minor amendments to the age of some Tethyan Beds, and remarks on <i>Lepidocyclusina</i> . . . . .	15
VIII	The dating of marine so-called 'Oligocene' beds in the Central American Region . . . . .	22
	A Introduction . . . . .	22
	B Historical . . . . .	22
	C The type Vicksburgian . . . . .	23
	D The fauna of the Antigua Limestone . . . . .	26
	E Other closely related coral faunas in the Central American Region . . . . .	27
	F The evidence of the Miogypsinids and <i>Orbulina</i> . . . . .	27
	G Brief review of significant faunas from important parts of the Central American Region . . . . .	31
	H Palaeogeography and conclusions . . . . .	48
IX	General conclusions . . . . .	49

## CONTENTS

Appendix I. Recognition of the genus <i>Palaeonummulites</i> Schubert, 1908	page 50
Appendix II. Revision of the genus <i>Pliolepidina</i> H. Douvillé, 1915 . . . . .	50
Selected references . . . . .	52
Plates I-VII . . . . .	<i>following page</i> 60

**PART 2: THE MID-TERTIARY  
(UPPER EOCENE TO AQUITANIAN)  
GLOBIGERINACEAE**

BY W. H. BLOW & F. T. BANNER

I Introduction . . . . .	61
II Acknowledgments . . . . .	62
III The Tanganyikan Middle Tertiary (Upper Eocene to Lower Miocene) . . . . .	62
A General geology . . . . .	62
B General palaeontology . . . . .	66
C Planktonic foraminiferal biozones . . . . .	67
IV Comparison of the Globigerinaceae of the Oligocene and Aquitanian in the Tethyan, European and Caribbean Regions . . . . .	69
V The stratigraphical position of the 'San Fernando Formation', Southern Trinidad . . . . .	77
VI Systematic palaeontology . . . . .	81
<i>Cassigerinella</i>	
<i>C. chipolensis</i>	
<i>Globigerina</i>	
<i>G. ampliapertura ampliapertura</i>	
<i>G. ampliapertura euapertura</i>	
<i>G. angulisuturalis</i>	
<i>G. angustumibilicata</i>	
<i>G. linaperta linaperta</i>	
<i>G. linaperta pseudoeocaena</i>	
<i>G. officinalis</i>	
<i>G. oligocaenica</i> sp.nov.	
<i>G. ouachitaensis ouachitaensis</i>	
<i>G. ouachitaensis ciperoensis</i>	
<i>G. ouachitaensis gnaucki</i> subsp.nov.	
<i>G. praebulloides praebulloides</i>	
<i>G. praebulloides leroyi</i> subsp.nov.	
<i>G. praebulloides occlusa</i> subsp.nov.	
<i>G. pseudoampliapertura</i> sp.nov.	
<i>G. senilis</i>	
<i>G. tripartita tripartita</i> emended	
<i>G. tripartita tapuriensis</i> subsp.nov.	
<i>G. turritilina turritilina</i> sp.nov. et subsp. nov.	
<i>G. turritilina praeturritilina</i> subsp.nov.	
<i>G. yeguaensis yeguaensis</i>	
<i>G. yeguaensis pseudovenezuelana</i> subsp. nov.	
<i>G. aff. yeguaensis</i>	
<i>Globigerinita</i> emended	
<i>G. africana</i> sp.nov.	
<i>G. dissimilis dissimilis</i>	
<i>G. dissimilis ciperoensis</i> subsp.nov.	
<i>G. globiformis</i> sp.nov.	
<i>G. howei</i> sp.nov.	
<i>G. martini martini</i> sp.nov. et subsp.nov.	
<i>G. martini scandretti</i> subsp.nov.	
<i>G. pera</i>	
<i>G. unicava unicava</i>	
<i>G. unicava primitiva</i> subsp.nov.	
<i>Globigerinoides</i>	
<i>G. quadrilobatus primordius</i> subsp.nov.	
<i>Globoquadrina</i>	

## CONTENTS

<i>G. dehiscens praedehiscens</i> subsp.nov.	<i>G. suteri</i>
<i>Globorotalia (Turborotalia)</i>	<i>Globigerapsis</i>
<i>G. (T.) centralis</i>	<i>G. tropicalis</i> sp.nov.
<i>G. (T.) cerro-azulensis</i>	<i>G. semi-involuta</i>
<i>G. (T.) hansbollii</i> sp.nov.	<i>Globigerinatheka</i>
<i>G. (T.) increbescens</i>	<i>G. lindiensis</i> sp.nov.
<i>G. (T.) opima opima</i>	<i>Hantkenina</i>
<i>G. (T.) opima nana</i>	<i>H. alabamensis</i>
<i>G. (T.) permicra</i> sp.nov.	<i>H. primitiva</i>
<i>G. (T.) postcretacea</i>	<i>Cribrohantkenina</i>
<i>Truncorotaloides</i>	<i>C. danvillensis</i>
<i>T. rohri</i>	<i>Pseudohastigerina</i>
<i>Turborotalita</i> gen.nov.	<i>P. micra</i>
<i>Globorotaloides</i>	
VII The evolution of some Upper Eocene to Lower Miocene	
Globigerinaceae . . . . . page 130	
References . . . . .	146
Plates VIII–XVII . . . . .	following page 152
Indexes (to both Parts) . . . . .	153

## FIGURES

1 Correlation of Oligocene and Lower Miocene stages of Europe and the Mediterranean with the East . . . . .	12
2 Ranges of some stratigraphically restricted microfossils in the Tertiary of the Far East . . . . .	14
3 Correlation of some mid-Tertiary beds of the Gulf States . . . . .	32
4 Correlation of some Tertiary beds in the Indo-Pacific province*	
5 Correlation of some Miocene Formations*	
6 Map of East Africa . . . . .	64
7 Geological map of the Lindi area showing type localities of the new biozones . . . . .	65
8 Correlation of the biozones and the geological successions in East Africa and Trinidad . . . . .	70
9 Some stratigraphically significant species from Europe, East Africa and America . . . . .	74

\* Figures 4 and 5 are available for download from  
[www.cambridge.org/9780521172295](http://www.cambridge.org/9780521172295)

## CONTENTS

10 Diagrams illustrating the descriptive morphological terminology of the Globigerinidae . . . . .	page 82
11 Variation in the bullae and apertural modifications in some Globigerinidae . . . . .	103
12 The evolution of the <i>Globorotalia opima</i> → <i>Globigerina ampliapertura</i> and the <i>Globorotalia centralis</i> → <i>Globigerina pseudoampliapertura</i> lineages . . . . .	131–5
13 The evolution of the <i>Globigerina praebulloides</i> → <i>Globigerinoides quadrilobatus</i> lineage . . . . .	137
14 Specimens illustrating evolutionary transition from <i>Globigerina praebulloides occlusa</i> to <i>Globigerinoides quadrilobatus primordius</i> . . . . .	138
15 The evolution of the <i>Globigerina officinalis</i> → <i>Globigerina ouachitaensis</i> lineage . . . . .	140
16 Specimens illustrating the morphological transition from <i>Globigerina officinalis</i> to <i>G. angustumbilicata</i> . . . . .	141
17 The evolution of the <i>Globigerina yeguaensis</i> → <i>Globigerina tripartita</i> lineage . . . . .	143
18 Specimens illustrating the morphological transition from <i>Globigerina yeguaensis yeguaensis</i> to <i>G. tripartita tripartita</i> . . . . .	144
19 Specimens illustrating the development of complex cibrate accessory apertures within the porticus of <i>Cribrohantkenina danvillensis</i> . . . . .	145
20 Ranges of Upper Eocene to Lower Aquitanian Globigerinaceae . . . . .	<i>preceding page</i> 153