

Index

- Aβ afferents
 - activation in
 - cutaneous reflexes, 365
 - RII reflex, 365
 - RIII reflex, 346–347
 - withdrawal reflexes, 346–347
 - conduction velocity, 365
 - electrical threshold, 365
- Aδ fibres
 - activation in
 - RIII reflex, 346
 - withdrawal reflexes, 346
 - conduction velocity, 346
 - electrical threshold, 346
- abdominal reflexes
 - central delay, 347
 - methodology, 341
 - trunk skin reflexes, organisation, 349
 - upper motoneurone lesions, 380
- acetylcholine
 - neuromuscular junction, 138
 - recurrent collaterals, 138
- activity-dependent hyperpolarisation, fibres
 - Ia afferents, 11, 72, 298
 - motor axons, 8, 11
- activity-dependent hyperpolarisation, motoneurons
 - (afterhyperpolarisation, AHP)
 - paired H reflex technique, 144, 147, 161
 - PSTH, discharge-related stimulation, 30–31, 50, 402
 - silent period, ongoing EMG/PSF/PSTH, 27, 33–35
- afferent feedback
 - delays, deleterious effects, 128, 482, 484–485
 - FRA hypothesis, 338, 451

internal vs. external, 482, 485	afferent feedback, TA, 519
role in	feedforward control, 518–520
cortical reciprocal inhibition, 493	visco-elastic stiffness, ankle, 517
corticospinal neurone excitability, 486	balance, standing
propriospinal (C3–C4) transmission, 422–425	perturbations, role of
quiet standing, multi-sensory, 516–517	cutaneous reflexes, 376
SICI, 495	feedback reflexes, 520–522
voluntary contraction	functional stretch reflex, 521–522
deafferentation, 128, 451	group II excitation (MLSR), 271–273, 521
multi-sensory, 453–454	Ia excitation (SLSR), 521
afferent feedback (spindle) in	long-latency responses, antagonist, 522
clonus, 129	presynaptic inhibition of Ia terminals, 319
concentric contraction, 127	reciprocal Ia inhibition, 202–204
kinaesthetic sensations, role in, 128	transcortical Ia excitation (LLSR), 521
lengthening contraction, 127	plasticity
motor learning, role in, 128	childhood, 547
quiet standing, 519	elderly, 524–525, 548–549
reciprocal Ia inhibition, 458–459	restoration after
steady contraction, 125, 128, 450–451	pitch perturbations, 520–523
tremor, Parkinson’s disease, 130	pitch-roll perturbations, 523–524
unloaded concentric contraction, antagonists, 127	training, changes in
unstable stance, 520	corticospinal excitability, 555
walking, 82, 128, 274–276, 279, 527–528	feedback reflexes, 548–549
alternative pathways in	presynaptic inhibition of Ia terminals, 550
FRA (short-latency) responses, walking, 362	transcortical Ia excitation, 524
FRA pathways (cat), 338	balance, walking
muscle synergies, 470–471	perturbations, changes in
amputation, corticospinal plasticity, 555–556	group II excitation (MLSR), 276–277, 530
amyotrophic lateral sclerosis (ALS), changes in	Ia excitation (SLSR), 82, 529–530
presynaptic inhibition of Ia terminals, 322	transcortical cutaneous excitation, 377–379, 531
recurrent inhibition, 170	transcortical Ia excitation (LLSR), 530
anastomosis, peripheral nerve, corticospinal plasticity, 556	restoration after
ankle strategy, standing, pitch perturbations, 522–523	slipping, 532
anticipatory responses	stepping down, 532
efference copy, possible role of, 482	stumbling, 277, 531–532
postural LLSRs, 477–478	ballistic contraction
arm cycling, modulation of	corticospinal involvement, 466, 492, 493
corticospinal neurone excitability, 490	motoneurons, orderly recruitment, 448
cutaneomuscular responses (deltoid), 374–376	preparatory period, presynaptic inhibition of Ia terminals, 453
presynaptic inhibition of Ia terminals, 316	quiet standing, predictive ballistic burts, 518–519
attention-demanding task, changes in	recurrent inhibition, 160, 161
gamma (γ) activity, 124–125	reticulospinal involvement, initiation, 492
non-reciprocal group I inhibition (wrist), 464	triphasic EMG pattern, 466
autonomic dysreflexia, SCI, 552, 573	beta (β) motoneurons, 113
Babinski response	contraction agonist, 113, 127
methodology, 341	bicycling, modulation of
upper motoneurone lesions, 381	corticospinal neurone excitability, 490
balance, quiet standing, role of	group II excitation (MLSR), 277
	presynaptic inhibition of Ia terminals, 321

582 Index

- biomechanics, elements of, in
 - multijoint movements, 447–448
 - posture, pitch-roll perturbations, 524
 - quiet standing, 517
 - walking, 525–526
- C3–C4 propriospinal neurones. *See* propriospinal relay
- caveats and limitations in studies of
 - corticospinal neurone excitability, 486, 491
 - cutaneous reflexes, 343–344
 - gamma (γ) drive, 114–115, 117–118
 - group II excitation, 260–261
 - Ia excitation (monosynaptic), 69, 71
 - Ib inhibition, 224–225
 - presynaptic inhibition of Ia terminals, 295–298, 299–300, 302, 321–322
 - propriospinal excitation (C3–C4), 398–400
 - propriospinal excitation (lumbar), 432
 - reciprocal Ia inhibition, 185–187
 - recurrent inhibition, 140–141, 146–147, 152
 - SICI, 38, 45, 494
 - withdrawal reflexes, 343–344
- caveats and limitations in techniques
 - coherence analyses, 51
 - F wave, 22–23
 - H reflex, 8–17
 - microneurography, 117–118
 - motor cortex stimulation, 38–39
 - ongoing EMG modulation, 27
 - paired H reflex technique, 146–147
 - PSF, 35
 - PSTH, 33–34
 - spatial facilitation, 50
 - thixotropy, 118
 - unitary H reflex, 36
- central fatigue. *See* muscular fatigue
- central latency, assessment of, changes in
 - H reflex, 8–9
 - on-going EMG modulation, 26
 - on-going EMG vs. H reflex, 26
 - PSTH, 32
- cerebellum
 - internal models, putative role in, 484
 - withdrawal reflexes, responses evoked by, 383
- cerebral palsy
 - changes in
 - group II excitation, 573
 - H_{\max}/M_{\max} , 572
 - homosynaptic depression, 573
 - reciprocal Ia inhibition, 206, 573
 - recurrent inhibition, 170
 - contracture, 567
 - motor impairment, spasticity, 567
 - reciprocal Ia excitation, 547
- Charcot-Marie-Tooth disease type 1A, 82, 271, 279
- childhood, plasticity
 - short-long thumb abductor synchrony, 556
 - standing, walking, 547
- clasp-knife phenomenon, 566
- clenching
 - changes in
 - corticospinal neurone excitability, 487
 - non-reciprocal group I inhibition (wrist), 464
 - presynaptic inhibition of Ia terminals, 464
 - synergy ECR–finger flexors, 463
- clonidine, group II excitation, SCI, 280
- clonus
 - prevention, movement, low gain SLSR
 - homosynaptic depression, 450
 - presynaptic inhibition of Ia terminals, 450, 475
 - recurrent inhibition, 165, 475
- spasticity
 - afferent feedback (spindle), 129
 - recurrent stretch reflex activity, 566
- CMEP, cervicomedullary MEP, 40, 43
- coherence analyses
 - evidence for
 - cortical control, co-contraction of antagonists, 475
 - corticospinal plasticity, childhood, 556
 - parallel bilateral control, quiet standing, 519
 - motor system investigations, use in, 51
- collision in motor axons, paired H reflex technique, 142–144
- concentric contraction, changes in
 - H reflex, MEP, 456–457
 - presynaptic inhibition of Ia terminals, 312
 - spindle inflow, 127, 453–454
- contraction remote muscle (Jendrassik manoeuvre)
 - α motoneurons, excitation, 124
 - elderly, 548
 - gamma (γ) activity, 123–124
 - presynaptic inhibition of Ia terminals
 - lower limb, 316
 - wrist, 317
 - reciprocal Ia inhibition, 202
- contracture, in spasticity
 - disuse, role of, 566

motor impairment, role in, 566–567	Ib interneurons, 232
muscle stiffness, contribution to, 566	lumbar propriospinal neurones, 434
convergence on interneurons mediating	non-reciprocal group I inhibitory interneurons (wrist), 462
C3-C4 propriospinal excitation, 403, 408, 411	PAD interneurons, 304–307
feedback (C3-C4 propriospinal) inhibition, 405, 408	propriospinal neurones (C3-C4), 402–405
group II excitation, 265–270	Renshaw cells, 157
Ib inhibition, 227–234	corticospinal neurone excitability, assessment
lumbar propriospinal excitation, 50, 433, 434	descending corticospinal volleys, 488
non-reciprocal group I inhibition (wrist), 192–193, 462	MEP vs. H reflex/ongoing EMG, 491–492
reciprocal Ia inhibition, 184, 188–190	TMS- vs. TES-induced MEP/CMEP, 43, 487
recurrent inhibition, 157–158	TMS-induced H reflex facilitation, 487–488
withdrawal reflexes, 335, 352, 380	corticospinal neurone excitability, modulation with
cooling	agonist contraction
afferent fibres, conduction velocity, 257	initiation, 491–492
spinal group II excitation, 257	phasic, 488–490
transcortical Ia excitation, 93	relaxation, 490–491
corollary discharge. <i>See</i> efference copy	tonic, 487–488
cortical map, changes in	antagonist contraction
amputation, 555–556	ballistic, 493
badminton players, 555	initiation, 492–493
Braille readers, 555	tonic, 492
hemispherectomy, 556	arm cycling, 490
peripheral nerve anastomosis, 556	bicycling, 490
peripheral stimulation, 554	co-contraction of antagonists (hinge joint), 474–475
string players, 555	disuse, 555
cortical reciprocal inhibition	inactivity, 486–487
evidence for, 493	lengthening contraction, 491
plasticity, elderly, 493, 556	motor cortex stimulation
cortical silent period, following	PAS (paired associative stimulation), 553–554
contralateral subthreshold TMS	rTMS (repetitive TMS), 47, 554
LLSR and long-latency cutaneomuscular reflex, differential	TBS (theta burst stimulation), 47, 554
effect on, 372	tDCS (transcranial direct current stimulation), 47
timing, 45	motor resonance, 490
transcortical Ia excitation, evidence for, 98	peripheral stimulation, 554
contralateral suprathreshold TMS	skilled tasks, 493–494
spinal fatigue, evidence for, 497	corticospinal peak, PSTH, 41, 94, 298, 369–371, 402–407, 488
timing, 45	corticospinal plasticity
ipsilateral TMS	age-related
interhemispheric inhibition, 46	childhood
transcortical Ia excitation, evidence for, 94	short-long thumb abductor synchrony, 556
triphasic pattern, evidence for corticospinal involvement,	string players, 556
466	elderly,
cortical stimulation. <i>See</i> motor cortex stimulation	cortical reciprocal inhibition, 493, 556
corticospinal control of	cortical use-dependent plasticity, 556
CPG, 471	long-term
feedback inhibitory interneurons (C3-C4 propriospinal	amputation, 555–556
system), 407–408	badminton players, cortical map, 555
gamma (γ) motoneurons, 123	Braille readers, cortical map, 555
Ia inhibitory interneurons, 190, 474, 492	hemispherectomy, 556

584 Index

- corticospinal plasticity (cont.)
 - motor learning, 555
 - motor skill, 555
 - peripheral nerve anastomosis, 556
 - string players, cortical map, fMRI, 555
 - stroke, 556–557
- stimulation-induced,
 - PAS, 553–554
 - peripheral stimulation, 554
 - rTMS, 47, 554
 - TBS, 47, 554
 - tDCS, 47
 - TMS, 553
- training-induced
 - balance, 555
 - co-contraction, 555
 - disuse, 555
 - skill vs. strength, 555
 - use-dependent plasticity, 553
- CPG (central pattern generator), locomotion
 - cortical control, 471
 - evidence for, 526
 - half-centre organisation, 526
 - long-latency FRA pathways, 339, 526
 - muscle synergies, role in, 471
 - rhythmic arm movements, 526
- cremasteric reflexes, 380
- CUSUM
 - EMG, 26
 - PSTH, 31–32
- cutaneomuscular reflexes
 - E1 (early excitation), 363
 - E2 (long-latency excitation), 363
 - I1 (inhibition), 363
 - maturation, 363, 369
 - methodology
 - electrical stimulation, 340
 - on-going EMG modulation, 343
 - spike-triggered averaging, 341, 363, 367
 - mirror movements, 369
 - plasticity
 - elderly, E1, 363
 - habituation, 363
 - newborn, E2 vs. E1, 369, 546
 - upper motoneurone lesions, 369, 382
- cutaneous afferents (tactile)
 - activation in
 - cutaneomuscular reflexes, 363
 - LLSRs, 95–96
 - movement, 334
 - RII reflex, 362–363
 - spinal cutaneous effects, 365–368
 - transcortical cutaneous responses, 368–372
- motoneurone recruitment order, reversal, 372–374
- projection to
 - C3–C4 propriospinal neurones, 402
 - feedback inhibitory interneurons (C3–C4 propriospinal system), 405, 414–417, 423–424, 454
 - gamma (γ) motoneurons, 121–122
 - Ia inhibitory interneurons, 188–190
 - Ib interneurons, 227–230, 454
 - interneurons mediating withdrawal reflexes, 352
 - PAD interneurons, 304
- suppression of
 - C3–C4 propriospinal transmission, 414–417, 423
 - early withdrawal reflexes, 359
 - Ib inhibition, 227–229
- cutaneous reflexes
 - background from animal experiments, 335–337
 - methodology
 - caveats and limitations, 343–344
 - cutaneomuscular reflexes, 340, 363
 - monosynaptic reflexes, modulation of, 363–365
 - on-going EMG modulation, 343
 - PSTH, 343
 - responses at rest, 341–343
 - stimulation, 340–341
- organisation
 - afferent pathway, A β afferents, 365
 - cutaneomuscular reflexes, 363
 - interlimb responses, 377
 - motoneurone recruitment order, 372–374
 - RII reflex, 362–363
- pathway
 - spinal early responses, 365–368
 - transcortical late responses, 368–372
- physiological tasks
 - grip, manipulation, 374, 451, 454
 - standing, unstable stance, 376
 - walking
 - muscle activation, 374
 - perturbations, swing phase, 377–379, 531
- presynaptic inhibition of Ia terminals, changes
 - in, 368
- cutaneous silent period (CSP)
 - central pathway, upper limb, 347
 - organisation, 353
 - protective function, 353

- D wave. *See* descending corticospinal volleys
- D1, D2 inhibition, H reflex, 298–300
- deafferented patients, evidence for
 - afferent feedback control, contraction, 128, 451
 - central plan in triphasic pattern, 466
 - recurrent inhibition, antidromic volleys, 141, 152
- decerebrate rigidity vs. spasticity, 566
- descending command, transmission through C3–C4 propriospinal relay
 - evidence for, 414–416
 - group I-induced facilitation, 422, 453
 - possible function
 - control of force, 423, 452
 - focussing action, 419, 423, 452
 - grip, 420
 - grip–lift task, 422
 - multi-joint movements, 419, 471–472
 - muscle synergies, 419, 471–472
 - muscular fatigue, compensation for, 417–418
 - reaching, 420–422, 472, 485–486
 - termination movement, 423–424, 454
 - quantitative part, 416–417
- descending corticospinal volleys (D and I waves), 41–43
 - cervicomedullary stimulation, 43
 - TES, 41
 - TMS, 41–43
- disfacilitation vs. inhibition, on-going EMG, 24
- disfacilitation, suppression of
 - C3–C4 propriospinal transmission, 365, 400, 405–407, 414–416, 423–424
 - group II excitation, 266, 267–270
 - lumbar propriospinal transmission, 267–270, 434
- distal movements
 - afferent feedback delay, deleterious effects, 128
 - feedforward control, 128
 - gamma (γ)-driven spindle inflow, 128
 - kinaesthetic sensations, stretched antagonists, 128
 - monosynaptic corticospinal excitation, 465
 - monosynaptic Ia excitation, 465
 - propriospinal excitation, absence of, 402
 - recurrent inhibition, absence of, 153
 - transcortical Ia excitation, 96
- disuse, changes in
 - corticospinal neurone excitability, 555
 - H_{\max}/M_{\max} ratio, 549
 - homosynaptic depression, 89, 90, 549
 - presynaptic inhibition of Ia terminals, 313, 549
 - reciprocal Ia inhibition, 198, 549
- disuse, role in spasticity
 - contracture, 566
 - transmission in spinal pathways, changes in, 90, 570
- DOPA
 - group II excitation, spinal cord injury, 280
 - long-latency FRA pathways, cat, 337, 339
 - Parkinson’s disease, 241, 282, 324, 382, 430–431, 575
- dynamics (limb’s intersegmental), 447–448, 479–480, 483
- dystonia, changes in
 - C3–C4 propriospinal excitation, 431
 - group II excitation, 284
 - non-reciprocal group I inhibition (wrist), 324
 - presynaptic inhibition of Ia terminals, 324
- dystonia, unaffected side, 324
- E1, early cutaneomuscular excitation, 363
- E2, long-latency cutaneomuscular excitation, 363
- eccentric contraction. *See* lengthening contraction
- efference copy
 - evidence for, 482–484
 - generation
 - C3–C4 propriospinal relay, 485
 - upstream primary motor cortex, 484
 - possible function
 - afferent feedback delays, compensation for, 482
 - distinction external-internal feedback, 482
 - length signal vs. gamma (γ) drive, 482
 - motor learning, 482
 - reaching, corrective movements, 483–484
- elderly
 - balance, deficit, 524–525, 548–549
 - changes in
 - cortical reciprocal inhibition, 493, 556
 - cutaneomuscular reflexes, 363
 - Ia excitation, monosynaptic reflexes, 548
 - reciprocal Ia inhibition, 188, 548
 - use-dependent corticospinal plasticity, 556
 - presynaptic inhibition of Ia terminals, methodology, 321–322, 548
 - recurrent inhibition, 153, 548
 - walking, 548
- EMG. *See* ongoing EMG modulation
- external feedback vs. reafference, 482, 485
- F wave, 21
 - caveats and limitations, 21–24
 - motoneurons involved, 21–22
 - studies in patients, 24

586 Index

- fatigue. *See* muscular fatigue
- feedback inhibition. *See* propriospinal relay (C3-C4), organisation
- feedback inhibitory interneurons (C3-C4 propriospinal system)
- organisation
- corticospinal control, 407–408
- cutaneous input, 405, 423, 454
- group I input, 405
- possible function
- control of force, 423, 452
- focussing action, 423, 452
- termination movement, 423–424
- feedforward control
- distal movements, 128, 465
- propriospinal relay (C3-C4), feed forward inhibition, 396–397, 410, 412, 423, 425
- quiet standing, 518–519
- flexor spasms (SCI), PICs in motoneurons, 381
- fMRI (functional magnetic resonance imaging)
- corticospinal plasticity, string players, 555
- dynamic cortical changes, stroke recovery, 557
- efference copy
- generation upstream M1, evidence for, 484
- possible role of, evidence for, 482–483
- plantar responses, cortical control of, 381
- propriospinal relay (C3-C4), evidence for, 410
- FRA (flexion reflex afferent) pathways, background from animal experiments, 337–339
- long-latency FRA pathways, 339
- short-latency FRA pathways, 338–339
- FRA hypothesis, 338
- group II excitation, 274, 451, 470
- FRA pathways
- long-latency, SCI patients
- contralateral inhibition, 357
- half-centre organisation, 526
- inhibition from short-latency FRA, 355, 380
- long-latency FRA responses, 355, 380
- presynaptic inhibition of Ia terminals, 355–357
- short-latency
- functional organisation, 349–353
- modulation during
- standing, 362
- voluntary contraction, 361–362
- walking, alternative FRA pathways, 362
- functional stretch reflex, standing, perturbations, 98, 521–522
- fusimotor. *See* gamma (γ)
- gain hypothesis
- recurrent inhibition, possible function
- control of force, 162–163
- power vs. skill, 162–163, 452–453
- gait. *See* walking
- gamma (γ) activity
- efference copy, requirement of, 482
- kinaesthetic sensations, 128
- relaxed muscles, 118–119
- gamma (γ) drive
- evidence for
- microneurography, 115–117
- thixotropy, 117
- methodology, 114–118
- caveats and limitations, 117–118
- microneurography, 115–117
- nerve blocks, 115
- tendon jerk vs. H reflex, 114–115
- thixotropy, 117
- motor tasks, involvement in
- agonist contraction, 113–114, 125–127
- attention-demanding task, 124–125
- co-contraction of antagonists, 127
- motor imagery, 124
- motor learning, 128–129
- motor preparation, 124
- muscular fatigue, 497
- muscular fatigue, compensation for (not involved), 125
- remote contraction (Jendrassik manoeuvre), 123–124
- servo-assistance, 110, 450–451
- standing, task-dependence, 127–128
- steady contraction, unloading reflex, 125, 128
- studies in patients
- Parkinson's disease, 130
- spasticity, 129–130, 568
- spinal cord injury, 130
- stroke, 129
- gamma (γ) motoneurons, 112
- corticospinal control, 123
- cutaneous control, task-dependence, 121–123
- dynamic (γ_d) motoneurons, 112, 127, 129
- static (γ_s) motoneurons, 112, 127, 129
- vestibular control, 123
- Golgi tendon organs (GTO), 214
- prolonged vibration, selective electrical stimulation of Ib afferents, 193, 215
- vibration, sensitivity to, 120, 214
- grasp reflex, 382

grip, role of	
cutaneomuscular reflexes, 374, 451, 454	
feedback inhibition (C3-C4 propriospinal system),	
suppression of, 420	
heteronymous Ia excitation, hand to proximal muscles, 84, 469	
grip-lift task, role of	
feedback inhibition (C3-C4 propriospinal system),	
suppression of, 422	
grip-load force coupling, efference copy, possible role of, 483	
group II afferents (spindle)	
activation in	
concentric contraction, 127, 453–454	
lengthening contraction, 127, 453–454	
MLSR, 257–260	
steady contraction, γ -driven discharge, 125–127, 453	
conduction velocity, 261–262	
electrical threshold, 262	
identification, microneurography, 115–116	
origin from spindle secondaries, 111–112	
projection to lumbar propriospinal neurones, 264, 433	
group II excitation	
background from animal experiments	
extensor motoneurons, inhibition of, 251	
lumbar propriospinal neurones, 249–251	
homosynaptic depression, absence of, 252	
input, 251–252	
output, 251	
monoaminergic gating, 252	
presynaptic gating, 252	
evidence for	
cooling, 257	
late high-threshold response, 254–257	
MLSR, 252–254	
spindle afferents (not cutaneous/joint), 260	
tizanidine, 260	
vibration, 257–260	
methodology	
caveats and limitations, 260–261, 270	
H reflex, 254	
ongoing EMG modulation, 254–257	
PSTH, 254	
stretch, reclining, 254	
stretch, standing, 252	
organisation	
central delay, 262–263	
convergence, Ia afferents, 265–266	
corticospinal facilitation of	
feedback inhibitory interneurons, 267–270	
lumbar propriospinal neurones, 267	
crossed responses, 264	
feedback inhibition, 266	
heteronymous, pattern	
lower limb, 264	
upper limb, 264–265	
homonymous responses, 252–254, 264	
propriospinal neurones, rostral location of, 263–264	
physiological tasks	
bicycling, 277	
standing, 271–273, 520, 521	
voluntary contraction, 270–271	
walking, 274–279, 528	
possible function	
standing, balance, MLSR, 271, 521	
unstable standing, muscle activation, 273–274,	
470, 520	
voluntary contraction	
FRA hypothesis, 451, 470	
muscle synergies, 470	
servo-assistance, 451	
walking	
muscle activation, 274–276, 470, 528	
perturbations, 276–277, 530	
studies in patients	
cerebral palsy, 573	
dystonia, 284	
Parkinson's disease, 282, 574–575	
peripheral neuropathies, 279	
spasticity, 279–282, 569	
spinal cord injury, 280	
stroke, 279–280	
group II inhibition, absence of evidence for,	
266–267	
group III-IV muscle afferents, 157–158, 451, 566	
H reflex	
assessing motoneurone pool excitability, 1–21	
basic methodology, 4–9	
caveats and limitations, 8–9, 11–13, 15–17	
gating of monosynaptic Ia EPSP through	
activity-dependent hyperpolarisation, 11	
curtailment by disynaptic Ib IPSP, 12–13, 270, 432	
homosynaptic depression, 12, 87, 297–298, 322	
presynaptic inhibition, 11–12, 295–297, 307–308	
H and M recruitment curve, 7–8	
motoneurons, orderly recruitment of, 3	
normalisation, 6, 7, 15	
PICs in motoneurons, unchanged with, 20	
pool problems	

H reflex (cont.) <ul style="list-style-type: none">non-linearity, size-related sensitivity, 15–16recruitment gain, 16–17, 302, 374	Parkinson’s disease, 574
quantitative data, 20–21	power training, 549, 551–552
recovery cycle, 9–11	spasticity, 567
spatial facilitation technique, 50	spinal cord injury (SCI), 573
studies in patients <ul style="list-style-type: none">Parkinson’s disease, 574peripheral neuropathies, nerve lesions, 21, 84spasticity, 21, 85, 567	standing, 318
threshold tracking, 11, 87	stroke, affected side, 572
time resolution, 8–9	homosynaptic depression (Ia–motoneurone synapse) <ul style="list-style-type: none">background from animal experiments, 85evidence for<ul style="list-style-type: none">H reflex, stimulus rate, 12, 87passive stretch, 85threshold tracking, 87
half-centres, locomotion <ul style="list-style-type: none">CPG, 526long-latency FRA pathways, 339, 526	motoneurons of different type, sensitivity of, 87
handedness-related asymmetry <ul style="list-style-type: none">H reflex, 21homosynaptic depression, 21MEP threshold, 40–41proprio-spinal relay (C3–C4), voluntary contraction, group I<ul style="list-style-type: none">facilitation of, 419–420withdrawal reflexes (early), 353	plasticity <ul style="list-style-type: none">disuse, 89, 549functional significance, 89–90skill training, 89, 550
heteronymous connections <ul style="list-style-type: none">group II excitation<ul style="list-style-type: none">muscle synergies, 470pattern, 264–265Ia (monosynaptic) excitation<ul style="list-style-type: none">muscle synergies, 469pattern, 77–80recurrent inhibition<ul style="list-style-type: none">pattern, 153–155possible function in<ul style="list-style-type: none">flexible muscle synergies, 83–84, 165–167, 470standing, 167walking, 169	prolonged homonymous vibration, activated by, 297, 322
heteronymous Ia facilitation, H reflex <ul style="list-style-type: none">change in recruitment gain, 16–17, 302, 374presynaptic inhibition of Ia terminals, 300–302	studies in patients <ul style="list-style-type: none">cerebral palsy, 573multiple sclerosis, 90, 573Parkinson’s disease, 90, 575spasticity, 90, 568spinal cord injury, 90, 573stroke, 90, 572
hip strategy, standing, pitch perturbations, 523	voluntary contraction, attenuation during, 88, 450–451
H _{max} /M _{max} ratio <ul style="list-style-type: none">balance training, 550ballet dancers, 551cerebral palsy, 572disuse, 549elderly, 548endurance training, 549, 551–552healthy subjects, symmetrical, 20multiple sclerosis, 573newborn, 547	homosynaptic depression, absence of, on <ul style="list-style-type: none">lumbar proprio-spinal neurones, 85, 88, 217, 252, 270non-reciprocal group I interneurons (wrist), 193, 462
	homosynaptic depression, Ia fibre–Ia interneurone synapse, <ul style="list-style-type: none">193, 197, 459
	hopping, role of <ul style="list-style-type: none">cutaneous reflexes, 379heteronymous Ia excitation, 83–84Ia excitation, SLSR, 81
	Huntington’s disease, transcortical Ia excitation, absence <ul style="list-style-type: none">of, 94
	hyperekplexia <ul style="list-style-type: none">Ib inhibition, 241reciprocal Ia inhibition, 206recurrent inhibition, 170
	I waves. <i>See</i> descending corticospinal volleys
	I1, cutaneomuscular inhibition, 363
	Ia afferents <ul style="list-style-type: none">activation in<ul style="list-style-type: none">concentric contraction, 127, 453–454H reflex, tendon jerk, 3lengthening contraction, 127, 453–454

LLSR, 91, 95–96	manipulatory movements (hand to proximal muscles), 84, 469
monosynaptic reflex, 3–4	muscle activation, hopping, landing, running, standing, walking, 80–82, 532
SLSR, 80	muscle synergies, 83, 469
steady contraction, γ -driven discharge, 125–127, 450	voluntary contraction
activity-dependent hyperpolarisation, 72, 298	muscle synergies, 469
conduction velocity, 75–76	servo-assistance, 450–451, 452
conduction velocity, assessment of, 33	walking
discharge rates, 118–119	muscle activation, 82, 527–528
electrical threshold, 69, 72	perturbations, SLSR, 82, 529–530
identification, microneurography, 115–116	studies in patients
origin from spindle primaries, 111	Parkinson’s disease, H_{\max}/M_{\max} , 574
projection to	peripheral neuropathies, nerve lesions, 84
C3–C4 propriospinal neurones, 401–402	spasticity, H_{\max}/M_{\max} , 85, 567
feedback inhibitory interneurones (C3–C4 propriospinal system), 405	Ia excitation (monosynaptic), plasticity
group II interneurones, 265–266	elderly, monosynaptic reflexes, 548
Ia inhibitory interneurones, 181–184	newborn
Ib interneurones, 227	monosynaptic reflexes, 547
lumbar propriospinal neurones, 432–433	reciprocal Ia excitation, 80, 547
non-reciprocal group I interneurones (wrist), 192–193	PTP (post-tetanic potentiation), 542
PAD interneurones, 304	Ib afferents
Ia excitation (monosynaptic)	activation in
background from animal experiments, 65–66	LLSR (possibly), 96
evidence for	voluntary contraction, 234, 454
Ia afferents, 67–69, 72	conduction velocity, 214
monosynaptic linkage	electrical threshold, 214–215, 218–221
bidirectional connections, 71–72	identification, microneurography, 115–116
heteronymous connections, 69–72	origin from Golgi tendon organs, 214
homonymous connections, 67	projection to
initial part PSTH, 12–13, 67, 302	C3–C4 propriospinal neurones, 402
methodology	feedback inhibitory interneurones (C3–C4 propriospinal system), 405
caveats and limitations, 69, 71	group II interneurones, 251, 265–266
heteronymous connections, 69–74	Ib interneurones, 225
homonymous connections, 67–69	lumbar propriospinal neurones, 432–433
organisation	non-reciprocal group I interneurones (wrist), 192–193
flawed connections, tendon tap, 80	PAD interneurones, 304
heteronymous, pattern	Ib excitation
lower limb, 77–78	evidence for, 224
upper limb, 78–80	organisation, 226
homonymous, 76–77	spasticity, 242, 570
motoneurone recruitment order, 3, 76–77	Ib inhibition,
phylogenetic evolution, 78, 79	background from animal experiments. <i>See</i> Ib pathways, 214–217
reciprocal Ia excitation, newborn, 80, 547	evidence for
physiological tasks	disynaptic linkage, 221–223
hopping, landing, running, standing, walking, 80–84	Ib afferents, 218–221
manipulatory movements, support to hand	recurrent inhibition, absence of, 223
muscles, 84	
voluntary contraction, 450–451	
possible function	

590 Index

- Ib inhibition (cont.)
 - short duration, 223–224
- methodology
 - caveats and limitations, 224–225, 241
 - H reflex, 218
 - PSTH, 218
 - tendon stimulation, 224–225
- organisation
 - convergence of
 - corticospinal input, 232
 - cutaneous afferents, 227–230
 - Ia afferents, 227
 - Ib afferents, different muscles, 226
 - joint afferents, 230
 - multiple inputs, on the same interneurons, 232–234
 - nociceptive afferents, 230
 - vestibulo-spinal input, 232
 - heteronymous, pattern, 225
 - homonymous, strength, 225
 - subsets, organised with respect to target motoneurons, 226
- physiological tasks
 - agonist contraction, 234–238, 451–454
 - underlying mechanisms, 234–236
 - antagonist contraction, 238
 - inactive synergists, 238
 - lengthening contraction, 457–458
 - see also* Ib pathways, walking, 238–240
- possible function
 - agonist contraction, 237
 - antagonist contraction, 238, 460
 - control of force, 234, 451–453
 - curtailment movement, convergence cutaneous and joint afferents, 237–238, 454
 - dynamic sensitivity, convergence Ia afferents, 238
 - focussing action, 238–240, 452, 528, 528, 238–240
- Ib inhibition, studies in patients
 - hyperekplexia, 241
 - Parkinson's disease, 241, 575
 - restless leg syndrome, 241
 - spasticity, 241, 569–570
 - spinal cord injury, 241
 - stroke, 241
 - supranuclear palsy, 241
- Ib pathways
 - background from animal experiments
 - Ib interneurons, 215
 - input, 216–217
 - mutual inhibition, 216
 - output, 215–216
 - presynaptic gating, 217
 - reversal, locomotion, 217
 - methodology, 217–225
 - walking, 238–240, 528
- immobility *See* disuse
- inactivity, corticospinal neurone excitability, changes in, 486–487
- initial part, Ia excitation, PSTH
 - monosynaptic excitation, evidence for, 32, 67, 302
 - presynaptic inhibition of Ia terminals, 224, 302, 303, 304
- interlimb connections
 - cutaneomuscular responses, 377
 - presynaptic inhibition of Ia terminals, 316
 - walking, cutaneomuscular responses, 377, 531
- internal models
 - optimal feedback control, 485
 - possible function
 - afferent feedback delays, compensation for, 485
 - distinction external-internal feedback, 484–485
 - motor learning, 485
 - quiet standing, 518
 - stepping down, 532
 - putative neural network(s)
 - C3-C4 propriospinal relay, 485
 - cerebellum, 484
- interneuronally mediated effect, initial sparing of
 - corticospinal peak, PSTH, 404, 405–407
 - Ia excitation, PSTH, 12–13, 224, 230, 232–234, 407
- intracortical excitability
 - IHI (interhemispheric inhibition), 46
 - LAI (long-latency afferent inhibition), 46–47
 - LICF (long-interval intracortical facilitation), 46
 - LICI (long-interval intracortical inhibition), 45
 - PAS (paired associative stimulation), 47
 - SAI (short-latency afferent inhibition), 46–47
 - SICF (short-interval intracortical facilitation), 45–46
 - SICI (short-interval intracortical inhibition), 44–45
- inverted pendulum, body at the ankle joint
 - pitch perturbations, ankle strategy, 522–523
 - quiet standing, 515–516
- ipsilateral motor cortex, recovery from
 - hemispherectomy, 556
 - stroke, 556
- ipsilateral premotor cortex, recovery from stroke, 557
- IPSP, motoneurons
 - decay phase, ongoing EMG vs. H reflex, 26–27
 - ongoing EMG, reciprocal Ia inhibition, 181
- ischaemic blockade, evidence for
 - efference copy, role of, 483

Ia afferents, 69, 80, 82, 85, 89, 123, 194, 196, 204, 266, 274, 276, 277, 313, 316	stumbling, 531–532
Ib afferents, 221	upper limb movements, 96, 476–481
Jendrassik manoeuvre. <i>See</i> contraction remote muscle	voluntary movement, coordination, 476–481
joint afferents	walking, perturbations, 98, 530, 532
activation, extremes of movement, 237–238, 454	Parkinson's disease, changes in, 574
excitation quadriceps motoneurons, 230	sensory pathway, 95–96
late withdrawal reflexes (SCI), contribution to, 380	standing, balance training, changes with, 524
possible function, curtailment movement, 237–238, 454	local sign in
projection to	cutaneous facilitation of
Ib interneurons, 230, 237–238, 241, 454	Ib inhibition, 237, 454
Renshaw cells, 158	PAD interneurons, lower limb, 304
joint inflammation, Ib inhibition, 241	cutaneous responses, gait, 377
kinaesthetic sensations	cutaneous suppression, C3–C4 propriospinal transmission, 423, 454
active vs. passive movements, 482	early withdrawal reflexes, 349, 353, 361
afferent feedback (spindle), stretched antagonists, 127	local sign, loss of, in
efference copy, possible role of, 483	cutaneous reflexes, upper motoneurone lesions, 380
γ -driven spindle inflow, 128	withdrawal reflexes
L-acetylcarnitine (L-Ac). <i>See</i> recurrent inhibition, evidence for	spinal cord injury, 380
landing, role of	upper motoneurone lesions, 381
heteronymous Ia excitation, 83–84	locomotion
Ia excitation, SLSR	CPG (central pattern generator), 471, 526
lower limb, 81	long-latency FRA pathways, 339, 526
upper limb, 81–82	muscle activation, afferent feedback, 527–529
lengthening contraction	perturbations, responses to, 529–532
changes in	long-loop reflex
corticospinal excitability, 491	long-latency cutaneous facilitation, 368
EMG activity, 454–455	presynaptic inhibition of Ia terminals, median to soleus, 316–317
H reflex, 312, 456	LRN (lateral reticular nucleus), 396
Ia–Ib-induced non-reciprocal group I inhibition, 457–458	lumbar propriospinal excitation
MEP, CMEP, 456–457	group I excitation, methodology, 431–432, 249–291
motoneurone recruitment order, reversal, 456	lumbar propriospinal neurones
presynaptic inhibition of Ia terminals, 312, 456–457	background from animal experiments
spindle inflow, 127, 453–454	binary system, 431
walking, occurrence in, 128, 527	group I–group II effects, transmission of, 249–251
LICF (long-interval intracortical facilitation), 46	corticospinal input
LICI (long-interval intracortical inhibition), 45	disfacilitation, 267–270, 434
LLSR (transcortical stretch reflex)	excitation, 267, 434
evidence for	homosynaptic depression, absence of, 85, 88, 217, 270
lower limb, 96–98	peripheral input
upper limb, 91–94	disfacilitation
function in	group I afferents, 433
flexible muscle synergies, 471, 476–478, 481	group II afferents, 266
postural responses, 476–478	excitation
servo-assistance, 96	DPN–quadriceps connection, 433
standing, pitch perturbations, 521	group I afferents, different muscles, 265–266, 432–433
	group II afferents, 264, 433
	rostral location, 432–432

- M wave, M_{max} , 7–8
- manipulatory movements, possible role of
 - cutaneomuscular reflexes, 374, 451, 454
 - heteronymous Ia excitation, hand to proximal muscles, 84, 469
 - Ib inhibition, convergence Ia afferents, dynamic sensitivity, 238
 - transcortical Ia excitation, 96
- medium-latency group I inhibition from foot muscles
 - GS and quadriceps, projection to, 433
 - walking, muscle activation, role in, 436, 528
- medium-latency reciprocal inhibition
 - evidence for, 194–196, 433
 - voluntary antagonist contraction, 459, 460
- MEP (motor evoked potential)
 - motoneurones, orderly recruitment, 44, 448
 - motor cortex stimulation, 40–44
- microneurography
 - afferent fibres, identification, 115–116
 - caveats and limitations, 117–118
 - methodology, 115–118
- minimal intervention hypothesis
 - evidence for, 468
 - optimal feedback control, 485
- mirror movements, patients with, evidence for
 - transcortical cutaneous excitation, 369
 - transcortical Ia excitation, 93, 94
- MLSR (medium-latency stretch response)
 - evidence for
 - group I contribution, 265–266
 - group II (spindle) excitation, 257–260
 - function in
 - standing perturbations, 271, 521
 - stumbling, 530, 532
 - walking perturbations, 276–279, 530
 - studies in patients
 - Parkinson's disease, 282, 574, 575
 - stroke, 280, 569
- monosynaptic reflex
 - method, principles, 3–4
 - motoneurones, orderly recruitment of, 3
 - pathway, 3
 - see also* H reflex; tendon jerk
- motoneurone excitability, assessment, pool
 - F wave, 21–24
 - H reflex, 1–21
 - ongoing EMG modulation, 24–28
 - tendon jerk, 8
- motoneurone excitability, assessment, single units
 - paired motor unit techniques, 19
 - PSF, 34–35
 - PSTH, 28–34
 - unitary H reflex, 35–36
- motoneurones, recruitment order
 - orderly
 - ballistic contraction, 448, 465–466
 - H reflex, 3
 - monosynaptic reflex, 3
 - TMS-induced MEP, 44, 448
 - voluntary contraction, 448
 - reversal with
 - cutaneous stimulation, 372–374
 - Ia excitation (monosynaptic), distal muscles, 77
 - lengthening contraction, 456
 - presynaptic inhibition of Ia terminals, 302–303
 - propriospinally mediated corticospinal excitation, 412
- motor axon excitability, changes in
 - multiple sclerosis, spinal cord injury, stroke, 542
 - spasticity, 568
- motor cortex excitability. *See* corticospinal neurone excitability; intracortical excitability
- motor cortex stimulation
 - caveats and limitations, 38–39, 43–44, 486, 491
 - cervicomedullary junction, 40, 43
 - conditioning spinal neurones, 43
 - C3–C4 propriospinal neurones, 402–405
 - feedback inhibitory interneurones (C3–C4 propriospinal system), 407–408
 - feedback inhibitory interneurones (lumbar propriospinal system), 267–270, 434
 - gamma (γ) motoneurones, 123
 - Ia inhibitory interneurones, 190
 - Ib interneurones, 232
 - lumbar propriospinal neurones, 434
 - non-reciprocal group I interneurones, 462
 - PAD interneurones, 304–307
 - Renshaw cells, 157
- corticospinal neurone excitability assessment
 - descending corticospinal volleys, 488
 - MEP vs. H reflex/ongoing EMG, 491–492
 - TMS- vs. TES-induced MEP/CMEP, 43, 487
 - TMS-induced H reflex facilitation, 487–488
- descending corticospinal volleys (D and I waves), 41–43
- electrical (TES), methodology, 39
- magnetic (TMS), methodology, 39–40
- MEP vs. H reflex, evidence for
 - presynaptic inhibition of Ia terminals, 43, 295–297
 - propriospinal (C3–C4) corticospinal transmission, 43, 414–416

- motor evoked potential (MEP), 40–41
- motor evoked potential, cervicomedullary junction (CMEP), 40, 43
- PSTH, corticospinal peak, 41
- motor cortex stimulation, combined peripheral and cortical stimulation
 - paired associative stimulation (PAS) protocol. *See* PAS
 - paired pulse protocol, 46–47
- motor cortex stimulation, paired-pulse
 - IHI (interhemispheric inhibition), 46
 - LICF (long-interval intracortical facilitation), 46
 - LICI (long-interval intracortical inhibition), 45
 - SICF (short-interval intracortical facilitation). *See* SICF
 - SICI (short-interval intracortical inhibition). *See* SICI
- motor cortex stimulation, prolonged
 - repetitive TMS (rTMS), 47
 - theta burst stimulation (TBS), 47
 - transcranial direct current stimulation (tDCS), 47
- motor imagery
 - centrally programmed SICI, evidence for, 495
 - gamma (γ) activity, 124
- motor learning
 - co-contractions of antagonists
 - production of, 472
 - long-term plasticity
 - activity-dependent, 541
 - corticospinal, 555
 - spinal, 550–552
 - possible contribution of
 - afferent feedback (spindle), 128
 - efference copy, 482
 - gamma (γ) activity, 128–129
 - internal models, 485
 - presynaptic inhibition of Ia terminals, 313, 550
 - propriospinal relay (C3–C4), 485
 - PTP (post-tetanic potentiation), 542
 - role in
 - muscle synergy selection, 469
 - quiet standing (childhood), 520
 - withdrawal reflexes organisation (newborn), 546
- motor preparation
 - gamma (γ) activity, 124
 - presynaptic inhibition of Ia terminals, changes in, 453
- motor resonance, changes in
 - corticospinal neurone excitability, 490
 - motoneurone excitability, 490
- motor skill
 - changes in
 - corticospinal neurone excitability, 493–494
 - homosynaptic depression, 89–90, 550
 - non-reciprocal group I inhibition (wrist), 464
 - presynaptic inhibition of Ia terminals, 313
 - SICI, 494
 - long-term corticospinal plasticity, 555
 - possible role of
 - gamma (γ) activity, 128–129
 - transcortical cutaneous excitation, 369
 - transcortical Ia excitation, 96
 - tDCS-induced, 554
- movement
 - distal vs. proximal movements, 465
 - FRA hypothesis, 338, 451, 470
- multijoint movements, biomechanics, 447–448
- multiple sclerosis
 - changes in
 - homosynaptic depression, 90, 573
 - Ib excitation, 242
 - presynaptic inhibition of Ia terminals, 322
 - reciprocal Ia inhibition, 205–206
 - H_{\max}/M_{\max} , 573
 - motor impairment, spasticity, 567
- muscle spindles, 110, 112–113
- muscle synergies
 - alternative pathways, 470–471
 - arguments for and against, 467–468
 - clenching/grasping, ECR–finger flexor synergy, 463
 - flexibility through
 - heteronymous recurrent inhibition, 83–84, 165–167, 470
 - presynaptic inhibition of Ia terminals, 83–84, 314, 470
 - transcortical Ia excitation, 476–478, 481
 - hierarchical control schema, 469
 - motor learning, formation during, 469
 - spinal implementation
 - CPG (central pattern generator), 471
 - cutaneomuscular responses, 470
 - heteronymous group II connections, 470
 - heteronymous Ia connections, 469
 - propriospinal relay (C3–C4), 471–472
 - withdrawal reflexes, 470
 - transcortical pathways, 476–481
- muscle tone, abnormal. *See* stiffness
- muscular fatigue
 - central fatigue, 496–497
 - disfacilitation, γ -driven spindle inflow, 497
 - spinal fatigue, 497
 - supraspinal fatigue, 496
 - compensation for
 - gamma (γ) drive, motor tasks (not involved), 125

594 Index

- muscular fatigue (cont.)
 - propriospinal relay (C3-C4), 417–418
 - recurrent inhibition, 158, 162
 - evidence for, twitch interpolation, 496
 - peripheral fatigue, 496
 - reciprocal Ia inhibition, changes in, 198
- newborn
 - cutaneomuscular reflexes, 363
 - Ia excitation, monosynaptic reflexes, 547
 - plantar response, 353, 546
 - reciprocal Ia excitation, 80, 547
 - recurrent inhibition, 153
- nociceptive afferents
 - activation in
 - RIII reflex, 344–347
 - withdrawal reflexes, 344–347
 - modulation of withdrawal reflexes (early), 358–359
 - projection to
 - Ib interneurons, 230
 - PAD interneurons, 304
 - Renshaw cells, 157–158
- non-reciprocal group I inhibition (hinge joints). *See* Ib inhibition
- running, suppression in, 321
- non-reciprocal group I inhibition (wrist)
 - changes in
 - agonist contraction, 462–463
 - antagonist contraction, 463
 - attention-demanding task, 464
 - clenching, 464
 - co-contraction antagonists, 473, 475–476
 - organisation, 190–193, 460–462
 - corticospinal control, 462
 - homosynaptic depression, absence of, 193
 - multi-excitatory convergence group I, 192–193
 - mutual inhibition, ‘opposite’ interneurons, 193
 - recurrent inhibition, absence of, 185, 192
- studies in patients
 - dystonia, 324
 - Parkinson’s disease, voluntary contraction, 575
 - stroke, 572
- oligosynaptic group I excitation
 - DPN–quadriceps connection. *See* lumbar propriospinal neurones
 - See* Ib excitation, 224, 226, 242
- ongoing EMG modulation, 24–28
 - basic methodology, 24–26
 - caveats and limitations, 27
 - central delay, estimate, 26
 - CUSUM, 26
 - inhibition vs. disfacilitation, 414–416
 - inhibition, IPSP, decay phase, 26–27
 - latency, changes in EMG vs. H reflex, 26
 - silent period, motoneurone discharge-related, 27
- operant conditioning, H reflex, humans
 - cortical plasticity, 544–546
 - spinal plasticity, 544–546
- operant conditioning, H reflex, rat
 - cortical plasticity, 543–544
 - spinal plasticity, 543–544
- optimal feedback control, 485
- PAD (primary afferent depolarisation) interneurons. *See* presynaptic inhibition of Ia terminals
- paired H reflex technique, 141–147
 - AHP vs. recurrent inhibition, 144
 - caveats and limitations, 146–147
 - collision in motor axons, 142
 - evidence for recurrent inhibition, 142–146
 - reference H reflex, 147
 - validation
 - animal experiments, 146
 - L-acetylcarnitine, 146
- paired motor unit techniques
 - evidence for PICs in motoneurons, 19
- Parkinson’s disease
 - changes in
 - C3-C4 propriospinal excitation, 430–431
 - gamma (γ) drive, 130, 574
 - group II excitation, 282, 575
 - H_{\max}/M_{\max} , 574
 - homosynaptic depression, 90, 575
 - Ib inhibition, 241, 575
 - presynaptic inhibition of Ia terminals, 323–324, 575
 - reciprocal Ia inhibition, 206, 575
 - recurrent inhibition, 170, 575
 - transcortical cutaneous inhibition, 382, 575
 - withdrawal reflexes, 382, 575
- motor tasks
 - MLSR, standing, postural set, 282, 575
 - non-reciprocal group I inhibition (wrist), 575
 - reciprocal Ia inhibition, 206, 575
- rigidity
 - group II excitation, 574
 - LLSR, 574
 - muscle stiffness, 573–574
- tremor

afferent feedback, 130	hip strategy, 523
F wave, 574	pitch vs. pitch-roll control, 524
PAS (paired associative stimulation), changes in	pitch-roll perturbations
corticospinal neurone excitability, 553–554	biomechanics, 524
presynaptic inhibition of Ia terminals, 306, 553–554	strategy, 523
SICI, 553–554	presynaptic inhibition of Ia terminals
peripheral neuropathies, nerve lesions, changes in	background from animal experiments
cutaneous reflexes, walking, 383	PAD interneurons, 292–293
F wave, 24	input, 293–294
H reflex, 21, 84	output, 294
MLSR (group II excitation), 279	primary afferent depolarisation (PAD), 292
SLSR (Ia excitation), 279	evidence for
PICs (persistent inwards currents) in motoneurons	GABA _A potentiation, 300–301
amplification	H reflex
reciprocal Ia inhibition, 179	H vs. MEP, 295–297
recurrent inhibition, 138	Ia vs. corticospinal facilitation, 300
synaptic input, 17–19	opposite changes in Ia facilitation vs. D1 inhibition, 302
background from animal experiments, 17	PSTH
evidence for	Ia peak, initial part, 302
paired motor unit techniques, 19	Ia vs. corticospinal peak, 298
PICs vs. changes in AHP, 19	methodology
prolonged Ia fibre stimulation	ageing, 321–322
electrical, 20, 542	caveats and limitations, 295–298, 299–300, 302, 321–322
vibration, TVR, 20	flawed technique, prolonged homonymous vibration, 297–
H reflex, input–output relationship, 20	298, 322
possible role in voluntary contraction	H reflex
relaxation, 453	D1, D2 inhibition, 298–300
steady, 450	H reflex vs. ongoing EMG/MEP/F wave, 295–297
studies in patients	heteronymous Ia facilitation, 300–302
flexor spasms, 381	heteronymous vibration/tendon tap, 298
spasticity, 568	PSTH, initial part, 302
spinal cord injury, 568	organisation
stroke, 568	corticospinal control
placing reaction, RII reflex, 376	lower limb, 304–306, 307
plantar responses	upper limb, 306, 307
pattern, 351–352	long-loop reflex, 316–317
plasticity, newborn, 353, 546	motoneurone recruitment order, reversal, 302–303
upper motoneurone lesions, 381	peripheral input
plasticity. <i>See</i> corticospinal plasticity; spinal plasticity	cutaneous (tactile) afferents, 304
plateau potentials. <i>See</i> PICs	group I afferents, 304
post-activation depression. <i>See</i> homosynaptic depression	nociceptive afferents, 304
postural set, influence on	sensitivity of
MLSR, 271, 521	H reflex, SLSR, tendon jerk, 307–308
MLSR, Parkinson’s disease, 282, 575	stretch-induced Ia volleys, 307–308
SLSR, 271, 521	subsets (target motoneurons), 303–304
posture, perturbations	tonic level, 307
pitch perturbations	vestibular control, 307
ankle strategy, 522–523	physiological tasks
changes in support strategies, 523	agonist contraction

presynaptic inhibition of Ia terminals (cont.)	PAS (paired associative stimulation), 306
concentric, 312	rTMS (repetitive TMS), 306
lengthening, 312, 456–457	training
lower limb, 308–313	arm cycling, 316
termination, 312	balance training, 550
tonic, 310	ballet dancers, 551
underlying mechanisms, 312–313	hopping training, 550
wrist, 317	locomotor training, 550
antagonist contraction	skill training, 313, 550
lower limb, 314, 459	presynaptic inhibition of Ib terminals
wrist, 317, 464	voluntary contraction of agonist, 236
arm cycling, 316	primary ending, 111
bicycling, 321	primary spindle ending
co-contraction of antagonists, ankle, 314–316, 473	sensitivity to stretch, 112
co-contraction of antagonists, wrist, 473	sensitivity to vibration, 112, 119
inactive synergists, 313–314	progressive spastic paraparesis
remote contraction	motor impairment, spasticity, 567
Jendrassik manoeuvre, 316	reciprocal Ia inhibition, 205
wrist, contralateral arm, 317	recurrent inhibition, 169–170
running, 320–321	withdrawal reflexes, walking, 382
standing, 317–319	propriospinal relay (C3–C4)
walking, 319–320	background from animal experiments
possible function	cat, 395–397
agonist contraction, 313	monkey, 397–398
antagonist contraction, 316, 460	conflicting conclusions
bicycling, 321	humans, 401, 410–411
co-contraction of antagonists, hinge joint, 316, 475	monkey, 397–398
control of force, 310	segmental interneurons, evidence against, 408–410
focusing action, 314, 452	evidence for
motor learning, 313, 550	BOLD fMRI, cervical spinal cord, 410
running, 321	corticospinal excitation, disclosure of, 411
standing	disfacilitation of
quadriceps, 319	corticospinal/peripheral excitation, by increasing
soleus, 319, 521	excitatory input, 405–408
termination contraction, suppression Ia burst, 312	MEP/EMG vs. H reflex, by cutaneous afferents, 414–416
walking, 320, 527–528	initial sparing of
studies in patients	corticospinal peak PSTH, with peripheral group I
amyotrophic lateral sclerosis (ALS), 322	disfacilitation, 408
dystonia, 324	facilitation, 404
multiple sclerosis, 322	MEP, with cutaneous disfacilitation, 416
Parkinson’s disease, 323–324, 575	lesion at C6–C7 spinal level, 425–427
spasticity, 322–323, 569	long-latency of
spasticity, motor tasks, 323, 571–572	cutaneous suppression, descending command, 400
spinal cord injury, late FRA-induced changes, 355–357	group I excitation, 398
spinal cord lesions, 322	rostral location (with respect to motoneurons), 400–401,
stroke, 322–323	405, 407, 408
presynaptic inhibition of Ia terminals, plasticity	methodology
disuse, 313, 549	caveats and limitations, 398–400
motor cortex stimulation	ongoing EMG modulation, 400

- PSTH, 398
- TMS-induced MEP, 398
- organisation
 - afferent input, 401–402
 - corticospinal input, 402–405
 - cutaneous inhibition, pattern, 423
 - diffuse distribution, 401–402
 - divergent projections, 411–412, 419
 - efference copy, possibility of, 485
 - feedback inhibition
 - afferent input, 405–407
 - convergence on inhibitory interneurons, 405, 408
 - corticospinal input, 407–408
 - group I inhibition, pattern, 405
 - hand muscles, absence of projections to, 402
 - inhibitory propriospinal neurones, 409
 - interaction excitatory vs. inhibitory inputs, 408
 - internal models, possibility of, 485
 - motoneurone recruitment order, 412
 - multi-excitatory convergence, 403, 408, 411
 - rostral location, 400–401, 405, 407, 408
 - subsets, organised with respect to muscle afferent input, 411, 419, 423
- physiological tasks
 - agonist contraction
 - muscular fatigue, 417–418
 - offset, 423–424, 454
 - onset, 418–419
 - tonic, 414–417, 418–419
 - grip, 420
 - grip–lift task, 422
 - reaching, 420–422
- possible function
 - co-contraction of antagonists, 474
 - control of force, 423
 - descending command
 - group I facilitation, 422, 453
 - modulation, 422
 - transmission, 414–425
 - focussing action, 419, 423, 452
 - grip, feedback inhibition, suppression of, 420
 - grip–lift task, feedback inhibition, suppression of, 422
 - locomotion (arm movements), 425
 - motor learning, 485
 - multi-joint movements, 419, 422–423
 - muscular fatigue, compensation for, 417–418
 - reaching, descending command, transmission of, 420–422, 472, 485–486
 - servo-assistance, complex movements, 422–423
 - synergies, multi-joint movements, 471–472
 - termination movement, 423–424, 454
- studies in patients
 - dystonia, 431
 - lesion at C6–C7 spinal level, 425–427
 - Parkinson’s disease, 430–431
 - stroke, recovery, 427–430
 - synkinetic movements, 430, 572
- propriospinally mediated reciprocal inhibition. *See* medium-latency reciprocal inhibition
- PSF (post-stimulus frequencygram)
 - caveats and limitations, 35
 - methodology, 34–35
 - multiple peaks, PSTH, 34–35
 - silent period, motoneurone discharge-related, 35
- PSTH (post-stimulus time histogram)
 - caveats and limitations, 33–34
 - CUSUM, 31–32
 - firing probability, motoneurone, 28–30
 - latency, assessment of, 31–32
 - methodology, 30–31
 - monosynaptic excitation, initial part, 12, 32, 67, 230, 232, 302, 303, 304, 404, 405–407
 - multiple peaks, 34
 - normalisation, 32
 - recurrent inhibition vs. AHP, 151
 - silent period, motoneurone discharge-related, 33
 - spatial facilitation technique, 48–50
 - stimulation, random vs. discharge-related, 30–31
- PTP (post-tetanic potentiation)
 - Ia fibre–motoneurone synapse, 542
 - motor learning, 542
- quadriceps
 - group II excitation, MLSR, walking, 277
 - H reflex, truncation of (disynaptic Ib IPSP), 12–13
- heteronymous
 - group II excitation (from TA)
 - standing (leaning backwards), 273–274
 - walking, muscle activation, 276
 - Ia excitation, pattern, 78
 - recurrent inhibition, changes in
 - standing (from TA), 165–167
 - voluntary contraction (from GS), 161–162
 - walking (to GS), 167–169
 - recurrent inhibition, pattern, 153–154
- Ib inhibition
 - convergence of
 - cutaneous afferents, 227–230

598 Index

- quadriceps (cont.)
 - Ib afferents, quadriceps and GS, 226
 - joint afferents, 230
 - inactive synergist contraction, 238
- lumbar propriospinal excitation, 433
- presynaptic inhibition of Ia terminals
 - agonist contraction, 310
 - inactive synergist contraction, 313–314
 - standing, 317–319
- quiet standing
 - evidence against a role of
 - GS spring-like properties, 517
 - short-latency stretch reflex (SLSR), 517–518
 - simple feedback system, 519
 - evidence for
 - paradoxical GS movements, 518
 - predictive controller, 518
 - feedforward control, 518–519
 - inverted pendulum, body at the ankle joint, 515–516
 - motor learning, requirement in childhood, 520
 - multi-sensory afferent feedback, 516–517
 - parallel bilateral control, 519
 - role of
 - afferent feedback, TA, 519
 - internal model, 518
 - predictive ballistic GS contractions, 518–519
 - visco-elastic stiffness, ankle, 517
- reaching
 - cat, monkey, 396–398
 - propriospinal relay (C3–C4), role in
 - coordinated synergies, 472
 - efference copy, 483–484
 - fast corrections, 485–486
 - transmission descending command, 420–422
- reafference vs. external feedback, distinction, 482, 485
- reciprocal Ia excitation
 - cerebral palsy, 547
 - newborn, 80, 547
- reciprocal Ia inhibition
 - background from animal experiments
 - Ia inhibitory interneurons, 178–179
 - inhibition, ‘opposite’ interneurons, 179
 - input, 179
 - output, 179
 - presynaptic gating, 179–180
 - evidence for
 - disynaptic linkage, 184–185
 - Ia afferents, 181–184
 - recurrent suppression, 155–157, 185
- methodology
 - caveats and limitations, 185–187
 - H reflex, 180–181
 - ongoing EMG modulation, 181
 - PSTH, 181
 - tendon jerk, 181
- organisation
 - ankle level, 187–188
 - corticospinal control, 190
 - cutaneous control, 188–190
 - elbow level, 188
 - knee level, 188
 - vestibular control, 190
- physiological tasks
 - ‘natural’ reciprocal Ia inhibition, 194
 - agonist contraction, 199–200
 - antagonist contraction, 194–198, 458–460
 - dynamic, 196
 - homosynaptic depression (Ia interneurone level), 196–197
 - tonic, 194–196
 - underlying mechanisms, 197
 - co-contraction of antagonists, 200–202, 472
 - co-contraction of antagonists, underlying mechanisms, 202, 473–475
 - initiation stepping, 202–204
 - remote contraction (Jendrassik manoeuvre), 202
 - standing, unstable, 202
 - walking, 204, 528
- possible function
 - agonist contraction, 200
 - antagonist contraction, 198, 460
 - walking, 204, 528
- studies in patients
 - cerebral palsy, 206, 573
 - hyperekplexia, 206
 - multiple sclerosis, 205–206
 - Parkinson’s disease, 206, 575
 - progressive spastic paraparesis, 205
 - spasticity, 205–206, 570
 - spinal cord injury, 205
 - stroke, 205
- studies in patients, motor tasks
 - Parkinson’s disease, 206, 575
 - spasticity, 206, 571
- reciprocal Ia inhibition, plasticity
 - daily training, 198, 551
 - elderly, 188

- explosive training, 198, 549–550
- patterned stimulation, 199
- spasticity, external stimulator, 199, 551
- reciprocal inhibition between antagonists
 - ankle, medium-latency reciprocal inhibition, 433, 460
 - ankle/elbow/knee, reciprocal Ia inhibition, 458–460
 - cortical reciprocal inhibition, 473
 - wrist, non-reciprocal group I inhibition, 190–193, 460–464
- recovery cycle, H reflex, 9–11
- recruitment curve, H reflex, 7
- recruitment gain, H reflex
 - cutaneous stimuli, effect of, 374
 - heteronymous Ia facilitation, effect on, 16–17, 302
- recruitment order motoneurons. *See* motoneurons
- recurrent inhibition
 - background from animal experiments
 - Renshaw cells, 138–140
 - input, 139
 - output, 139–140
- evidence for
 - heteronymous
 - antidromic motor volley, 151–152
 - orthodromic motor discharge, 148, 151
 - homonymous, paired H reflex technique, 142–146
 - L-acetylcarnitine, 141, 146, 152, 185
- methodology
 - AHP vs. recurrent inhibition, 144, 151, 161
 - antidromic motor stimulation, deafferented patients, 141, 152
 - arguable techniques, 140–141
 - caveats and limitations, 146–147, 152
 - heteronymous, H reflex/MEP/PSTH/ongoing EMG modulation, 147–152
 - homonymous, paired H reflex technique, 141–147
- organisation
 - corticospinal control, 157
 - heteronymous, pattern
 - lower limb, 153–154
 - upper limb, 154–155
 - homonymous, pattern
 - distal muscles, 153
 - proximal muscles, 153
 - Ia inhibitory interneurons, projection to, 155–157, 185
 - peripheral control
 - joint afferents, 158
 - nociceptive afferents, 157–158
 - tonic descending drive, 157
- physiological tasks
 - agonist contraction, 158–163, 451
 - agonist contraction, underlying mechanisms, 160–161
 - antagonist contraction, 163, 464–465
 - co-contraction of antagonists
 - ball joint, 464–465
 - hinge joint, 163–165, 472–473
 - standing
 - heteronymous, 165–167
 - homonymous, 165
 - walking, 167–169, 528
- possible function
 - agonist contraction, 162–163, 452–453
 - antagonist contraction, 163, 460
 - co-contraction of antagonists
 - ball joint, 475–476
 - hinge joint, 165, 472–473
 - control of force, 162–163
 - gain hypothesis, power vs. skill, 162–163
 - muscular fatigue, compensation for, 162
 - standing
 - heteronymous, 167
 - homonymous, 165
 - walking, 169, 528
- studies in patients
 - cerebral palsy, 170
 - hyperekplexia, 170
 - Parkinson's disease, 170
- spasticity
 - motor tasks, 170
 - progressive spastic paraparesis, ALS, 169–170
 - spinal cord injury, stroke, 169
- recurrent inhibition, plasticity, endurance training, 162
- reference H reflex, paired H reflex technique, 147
- rehabilitation
 - cortical plasticity, 553
 - spinal cord injury, locomotion, 552–553
 - spinal plasticity, 552–553
 - stroke, corticospinal plasticity, peripheral-induced, 554
- Renshaw cells. *See* recurrent inhibition
- restless leg syndrome, changes in
 - Ib inhibition, 241
 - withdrawal reflexes (cutaneous silent period, EDB), 382
- rigidity, Parkinson's disease, changes in
 - group II excitation, 574
 - LLSR, 574
 - muscle stiffness, 573–574
- RII reflex, 341–343, 362–363, 365, 376
- RIII reflex, 341, 344–347
- rTMS (repetitive TMS), changes in
 - corticospinal neurone excitability, 47, 94, 98, 299, 554

600 Index

- rTMS (repetitive TMS), changes in (cont.)
 - presynaptic inhibition of Ia terminals, 306, 554
 - SICI (short-interval intracortical inhibition), 554
- running, role of
 - cutaneous reflexes, 379
 - heteronymous Ia excitation, 83–84
 - Ia excitation, SLSR, 80–81
 - non-reciprocal group I inhibition, suppression of, 321
 - presynaptic inhibition of Ia terminals, 320–321
- SCI. *See* spinal cord injury
- secondary spindle ending, 111–112
 - sensitivity to stretch, 112
 - sensitivity to vibration, 112, 119–120
- servo-assistance, through
 - gamma (γ)-driven group II excitation, 451
 - gamma (γ)-driven Ia excitation, 450–451, 452
 - propriospinal (C3–C4) relay, 422–423
 - reciprocal Ia inhibition, 460
 - transcortical Ia excitation, 96
- shortening contraction. *See* concentric contraction
- SICF (short-interval intracortical facilitation)
 - evidence for, 45
 - methodology, caveats and limitations, 38
 - stroke, recovery, 556
 - threshold tracking, 40
- SICI (short-interval intracortical inhibition)
 - changes with
 - contralateral suprathreshold cortical silent period, 45
 - cutaneous stimulation, 372
 - initiation contraction, 495
 - LICI (long-interval intracortical inhibition), 45
 - motor imagery, 495
 - movement prevention, 495
 - PAS (paired associative stimulation), 47, 553–554
 - termination of contraction, 495
 - TMS (repetitive TMS), 554
 - vibration-induced Ia discharge, 495
 - voluntary contraction, focussing action, 45, 494–495
 - characteristics, 44
 - diagnostic significance, 45
 - methodology
 - caveats and limitations, 38, 45, 494
 - normalisation, 44–45
 - threshold tracking, 40
 - underlying mechanism, binary process, 45
- silent period, ongoing EMG
 - contraction onset, cause of delay in, 491
 - cortical silent period, 45, 46
 - cutaneous silent period (CSP), 347, 353
 - initiation of stepping, reciprocal Ia inhibition, 202–204
 - motoneurone discharge-related, 27
 - unloading, 125
- single motor units, recording of
 - PSF, 34–35
 - PSTH, 28–34
 - unitary H reflex, 35–36
- skeleto-fusimotor. *See* beta (β)
- slipping, restoration of balance, 532
- SLSR (short-latency stretch reflex)
 - function in
 - hopping, 81
 - landing, 81
 - running, 80–81
 - standing, pitch perturbations, 82, 271, 521
 - stumbling, 529, 532
 - walking, perturbations, 82, 529–530
 - gain during
 - co-contraction of antagonists
 - hinge joint, 475
 - wrist, 476
 - quiet standing, 517–518
 - presynaptic inhibition of Ia fibres, sensitivity to, 307–308
 - spasticity, exaggerated SLSR, 565–566
- spasticity, pathophysiology
 - clinical features
 - clasp-knife phenomenon, 566
 - clonus, 566
 - contracture, 566
 - stretch reflex (SLSR), exaggeration, 565–566
 - decerebrate rigidity, differences with, 566
 - definition, 565–566
 - underlying mechanisms
 - abnormal descending control, 570
 - disuse, 90, 570
 - plastic changes, 570
 - spinal fixation, 552, 570
 - sprouting, 570
- spasticity, spinal pathways
 - changes in
 - F wave, 567
 - gamma (γ) drive, 129–130, 568
 - group II excitation, 279–282, 569
 - H_{\max}/M_{\max} , 85, 567
 - homosynaptic depression, 90, 568
 - Ib excitation, 242, 570
 - Ib inhibition, 241, 569–570
 - motoneurone excitability, PICs, 568

motor axon excitability, 542, 568	local sign, loss of, 380
presynaptic inhibition of Ia terminals, 322, 569	multi-afferent, 380
spinal cord lesions, 322	presynaptic inhibition of Ia terminals, late FRA-induced
stroke, 322–323	changes, 355–357
reciprocal Ia inhibition	rehabilitation, locomotion, 552–553
soleus, 205–206, 570	spinal fixation, 552
TA, 205	spinal fixation, 543
recurrent inhibition	spinal cord injury, 552
progressive spastic paraparesis, ALS, 169–170	stroke, 552
stroke, SCI, 169, 570	spinal plasticity
mechanisms underlying changes in	H reflex, operant conditioning
group II excitation, 280	humans, 544–546
homosynaptic depression, 90	rat, 543–544
Ib excitation, 242	training and disuse, 543
Ib inhibition, 241	spinal plasticity, age-related
non-monosynaptic group I excitation, 280	childhood
presynaptic inhibition of Ia terminals, 323	standing, feedback reflexes, 547
reciprocal Ia inhibition, 206	walking, 547
recurrent inhibition, 169, 170	elderly
motor tasks, changes in	cutaneomuscular reflexes, 363
presynaptic inhibition of Ia terminals, 323, 571–572	Ia excitation, monosynaptic reflexes, 548
reciprocal Ia inhibition, 206, 571	Jendrassik manoeuvre, 548
recurrent inhibition, 170	reciprocal Ia inhibition, 188, 548
spatial facilitation technique	recurrent inhibition, 153
assessment with	newborn
H reflex, 50	cutaneomuscular reflexes, 363, 369
PSTH, 48–50	Ia excitation, monosynaptic reflexes, 547
background from animal experiments, 48	plantar response, 353, 546
caveats and limitations, 50, 51	reciprocal Ia excitation, 80, 547
convergence on interneurons, 48, 184, 192–193, 230, 232,	spinal plasticity, CNS lesions
267, 304, 403, 434	interlimb connections, 380, 552
spinal cord injury (SCI)	motor axon excitability, 542
autonomic dysreflexia, 552, 573	spinal plasticity, TMS-related, presynaptic inhibition of Ia
changes in	terminals
gamma (γ) drive, 130	PAS, 306
group II excitation, 280	rTMS, 306
H_{\max}/M_{\max} , 573	spinal plasticity, training-related
homosynaptic depression, 90	homosynaptic depression
Ib excitation, 242	disuse, 89, 549
Ib inhibition, 241	skill training, 89–90, 550
PICs, 568	Ia excitation (monosynaptic)
presynaptic inhibition of Ia terminals, 322	power vs. endurance training, 549, 551–552
reciprocal Ia inhibition, 205	presynaptic inhibition of Ia terminals
recurrent inhibition, 169, 570	arm cycling, 316
flexor spasms, 381	balance training, 550
interlimb connections, 380, 552	ballet dancers, 551
late withdrawal reflexes, 355–357, 380	disuse, 313, 549
early reflexes, replacement of, 355, 380	hopping training, 550
inhibition from short-latency FRA, 355	locomotor training, 550

602 Index

- spinal plasticity, training-related (cont.)
 - skill training, 313, 550
- reciprocal Ia inhibition
 - daily training, 198, 551
 - explosive training, 198, 549–550
 - patterned stimulation, 199
 - spasticity, external stimulator, 199, 551
- recurrent inhibition
 - endurance training, 162, 552
- sprouting, spinal cord injury
 - autonomic dysreflexia, 552
 - interlimb connections, 552
 - recovery, 552
- stair climbing, cutaneous reflexes, role of, 379
- standing
 - corrective function of
 - cutaneomuscular reflexes, 376–377
 - group II excitation (MLSR), 271, 521
 - Ia excitation (SLSR), 82, 271, 521
 - presynaptic inhibition of Ia terminals, 319, 521
 - transcortical Ia excitation (LLSR), 521
 - transcortical Ia excitation, training, 524
 - responses to pitch perturbations
 - ankle strategy, 522–523
 - changes in support strategies, 523
 - conflicting results, 523
 - cutaneous reflexes, contribution of, 522
 - functional stretch reflex, ankle muscles, 521–522
 - hip strategy, 523
 - LLSR, ankle muscles, 521
 - long-latency responses, antagonist, 522
 - MLSR, ankle muscles, 521
 - SLSR and MLSR, flexor digitorum brevis, 522
 - SLSR, ankle muscles, 521
 - responses to pitch–roll perturbations
 - neural pathways, 524
 - strategy, 523–524
 - standing, role of
 - gamma (γ) activity, 127–128
 - presynaptic inhibition of Ia terminals
 - quadriceps, 319
 - soleus, 319
 - withdrawal reflexes (early), modulation, 362
- standing, unstable, muscle activation, role of
 - heteronymous
 - group II excitation, 273–274, 470, 520
 - Ia excitation, 82–83, 469–520
 - recurrent inhibition, 165–167
 - homonymous recurrent inhibition, 165
 - reciprocal Ia inhibition, 202
- steady contraction
 - changes in
 - afferent feedback, 125, 128, 451, 452
 - H reflex, MEP, CMEP, 456–457
 - presynaptic inhibition of Ia terminals, 310
 - spinal pathways contributing to
 - control of speed and force, 452–453
 - maintenance, 450–452
 - selectivity, 452–453
 - timing, 453
- stepping down, restoration of balance
 - braking reaction, 532
 - internal model, possible role of, 532
- stiffness
 - co-contraction of antagonists
 - ball joint, 475–476
 - hinge joint, 475
 - muscle tone, abnormal
 - fusimotor drive, role of, 129
 - Parkinson's disease, 573–574
 - spasticity, contracture, 566
 - spasticity, stretch reflex, 565–566
 - visco-elastic, ankle, quiet standing, 517
- stretch reflex, long-latency (LLSR). *See* LLSR
- stretch reflex, medium-latency (MLSR). *See* MLSR
- stretch reflex, short-latency. *See* SLSR
- stroke, motor impairment, role of
 - contracture, 566–567
 - gamma (γ) drive, 129
 - spasticity, 566–567
 - synkinetic movements, 430, 572
- stroke, recovery, role of
 - bilateral cortical overactivation, 557
 - C3–C4 propriospinal excitation, 427–430
- contralateral
 - corticospinal tract, 557
 - intracortical disinhibition, 557
- dynamic cortical changes, fMRI, 557
- ipsilateral
 - motor cortex, 556
 - premotor cortex, 557
 - SICF, 556
- stroke, spasticity
 - changes due to spinal fixation, 552
 - changes in
 - H_{\max}/M_{\max} , 572
 - PICs, 568
 - changes in spinal pathways

- gamma (γ) drive, 129
- group II excitation, 279–280
- homosynaptic depression, 90
- Ib excitation, 242
- Ib inhibition, 241
- presynaptic inhibition of Ia terminals, 322–323
- reciprocal Ia inhibition, 205
- recurrent inhibition, 169, 570
- withdrawal reflexes, walking, 382
- stroke, unaffected side, changes in
 - C3–C4 propriospinal excitation, 429, 572
 - group II excitation, 280
 - presynaptic inhibition of Ia terminals, 572
 - reciprocal Ia inhibition, 205, 572
 - stretch responses, 572
- stumbling, corrective responses, involvement of
 - compensatory arm responses, 531
 - diffuse SLSR, 529, 532
 - LLSR, 531–532
 - MLSR, 530
- subsets of interneurons, organised with respect to
 - muscle afferent input, C3–C4 propriospinal neurones, 411, 419, 423
- target motoneurons
 - Ia inhibitory interneurons, 179
 - Ib interneurons, 226
 - PAD interneurons, 303–304
- supranuclear palsy, Ib inhibition, 241
- synkinetic movements, C3–C4 propriospinal relay, 430, 467, 572
- task-related changes in
 - corticospinal neurone excitability, 487–494
 - cutaneous reflexes, 374–379
 - gamma (γ) drive, 125–129
 - group II excitation, 270–279
 - Ib inhibition, 234–240
 - non-reciprocal group I inhibition (wrist), 462–464
 - presynaptic inhibition of Ia terminals, 308–321
 - propriospinal (C3–C4) transmission, 420–422, 465
 - reciprocal Ia inhibition, 193–204
 - recurrent inhibition, 158–169
 - transcortical stretch reflexes (LLSRs), 476–478
 - withdrawal reflexes, 361–362
- TBS (theta burst stimulation)
 - corticospinal neurone excitability, changes in, 47, 554
 - efference copy, lack of effect on, 484
- tDCS (transcranial direct current stimulation)
 - changes in
 - corticospinal neurone excitability, 47, 474
 - transmission in spinal pathways, 47, 474
 - motor skill, effect on, 554
- tendon jerk
 - assessing motoneurone pool excitability, 8
 - comparison with H reflex, flawed technique for γ drive assessment, 114–115
 - difference with H reflex in
 - afferent volleys, 114
 - EPSPs in motoneurons, 114–115
 - sensitivity to presynaptic inhibition of Ia terminals, 307–308
 - pathway, 65
- tendon tap
 - evidence for Ia afferents, 67, 72, 184
 - evidence for Ib afferents, absence of effect, 221
- termination of movement, via
 - Ib inhibition, convergence cutaneous and joint afferents, activation through, 237–238, 454
 - propriospinal transmission, cutaneous suppression of, 423–425, 454
 - SICI, activation of, 495
- TES (transcranial electrical stimulation). *See* motor cortex stimulation
- thixotropy, gamma (γ) drive
 - caveats and limitations, 118
 - methodology, 117
- threshold tracking, assessment of
 - H reflex, 11
 - homosynaptic depression, 87
 - MEP (SICI, SICF), 40
 - unitary H reflex, 35–36
- tizanidine, group II excitation
 - evidence for, 260
 - stroke, 280
- TMS (transcranial magnetic stimulation). *See* motor cortex stimulation
- tonic level of
 - Ib inhibition, contraction of agonist, 236–237
 - presynaptic inhibition of Ia terminals, 307
 - reciprocal Ia inhibition, 179, 206
 - recurrent inhibition, 157
- training, changes in, with
 - balance training
 - corticospinal neurone excitability, 555
 - feedback reflexes, 548–549
 - presynaptic inhibition of Ia terminals, 550
 - transcortical Ia excitation, 524
 - ballet dancer training
 - presynaptic inhibition of Ia terminals, 551

604 Index

- training, changes in, with (cont.)
 - co-contraction training
 - corticospinal neurone excitability, 555
 - daily physical activity
 - reciprocal Ia inhibition, 198, 551
 - endurance training
 - Ia excitation (monosynaptic), 549, 551–552
 - recurrent inhibition, 162, 552
 - explosive training, 549–550
 - reciprocal Ia inhibition, 198, 549
 - hopping
 - presynaptic inhibition of Ia terminals, 550
 - locomotor training
 - presynaptic inhibition of Ia terminals, 550
 - power training
 - Ia excitation (monosynaptic), 549, 551–552
 - skill training
 - corticospinal neurone excitability, 555
 - homosynaptic depression, 89, 550
 - presynaptic inhibition of Ia terminals, 313, 550
 - TMS-coupled training
 - use-dependent corticospinal plasticity, 553
- transcortical cutaneous excitation
 - contralateral subthreshold TMS, effect of, 372
 - evidence for
 - lower limb, 368–371
 - upper limb, 368–369, 371–372
 - walking, swing phase perturbations, 377–379, 531, 532
- transcortical cutaneous inhibition
 - mirror movements, evidence for transcortical, 369
 - Parkinson's disease, 382
- transcortical Ia excitation
 - flexible muscle synergies, 476–478, 481
- lower limb
 - evidence for, 96–98
 - standing
 - balance training, 524
 - pitch perturbations, function in, 521
 - walking, function in, 98, 530, 532
- sensory pathway, 95–96
- task-dependence, 476–478
- upper limb
 - contralateral subthreshold TMS, effect of, 372
 - evidence for, 93–94
 - function, 96, 476–481
 - pattern multijoint, 478–481
 - proximo-distal pattern, 94
 - voluntary intent, overlap with, 94, 478
- transcortical stretch reflex. *See* LLSR
- transcranial stimulation. *See* motor cortex stimulation
- tremor, Parkinsonian, afferent feedback, 130
- triphasic EMG pattern, 466
- TVR (tonic vibration reflex)
 - PICs in motoneurons, 20
 - unloading-induced decreased spindle discharge, 121
 - vibration paradox, 297
- twitch interpolation, muscular fatigue, 496
- unaffected side, changes in spinal pathways
 - dystonia, 324
 - stroke, 572
- unitary H reflex, 35–36
- unloading reflex, 125, 128
- upper motoneurone lesions
 - cutaneous reflexes, changes in
 - abdominal and cremasteric reflexes, 380
 - cutaneomuscular reflexes, 382
 - early withdrawal reflexes, 381
 - local sign, loss of, 380
 - nociceptive extensor inhibition, 381–382
 - plantar responses, Babinski response, 381
 - transcortical cutaneous excitation, 369
 - spasticity, 565–573
- use-dependent corticospinal plasticity
 - D-amphetamine, effect of, 553
 - deafferentation, effect of, 553
 - elderly, 556
 - evidence for, 553
 - motor training, induced by, 553
- V1 response, 158
- vestibular control of
 - gamma (γ) motoneurons, 123
 - Ib inhibition, 232
 - presynaptic inhibition of Ia terminals, 307
 - reciprocal Ia inhibition, 190
- vibration
 - activation of
 - cutaneous mechanoreceptors, 120
 - Golgi tendon organs (GTO), 120
 - primary endings, 119
 - secondary endings, 119–120
 - evidence for
 - group II excitation, 257–260
 - Ia afferent transmission, 72, 184
 - presynaptic inhibition of Ia terminals, homonymous,
 - flawed technique, 297–298
 - production of

- activity-dependent hyperpolarisation, Ia afferents, 72, 298
- homosynaptic depression (Ia-motoneurone), 297, 322
- SICI (short-interval intracortical inhibition), 495
- vibration paradox, 297
- voluntary contraction, agonist
 - changes in (hinge joint)
 - afferent discharges, 453–454
 - beta-gamma (β - γ) activation, 125–127
 - corticospinal neurone excitability, 487–492
 - group II excitation, 125–127, 270–271, 451, 470
 - homosynaptic depression (Ia-motoneurone), 88–89
 - Ia afferent hyperpolarisation, 11
 - Ia excitation (monosynaptic), 125–127, 450–451, 452, 469
 - Ib inhibition, 234, 451–452
 - PICs in motoneurons, 450
 - presynaptic inhibition of Ia terminals, 308–313
 - proprio-spinal transmission, 416–425
 - reciprocal Ia inhibition, 199–200
 - recurrent inhibition, 158–163, 451, 452
 - SICI, 494–495
 - transcortical Ia excitation, 476–481
 - withdrawal reflexes (early), 361–362
 - changes in (wrist)
 - non-reciprocal group I inhibition, 462–463
 - presynaptic inhibition of Ia terminals, 317
 - control of force through
 - feedback inhibition (C3–C4 proprio-spinal system), 423
 - Ib inhibition, 234, 451–452
 - presynaptic inhibition of Ia terminals, 310, 452
 - recurrent inhibition, 162–163, 451
 - focussing action through
 - feedback inhibition (C3–C4 proprio-spinal system), 423, 452
 - Ib inhibition, 238, 452
 - presynaptic inhibition of Ia terminals, 314, 452
 - proprio-spinal (C3–C4) transmission, 419
 - SICI, 45, 494–495
 - mechanisms involved in relaxation
 - PICs in motoneurons, 453
 - presynaptic inhibition of Ia terminals, 312
 - SICI, 495
 - servo-assistance through
 - group II excitation, 451
 - Ia excitation (monosynaptic), 450–451, 452
 - presynaptic inhibition of Ia terminals, 313
 - proprio-spinal transmission, 422–423
 - transcortical Ia excitation, 96
 - voluntary contraction, antagonist
 - changes in
 - cortical reciprocal inhibition, 493
 - corticospinal neurone excitability, 492–493
 - changes in (hinge joint)
 - Ib inhibition, 238, 460
 - presynaptic inhibition of Ia terminals, 314, 460
 - reciprocal Ia inhibition, 194–198, 458–459, 460
 - recurrent inhibition, 163, 459, 460
 - servo-assistance, reciprocal Ia inhibition, 460
 - changes in (wrist)
 - non-reciprocal group I inhibition, 462–464
 - presynaptic inhibition of Ia terminals, 317, 464
 - voluntary contraction, co-contraction of antagonists
 - hinge joint, changes in
 - gamma (γ) activation, 127
 - corticospinal neurone excitability, 474–475
 - presynaptic inhibition of Ia terminals, 314–316, 473
 - reciprocal Ia inhibition, 200–202, 472
 - recurrent inhibition, 163–165, 472–473
 - stretch reflex (SLSR), 475
 - wrist, changes in
 - non-reciprocal group I inhibition, 473
 - presynaptic inhibition of Ia terminals, 473
 - recurrent inhibition, 464–465, 475–476
 - stretch reflex (SLSR), 476
 - voluntary contraction, corticospinal control of
 - agonist contraction, 487–492
 - antagonist contraction, 492–493
 - co-contraction of antagonists (hinge joint), 474–475
 - skilled tasks, 493–494
 - voluntary movement, coordination
 - LLSRs (transcortical stretch reflexes), 476–481
 - muscle synergies, 466–472
 - walking
 - general features
 - biomechanics, 525–526
 - gamma (γ)-driven spindle inflow, 128, 527
 - CPG (central pattern generator), 471, 526
 - lengthening loaded contractions, 128, 527
 - muscle activation, pattern of, 527
 - muscle activation, role of
 - afferent feedback (spindle), 128
 - cutaneous feedback, motoneurone recruitment order, 374
 - group I inhibition from foot muscles, 436, 528
 - group II excitation
 - heteronymous, 276, 528
 - homonymous, 274–276, 528
 - Ib pathways, reflex reversal, 238–240, 528

walking (cont.)	descending effects, 359
monosynaptic Ia excitation	nociceptive afferents, 358–359
heteronymous, 83–84	non-nociceptive cutaneous afferents, 359
homonymous, 82, 527	modulation during
presynaptic inhibition of Ia terminals	standing, 362
quadriceps, 320	voluntary contraction
soleus, 319–320, 527–528	antagonist, inhibition of, 361–362
reciprocal Ia inhibition, 204, 528	responses, 361
recurrent inhibition, heteronymous, 167–169, 528	walking, alternative FRA pathways, 362
perturbations, role of	modulation with motor learning, 546
alternative FRA pathways, 362, 531	organisation
compensatory arm responses, 531	hand muscles, cutaneous silent period, 353
cutaneous reflexes, 379, 531	handedness-related asymmetry, 353
group II excitation (MLSR)	local sign, 349, 353, 361
GS, TA, 277, 530	lower limb, 351–353
quadriceps, 277	plantar responses, 351–353
heteronymous Ia connections, ankle, 83, 528	protective function, 349, 352–353
Ia excitation (SLSR), GS, 82, 529	RIII reflex, 341, 344–347
interlimb cutaneomuscular responses, 377, 531	trunk skin reflexes, 349–351
reciprocal Ia inhibition, 528	upper limb, 353
transcortical cutaneous responses, 377–379, 532	pathway
transcortical Ia excitation, 98, 530, 532	A β afferents, convergence of, 346–347, 352
restoration of balance after	A δ fibres, 346
slipping, 532	central delay, 347
stepping down, 532	central pathway, 347–349
stumbling, 531–532	plasticity, newborn, plantar response, 353, 546
walking, plasticity	withdrawal reflexes (late)
childhood, 547	normal subjects, 357–358
elderly, 548	spinal cord injury
walking, studies in patients	contralateral inhibition, 357
cutaneous reflexes, 383	inhibition from short-latency FRA, 355, 380
presynaptic inhibition of Ia terminals, 323, 572	long-latency FRA responses, 355, 380
withdrawal reflexes (early)	presynaptic inhibition of Ia terminals, 355–357
background from animal experiments, 335–339	replacement of early withdrawal reflexes, 355
methodology	withdrawal reflexes, studies in patients
caveats and limitations, 343–344	cerebellar lesions, 383
motoneurone excitability modulation, 343	Parkinson’s disease, 382, 575
responses at rest, 341	restless leg syndrome, 382
stimulation, 340–341	spinal cord injury, 380
modulation by	upper motoneurone lesions, 381