GRAPH SPECTRA FOR COMPLEX NETWORKS

PIET VAN MIEGHEM Delft University of Technology



> CAMBRIDGE UNIVERSITY PRESS Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore, São Paulo, Delhi, Dubai, Tokyo, Mexico City

> > Cambridge University Press The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org Information on this title: www.cambridge.org/9780521194587

© Cambridge University Press 2011

This publication is in copyright. Subject to statutory exception and to the provisions of relevant collective licensing agreements, no reproduction of any part may take place without the written permission of Cambridge University Press.

First published 2011

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this publication is available from the British Library

Library of Congress Cataloguing in Publication data

ISBN 978-0-521-19458-7 Hardback

Cambridge University Press has no responsibility for the persistence or accuracy of URLs for external or third-party internet websites referred to in this publication, and does not guarantee that any content on such websites is, or will remain, accurate or appropriate.

to Saskia

who tolerated with love that science is a passionate and most demanding mistress Cambridge University Press 978-0-521-19458-7 - Graph Spectra for Complex Networks Piet van Mieghem Frontmatter More information

Contents

Preface		ix	
Ac	knowled	gements	xiii
Syn	mbols		XV
1	Intro	duction	1
	$1.1 \\ 1.2 \\ 1.3 \\ 1.4$	Interpretation and contemplation Outline of the book Classes of graphs Outlook	2 5 7 10
	Part	I Spectra of graphs	11
2	Algeb	braic graph theory	13
	$2.1 \\ 2.2$	Graph related matrices Walks and paths	13 25
3	Eigenvalues of the adjacency matrix		29
	3.1 3.2 3.3 3.4 3.5 3.6 3.7	General properties The number of walks Regular graphs Bounds for the largest, positive eigenvalue λ_1 Eigenvalue spacings Additional properties The stochastic matrix $P = \Delta^{-1}A$	29 33 43 46 55 58 63
4	Eigen	values of the Laplacian Q	67
	4.1	General properties	67

CAMBRIDGE

Cambridge University Press	
978-0-521-19458-7 - Graph S	Spectra for Complex Networks
Piet van Mieghem	
Frontmatter	
More information	

vi	Contents	
	 4.2 Second smallest eigenvalue of the Laplacian Q 4.3 Partitioning of a graph 4.4 The modularity and the modularity matrix M 4.5 Bounds for the diameter 4.6 Eigenvalues of graphs and subgraphs 	80 89 96 108 109
5	Spectra of special types of graphs	115
	5.1The complete graph5.2A small-world graph5.3A circuit on N nodes5.4A path of $N-1$ hops5.5A path of h hops5.6The wheel W_{N+1} 5.7The complete bipartite graph $K_{m,n}$ 5.8A general bipartite graph5.9Complete multi-partite graph5.10An m-fully meshed star topology5.11A chain of cliques5.12The lattice	$ \begin{array}{r} 115 \\ 115 \\ 123 \\ 124 \\ 129 \\ 129 \\ 129 \\ 129 \\ 131 \\ 135 \\ 138 \\ 147 \\ 154 \\ \end{array} $
6	Density function of the eigenvalues	159
	6.1Definitions 6.2 The density when $N \to \infty$ 6.3 Examples of spectral density functions 6.4 Density of a sparse regular graph 6.5 Random matrix theory	159 161 163 166 169
7	Spectra of complex networks	179
	 7.1 Simple observations 7.2 Distribution of the Laplacian eigenvalues and of the degree 7.3 Functional brain network 7.4 Rewiring Watts-Strogatz small-world graphs 7.5 Assortativity 7.6 Reconstructability of complex networks 7.7 Reaching consensus 7.8 Spectral graph metrics 	179 181 184 185 187 196 199 200
	Part II Eigensystem and polynomials	209
8	Eigensystem of a matrix	211
	8.1 Eigenvalues and eigenvectors8.2 Functions of a matrix	211 219

CAMBRIDGE

Cambridge University Press	
978-0-521-19458-7 - Graph Spectra for Complex Ne	etworks
Piet van Mieghem	
Frontmatter	
More information	

	Contents	vii
8.3	Hermitian and real symmetric matrices	222
8.4	Vector and matrix norms	230
8.5	Non-negative matrices	235
8.6	Positive (semi) definiteness	240
8.7	Interlacing	243
8.8	Eigenstructure of the product AB	252
8.9	Formulae of determinants	255
9 Polyn	nomials with real coefficients	263
9.1	General properties	263
9.2	Transforming polynomials	270
9.3	Interpolation	274
9.4	The Euclidean algorithm	277
9.5	Descartes' rule of signs	282
9.6	The number of real zeros in an interval	292
9.7	Locations of zeros in the complex plane	295
9.8	Zeros of complex functions	302
9.9	Bounds on values of a polynomial	305
9.10	Bounds for the spacing between zeros	306
9.11	Bounds on the zeros of a polynomial	308
10 Orth	ogonal polynomials	313
10.1	Definitions	313
10.2	Properties	315
10.3	The three-term recursion	317
10.4	Zeros of orthogonal polynomials	323
10.5	Gaussian quadrature	326
10.6	The Jacobi matrix	331
References		339
Index		345

Preface

During the first years of the third millennium, considerable interest arose in complex networks such as the Internet, the world-wide web, biological networks, utility infrastructures (for transport of energy, waste, water, trains, cars and aircrafts), social networks, human brain networks, and so on. It was realized that complex networks are omnipresent and of crucial importance to humanity, whose still augmenting living standards increasingly depend on complex networks. Around the beginning of the new era, general laws such as "preferential attachment" and the "power law of the degree" were observed in many, totally different complex networks. This fascinating coincidence gave birth to an area of new research that is still continuing today. But, as is often the case in science, deeper investigations lead to more questions and to the conclusion that so little is understood of (large) networks. For example, the rather simple but highly relevant question "What is a robust network?" seems beyond the realm of present understanding. The most natural way to embark on solving the question consists of proposing a set of metrics that tend to specify and quantify "robustness". Soon one discovers that there is no universal set of metrics, and that the metrics of any set are dependent on each other and on the structure of the network.

Any complex network can be represented by a graph. Any graph can be represented by an adjacency matrix, from which other matrices such as the Laplacian are derived. These graph related matrices are defined in Chapter 2. One of the most beautiful aspects of linear algebra is the notion that, to each matrix, a set of eigenvalues with corresponding eigenvectors can be associated. The physical meaning of an "eigen" system is best understood by regarding the matrix as a geometric transformation of "points" in a space. Those "points" define a vector: a line segment from an origin that ends in the particular point and that is directed from origin to end. The transformation (rotation, translation, scaling) of the vector is again a vector in the same space, but generally different from the original vector. The vector that after the transformation turns out to be proportional with itself is called an eigenvector and the proportionality strength or the scaling factor is the eigenvalue. The Dutch and German adjective "eigen" means something that is inherent to itself, a characteristic or fundamental property. Thus, knowing that each graph is represented by a matrix, it is natural to investigate the "eigensystem", the set of all eigenvalues with corresponding eigenvectors because the "eigensystem" characterizes the graph. Stronger even, since both the adjacency and Laplacian matrix are symmetric, there is a one-to-one correspondence between the matrix and the "eigensystem", established in art. 151.

In a broader context, transformations have proved very fruitful in science. The

х

Preface

most prominent is undoubtedly the Fourier (or Laplace) transform. Many branches of science ranging from mathematics, physics and engineering abound with examples that show the power and beauty of the Fourier transform. The general principle of such transforms is that one may study the problem in either of two domains: in the original one and in the domain after transformation, and that there exists a one-to-one correspondence between both domains. For example, a signal is a continuous function of time that may represent a message or some information produced over time. Some properties of the signal are more appropriately studied in the time-domain, while others are in the transformed domain, the frequency domain. This analogy motivates us to investigate some properties of a graph in the topology domain, represented by a graph consisting of a set of nodes connected by a set of links, while other properties may be more conveniently dealt with in the spectral domain, specified by the set of eigenvalues and eigenvectors.

The duality between topology and spectral domain is, of course, not new and has been studied in the field of mathematics called *algebraic graph theory*. Several books on the topic, for example by Cvetković et al. (1995); Biggs (1996); Godsil and Royle (2001) and recently by Cvetković et al. (2009), have already appeared. Notwithstanding these books, the present one is different in a few aspects. First, I have tried to build-up the theory as a connected set of basic building blocks, called articles, which are abbreviated by **art**. The presentation in article-style was inspired by great scholars in past, such as Gauss (1801) in his great treatise Disquisitiones Arithmeticae, Titchmarsh (1964) in his Theory of Functions, and Hardy and Wright (1968) in their splendid Introduction to the Theory of Numbers, and many others that cannot be mentioned all. To some extent, it is a turning back to the past, where books were written for peers, and without exercise sections, which currently seem standard in almost all books. Thus, this book does not contain exercises. Second, the book focuses on general theory that applies to all graphs, and much less to particular graphs with special properties, of which the Petersen graph, shown in Fig. 2.3, is perhaps the champion among all. In that aspect, the book does not deal with a zoo of special graphs and their properties, but confines itself to a few classes of graphs that depend at least on a single parameter, such as the number of nodes, that can be varied. Complex networks all differ and vary in at least some parameters. Less justifiable is the omission of multigraphs, directed graphs and weighted graphs. Third, I have attempted to make the book as self-contained as possible and, as a peculiar consequence, the original appendices consumed about half of the book! Thus, I decided to create two parts, the main part I on the spectra, while part II overviews interesting results in linear algebra and the theory of polynomials that are used in part I. Since each chapter in part II discusses a wide area in mathematics, in fact, separate books on each topic are required. Hence, only the basic theory is discussed, while advanced topics are not covered, because the goal to include part II was to support part I. Beside being supportive, part II contains interesting theory that opens possibilities to advance spectral results. For example, Laguerre's beautiful Theorem 51 may once be applied to the characteristic

Preface

polynomials of a class of graphs with the same number of negative, positive and zero eigenvalues of the adjacency matrix.

A drawback is that the book does not contain a detailed list of references pointing to the original, first published papers: it was not my intention to survey the literature on the spectra of graphs, but rather to write a cohesive manuscript on results and on methodology. Sometimes, different methods or new proofs of a same result are presented. The monograph by Cvetković *et al.* (1995), complemented by Cvetković *et al.* (2009), still remains the invaluable source for references and tables of graph spectra.

The book is a temporal reflection of the current state of the art: during the process of writing, progress is being made. In particular, the many bounds that typify the field are continuously improved. The obvious expectation is that future progress will increasingly shape and fine-tune the field into – hopefully – maturity. Hence, the book will surely need to be updated and all input is most welcome. Finally, I hope that the book may be of use to others and that it may stimulate and excite people to dive into the fascinating world of complex networks with the rigorous devices of algebraic graph theory offered here.

 $Ars\ mathematicae$

October 2010

PIET VAN MIEGHEM

xi

Acknowledgements

I would like to thank Huijuan Wang, for her general interest, input and help in pointing me to interesting articles. Further, I am most grateful to Fernando Kuipers for proofreading the first version of the manuscript, to Roeloef Koekoek for reviewing Chapter 10 on orthogonal polynomials, and to Jasmina Omic for the numerical evaluation of bounds on the largest eigenvalue of the adjacency matrix. Javier Martin Hernandez, Dajie Liu and Xin Ge have provided me with many nice pictures of graphs and plots of spectra. Stojan Trajanovski has helped me with the *m*-dimensional lattice and **art.** 105. Wynand Winterbach showed that the assortativity of regularly graphs is not necessarily equal to one, by pointing to the example of the complete graph minus one link (Section 7.5.1.1). Rob Kooij has constructed Fig. 4.1 as a counter example for the common belief that Fiedler's algebraic connectivity is always an adequate metric for network robustness with respect to graph disconnectivity. As in my previous book (Van Mieghem, 2006b), David Hemsley has suggested a number of valuable textual improvements.

Symbols

Only when explicitly mentioned, will we deviate from the standard notation and symbols outlined here.

Random variables and matrices are written with capital letters, while complex, real, integer, etc., variables are in lower case. For example, X refers to a random variable, A to a matrix, whereas x is a real number and z is complex number. Usually, i, j, k, l, m, n are integers. Operations on random variables are denoted by [.], whereas (.) is used for real or complex variables. A set of elements is embraced by $\{.\}$.

Linear algebra

A

 $n \times m \text{ matrix} \begin{bmatrix} a_{11} & \cdots & a_{1m} \\ \vdots & & \\ a_{n1} & \cdots & a_{nm} \end{bmatrix}$

$\det A$	determinant of a square matrix A ; also denoted by	÷			
		a_{n1}		a_{nn}	
$\operatorname{trace}(A)$	$=\sum_{j=1}^{n} a_{jj}$: sum of diagonal elements of A				
$\operatorname{diag}(a_k)$	$=$ diag (a_1, a_2, \ldots, a_n) : a diagonal matrix with diagonal	onal el	ement	ts liste	d,
	while all off-diagonal elements are zero				
A^T	transpose of a matrix, the rows of A are the column	s of A	Т		
A^*	matrix in which each element is the complex conjugate of the				
	corresponding element in A				
A^H	$= (A^*)^T$: Hermitian of matrix A				
$c_A(x)$	$= \det (A - xI)$: characteristic polynomial of A				
$\mathrm{adj}A$	$= A^{-1} \det A$: adjugate of A				
$Q\left(\lambda ight)$	$=\frac{c_A(\lambda)}{\lambda I-A}$: adjoint of A				
e_j	basic vector, all components are zero, except for cor	npone	nt j t	hat is	1
δ_{kj}	Kronecker delta, $\delta_{kj} = 1$ if $k = j$, else $\delta_{kj} = 0$				

 $|a_{11} \cdots a_{1n}|$

CAMBRIDGE

Cambridge University Press 978-0-521-19458-7 - Graph Spectra for Complex Networks Piet van Mieghem Frontmatter <u>More information</u>

xvi	Symbols	
Probability theory		
$\Pr\left[X ight]$	probability of the event X	
$E\left[X ight]$	$= \mu$: expectation of the random variable X	
$\operatorname{Var}[X]$	$=\sigma_X^2$: variance of the random variable X	
$f_{X}\left(x\right)$	$=\frac{dF_X(x)}{dx}$: probability density function of X	
$F_X(x)$	probability distribution function of X	
$\varphi_{X}\left(z ight)$	probability generating function of X	
	$\varphi_X(z) = E[z^X]$ when X is a discrete r.v.	
	$\varphi_X(z) = E\left[e^{-zX}\right]$ when X is a continuous r.v.	
$\{X_k\}_{1 \le k \le m}$	$= \{X_1, X_2, \dots, X_m\}$	
$X_{(k)}$	k-th order statistics, k-th smallest value in the set $\{X_k\}_{1 \le k \le m}$	
P	transition probability matrix (Markov process)	
$1_{\{x\}}$	indicator function: $1_{\{x\}} = 1$ if the event or condition $\{x\}$ is true,	
	else $1_{\{x\}} = 0$. For example, $\delta_{kj} = 1_{\{k=j\}}$	

Graph theory

L	set of links in a graph
\mathcal{N}	set of nodes in a graph
L	$= \mathcal{L} $: number of links in a graph
N	$= \mathcal{N} $: number of nodes in a graph
K_N	the complete graph with N nodes
$K_{n,m}$	the complete bi-partite graph with $N = n + m$
d_j	degree of node j
$d_{(j)}$	the j -th largest degree of node in a graph
d_{\max}	maximum degree in the graph
d_{\min}	minimum degree in the graph
D	degree in a graph (random variable)
$\kappa_{\mathcal{N}}\left(G\right)$	vertex (node) connectivity of graph G
$\kappa_{\mathcal{L}}\left(G\right)$	edge (link) connectivity of graph G
Η	hopcount in a graph (random variable)
A	adjacency matrix of graph G
В	incidence matrix of graph G
Q	$= BB^T$ Laplacian matrix of graph G
J	all-one matrix
u	all-one vector
\blacktriangle_G	the number of triangles in G
Δ	$=$ diag (d_1, d_2, \ldots, d_N) : diagonal matrix of the nodal degrees
$\{\lambda_k\}_{1 \le k \le N}$	set of eigenvalues of A ordered as $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_N$
$\{\mu_k\}_{1 \le k \le N}^{}$	set of eigenvalues of Q ordered as $\mu_1 \ge \mu_2 \ge \cdots \ge \mu_N$
N_k – –	total number of walks with length k
W_k	number of closed walks with length k