

Index

- abstinence violation effect (AVE) 341
- abstract techniques 490
- Adolescent Transition Program 463–4
- adolescents
 - aggression in 443, 447–9, 453–4
 - attention deficit/hyperactivity disorder 423, 424
 - suicidal behavior 359
 - see also* child and adolescent depression; childhood
 - anxiety; conduct disorders (CD); oppositional defiant disorder (ODD)
- aggression, in children and adolescents 443, 447–9, 453–4
 - see also* conduct disorders (CD); oppositional defiant disorder (ODD)
- alcohol dependence syndrome (ADS) 334
- alcohol-use disorders 333–53
 - alcohol abuse 334
 - alcohol dependence 333–4
 - cognitive processes in initiation and maintenance 344–5
 - cognitive-behavioral treatments 335–50
 - Beck's cognitive therapy 342–4, 348
 - comprehensive functional analysis 339
 - coping skills development 340
 - effectiveness 349–50, 522
 - empirical evidence 346–8
 - history of 335–7
 - mechanism of change 345–6, 348–9
 - motivation to change 337–8
 - predictors of response 346
 - rational emotive behavior therapy (REBT) 342, 343–4, 348
 - relapse prevention 336, 340–1, 343–4, 347
 - therapeutic relationship development 338–9
 - treatment comparisons 343–4
 - treatment goals development 340
- etiology 334–5
- future research 350–2
 - broad-spectrum CBT 352
 - identification of mediators of change 350–1
 - treatment process analyses 350–1
- history of classification 333–4
- pathological effects 334
- prevalence 334
- alliance–outcome relationship 484–6
 - see also* therapeutic relationship
- anger arousal model, conduct problems 446–7
- Anger Control Program 446
- anger control training 463
- Anger Coping Program 441, 457–8, 465, 466
 - outcomes 459–60
- antidepressants, bulimia nervosa treatment 303–4
- antisocial personality disorder (APD) 443
- anxiety disorders 60, 61, 234–5
 - childhood *see* childhood anxiety disorders
 - cognition in 317, 514–15
 - similarities to schizophrenia 317–18
 - cognitive versus behavioral therapy 489
 - see also* generalized anxiety disorder (GAD); panic disorder; social phobia
- anxiety management training 74
- anxiety sensitivity 128–9
- Anxiety Sensitivity Index (ASI) 128–9, 519
- applied science approach 510–12
- appraisal, in obsessive-compulsive disorder 143–4
- Articulated Thoughts in Simulated Situations (ATSS) 351
- articulated treatment protocols 517
- assessment
 - attention deficit/hyperactivity disorder 431–2, 436
 - child and adolescent depression 371–5
 - of cognitive operations 374–5
 - of cognitive products 372–3
 - of schema 373–4
 - obsessive-compulsive disorder 150–1
 - self theory and research applications 214–16
 - case illustration 220–1
 - conceptual orientation 214–15
 - instruments and strategies 215–16
- attention deficit/hyperactivity disorder (AD/HD) 418–37
 - assessment 431–2, 436
 - cognitive interventions 430–5
 - addressing comorbid concerns 433
 - distress over diagnostic feedback 432–3
 - efficacy 436
 - facilitating home- and school-based interventions 434–5
 - misperceptions of clinical presentation 430–1

530 Index

- attention deficit (*cont.*)
 - past and future perspectives 432
 - preparation for medication trials 433–4
- cognitive models 426–30, 510
- comorbidity 424, 433
- developmental course 423
- diagnostic criteria 419–21
- etiology 424–6
 - biological accounts 424–5
 - psychological conceptualizations 425–6
 - psychosocial conceptualizations 425
- future directions 435–7
- prevalence 422
- psychosocial impact 423–4
- situational variability 421–2
- subtypes 419–20, 422
- attentional bias
 - in depression 18
 - in panic disorder 124
 - in social phobia 263, 265–8, 282–3
- attentional training 280
- attribution–emotion linkages 454
- attributional style 365–7
 - as predictor of cognitive therapy response 480–1
- Attributional Style Questionnaire 373, 480
- autonomic inflexibility 65
- Autonomic Thoughts Questionnaire 494
- autonomy 14, 22–3, 363, 520
- aversion therapy, alcohol-use disorders and 336, 346
- avoidant personality disorder (AvPD), cognitive therapy 194
- Beck Depression Inventory 46, 77, 493, 495
- behavioral inhibition deficits 428, 429
- behavioral marital therapy, alcohol-use disorders and 336
- behavioral therapy
 - anxiety 489
 - bulimia nervosa 300, 305
 - depression 488–9
 - see also* cognitive-behavioral therapy
- binge eating 297
 - medical complications 293–4
 - see also* bulimia nervosa
- bipolar disorder (BP) 40–55, 358
 - cognitive models 40–7
 - early descriptions 40–2
 - empirical evidence 42–7
 - interaction of life events and cognitive style 45–7
 - cognitive therapy 47–51
 - group therapy 47–8
 - individual therapy 48–51
 - future developments 52–5
- blushing, fear of 272–3
 - see also* social phobia
- Body Sensations Questionnaire 129
- borderline personality disorder (BPD) 183–4, 193
- bulimia nervosa 293–309
 - cognitive model 295–7, 513
 - evidence for 299
 - cognitive-behavioral therapy 295, 297–305
 - comparison with other psychotherapies 300–3
 - effective components of 304–5
 - effectiveness of 299–304, 306
 - relapse prevention 298–9
 - with antidepressant medication 303–4
 - comorbidity 293
 - future research directions 306–8
 - medical complications 293–4
 - prevalence 294
- California Psychotherapy Alliance Scale (CALPAS) 196
- catastrophic misinterpretation
 - in obsessive-compulsive disorder 98
 - in panic attacks 120–1, 123–4, 127, 129, 510, 512
 - in schizophrenia 319
- child and adolescent depression 358–86
 - assessment 371–5
 - of cognitive operations 374–5
 - of cognitive products 372–3
 - of schema 373–4
 - cognitive models 362–70, 509–10, 512
 - cognitive distortion models 362–4
 - helplessness, hopelessness, and attributional style 365–7
 - integrative models 369–70
 - problem-solving deficits 367–8
 - schema-based models 368–9
 - self-regulation deficits 364–5
- cognitive therapy 375–81
 - cognitive restructuring approaches 375–6
 - future research directions 378–81
 - integrative treatment approaches 377–8
 - methodological issues 378
 - self-control training 376–7
- comorbidity 371
 - consequences of 358–9
 - etiology 359–62
 - prevalence 348
 - prevention 381–4
 - selective and indicated prevention 383–4
 - universal prevention 382–3
 - unresolved conceptual issues 370–1
 - see also* depression
- childhood anxiety disorders 396–411
 - behavioral treatments 402–4
 - contingency management 403–4
 - modeling 403
 - systematic desensitization 402–3
- cognitive constructs 400–2, 509
- cognitive treatment 404–5
- cognitive-behavioral treatments 397, 405–8
 - family therapy 406–7
 - group therapy 407–8
 - individual therapy 405–6
 - versus education support 408

531 Index

- etiological models 398–400
 - ethological model 398
 - familial determinants 399–400
 - temperament 399
 - tripartite model 399
- future directions 408–11
- impact of 396–7
- prevalence 396, 397
- see also* anxiety disorders
- Children's Attributional Style Questionnaire (CASQ) 373
- Children's Cognitive Error Questionnaire 373
- cholecystokinin (CCK) 126
- circadian rhythm disruption 52, 54–5
- classical conditioning theory, alcohol-use disorders and 335
- client factors in cognitive therapy 479–82
- cognitive deficits, in obsessive-compulsive disorder 93–6
- cognitive distortions 13
 - in child and adolescent depression 362–4
 - in childhood anxiety disorders 401–2
 - in parents of children with attention deficit/hyperactivity disorder 434
- cognitive operations 362
 - assessment of in children and adolescents 374–5
 - distortions in childhood anxiety disorders 401–2
- cognitive processing therapy (CPT)
 - components of 247–8
 - posttraumatic stress disorder 247, 250–1
- cognitive products 362
 - assessment of in children and adolescents 372–3
- cognitive research 508
 - advances in 509–17
 - future directions 519–21
 - see also* cognitive theory; *specific disorders*
- cognitive restructuring 80
 - in bulimia nervosa 307
 - in depression 363–4
 - child and adolescent depression 375–6
 - in panic disorder 127
 - in parents of children with attention deficit/hyperactivity disorder 434
 - in social phobia 277
- cognitive self-instructional training (CSIT)
 - program 426, 429
- Cognitive Taxonomic System 452
- cognitive theory
 - advances in 509–17
 - applied science approach 510–12
 - cognitive nature of clinical disorders 513–15
 - cognitive vulnerability and recovery 516–17
 - empirically based methodology 512–13
 - of therapeutic change 478–9
 - see also* cognitive research; *specific disorders*
- cognitive therapy 1–2, 315, 477–99, 507–9
 - active ingredients of 483–93
 - behavioral interventions 488–90
 - cognitive interventions 490–3
 - homework compliance 487
 - provision of a treatment rationale 486–7
 - therapeutic relationship 484–6
- advances in 517–19
 - articulated treatment protocols 517
 - combined treatments 519
 - relapse prevention 518
 - symptom reduction 517–18
- client factors 479–82
- cognitive theory of therapeutic change 478–9
- comparative efficacy 522–3
- mechanisms of action 29–30, 493–7, 523–4
 - cognitive and behavioral components 29
 - future research 497–9
 - role of cognition 494–7
 - role of noncognitive factors 493–4
- technical fidelity 30
- therapist empathy 30
- prescriptive and prognostic indicators 524–5
- self theory and research applications 214–22
 - assessment 214–16
 - case illustration 219–22
 - strategies 217–19
 - treatment planning 216–17
- therapist factors 482–3
- see also* cognitive-behavioral therapy; *specific disorders*
- Cognitive Therapy Scale (CTS) 31
- cognitive vulnerability 516–17, 520–1
- cognitive-behavioral models 3–4, 8
- cognitive-behavioral therapy (CBT) 2, 7–9, 507
 - bounds of 6–7
 - definition of 2–4
 - efficacy of 4–6, 522–3
 - guidelines 4–6
 - prescriptive and prognostic factors 524–5
 - see also* behavioral therapy; cognitive therapy; *specific disorders*
- Collaborative Study Psychotherapy Rating Scale (CSPRS) 482, 490
- combination treatment 519, 524
 - conduct disorders 464–5
 - generalized anxiety disorder 74–5
 - social phobia 281
- community reinforcement approach, alcohol-use disorders and 336
- completion tendency 235
- compulsions 90–1, 92–3
 - see also* obsessive-compulsive disorder (OCD)
- conceptual entropy 511–12
- Concern over Failures in Thought Control Questionnaire 102
- concrete techniques 490
- conduct disorders (CD) 424, 441–69
 - cognitive interventions 456–67
 - adolescent transition program 463–4
 - anger control training 463
 - Anger Coping Program 457–8, 459–60
 - applications to school, outpatient, and residential settings 466–7

532 Index

- conduct disorders (*cont.*)
 - Coping Power Program 458, 460–1
 - developmental appropriateness 466
 - Dinosaur School 461
 - ethnic and community context 465
 - group process issues 467
 - history of 444–6
 - individualizing intervention 465
 - Montreal Delinquency Prevention Program 461–2
 - multicomponent interventions 464–5
 - outcomes 458–64
 - problem-solving skills training 462–3
- comorbidity 443
- prevalence 443
- research issues 468–9
 - dissemination research 469
 - maintenance of change 468
 - mediational processes 468
 - moderator effects 468
- social-cognitive models 446–56, 510
 - anger arousal model 446–7
 - role of schemata 451–6
 - social information-processing model 447–51
- contingency management, childhood anxiety 403–4
- Coping Cat program 405–6
- Coping Power Program 441, 458
 - outcomes 459–61
- Coping Responses Inventory 351
- coping skills development, alcohol-use disorders 340
- covert sensitization, alcohol-use disorders and 336
- cue exposure, alcohol-use disorders and 335
- culture 7
- cyclothymia 42
- Daily Activity Schedule 189
- depression 12–32
 - adolescents *see* child and adolescent depression
 - behavioral interventions 488–9
 - causal hypotheses 18–23
 - dysfunctional beliefs 18–22
 - personality styles and stressful events 22–3
 - children *see* child and adolescent depression
 - cognitive theory of 12–23, 362–70, 510, 512, 514–15
 - cognitive vulnerability 21, 516
 - future directions 30–1
 - similarities to schizophrenia 317–18
 - cognitive therapy 23–30, 375–81
 - active ingredients of 484–5, 486, 488–9, 490
 - acute efficacy 24
 - client factors 479–82
 - comorbidity and 27
 - future directions 30–2
 - maintenance 24–5
 - mechanisms of action 29–30, 493–7
 - prescriptive factors 27–9
 - prognostic factors 26–7
 - specificity 25
 - therapist factors 482–3
- comorbidity 27, 77, 371
- descriptive hypotheses 16–18
 - thoughts of failure and loss 16–18
- remitted-depressed studies 19–21
- self-esteem and 212
- versus social phobia 283
 - see also* bipolar disorder
- depressotypic thoughts 16–17
- Diagnostic Interview for Narcissism (DIN) 176
- dichotomous thinking 13
- Dinosaur School 461
- dispositional resistance, as prescriptive factor 27–8
- Dysfunctional Attitude Scale 22, 495
- dysfunctional beliefs 12–13
 - as prognostic factor 26–7
 - depression onset and 18–22
- Dysfunctional Thought Record modifications 162
- dysphoria 358
- dysthymia 42, 358
- eating disorders 293, 295
 - see also* bulimia nervosa
- education support, childhood anxiety and 408
- emotion 454, 493, 520
- emotion dysregulation 68, 71, 513
- emotion regulation
 - in generalized anxiety disorder 67–71, 513
 - future directions 71–2, 81–2
 - incorporation into cognitive-behavioral therapy 78–82
- emotional competence 69
- emotional deprivation, in narcissistic personality disorder 182
- emotional-processing theory 66, 67
 - posttraumatic stress disorder 236–43
 - core cognitions that maintain PTSD 239–40
 - pathological fear structures 236–9
 - rigidity and extremity of beliefs 240–1
 - treatment implications 241–3
- empathy deficiency, in narcissistic personality disorder 178–9
 - empathy training 190–1
- ethnicity, cognitive therapy and 465, 481
- expectancy theory 128–9
- exposure and response prevention (ERP),
 - obsessive-compulsive disorder 91–2, 103–7, 109, 138, 149, 161–2
- exposure therapy
 - components of prolonged exposure therapy 245–6
 - panic disorder 118
 - posttraumatic stress disorder 243–6, 249–51, 252–3
 - case example 245
 - mechanisms 253

533 Index

- failed thought control model, obsessive-compulsive disorder 100–3
- failure, prominent thoughts of 16–17
 - specificity to depression 17–18
- family cognitive-behavioral therapy, childhood anxiety 406–7
- Fast Track program 442
- fear structure 66
 - pathological 236–9
- Functional Analytic Enhanced Cognitive Therapy (FECT) 485
- General Behavioral Index 46
- general contingent reinforcement 339
- generalized anxiety disorder (GAD) 60–83
 - childhood 397
 - clinical characteristics 62–3
 - cognitive-behavioral therapy 61, 72–82
 - active components of 489
 - anxiety management training 74
 - efficacy 73–4, 75–7
 - emotion regulation perspective incorporation 78–82, 513
 - future directions 81–2
 - multicomponent therapy 74–5
 - relaxation training 72–3, 74, 75
 - self-control desensitization 74–5
 - comorbidity 63, 77–8
 - diagnosis 60–1
 - prevalence 62
 - theoretical models of 64–72, 509, 515
 - avoidance of worry 64–7
 - emotion regulation perspective 67–72
 - future directions 71–2
- goal setting
 - in alcohol-use disorders 340
 - in obsessive-compulsive disorder treatment 152
- goals 205–6
- gross stress reaction 231
- group cognitive therapy
 - bipolar disorder 47–8
 - childhood anxiety 407–8
 - conduct disorders 467
- Hamilton Rating Scale for Depression 495
- helplessness model, depression 365
- homework compliance 487
- Hopelessness Scale for Children 374
- hopelessness theory, depression 365
- hyperactivity 430
 - see also* attention deficit/hyperactivity disorder (AD/HD)
- hyperkinetic disorder 420, 421
- imaginal exposure, posttraumatic stress disorder 243–4, 245, 246, 249
- information-processing deficits 321
 - in schizophrenia 316–17, 320–2, 323–4
 - limits to information-processing models 321–2
- social information-processing model, conduct problems 447–51
 - schemata effects on information processing 453–5
 - see also* cognitive theory
- instrumental emotional responses 80–1
- intelligence, as prognostic factor 26
- Internal State Scale (ISS) 49
- internalization of speech 428–9
- interoceptive conditioning model, panic disorder 129–30
- interpersonal problems 480
- interpersonal therapy (IPT), bulimia nervosa 300–3, 307–8
- interpersonal–social rhythms therapy (IPSRT) 55
- Interpretation of Intrusions Inventory (III) 99
- intrusive thoughts, in obsessive-compulsive disorder 139–42
 - normalization 154–5
- learned resourcefulness 28
- learning theory
 - of panic disorder 130
 - of posttraumatic stress disorder 233
- life events
 - bipolar disorder and 45–7
 - see also* stressful events
- Life Goals program 48
- loss, prominent thoughts of 16–17
 - specificity to depression 17–18
- magnification 13
- mania 41, 43–5, 51, 53, 510
 - see also* bipolar disorder
- marital discord, as prescriptive factor 27
- Means–Ends Problem Solving (MEPS) inventory 374
- memory impairment
 - bias in depression 17–18
 - in obsessive-compulsive disorder 94–5
- Millon Clinical Multiaxial Inventory (MCMI) 176
- Minnesota Multiphasic Personality Inventory (MMPI) 176
- mistrust, in narcissistic personality disorder 182–3
- modeling, childhood anxiety treatment 403
- modified Stroop test 124, 266
- Montreal Delinquency Prevention Program 461–2
- mood-congruent memory 42–3
- mood-dependent memory 42–3
- motivational enhancement therapy (MET), alcohol-use disorders and 338, 347
- motivational interviewing 338
- multicomponent treatment *see* combination treatment
- multidimensional, hierarchical models, of self-concept 203–5
- mutual perception 322

534 Index

- narcissistic personality disorder (NPD) 172–96, 510
- cognitive characteristics of 179–83
 - Personality Beliefs Questionnaire (PBQ) 180–1
 - Schema Questionnaire (SQ) 181–3
 - comorbidity 176, 183–4, 193
 - construct of 175–7
 - developmental factors 177–9
 - future studies 193–6
 - prevalence 176–7
 - treatment issues 183–93, 194–6
 - empathy training 190–1
 - moderating extreme opinions of self and others 191–2
 - presenting problems 183–5
 - self-correction skills 188–90
 - therapeutic relationship 185–8
 - tolerating frustration and limits more gracefully 192–3
- Narcissistic Personality Inventory (NPI) 176
- negative activation or affect (NA) 53–4
- network therapy, alcohol-use disorders and 336
- obsessions 90, 92–3, 100–1, 138, 139
- Obsessive Beliefs Questionnaire (OBQ) 99
- Obsessive Compulsive Cognitions Working Group (OCCWG) 99, 165
- obsessive-compulsive disorder (OCD) 90–109, 138–67
- assessment 150–1
 - clinical course 91
 - cognitive-behavioral theory and research
 - 92–103, 139–45, 194, 509, 512
 - behavioral theory 92–3, 107
 - effectiveness 165
 - evaluation 164
 - failed thought control model 100–3
 - general cognitive deficit theory 93–6
 - Rachman's significance model 98–100
 - responsibility theory 96–8, 513
 - treatment implications 146–8
 - why obsessional patients try too hard 143–5
 - cognitive-behavioral treatment 103–7, 138–9, 148–9, 152–64
 - challenging responsibility appraisals 160–1
 - evaluation 164, 522
 - exposure and response prevention (ERP) 91–2, 103–7, 109, 138, 149, 161–2
 - linking history to formulation 155–6
 - normalizing intrusions 154–5
 - reappraisal and belief change strategies 162–4
 - style and format of 148–9
 - understanding and testing counterproductive strategies 156–60
 - future directions 107–9, 165–7
 - goal setting 152
 - phenomenology 166
 - prevalence 91
- omission bias, in obsessive-compulsive disorder 144–5
- Open Middle Interview (OMI) 374
- operant conditioning, alcohol-use disorders and 336
- oppositional defiant disorder (ODD) 424, 441–69
- comorbidity 443
 - history of cognitive interventions 444–6
 - prevalence 443
 - social-cognitive models 446–56
 - anger arousal model 446–7
 - role of schemata 451–6
 - social information-processing model 447–51
 - see also* conduct disorders
- overgeneralization 13
- panic attacks 117, 120–1, 125, 129
- catastrophic misinterpretation and 120–1, 123–4, 129, 510, 512
- panic control treatment (PCT) 118
- panic disorder 61, 117–32
- anxiety sensitivity and expectancy theory 128–9
 - cognitive model 117–18, 120–1, 132, 509
 - critiques 123–7
 - epistemological considerations 122–3
 - cognitive-behavioral therapy 117–20, 130–1
 - efficacy 119–20, 126–7
 - significance for cognitive model 126–7
 - future research directions 130–2
 - interoceptive conditioning model 129–30
 - modern learn-theory perspective 130
 - prevalence 117, 125
 - schizophrenia comorbidity 125–6
- parallel distributed processing (PDP) 316
- Patient Cognitive Change Scale 497
- perceived self-efficacy 206–7
- person–environment fit 213–14
- Personality Beliefs Questionnaire (PBQ) 180–1
- Personal Style Inventory (PSI) 23
- personality disorder 77–8
- see also* antisocial personality disorder; avoidant personality disorder; borderline personality disorder; narcissistic personality disorder
- personality styles, depression onset and 22–3
- personalization 13
- phobia formation 233
- positive activation or affect (PA) 53–4
- posttraumatic stress disorder (PTSD) 231–53
- cognitive theory 233–6, 509, 511
 - cognitive model of emotional disorders 234–5
 - cognitive-behavioral interventions 243–51, 252–3
 - cognitive therapy 246–8, 250–1
 - exposure therapy 243–6, 249–51, 252–3
 - outcome studies 248–51
 - stress inoculation training 248, 249
- diagnostic criteria 232
- emotional-processing theory 236–43
- core cognitions that maintain PTSD 239–40
 - pathological fear structures 236–9
 - rigidity and extremity of beliefs 240–1
 - treatment implications 241–3

535 Index

- learning theory 233
- prevalence 232
- prescriptive factors 524–5
 - depression therapy 27–9
- primal loss mode 15
- problem-solving deficits
 - in child and adolescent depression 367–8
 - in conduct problems 449
- problem-solving training
 - conduct disorder treatment 462–3
 - depression treatment 368
- prognostic factors 524–5
 - depression therapy 26–7
- psychodynamic psychotherapy 491
 - bulimia nervosa 306–7
- psychological reactance 319
- Purdue Elementary Problem-Solving Inventory (PEPSI) 374
- purging, in bulimia nervosa 297
 - medical complications 293–4
- Rachman's significance model, obsessive-compulsive disorder 98–100
- rational emotive behavior therapy (REBT) 342, 348
 - comparisons 343–4
- rational-emotive therapy (RET) 105
- reality monitoring, in obsessive-compulsive disorder 95
- relapse
 - prevention 518
 - alcohol-use disorders 336, 340–1, 343–4, 347
 - bulimia nervosa 298–9
 - risk 521
- relaxation therapy
 - alcohol-use disorders 336
 - generalized anxiety disorder 72–3, 74, 75, 489
 - efficacy 73
- Responsibility Attitudes scale 163
- responsibility, in obsessive-compulsive disorder
 - 96–8, 141–3, 146, 164, 513
 - challenging responsibility appraisals 160–1
- role functioning, in schizophrenia 323–4
- safety behaviors, in social phobia 272, 273
 - reduction of 280–1
- Schedule for Affective Disorders and Schizophrenia 46
- Schema Questionnaire (SQ) 181–3, 216
- Schema Self-Monitoring Record (SSMR) 189–90
- schema-based models, child and adolescent depression 368–9
- schemata 362, 451, 454, 492
 - assessment of in children and adolescents 373–4
 - role in conduct problems 451–6
 - assessment issues 455–6
 - effects on information processing 453–5
- schizophrenia 315–27
 - cognition in 316–17, 509
 - cognitive deficits versus role functioning 323–4
 - similarities to depression and anxiety 317–18
- cognitive therapy 315, 318–20, 325–6
 - stereotype threat and 325–6
- information processing versus social cognition deficits 320–3
 - limits to information-processing models 321–2
 - social cognition defined 322–3
- panic disorder comorbidity 125–6
- seemingly irrelevant decisions (SIDs) 345
- self-concept 202–23
 - applications to cognitive therapy 214–22
 - assessment 214–16
 - case illustration 219–22
 - intervention 216–19
 - motivation for self-consistency 207
 - multidimensional, hierarchical models 203–5
 - perceived self-efficacy 206–7
 - self in context 212–14
 - person–environment fit 213–14
 - self as mediator of contextual influences 212–13
 - self as moderator of environmental risk 213
 - social phobia and 262–3, 264–5
 - values, standards and goals 205–6
 - see also* self-esteem
- self-control desensitization 74–5
- self-control therapy 364
 - child and adolescent depression 376–7
 - see also* self-regulation treatment
- self-correction skills 188–90
- Schema Self-Monitoring Record (SSMR) 189–90
- self-esteem 172, 206, 210–12
 - bipolar disorder and 43–5
 - depression and 212
 - motivation for 207
 - implications for persons with low self-esteem 207–9
 - individuals with high self-esteem 209–10
 - negative self-evaluations in social phobia 261–2
 - obsessive-compulsive disorder and 146
 - see also* self-concept
- self-instruction *see* cognitive self-instructional training (CSIT) program
- self-interest 172
 - see also* narcissistic personality disorder
- self-monitoring 427
- self-presentation theory, of social phobia 260–1
- self-regulation deficits
 - in attention deficit/hyperactivity disorder 429
 - in child and adolescent depression 364–5
- self-regulation treatment
 - attention deficit/hyperactivity disorder 426–7, 429–30, 435
 - see also* self-control therapy
- separation anxiety disorder 397
- serotonin, bulimia nervosa and 304
- Situational Competency Test 351

536 Index

- social cognition 322–3
- social information-processing model, conduct problems 447–51
 - schemata effects on information processing 453–5
- social learning theory 336, 453
- social phobia 61, 258–84
 - childhood 397
 - cognitive models 259–74, 512, 513
 - appraisals of the consequences of negative evaluation 269–72, 283
 - attentional bias 265–8, 282–3
 - current models 260–3
 - history of 259–60
 - mental representation of self 264–5
 - performance-standard discrepancies 268–9
 - symptoms of social anxiety 272–3
 - comorbidity 63, 258
 - treatment response and 278
 - etiological factors 274–6
 - future research directions 281–4
 - prevalence 258–9
 - treatment considerations 276–81
 - attentional training 280
 - cognitive restructuring 277
 - combination treatment 281
 - safety behavior reduction 280–1
 - social skills training 277–8
 - video feedback regarding social performance 279
 - versus depression 283
- Social Problem-Solving Inventory–Revised (SPSI-R) 374
- social rhythm disruption 54–5
- social skills training
 - alcohol-use disorders and 336
 - social phobia and 277–8
- sociotropy 14, 22–3, 363, 520
- Sociotropy–Autonomy Scale 23
- standards 205–6
- stereotype threat, in schizophrenia 325–6
 - changing strategies 326
- stimulus control techniques, alcohol-use disorders and 335
- stress inoculation training (SIT), posttraumatic stress disorder 248, 249
- stressful events
 - bipolar disorder and 45–7
 - depression onset and 22–3
- Stroop test 124
- Subjective Units of Discomfort (SUDS) 242
- substance abuse 334
 - comorbidity 184
 - rational emotive behavior therapy (REBT) 342
 - see also* alcohol-use disorders
- suicidal behavior, adolescents 359
- supportive-expressive therapy, bulimia nervosa 300
- symptom reduction 517–18
- systematic desensitization, anxiety treatment 489
 - childhood anxiety 402–3
- Temple-Wisconsin Cognitive Vulnerability to Depression (CVD) project 21, 516, 521
- theories 122
 - see also* cognitive theory
- therapeutic relationship 484–6
 - in alcohol-use disorders 338–9
 - in narcissistic personality disorder 185–8
- therapist factors in cognitive therapy 30, 482–3
- thinking errors, in obsessive-compulsive disorder 144
- thought listing 373
- thought–action fusion (TAF) 97, 98–100, 102, 144
- Treatment of Depression Collaborative Research Program (TDCRP) 482
- twelve-step facilitation (TSF), alcohol-use disorder treatment 347–8, 349
- values 205–6
- video feedback, regarding social performance 279
- videotaped cued recall 373
- worry 63
 - avoidance theory 64–7
 - definition 64
 - see also* anxiety disorders; generalized anxiety disorder (GAD)
 - worry exposure techniques 79–80