

Wind as a Geomorphic Agent in Cold Climates

Matti Seppälä

University of Helsinki



CAMBRIDGE
UNIVERSITY PRESS

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS

The Edinburgh Building, Cambridge CB2 2RU, UK
40 West 20th Street, New York, NY 10011-4211, USA
477 Williamstown Road, Port Melbourne, VIC 3207, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

© Matti Seppälä 2004

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2004

Printed in the United Kingdom at the University Press, Cambridge

Typeface Times 11/14 pt System L^AT_EX 2_ε [TB]

A catalogue record for this book is available from the British Library

Library of Congress Cataloguing in Publication data

Seppälä, Matti.

Wind as a geomorphic agent in cold climates / Matti Seppälä.

p. cm. – (Studies in polar research)

Includes bibliographical references and index.

ISBN 0 521 56406 9

1. Eolian processes—Polar Regions. I. Title. II. Series.

QE597.S46 2004 551.3'7'0911—dc22 2003066159

ISBN 0 521 56406 9 hardback

Contents

Acknowledgements *page ix*

1	Introduction	1
2	Delimitation and characterization of cold environments	8
2.1	Climate	9
2.2	Seasons	26
2.3	Seasonal snow cover	27
2.4	Glaciers	28
2.5	Soils	32
2.6	Permafrost	33
2.7	Seasonal frost	36
2.8	Vegetation	36
2.9	Some relevant ecological factors	49
3	General wind patterns in cold regions	56
3.1	Arctic and Subarctic regions	57
3.2	Antarctica	65
3.3	Local winds	69
3.4	Winds and temperature	79
4	Wind drift of mineral material	80
4.1	Sand drift mechanisms	80
4.2	Basic elements affecting sand drift	85
4.3	Aeolian sand	106
4.4	Aeolian dust and loess	115
4.5	Salt drift and weathering	117

5	Abrasion	119
5.1	Signs of abrasion on rocks	119
5.2	Rock varnish	136
5.3	Frosted quartz grains	137
5.4	Abrasion of peat layers	139
5.5	Practical meaning of abrasion	140
6	Deflation	142
6.1	Definitions	142
6.2	Activation of deflation	146
6.3	Blowouts in the Subarctic	148
6.4	Deflation in Iceland	151
6.5	Wind channels in New Zealand	154
6.6	Arctic blowouts	154
6.7	Deflated sand and silt surfaces	155
6.8	Frost action and deflation	158
6.9	Seasonal variation of deflation	160
6.10	Sand transport and deflation measurements	162
6.11	Cessation of deflation	173
6.12	Pavements	174
6.13	Succession of vegetation on deflation surfaces	174
6.14	Ground water table and deflation lakes	177
7	The question of oriented lakes	182
7.1	General characteristics	182
7.2	Formation processes of oriented lakes	185
7.3	Origin of oriented lakes	187
7.4	The possible role of snow in the formation of oriented lakes	192
7.5	Wind is the key factor	194
8	Accumulation	195
8.1	Ripple marks	195
8.2	Aeolian accumulation among vegetation	196
8.3	Sand dunes	197
8.4	Dune migration	210
8.5	Loess formation and dust storms in Polar regions	212
8.6	Niveo-aeolian material	217

9	Wind directions interpreted from field evidence	222
9.1	Small indicators	222
9.2	Deflation basins and furrows	226
9.3	Dune orientation	226
10	Ice-wedge casts and sand wedges	227
10.1	Frost cracking	227
10.2	Ice-wedge casts	228
10.3	Sand wedges	229
10.4	Soil and sand wedges formed by seasonal frost	231
10.5	Sand wedges at Hietatievat	235
11	Snow	236
11.1	General distribution of snow cover	236
11.2	Physical characteristics of snow	240
12	Drift of snow	251
12.1	Quality of drifted snow	251
12.2	Transport	253
12.3	Conditions during the drift	255
12.4	Amount of snow drift	258
12.5	Practical implications of snow drift	260
13	Snow accumulation	262
13.1	Snow patches and nivation	262
13.2	Accumulation and redistribution of snow in the mountains	265
13.3	Orientation of cirques	268
13.4	Snow drift and glacier mass balance	273
13.5	Asymmetric valleys	275
13.6	Slushflows	276
14	Deflation of snow cover	279
14.1	Erosional features of snow	279
14.2	Deflation lakes on glaciers	282
15	Snow and frost formation	287
15.1	Effects on permafrost formation	291
15.2	Palsa experiment	292

16	Aeolian landforms indicating palaeowind conditions	295
16.1	Environmental conditions during the Pleistocene	295
16.2	Past wind directions	297
16.3	Pleistocene abrasion	298
16.4	Other evidence of past aeolian activities	300
16.5	How to make progress in palaeoenvironmental studies	301
<i>Bibliography</i>		303
<i>Index</i>		343

The colour plates are situated between pages 141 and 142.

Introduction

Sand dune formation has been studied for a long time (e.g. Sokolów 1894) and several excellent text books on that topic have been published recently (e.g. Greeley & Iversen 1985; Pye & Tsoar 1990; Nordstrom *et al.* 1990). Large sand seas and very high sand dunes in the deserts in low latitudes are such fascinating features that they have attracted students much more than cold environment features. Detailed surveys of aeolian landforms in tropical, subtropical and temperate environments are numerous. A very extensive literature exists on deserts, their morphology and climate (e.g. Cooke & Warren 1975; Mabbutt 1977; McKee 1979; Thomas 1989, 1997; Cooke *et al.* 1993; Abrahams & Parsons 1994) and coastal environments (e.g. Bakker *et al.* 1990; Carter *et al.* 1992).

Most presentations of aeolian landforms are limited to latitudes lower than 60°, with most close to the equator. In these studies the Polar world has been totally excluded. However, we find in the Polar regions real deserts with very limited or no vegetation. People seldom think of the regions with dry snow as white deserts (e.g. Giaever 1955). We can call Polar regions outside glaciers seasonal Polar deserts, without liquid water, with no vegetation, where snow cover melts away and with wind drifting granular material in an unlimited way. These seasonal cold deserts are limited to a zone where temperature in the winter is frequently fluctuating around 0 °C which means liquid water and hard pan in the snow cover. Occasionally wind there is also able to drift and pack snow especially when new. This book examines cold environments with respect to wind and its geomorphic impact in high latitudes. Much of what is presented here can be applied also in Alpine regions and in cold regions at lower latitudes (consider, for example, Mongolia). The topic has been studied earlier, for example by Hörner (1926) and Samuelsson (1926), but apparently since then there has been no review published of the geomorphic effects of wind in cold environments.

Extremely strong winds are well-known both from the Arctic and Antarctica. In South Victoria Land, Antarctica and on both sides of the peninsula of West Antarctica, 'where rock is exposed denudation is powerful as a result of summer

insolation, frost weathering, and strong winds. The violent wind acts as a formative agency on rock and snow surfaces. It transports snow, dust, and sand; creates small heaps of coarse detrital material like those in stony deserts; it polishes boulders and bed rock; it furrows the snow and shapes it into sharp-edged *sastrugi*' (Nordenskjöld & Mecking 1928: 287).

Hobbs (1931) called attention to the (according to him) erroneous conception regarding the conditions outside the Pleistocene continental glaciers of North America and Europe. He had observed in Greenland fierce storms which blew outward from the ice of continental glaciers and the restriction of drainage to a brief warmer season during which meltwater issued from beneath the glacier. In 1942 Hobbs wrote: 'As soon as the brief warm season has come to an end, the agent of extraglacial transportation is no longer running water but strong wind currents directed radially outward from the glacier, or nearly at right angles to the glacier front'. Wind takes over from the thaw water the work of transportation so totally that Hobbs (1942) considered wind as the dominant transportation agent within extra marginal zones to continental glaciers. He had found how 'dust, sand, and smaller pebbles are lifted and carried away, leaving behind an armour of pebble pavement to protect the material below, and in every way this pavement is similar to the *sêrir* of low-latitude deserts'.


Thus an overall presentation of aeolian geomorphology from the cold regions was missing. Hamelin and Cook (1967: 129) wrote in their glossary of periglacial phenomena: 'The place of wind action in periglacial geomorphology is not fully understood. It does, however, contribute to both depositional and erosional features. Lack of vegetation in periglacial areas greatly increases the efficacy of the wind. The accumulative or depositional aspect is by far the most important. Some loess is of periglacial origin and is deposited by wind action. . . . Abrasion is performed by wind, both by blowing sand and snow.'

However, the importance of wind as a geomorphic agent in cold climates has been known a long time (e.g. Sapper 1909; Enquist 1916; Samuelsson 1926); the High Arctic deserts are rather seldom mentioned in literature (e.g. Passarge 1921; Fristrup 1952a, 1952b, 1952–53; Smiley & Zumberge 1974). In 1993 it was still possible to publish a book about *Canada's Cold Environments* (French & Slaymaker 1993) without mentioning the term wind erosion or deflation, although strong winds are pointed out. Tricart (1970) credits deflation with a greater transporting role than running water in periglacial areas. Periglacial environments are characterized by strong wind action. Climatic dryness and the sparse plant cover are only partly important for the aeolian processes, but above all the frost and melting glacial ice produce large quantities of unconsolidated sediments favourable for wind transport (Troll 1948). Peltier (1950: 221) believes that wind has a secondary role that increases in the later part of the periglacial cycle.

Maximum wind action occurs in morphogenetic regions with low annual precipitation and high or low mean annual temperatures (Peltier 1950: Fig. 6; Thornbury 1954: 60–65). In cold morphogenetic environments Peltier (1950: Fig. 7) considers glacial, periglacial, boreal and partly savanna (sic! term used in unusual context) regions, which have from moderate to strong wind action (Fig. 1.1).

In the existing literature of the twentieth century we can find certain centres of periglacial aeolian studies. First a strong school of this aeolian geomorphology was established in Uppsala, Sweden (Enquist 1916, 1932; Hörner 1926; Högbom 1923; Samuelsson, 1921, 1925, 1926). The papers were published in a local journal: *Bulletin of the Geological Institution of the University of Upsala*. Then the centre of gravity moved to the central European sand belt and to Poland where many ancient late Pleistocene and Holocene sand dunes of cold climate origin appear (e.g. Dylikowa 1958; Galon 1958; Kobendza & Kobendza 1958;

Fig. 1.1 Morphoclimatic regions and the activity of aeolian processes according to mean annual air temperature and precipitation compiled after Peltier (1950: Figs. 6 and 7). © Association of American Geographers, Blackwell Publishing.



Mrózek 1958; Pernarowski 1959, 1966; Stankowski 1963; Wasylikowa 1964; Urbaniak 1969; Kozarski 1991). In Łódź the *Biuletyn Peryglacjalny*, founded by Jan Dylik, was published. Poser (1948, 1950), by continuing the old German tradition in late glacial sand dune studies (Sokolów 1894; Solger 1910), established in Göttingen, Germany, a study centre of periglacial processes in general and aeolian processes, too. Several studies were published in *Abhandlungen der Akademie der Wissenschaften* in Göttingen in the 1970s and 1980s. At present we have no very well specialized, leading cold environment research group studying cold environment aeolian processes but the interest seems to be split into several places (Seppälä 1975b; Niessen *et al.* 1984; Koster 1988, 1995). Recently a new journal in cold environment studies has been founded: *Permafrost and Periglacial Processes*, which contains some aeolian geomorphology, too.

Gary *et al.* (1972: 232) in the *Glossary of Geology* defined *aeolian* (= *aeolian*) as 'pertaining to the wind; especially said of rocks, soils, and deposits (such as loess, dune sand, and some volcanic tuffs) whose constituents were transported (blown) and laid down by atmospheric currents, or of landforms produced or eroded by the wind, or of sedimentary structures (such as ripple marks) made by the wind, or of geologic processes (such as erosion and deposition) accomplished by the wind'. Nevertheless, wind transport is a major part of sedimentology and perhaps it has received less attention than it deserves (Pettijohn *et al.* 1987: 310).

I would like to expand further the use of the term *aeolian* to the secondary effects of wind in cold environment. Then we are speaking of snow drift and accumulation, glacier formation, permafrost formation, deflation features which could later be lake basins, and deposition of mineral material together with snow, so called *niveo-haeolian* deposits. Deflation of snow cover can increase the thickness of permafrost, and form patterned ground which has connections with wind activity. Regions with glaciers and some other areas in middle latitudes have been included in this presentation as a part of the cold climate aeolian geomorphology.


Aeolian processes are environmentally controlled. The basic factors are the latitude which determines the amount of solar radiation and relief which both strongly affect climate (Fig. 1.2). Dominating meteorological factors from the aeolian point of view are temperature and precipitation which both affect the main limiting factors: vegetation and moisture of the soil surface and also soil formation. The special controlling factors of aeolian processes in the cold environments are frost, ice, snow, icing and meltwaters. Large quantities of meltwaters in spring and summer partly prevent the aeolian drift of mineral material and partly produce favourable sand surfaces to be deformed by wind. Most of the year snow cover and ground frost in certain parts of Polar regions can totally stop the deflation of the ground, but as contrast in some other regions the winter

activity is the predominating one. As a result of favourable climatic factors wind can transport in cold environments sand, silt and dust, in some cases also small stones and especially large amounts of snow. Particle drift causes abrasion on rock surfaces and deflation forms. Material is deposited. Sand dune formation is only a small portion of the total aeolian activity in the cold environment. Snow deposition is probably much more important for landforms. A secondary effect of the thinning of snow cover is deep frost formation. Many of these processes and events have some kind of feedback on aeolian processes, for example, wind action drifts snow forming glaciers, which then affect wind pattern (Fig. 1.2).

In extreme conditions as in the Antarctic cold deserts, in the absence of contemporary fluvial and other geological and biological processes capable of removing weathered material from rock surface, aeolian abrasion acts as a controlling factor in the rate and magnitude of erosion (Malin 1992: 27).

Wind plays an important role in cold climate geomorphology. Tricart (1970: 143) enumerates the main reasons for this: (1) high frequency of strong winds and storms in the Arctic; (2) low precipitation; (3) reduction of the vegetation cover by the cold, which is increased (4) by the wind blowing away the snow and exposing the ground surface, leading to the destruction of the vegetation; and to (5) the concentration of water in the ground by freezing which tends to dry

Fig. 1.2 Flowage diagram of the aeolian processes and formations combined with other limiting and supporting factors in general.



out the surface layer of silty soils, and to the production of abundant fine debris, especially silt, by frost weathering. This material is suitable for long distance transport by wind.

Wind is everywhere and in the circumpolar environments and on high mountains it is strong. There its force is not much limited by high vegetation like forests. When considering the importance of aeolian processes in cold environments we have to keep in mind how vast areas of the earth are covered by snow permanently or seasonally. Dry snow is relatively light material and easily drifted. Snow as such does not form geomorphic features but when it is accumulated in big amounts and forming glaciers and snow patches or melting, then it produces erosional and depositional landforms of nival origin and wind drift is the primary reason for them. In certain places snow cover is thin and in other places thick and the controlling factor is wind. Blowing snow itself is a major problem in cold environments, e.g. Canada (French & Slaymaker 1993: 9). Accumulation and redistribution of snow in the mountains has considerable effects on snow transport, glacier dynamics, snow drifts and avalanche danger (Dyunin & Kotlyakov 1980).

Fig. 1.3 Wind piled sea ice blocks on the shore of the northern Gulf of Bothnia, Hailuoto, Finland. Photo by H. Tabuchi.



We should consider the winter conditions much more when studying the periglacial phenomena and the frost action on landforms. In the literature we can find a lot of knowledge of snow and of its physical characteristics which should be integrated into the geomorphic processes. In the summer time the periglacial phenomena are mainly in rest phase and melting. For example, data of freezing on the active layer and the impact of accumulating snow are limited. This book partly tries to open that gate a little.

In this book the stress of wind on sea ice (see Sverdrup 1957) and coastal processes is largely excluded, although cold environments give several characteristic features on beaches, too, such as rafted ice walls pressed up on the shores by wind (e.g. Alestalo & Häikiö 1976) (Fig. 1.3), ice-pushed boulders (Philip 1990) and boulder barricades on shores (e.g. Rosen 1979; Dionne 1994), permafrost degradation by wave action on Arctic coasts, buried ice blocks melting in beach sand, drifting of icebergs, ice scouring and plucking of rock surfaces on tidal flats (e.g. Dionne & Quessy 1995), etc.

By knowing the present-day wind action in cold environments we have a better chance of understanding also the ancient aeolian features found in the vicinity of the glaciated regions. The palaeoenvironment reconstructions deal mainly with temperatures and humidity, interpreted through plant remains, especially pollens, and through soil formation. Some wind regimes relating to the aeolian landforms are described in the literature but the winter conditions are almost totally neglected, with the exception of the evidence of permafrost. Occurrence of snow is considered mainly for niveo-aeolian deposits. It seems to be obvious that today's aeolian processes in cold regions are considerably weaker and scattered compared with the Pleistocene activity in North America, central Europe, Asia, Argentina and New Zealand when the glaciers reached their maximum stage.