

Index

- aesthesiometers 193, 199
- age-related changes
 - postural stability 17, 18
 - sensorimotor function 18, 40–1
 - muscle strength 47
 - peripheral sensation 44–5
 - reaction time 48
 - vestibular sense 46
 - vision 41, 64
 - walking 26–9
- age-related macular degeneration 65
- aids *see* assistive devices; walking aids
- alcohol consumption 108
- Alzheimer’s disease 8, 45, 68
- amiloride 87
- amputation, fall risk and 8
- analgesics 87–8
- angina pectoris 86
- angiotensin-converting enzyme (ACE) inhibitors 86
- ankle strategy 24
- anti-inflammatory drugs 87–8
- antidepressants 84–5, 88
- antihypertensive agents 86–7, 88, 112–13
 - withdrawal of 209
- antipsychotics 84, 88
- anxiety, psychosocial therapies 209–10
- anxiolytics 83–4
- arthritis 87–8
- assistive devices 166, 170–1
 - hip protectors 172–3
 - in institutional settings 185
 - ‘long lie’ prevention 173–4
 - restraints 174–5, 185
 - spectacles 171–2
 - see also* walking aids
- balance
 - assessment of 201
 - future research directions 239
 - see also* postural stability; vestibular sense
- benzodiazepines 84, 89
- beta blockers 86
- bifocal spectacles 172
- blood pressure, variation in 60–1
 - see also* hypotension
- body sway *see* postural sway
- calcium channel blockers 86
- cardiovascular problems
 - drop attacks 61–3
 - stroke 8, 55
 - syncope 63–4
 - see also* hypotension
- cardiovascular system medications 86–7
 - antihypertensive agents 86–7
 - diuretics 87
- care institutions *see* hospitals; residential care facilities
- carotid sinus hypersensitivity 62, 64
- cataracts 64–5
- cerebellar disorders 56–7
- cervical spondylosis 56
- codeine 88
- cognitive factors 68–9
- consciousness, loss of 63–4
- contrast sensitivity 42–4
- coordinated stability test 21–2
- corticosteroids 87
- counselling of patients at risk 192
 - peripheral sensation counselling 232–4
 - vestibular system and visual field dependence counselling 234
- dementia 68
- demographic factors 107–8
- depression 68–9
 - psychosocial therapies 209–10
- depth perception 41, 42
- diabetes 46, 58
- digoxin 87
- diuretics 87, 88
- dizziness 57, 112
 - orthostatic hypotension and 61
 - see also* balance; vestibular sense
- drop attacks 61–3, 64
- education
 - falls risk awareness 186
 - group education sessions 217

246 Index

- endurance training 127–9
 - background 127–8
 - effects on falls and falls risk 128–9
 - effects on functional abilities 129
- environmental modification strategies 146–51
 - as individual interventions 146–9
 - barriers to 150
 - design issues 150–1
 - in institutional settings 184–5
 - in public places 150–1
 - multifactorial programmes 149–50
 - sleep disturbance management 211–12
- environmental risk factors 96–104, 113–14
 - evidence for 97–102
 - in private residences 97–101
 - in public places 101–2
 - in residential care facilities 101
 - interaction between individual and environment 102–4
 - proportion of falls involving environmental factors 96–7
 - suggested risk factors 97, 98
- epilepsy 62
- exercise 121–37, 182–4
 - endurance training 127–9
 - fall rates and 121–2
 - future research directions 241
 - physiotherapy intervention 129–31, 182–4
 - resistance training 123–6
 - setting 134–7
 - group exercise 132–3, 134–5, 182–4
 - home exercise 133–4, 135
 - sleep disturbance management 210–11
 - targeted intervention 230–1
- falls
 - ascertainment of 3–4
 - assessment of 187, 218
 - avoidance response 25
 - consequences of 9–12
 - cost of 12–13
 - definition of 3
 - incidence of in elderly persons 4–8
 - community-dwelling people 4–6
 - long-term care institutions 6–7
 - particular groups 8
 - location of 8–9
 - prevention of *see* prevention strategies
 - risk assessment *see* general practice assessment
 - of at-risk patients; physiological profile assessment (PPA)
- falls clinics, evaluation of 242
- fitness training *see* endurance training
- floor surfaces 151
- foot problems 66–7
- footwear 154–61
 - advice 192, 196
 - foot problems and 67
 - future research directions 241
 - heel collar height 158–60
 - heel height 155–6
 - midsole cushioning 156–7
 - midsole flaring 160–1
 - slip-resistance 157–8
- fractures 10
- frailty 8, 9
- frusemide 87
- functional reach test 21
- future research directions 239–43
 - balance 239
 - determination of best interventions 243
 - evaluation of falls clinics 242
 - exercise intervention programmes 240–1
 - falls prevention
 - in institutions 241
 - in population subgroups 240
 - home modifications 241–2
 - intervention compliance 242
 - modification of medication use 242
 - multifactorial targeted interventions 243
 - neuropsychological risk factors 240
 - sensorimotor systems 239
 - shoe designs 241
 - transient risk factors 239–40
- gait
 - age-related changes 26–8
 - problems following stroke 130
 - see also* walking
- general practice assessment of at-risk patients 192–202
 - balance 201
 - orthostatic hypotension 201–2
 - reaction time 200
 - sensation 195–9
 - strength 199–200
 - vision 194–5
- glasses *see* spectacles
- glaucoma 65–6
- group education sessions 217
- group exercise 132–3, 134–5, 182–4
- heel height 155–6
- hip fractures 10
- hip protectors 172–3
- hip strategy 24
- home exercise 133–4, 135
- homes *see* private homes; residential care facilities
- hospitals
 - admissions to 9–10
 - discharge planning 186
 - falls prevention *see* prevention strategies
 - incidence of falls 180–1
- hypertension 86
- hypnotics 83–4
- hypotension 58–61
 - drug-induced 88–9
 - orthostatic 58–61, 112
 - assessment of 201–2
 - post-prandial 60–1, 63
- incontinence 67–8
- injuries from falls 10
- insomnia *see* sleep disturbances

247 Index

- institutional settings *see* hospitals; prevention strategies; residential care facilities
- intervention strategies *see* prevention strategies
- joint position sense 45
- leaning, postural stability 20–2
- ‘long lie’ 10–11
 - preventative aids 173–4
- lorazepam 84
- lower extremity problems 66–7
- macular degeneration 65
- maximum balance range test 21
- medical factors 111–12
 - management of 190–2
 - see also specific medical conditions*
- medical management 190–203
 - conditions associated with falls 190–2
 - counselling of patients at risk 192
 - see also* general practice assessment of at-risk patients
- medications
 - as risk factors 82–91, 112–13, 206
 - physiological mechanisms 88–90
 - study design issues 82–3
 - future research directions 242
 - record cards 206–7
 - withdrawal of 206–9
 - see also specific drugs and classes of drugs*
- Menière’s disease 62
- monoamine oxidase inhibitors 84
- monofilament aesthesiometers 199
- multifocal spectacles 172
- muscle strength 47–8
 - age-related changes 47
 - assessment of 199–200
 - resistance training 123–6
- muscle synergies 24
- myelopathy 56–7
- near-tandem standing balance test 201, 202
- neurological problems 55–8
 - cerebellar disorders 56–7
 - myelopathy 56
 - neuropathy 45–6, 58
 - Parkinson’s disease 8, 56
 - stroke 8, 55
 - vestibular pathology 57
- neuropsychological risk factors 240
- nitrazepam 84
- nonsteroidal anti-inflammatory drugs (NSAIDs)
 - 87, 90, 112
 - reduction of use 208–9
- nursing homes
 - falls prevention *see* prevention strategies
 - incidence of falls 7, 181–2
 - see also* residential care facilities
- obstacle navigation 29–30
- occupational therapy intervention 146–8, 186
 - institutional settings 184
- orthostatic hypotension 58–61, 112
 - assessment of 201–2
- osteoarthritis 66
- Parkinson’s disease, fall risk and 8, 56
- peripheral neuropathy 45, 58
- peripheral sensation 44–6
 - assessment of 195–9
 - targeted counselling intervention 232–4
 - see also* sensorimotor function
- personal alarm systems 174
- physical activity 108
 - sleep disturbance management 211
 - see also* exercise
- physiological profile assessment (PPA) 221–34
 - randomized controlled trial 229–34
 - see also* targeted falls prevention strategies
- physiotherapy intervention 129–31, 182–4
 - background 129–30
 - effects on falls and falls risk 130
 - effects on functional abilities 131
- polypharmacy 82, 206, 208
 - see also* medications
- post-fall syndrome 12
- post-prandial hypotension 60–1, 63
- postural hypotension 58
 - drug-induced 88–9
- postural stability 17–31, 109–10
 - during leaning tasks 20–2
 - during normal walking 26–9
 - responses to external perturbations 22–5
 - when performing multiple tasks 30–1
 - when standing 18–20
 - see also* balance; vestibular sense
- postural strategies 24
- postural stress test 22
- postural sway
 - during leaning tasks 20–2
 - when standing 18–20
- presbyopia 172
- prevention strategies 119–20
 - future research directions 240–3
 - determination of best interventions 243
 - intervention compliance 242
 - in institutional settings 182–7, 241
 - discharge planning 186
 - environmental interventions 184–5
 - evaluation of prevention programmes 187
 - increasing awareness of falls risk 186
 - post-fall assessment 187
 - reducing falls injuries 186
 - risk factor modification 182–4
 - screening protocols 182
 - ‘long lie’ prevention 173–4
 - targeted prevention strategies 215–19, 229–34
 - exercise intervention 230–1
 - peripheral sensation counselling intervention 232–4
 - vestibular system and visual field dependence counselling intervention 234
 - visual intervention 231–2

248 Index

- prevention strategies (*cont.*)
 - see also* assistive devices; environmental modification strategies; exercise; physiological profile assessment (PPA)
- private homes
 - environmental modification strategies 146–9
 - future research directions 241–2
 - environmental risk factors 97–101
 - incidence of falls 8–9
- propoxyphene 88
- proprioception 45, 46
- psychoactive medications 83–6
 - alternatives to 209–12
 - behavioural strategies 210
 - environmental interventions 211–12
 - exercise 210–11
 - psychosocial therapies 209–10
- antidepressants 84–5
- antipsychotics 84
- anxiolytics 83–4
- hypnotics 83–4
- withdrawal of 207–8
- psychological factors 68–9
- psychosocial factors 107–8
- psychosocial therapies 209–10
- public places
 - environmental modification strategies 150–1
 - environmental risk factors 101–2
 - incidence of falls 8–9
- reaction time 48–9
 - assessment of 200–1
- rehabilitation programmes 186
- research directions *see* future research directions
- residential care facilities
 - environmental risk factors 101
 - falls prevention *see* prevention strategies
 - incidence of falls 6–7, 181–2
- resistance training 123–6
 - background 123–4
 - effects on falls and fall risk 124–5
 - effects on functional abilities 125–6
- restraints 174–5, 185
- retirement villages, incidence of falls 7
 - see also* private homes; residential care facilities
- Rhomberg test 201
- rod catch test 200–1
- rotational perturbation 22–4
- screening, in institutional settings 182
 - see also* general practice assessment of at-risk patients; physiological profile assessment (PPA)
- selective serotonin re-uptake inhibitors (SSRIs) 84–5
- sensorimotor function 40–9, 110–11
 - age-related changes 40–1
 - assessment of 195–9
 - future research directions 239
 - muscle strength 47–8
 - peripheral sensation 44–6
 - reaction time 48–9
 - vestibular sense 46–7
 - vision 41–4
- sensory organization test 22–3
- shoes *see* footwear
- sit-to-stand functional strength test 199–200
- sleep disturbances, nonpharmacological treatments 209–12
 - behavioural strategies 210
 - environmental interventions 211–12
 - exercise 210–11
 - psychosocial therapies 209–10
- slip-resistance of footwear 157–8
- spectacles 171–2
 - bifocal/multifocal spectacles 172
- stability limits 17
 - see also* postural stability
- standing, postural stability 18–20
- step width 27–8
- stepping responses 25–6
- stepping strategy 24–5
- strength *see* muscle strength
- strength training *see* resistance training
- stride length 28–9
- stroke
 - fall risk and 8, 55
 - gait problems and 130
- sway *see* postural sway
- syncope 63–4
- tactile sensitivity 44–5, 46
- Tai Chi exercise programme 122, 132
- tandem walking 29
- targeted falls prevention strategies 215–19
 - exercise intervention 230–1
 - future research directions 243
 - peripheral sensation counselling intervention 232–4
 - vestibular system and visual field dependence counselling intervention 234
 - visual intervention 231–2
 - see also* physiological profile assessment (PPA)
- tetracyclic antidepressants 84
- thiazide 87
- translational perturbation 24–5
- tricyclic antidepressants 84–5
- urinary incontinence 67–8
- vasodilators 87
- vestibular pathology 57, 112
- vestibular sense 46–7, 110
 - future research directions 239
 - targeted counselling intervention 234
 - see also* balance; postural stability
- vibration sense 44
- vision 41–4
 - assessment of 194–5
 - contrast sensitivity 42–4, 64
 - targeted visual intervention 231–2
 - visual acuity 41–2, 64, 194–5
- visual field dependence 44
 - targeted counselling intervention 234

Cambridge University Press
0521589649 - Falls in Older People: Risk Factors and Strategies for Prevention
Stephen R. Lord, Catherine Sherrington and Hylton B. Menz
Index
[More information](#)

249 **Index**

- visual problems 64–6
 - cataracts 64–5
 - glaucoma 65–6
 - macular degeneration 65
- voluntary stepping 25–6
- walking
 - age-related changes 26–9
 - exercise 127–8
 - see also* endurance training
 - navigating obstacles 29–30
 - postural stability 26–9
 - tandem walking 29
- walking aids 166–70
 - alternatives 170–1
 - indications 166–8
 - limitations 169–70
 - prescription principles 168–9
- withdrawal of medication 206–9