

Index

Note: page numbers in *italics* refer to figures and tables.

- ABC schizophrenia study 220
 Aboriginal peoples 327–8, 356–61
 alcohol use 359
 health status 357
 suicide 360
 abuse *see* alcohol-use disorders; child abuse;
 drug use/abuse; substance-abuse
 disorder
 access to treatment 110–11
 accountability
 financial restrictions 359
 mental health services 335–6
 acquired immunodeficiency syndrome
 (AIDS) 369, 372, 375
 acupuncture 175
 scientific evidence 183
 adaptive behavior impairment 346
 addictive disorders, prevalence 45
 adherence *see* compliance
 admission to psychiatric ward 400
 adolescents 32, 327, 330
 anxiety disorders 195–6
 autonomy 332
 depression 239, 337
 self-medication 734
 suicide 239
 treatment intervention outcomes 337
 affective disorder
 prevalence 17, 18, 20
 see also bipolar affective disorder
 Africa 238
 health systems 369
 age
 population structure 210–11
 public knowledge/attitudes 408–9
 aggressive behavior syndrome prevention
 1347
 evidence-based preventive interventions
 1357
 risk factors 1345
 aggressive syndromal patterns 134–7
 agoraphobia 263, 265, 267, 269
 agreement about need 271–2
 Aid for Families with Dependent Children
 (AFDC) 70
 alcohol
 dependence 73
 detoxification 297
 public education 295
 alcohol, drug and mental (ADM) disorders
 378
 alcohol-use disorders 196, 290
 Aboriginal peoples 359
 Composite International Diagnostic
 Interview (CIDI) 292–3
 comorbidity 290, 292, 293, 297–8
 detoxification 297
 Epidemiologic Catchment Area (ECA)
 study 291–2
 epidemiology 291
 hazardous use 295–6
 inpatient hospitalization 297
 matching hypothesis 297
 National Comorbidity Study (NCS)
 292–3
 outcome 291
 prevalence 292
 primary care 161
 psychiatric diagnoses 293–4
 public health 295
 public knowledge/attitudes 405
 screening 295–6
 specialist addiction treatment 296–8
 treatment 294–5
 Alcoholics Anonymous 297
 alternative medicine *see* complementary
 medicine
 alternative treatments, public
 knowledge/attitudes 405
 American Psychiatric Association (APA)
 146
 practice guidelines 148–50, 154

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)**430 Index**

- American Psychiatric Association (*cont.*)
 practice research network 154
 antidepressants 87
 medication frequency 93
 public perception 94
 antisocial behavior, suicide 251
 antisocial personality disorder, drug
 dependence co-occurrence 306
 antisocial spectrum condition prevention
 137
 antisocial syndromes 134–7
 anxiety
 cognitive-behavioral therapy 257
 complementary medicine 183–7
 need for treatment 902, 93
 postnatal 13940
 prenatal/postnatal 139–40
 predisposition 138
 prevention 138–42
 primary care 161
 primary prevention 32
 prognosis 406
 somatoform disorder overlap 318
 anxiety disorders 256–7
 budget allocation 29
 cognitive-behavioral therapy 257
 comorbidity 284
 counseling 272
 depression 236–7
 diagnostic criteria 256
 EDSP 270, 271
 exposure therapy 257
 inadequate treatment 268–70
 inappropriate treatment 272–3
 Munich Community Study 259–65, 266,
 267–73
 needs
 assessment 256–7
 evaluation 258–9
 pharmacological therapy 256–7, 286,
 287
 prevalence 17, 18, 20
 primary care 286
 psychiatrist treatment 286
 psychoeducation 257, 272
 psychotherapy 286
 self-help 257, 272
 self-perception of need 258
 service provision 257–8
 service use 284–6
 somatic symptoms 315
 specialist interventions 272
 spontaneous remission 257
 treatment 21, 256–7
 needs in community 257–8
 young people 195–6
 anxious disposition 314
 appearance in public, psychotic illness 385,
 387, 388
 attention deficit hyperactivity disorder
 (ADHD) 134
 audit of primary care 167
 Australia
 alcohol education 295
 budget allocation 30
 community-based surveys 422
 epidemiological survey 9, 151–7, 18,
 1922, 23, 245
 evidence-based medicine 13
 mental health expenditure 20
 National Mental Health Strategy 34
 Positive Parenting for Preschoolers
 135–6
 practice guidelines 154
 public knowledge/attitudes 399, 400,
 401, 402–9
 public perception of antidepressants 94
 somatoform disorders 316
see also Aboriginal peoples; National
 Survey of Mental Health and
 Well-Being (Australia)
 Austria, public knowledge/attitudes 399,
 400, 401, 402–9
 autogenic training for panic disorder 279
 automation of clinical psychiatry 329
 autonomy
 adolescents 332
 help-seeking 391
 Ayurvedic medicine 174, 175–6
 barriers to care 396, 397
 behavior therapy
 disability reduction 319
 public knowledge/attitudes 405
 somatoform disorders 318
 behavioral health problems
 low-income countries 375
see also conduct disorders
 Behaviorally Based Preventive Intervention
 137
 benefits
 economic 656, 121, 124
 health 121–3
 maximization with resource allocation
 66
 bereavement 1401, 409
 Berkeley Puppet Interview (BPI) 352
 bipolar affective disorder 26, 48
 black box of health care plans 146, 147
 Brief Disability Questionnaire 19
Bringing them Home 357
 Britain *see* United Kingdom
 British Medical Association (BMA) 173
 complementary medicine 178

- British National Survey of Psychiatric Morbidity 87, 88, 89, 93
- budget allocation 29–30, 32
- budget holders, complementary medicine 190
- budget-capping 127
- buffering factors 133
- bullying 136–7
- burden
 - depression on family 237–8
 - untreated illness 127
 - see also* disease burden; economic burden
- Burden of Disease Study, alcohol-use disorders 294
- Camberwell Assessment of Need 200–1, 203, 391
- Camberwell Needs for Care Survey 85, 90, 91, 93
- Canada
 - Aboriginal peoples 360
 - access to treatment 110–11
 - depression epidemiology 237
 - epidemiological survey 9, 141–5
 - International Consortium of Psychiatric Epidemiology (ICPE) 97, 989
 - mental health service system 99
 - service system 101
 - service use 1057
- cannabis dependence 303
- Canterbury Suicide Project 247–52
- capitation 127, 146
- Cardinal Needs Schedule 201–2, 391
- care
 - barriers to 396, 397
 - continuing 161
 - continuum 53
 - for children 337–8
 - cost 523, 54
 - demand 198–9
 - ethical imperative 6
 - hierarchy 420
 - need 1978
 - prioritizing 29–30, 420
 - quality monitoring 75
 - suicide attempts 252
 - supply 199
 - utilization 199
- cartoon-based assessment for children 351
- case identification 425–6
- causal chain 133
- causation web 366
- causes of mental illness, public knowledge/attitudes 4067
- cerebrovascular disease 236
- child abuse 135, 139
 - sexual 139, 141, 338
- childbirth care 135
- children 32, 327, 330
 - clinical interventions 336
 - continuum of care 337–8
 - continuum of mental health problems 332
 - coordination of treatment 337–8
 - depression 336
 - development 3312
 - emotional disorders 331
 - evaluation of interventions 341
 - features of mental disorders 330–2
 - four to eight year olds 345–53
 - cartoon-based assessment 350–2
 - informants 346–8
 - misclassification of service use/need 348–50
 - need assessment 345–6
 - parents 346–7
 - prevalence of unmet need 349–50
 - psychopathology assessment 350–3
 - puppet measures 352–3
 - self-reporting 353
 - teachers 346–7
 - identification of needs 332–5
 - inappropriate treatment 334, 336–7
 - ineffective treatment 334
 - measurement of needs 327
 - medical necessity 489
 - outcome of treatment 337–8, 340
 - prevention of disorders 338–40
 - primary care 331
 - resources allocation 332
 - risk factors 338
 - sub-syndromal disorders 336
 - treatment of disorders 335–8
 - vulnerability-focused interventions 339
 - vulnerable 142
- Children of Divorce Intervention Program 140
- children's global assessment scale (CGAS) 48
- Chinese Health Questionnaire 367
- Chinese medicine, traditional 174, 175
- chiropractic, scientific evidence 183
- chronic fatigue syndrome 312, 314
- clients, need 157
- clinical decisions 9
- Clinical Interview Schedule, Revised (CIS-R) 87
- Clinical Practice Guidelines (US) 235
- clinical trials 33, 152
 - complementary medicine 189
 - placebo response 257
- cocaine dependence 303
- Cochrane Collaboration 33
- cognitive impairment 17, 18, 20

432 Index

- cognitive-behavioral treatments
 - anxiety disorders 257
 - panic disorder 277–8, 279
 - past-therapy outcome 278
 - social phobia 277–8
 - somatiform disorders 318
 - techniques 166
- common mental disorders
 - disability 374
 - efficacious treatments 162–5
 - predictors of outcome 162
 - primary care 161, 162–5, 166
 - Primary Mental Health Care Project 373–4
- Commonwealth government 357
- community
 - adequacy of mental health provision 31
 - anxiety disorders treatment 257–8
 - disorganization 133
 - education 11, 131–4
 - health care procedures 13
 - institutions 133
 - medical necessity estimates 416
 - needs 3, 7
 - projects 42
 - specialist care for anxiety disorders 283–4, 285
- community care 391
- Community Care Act (UK; 1990) 201, 213
 - need definition 198
- community factors 133
 - depression 139
- community health workers, primary care 166
- Community Mental Health Centers (CMHC) legislation (US) 42, 44
- community-based surveys 422
- comorbidity
 - alcohol-use disorders 290, 292, 293, 297–8
 - anxiety disorders 284
 - drug dependence 304
 - drug use 306
 - physical disorders 19
 - utilization prediction 94
- complementary medicine 21, 115, 1723
 - attractions 188
 - beliefs of practitioners 179–80
 - budget holders 190
 - consumer spending 173
 - cost 190
 - developed after 1790 174–6
 - differences from orthodox 177
 - effectiveness studies 189
 - medical school curriculum 1901
 - popularity 177–8
 - psychiatric disorders 183–7
 - religious faith 175
 - respect 189
 - scientific evidence 180, 1813
 - self-selection 177
 - tradition before 1790 174
 - usage 172–3
 - varieties of treatments 174–6
 - views of orthodox doctors 178–9
- compliance 27983
- Composite International Diagnostic Interview (CIDI) 15, 105, 106, 256, 258
 - alcohol-use disorders 292–3
 - anxiety disorders 256, 284
 - perceived need assessment 392
 - see also* Munich CIDI
- computer technology 154, 419, 420
- concensus-based guidelines 166
- conduct disorders 134–7
 - primary prevention 32
 - see also* behavioral health problems
- cone of morbidity 423, 424
- consultation
 - behavior 212, 23, 245, 312
 - disability correlation 22, 24
 - effect 184–5, 188
 - healing power 186
 - rate 24
 - transactions 186
 - Consumers Association (UK) 173
- consumer movement 391
- consumers
 - attitudes to depression treatment 237
 - demand 159
 - empowerment 391
 - provider selection 111
 - role of individual 391
- contact with mental disorders, public
 - knowledge/attitudes 408–9, 410
- continuing care 161
- continuum of care 53
 - children 337–8
- ‘Coping with Stress’ intervention 336
- core responsibilities, defining 301
- cost offset approach 667
- cost-benefit analysis 121, 123
- cost-benefit ratio 65, 68
 - not treating treatable illness 127
 - treatment 1256
- cost-effectiveness 77, 78
 - analysis 121, 123
 - disease burden 165
 - interventions 70, 71
 - preventive interventions in children 340
 - research in low-income countries 372–3
 - studies 128

- cost-utility analysis 121, 123
- costs
- bearing 65–7
 - of care 52–3, 54
 - comparisons 128
 - complementary medicine 190
 - cutting 164
 - depression 238
 - direct 68
 - indirect 63–5, 67–70, 78
 - managed care 128–9
 - mortality/morbidity related 125
 - not treating treatable illness 125, 126–7
 - treatment of previously untreatable people 125
 - untreated cases 126
- counseling, anxiety disorders 272
- counselors 110
- public knowledge/attitudes 403
- criminal justice system, Indigenous peoples
- representation 356–7
- criminality, suicide association 2501
- cross-cultural investigations 365, 366, 367
- cross-national comparisons
- demographic variation 108
 - disorder type 107–8
 - provider availability 100
 - questionnaire design 112–16
 - sector use 103–5
 - service use 98, 101, 102, 103, 105–7
- cultural informants 379
- culture, health system 369–70
- culture-bound syndromes 110
- Defeat Depression Campaign (UK)
- 409–10, 411, 417
- definitions
- limited 49
 - one-year prevalence 50
- deinstitutionalization 391
- delusion of persecution/psychosis, Austrian
- public knowledge/attitudes survey 401, 4029
- demand 86
- care 198–9
 - for treatment/services 85
- dementia 6
- prognosis 406
 - public knowledge/attitudes 405
- demographic factors 209–11
- depression
- adolescents 239, 337
 - anxiety disorders 2367
 - awareness 235
 - budget allocation 29
 - children 336
 - community factors 139
 - community impact 238–9
 - complementary medicine 183–7
 - disability-adjusted life years (DALYs) 233
 - early intervention 141
 - early-onset 236
 - economic cost 238–9
 - environmental factors 138–9
 - epidemiology 237–8
 - evidence-based interventions 139–41
 - families 237–8
 - fundholder relevance 239–40
 - gender 239
 - general practitioner treatment 255
 - health planning 239–40
 - Irish public knowledge/attitudes 409
 - late-onset 236
 - low-income countries 3745
 - maternal 139–40
 - morbidity 138
 - need 233
 - for treatment 902
 - optimism enhancement 140
 - personality vulnerability 236
 - postnatal 139–40
 - predisposition 138
 - prenatal 139–40
 - primary care 161, 234–6
 - primary prevention 32
 - productivity loss 65
 - prognosis 405–6
 - protective factors 138, 139
 - psychotherapy 405
 - public knowledge/attitudes 410–11
 - pharmacological treatment 404
 - referral for specialist treatment 235
 - relapse potential 237
 - risk factors 138
 - schizophrenia 221, 224–7, 228, 229–30
 - significance 233
 - smoking 238–9
 - somatoform disorder overlap 318
 - specialist care provision 236–7
 - stressful life experiences 1401
 - substance abuse 239
 - suicide 250
 - terminology in Zimbabwe 367
 - tobacco use 238–9
 - treatment 235
 - undertreatment 234
 - unmet needs 92
 - workplace performance 67
- depressive disorders
- pain 315
 - prevention 136–42
 - treatment 87, 88, 89

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)**434 Index**

- depressive states, somatoform disorders 314–15
- depressive symptoms, subthreshold 239
- developing countries *see* low-income countries
- diagnosis 15
 - computer technology 419
 - medical necessity 49–50
 - need prediction 159–60
- diagnostic classification 86
- Diagnostic Interview Schedule for Children (DISC) 347
- Diagnostic Interview Schedule (DIS) 51, 293
 - somatoform disorders 311
- diagnostic related groups (DRGs) 52
- Diagnostic and Statistical Manual of Mental Disorders see* DSM
- diet manipulation, public
 - knowledge/attitudes 405, 408
- disability 17, 18, 19–20
 - behavioral strategies for reduction 319
 - common mental disorders 374
 - community setting 283–4, 285
 - consulting correlation 22, 24
 - days 19–20
 - health service use 284–6
 - measures 52
 - mental disorder classification 49
 - neurasthenia 316, 317
 - panic disorder 283–6
 - social phobia 283–6
 - specialist setting 283–4, 285
- Disability Assessment Schedule 220, 226
- disability-adjusted life years (DALYs) 257, 51, 160
 - depression 233
 - economic criteria 121, 122
 - primary care 165
 - rationing of health care 29
- disablement 328
 - ADM disorder 378
 - condition ranking 379
 - ranking 383
- discrete-time survival analysis 69
- disease 5
 - interventions 6
 - ranking 11
- disease burden 26, 27, 28, 334
 - cost-effectiveness analysis 165
 - economic 120
 - global 160, 162
 - measurement 69
 - priorities 27
- divorce 72, 139, 140
- doctorpatient relationship models 391
- drug dependence 73, 302–4
 - comorbidity 304
 - needs assessment 303–4
- drug use/abuse 32
 - altruistic sharing 304, 305
 - comorbidity 306
 - ‘honeymoon’ period 3045
 - initial opportunity 305–6
 - person-to-person spread 304–5
 - transition from initial opportunity 305–6
- DSM criteria, children 346
- DSM-III
 - alcohol abuse 291
 - anxiety 256
 - anxiety disorders 256
 - classification 14, 15
 - criteria-based disorders 467
 - syndrome prevalence 43
- DSM-III-R
 - attempted suicide 245
 - disorders 105, 106
- DSM-IV 146
 - anxiety disorders 256
 - evidence-based diagnostic guidelines 148
 - somatoform disorders 311
- dysthymia 239
 - complementary medicine 183–7
- Early Developmental Stages of Psychopathology (EDSP) 258
 - anxiety disorders 270, 271
- ecology of medical care 151–2
- economic burden 120–1
 - not treating treatable illness 125, 126–7
- economic costs of depression 238
- economic criteria 120–1
- economic effectiveness evaluation 121, 123, 124–5
- economic factors in somatization 316
- economic health 10
- economic outcome of treatment 124
- economic rationalism 426–7
- economists 119
- education 13, 312
 - alcohol 295
 - community 11, 13–14
 - literacy in mental health 328
 - mental health services 13–14
 - public knowledge/attitudes 408–9, 410
 - see also* medical school curriculum; medical training
- elderly people 139, 211
- emic approach 366, 367
- emotional disorders in children 331
- employment 65, 67, 69, 139
- environmental factors, depression 138–9

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)

- environmental influence 134
- Epidemiological Catchment Area (ECA)
 - study 39, 50–1, 291–2
 - costs 1267
 - diagnosis 293
 - severe mental illness prevalence 60
 - somatiform disorders 311
 - unmet needs 93–4
- epidemiological surveys 9, 11
- epidemiology 418, 422–3
 - alcohol-use disorders 291
 - health economics 56
 - health services 56
 - instruments 55
 - rates of prevalence of psychiatric disorder 204–5
- equivalence metrics 66
- estimates of unmet need, believable 425
- ethical imperative, care 6
- ethnicity 210
- etic approach 367
- Euro-American psychiatric diagnoses 366
- Europe, health systems 369
- evaluation services 4
- evidence
 - nature of 1802
 - objective 180, 181
 - scientific 180, 181–3
 - subjective 180, 181
 - systematic 180, 181
- evidence-based interventions 146–8
 - for depression 139–41
 - guideline development 154
 - limitations 151–2
 - preventive 135–7
 - quandaries 152–3
 - risk adjustment 153
 - see also* practice guidelines
 - evidence-based medicine 121–3, 21, 33, 147–8
 - primary care practice 166
 - resources for children 332
- exaggerated estimate (EE) problem 424–6
- expenditure
 - mental health 20
 - total 4
- explanatory styles, positive 140
- exposure therapy for anxiety disorders 257
- families
 - depression 237–8
 - needs 3, 5, 7
 - public knowledge/attitudes 403
- Family Bereavement Program 140–1
- family practice training, somatiform disorders 319
- family practitioners *see* general practitioners
- family therapy, public knowledge/attitudes 405
- fatigue
 - evaluation 315
 - somatiform disorders 314–15
 - syndromes 316
 - treatment 318
- fee-for-service system 32, 45, 127, 128, 146
- fibromyalgia 312, 318
- filter
 - between community and psychiatric admission 400
 - concept 423
- finance 119
 - barriers to treatment 127–8
 - evidence-based approaches 146
 - mechanisms 127
 - primary care 163–4
 - public 124–5
 - see also* funding
- Finland 93–4
- friends, public knowledge/attitudes 403
- fundholders
 - complementary medicine 190
 - depression relevance 239–40
- funding
 - government restrictions 9
 - types 127–8
 - see also* finance
- gender
 - aggressive behavior syndrome risk 134
 - depression 239
 - depressive symptoms 138
 - schizophrenia 230
 - social perception 138
- general practitioners 417–18, 423
 - abilities 417
 - comparison with public knowledge/attitudes 408
 - cone of morbidity 423, 424
 - depression treatment 255
 - low-income countries 372
 - Netherlands 109
 - Pathway to Care role 400
 - public knowledge/attitudes 403
- referral 400
 - role with children 333
 - see also* primary care
 - generalized anxiety disorders (GAD) 263, 265, 267–8
 - mental health service use 264
- generalized health questionnaire (GHQ) 89
- symptoms 48

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)**436 Index**

- genetic factors 133–4, 406
 - aggressive behavior syndromes 134
- Germany
 - community-based surveys 422
 - mental health service use 264–5
 - Munich Community Study 259–65, 266, 267–73
 - public knowledge/attitudes 399, 400, 401, 4029
 - public perception of antidepressants 94
- global assessment scale (GAS) 47–8
- Global Burden of Disease (GBD) study 689, 70
- goals 4
- ‘Good Behavior Game’ 338
- governments
 - funding restrictions 9
 - health status of population 7
 - needs 3
 - planning 334
- Griffith Early Intervention Project 141
- gross domestic product (GDP) 8
- group therapy, public knowledge/attitudes 405
- Handbook of Psychiatric Measures and Outcomes* 148
- handicap, role performance 425
- headache 314–15
- health gains, maximization 12–13
- health insurance 8
 - parity for mental health 55, 128–9
 - private in US 42, 43, 55, 100
- health maintenance organizations 128
- Health of the Nation* 204
- health professions, needs 34
- health status of population 7, 10
- health system
 - culture 369–70
 - families 370
 - India 369
 - low-income countries 368–9, 370
 - Zimbabwe 369
- health systems research (HSR) 363–4
 - dissemination 373
 - model 328
 - psychiatric research 370–3
- help, perception 22
- help-seeking 22
 - autonomy 391
 - perceived need assessment 395–6
- herbal medicines, public
 - knowledge/attitudes 405
- heroin dependence 303
- hierarchy of care 420
- homelessness 210
 - economic 200
 - multiproblem 200
 - needs assessment 199–200
- homeopathy 174, 176
 - scientific evidence 183
- hospitalization, inpatient 297
- housing 32, 133
- human capital 120–1, 122
- human resources 417
- human services sector 109–10
- ICD criteria, children 346
- ICD-10
 - anxiety 256
 - anxiety disorders 256
 - British National Survey of Psychiatric Morbidity 87
 - burden of disease 26
 - classification 15, 16
 - criteria-based disorders 467
 - psychiatric disorders with somatic symptoms 187–8
 - somatiform disorders 311
 - syndrome prevalence 43
 - ‘ice-smoking’ 304
- illness 5
 - behavior 400
 - course 49–51
 - supernatural 374
- impairment, mental disorder classification 478
- India
 - Ayurvedic medicine 174, 175–6
 - community-based surveys 422
 - health system 369
- Indigenous mental health 356–61
 - morbidity/mortality 357–8
 - professional interest 358–9
 - US 358
- individual, role of 391
- informants, child mental health 347–8
- information officers 167
- institutional care 391
- institutionalization 422
- internalizing syndromes, prevention 138–42
- International Consortium of Psychiatric Epidemiology (ICPE) 97–8
 - care providers 101, 103
 - contextual information 99–100, 101
 - demographic variation 108
 - disorder type 107–8
 - DSM-III-R disorders 105, 106
 - psychiatric need 105–7
 - questionnaire design 112–16
 - service system 101
 - service use 105–7
 - service utilization 102

- survey information 98–9
- interventions
 - appropriateness 60, 62
 - cost-effectiveness 70, 71
 - costs 612
 - delivery quality 74
 - effective 5, 7
 - expected benefits 426
 - follow-up 72
 - harmful 24
 - implementation 56
 - inadequate resources 63–70
 - physical illness 61
 - preventive 70–4
 - range used by professionals 411
 - resource allocation 61
 - universal 61
- interview for the retrospective assessment of the onset of schizophrenia (IRAOS) 218, 219–20
- invariance, cross-cultural investigations 365
- investment 4
- Ireland, public knowledge/attitudes 409
- iridology 176
- irritable bowel syndrome 312
 - treatment 318
- LAAM 303
- learned helplessness 138
- life course, major transitions 69
- Life Events and Difficulties Schedule 89
- life expectancy 13
- literacy in mental health 3289
- longevity, indirect cost 64
- longitudinal studies 50–1, 236–7
 - attempted suicide 246–7
 - panic disorder 278–9
- low-income countries 327, 328, 363–4
 - behavioral health problems 375
 - depression 374–5
 - etic-emic debate 365–8
 - general practitioners 372
 - health priorities 363
 - health system 368–9, 370
 - human resources 418
 - incidence of mental illness 364
 - nongovernmental organizations 370, 372, 373
 - primary care 419
 - sexually transmitted disease 369
 - universalist approach 365–6
- Luxembourg 328
 - psychotic disorder perception 378–89
- MacArthur Foundation 67, 353
- manageable symptoms 423
- managed care 456, 523, 55, 127, 129, 146
 - commercial 420
 - epidemiological data 45
- market-based model 147–8
- marriage 64, 69, 139
- Marshfield Primary Care Study 50
- ‘Mastery of Learning’ intervention 338
- maternal depression 139–40
- maternal overprotection 237
- medical anthropology 366
- medical necessity
 - children 489
 - community estimates 416
 - concept 46
 - cost of care 523
 - disability-adjusted life years (DALYs) 51
 - diagnosis 49–50
 - epidemiology of clinically significant disorders 469
 - illness course 49–51
 - planning 55–6
 - severe mental disorders 51–2
- medical practice 323
- Medical Research Council (UK)
 - Needs for Care Assessment 85, 89, 90, 91, 200, 202
 - Social Role Performance Schedule 89
- medical school curriculum
 - complementary medicine 172, 173, 189, 190–1
 - Zimbabwe 374
- medical students 173, 190, 374
- medical tests 310
- medical training 417
 - somatiform disorders 319
- meditation, public knowledge/attitudes 405
- mental disorders
 - recognition by public 403
 - role in care 400
 - somatic symptoms 187–8
 - see also* common mental disorders; severe mental illness
- mental health
 - care quality monitoring 75
 - expenditure 20
 - insurance parity 55, 128–9
 - literacy 328–9
- Mental Health Supplement (Ontario, Canada; 1990) 98–9
- Mental Illness Needs Index (MINI) 211
- met need 11–12
 - epidemiological survey 14, 15
- met un-need 12
 - epidemiological survey 14
 - incidence 21
- methadone 303

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)**438 Index**

- methamphetamine 304
- Midtown Manhattan study 55
- morbidity 132, 133
 - burden 11
 - cone 423, 424
 - generic risk factors 133
- mental disorder impact 132
- mortality burden 11
- motor function disorders 236
- moxibustion 175
- Munich Community Study
 - anxiety disorders 259–65, 266, 267–73
 - care system 262–3
 - counseling 272
 - diagnostic assessment 261
 - expert-rated need 263, 265, 266, 267–8, 271–2
 - health service use 262–3
 - inadequate treatment 268–70
 - inappropriate treatment 272–3
 - mental health service use 264–5
 - methods 261–3
 - need assessment 262–3
 - population affected by anxiety disorders 263–4
 - self-rated need 265, 266, 2678, 269, 271–2
 - service availability 271
 - services received 268–70
 - type of anxiety disorder 271
- Munich Prospective Community Survey 94
- Munich Composite International Diagnostic Interview (M-CIDI) 261–2
- National Comorbidity Study (NCS) 39, 98, 424
 - alcohol-use disorders 292–3
 - costs 127
 - diagnosis 293
 - indirect costs 64–5
 - preventive interventions 72, 73
 - severe mental illness prevalence 60
- national health insurance 8
- National Health Service (UK), budget holders 190
- National Institute of Mental Health (US) 199, 351
 - ICPE 97
- National Institutes of Health (US), funding 148
- National Primary Care Facilitation Programme (UK) 235
- National Survey of Mental Health and Well-Being (Australia) 15–25, 34, 391–2, 396–7
- needs
 - agreement 271–2
 - care 197–8
 - clients 157
 - comparative 333, 335
 - definition 4, 7, 85, 120, 159–60, 197–8
 - depression 233
 - deprivation-weighted approach 207–8, 209
 - determination 37
 - diagnosis 159–60
 - estimation 160
 - national patterns of use 205–7
 - expressed 333, 334
 - felt 333, 334
 - gradient 105
 - identification 86–7
 - individual-level assessment 199–202
 - management 159
 - manufacture 159
 - meetable 93
 - normative 333–4
 - perceived 86
 - population 87
 - prediction 159–60
 - providers 157
 - services 157
 - see also* met need; met un-need; unmet need
- needs assessment
 - anxiety disorders 256–7
 - individual-based 160
 - population-level 85, 160, 203–13
 - strategy 16–45
 - tools 391–2
 - Perceived Need for Care Questionnaire 392–7
- Needs for Care Assessment 85, 89, 90, 91, 200
 - modification 202
- Netherlands
 - access to treatment 110–11
 - formal care 108–9
 - ICPE 97, 99
 - mental health service system 99
 - service provision 104
 - service system 101
 - service use 105–7
- Netherlands Mental Health Survey and Incidence Study (NEMESIS) 99
- neurasthenia 17, 18, 311
 - disability 316, 317
- neurocognitive disorders 236
- neurotic disorder diagnosis 87, 88, 89
- neuroticism 138, 139, 237
- New Zealand
 - Canterbury Suicide Project 247–52
 - practice guidelines 154

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)

- suicide rate 245–6
- nicotine dependency 239
- nongovernmental organizations 370, 372
- nontreatment, inappropriate 623
- Norway
 - bully-victim interventions 136–7
 - mental disorder surveys 302
- nurses
 - primary care 166
 - terminology in Zimbabwe 367
- objectives 4
- OPCRIT (Operational Criteria Checklist) 202
- operant learning, pain 316
- opioids 303
- oppositional syndromes 134–7
- optimism 138
 - enhancement 140
- Oregon project 9, 28
- Organization for Economic Co-operation and Development (OECD) 8
- orthodox medicine, evidence 179–80, 181–2
- osteopathy, scientific evidence 183
- outcome
 - alcohol-use disorders 291
 - assessment 4, 76, 77
 - averages 3
 - children's treatment 337–8
 - comparisons 153
 - continuous 77
 - economic 124
 - expected 127
 - mental health treatment 46
 - past-therapy 278
 - predictors for common mental disorders 162
 - risk-adjustment 76
 - standards 756
- overprovision 199
 - rating 90
- overutilization measure 120
- pain
 - chronic disorder 311, 314–16
 - depressive disorders 315
 - operant learning 316
 - primary psychogenic disorder 315
- pain behavior, bizarre 315
- palliative care 323
- panic attacks, prevention 317–18
- panic disorder 265, 267–8, 277–9
 - autogenic training 279
 - cognitive-behavioral treatments 277–8, 279
 - disability 283–6
 - mental health service use 264
 - past-therapy outcome 278
 - pharmacological treatment 279
 - prevalence 28
 - psychodynamic therapy 279
 - treatment dropout 279–83
- parenting 69, 133, 135–6, 137, 237
 - adverse 139, 338
- parents
 - children with mental disorders 330–1, 335
 - children's mental health needs 346–7
- parity of treatment 55, 128–9, 164
- paternal care 237
- pathways to care 161, 328
 - model 400
- patients
 - with ADM disorder 378–89
 - needs 3, 455
- payers, fifth party 164
- payment mechanisms 127
- pediatricians 331
- role 333
 - Penn Prevention Program 336
- perceived need assessment 390–7
- Perceived Need for Care Questionnaire 392–7
- personality disorder 17, 18
- personality vulnerability to depression 236
- pharmaceutical industry 153
- pharmacological treatment
 - anxiety disorders 256–7, 286, 287
 - panic disorder 279
 - public knowledge/attitudes 404
- physical disorders, comorbidity 19
- physical illness, intervention 61
- physicians, primary care 166, 167
- placebo response 257
- planned child mental health services
 - research study, costs 127
- planning 4, 334, 328–9, 418–19, 421
 - depression 239–40
 - medical necessity 556
 - public knowledge/attitudes 410–11
- policy making 328–9
 - see also* planning
- population, health status 8, 10
- Positive Parenting for Preschoolers 135–6
- postnatal depression
 - with anxiety 139–40
 - support networks 238
- posttraumatic stress disorder 141
- poverty 133, 139
 - living conditions 42
- practice guidelines 148–50, 154
- practice research network 154
- practice-based research 154–5

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)**440 Index**

- Preferred Provider Organizations 128
- pregnancy, teenage 32, 70
- prenatal care 135
- prenatal depression/anxiety 139–40
- Prenatal/Early Infancy project 135
- President's Commission on Mental Health (US; 1978) 42
- prevalence of mental disorders 16–17
- community 44
 - epidemiological rates 204–5
 - treatment 44, 53
 - underestimates 424–5
- prevention 13, 32, 132–4
- aggressive behavior syndromes 134–7
 - antisocial spectrum conditions 137
 - anxiety 138–42
 - depressive disorders 136–42
 - evidence-based interventions 135–7
 - internalizing syndromes 138–42
 - interventions 704
 - mental health services 131–4
 - relapse 167
 - suicide 246
 - withdrawing syndromal patterns 138–42
- see also* primary prevention
- primary care 157–8
- anxiety disorders 286
 - audit 167
 - caseness 161–2
 - children 331
 - children's services 345–6
 - common mental disorders 161, 162–5, 166
 - community health workers 166
 - depression 234–6
 - disability-adjusted life years (DALYs) 165
 - education program 163
 - evidence-based practice 166
 - finance 163–4
 - low-income countries 419
 - mental health role 158
 - needs assessment strategy 164–5
 - nurses 166
 - physicians 166, 167
 - psychiatric disorders with somatic symptoms 187–8
 - public health 167–8
 - resource allocation 164
 - service provision 163
 - somatiform disorders 319–20
 - teams 167
 - teamwork 166
 - training 166
 - unmet need 160–2, 423
 - see also* general practitioners
- Primary Mental Health Care Project (Zimbabwe) 364, 373–4
- Primary Mental Health Toolkit 235
- primary prevention 11, 32
- prioritization 29–30, 420
- priority setting 33
- PRiSM Psychosis Study 202–3
- expected levels of service 208, 209
- private sector *see* health insurance
- process assessment 77
- productivity, loss 65
- professional help, public knowledge/attitudes 403–4
- professionals
- comparison with public knowledge/attitudes 408, 410, 411
 - disablement associated with chronic psychosis 378–89
 - range of interventions 411
- prognosis, public knowledge/attitudes 405–6
- Project MATCH 297
- protective influences 133
- providers
- availability 100
 - need 157
 - selection 111
- psychiatric services
- cone of morbidity 423, 424
 - use 42
- psychiatrists 417
- anxiety disorder treatment 286
 - comparison with public knowledge/attitudes 408
 - computer technology 419
 - outpatient treatment 418
 - public knowledge/attitudes 403, 404
 - specialist interventions 272
- psychoanalysis, public knowledge/attitudes 405
- psychodynamic therapy, panic disorder 279
- psychoeducation, anxiety disorders 257, 272
- psychological disorders, symptom presentation 185
- psychologists, public knowledge/attitudes 403, 404, 408
- psychopathology
- assessment in four to eight year olds 350–3
 - risk factors 32
- psychoses
- budget allocation 29
 - public order 31
- psychosis, prognosis 406
- psychosocial interventions 166
- psychosocial stress 406–7

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)

- psychotherapist interventions 272
 psychotherapy
 anxiety disorders 286
 depression 405
 public knowledge/attitudes 404–5
 psychotic disorders 210, 211
 psychotic illness
 appearance in public 385, 387, 388
 people's reactions 378–89
 social disapproval 385, 386
 stigma 388
 psychotropic medication, economic
 outcome 124
 public awareness programs 166–7
 public health 13
 alcohol-use disorders 295
 disease management 11
 education 295
 mental disorder impact 132
 primary care 167–8
 suicide 246
 public knowledge/attitudes 399–400
 public order 31
 public sector, managed care 55
 public services
 privatization 55
 provision 127
 Puerto Rico
 access to treatment 110–11
 ICPE 97, 99
 mental health service system 100, 101
 service use 105–7
 Puerto Rico Mental Health Care Utilization
 (MHCU) Project 99
 puppet measures 352–3
 purchasing health services 9

 quality assurance 167
 perceived need 397
 standards 74–7
 quality of life 3
 quality monitoring, mental health care 75
 quality-adjusted life years (QALYs) 66
 economic criteria 121, 122
 questionnaires
 alternative service assessment 115
 core bank design 11216
 covariates 115
 format 114
 outcomes 114
 priorities 113
 question placement 114
 question topics 113
 question types 114

 rationalism, economic 426–7
 rationing of health care 28–30

 strategies 163–4
 referral 424
 children by schools 335
 general practitioner 400
 reflexology 176
 reimbursement schemes 167
 relaxation 183
 public knowledge/attitudes 405
 religious faith 175
 research, HSR 370–3
 research and development (R & D) 27
 Research Diagnostic Criteria (RDC)
 disorders 47
 resource allocation 10, 59–62
 alcohol-use disorders 290–1
 benefit maximization 66
 children 332
 economic effectiveness 124–5
 inadequate 63–70
 intervention 61
 primary care 164
 societal decisions 68
 resources
 availability 120, 418
 utilization 11
 responsibility, assignment 6
 risk adjustment 767, 153
 risk factors 133
 generic 133
 psychopathology 32
 public knowledge/attitudes 406–7
 resource allocation 68
 somatoform disorders 313–14
 road accidents, single-vehicle 249
 role performance, handicap 425
 Royal Australia and New Zealand College of
 Psychiatrists 154
 Royal College of Psychiatrists (UK) 154
 rural areas 136, 239, 369, 370

 salaries 127
*Schedules for Clinical Assessment in
 Neuropsychiatry* (SCAN) 89
 schizophrenia 17, 18, 20, 195, 197
 age at onset of symptoms 220, 222
 Australian public knowledge/attitudes
 survey 401, 4029
 depression 221, 224–7, 228, 229–30
 duration of prephase 221, 224
 early course 218–20
 false-positive cases 218–19
 gender 230
 German public knowledge/attitudes
 survey 401, 402–9
 IRAOS 218, 219–20
 living condition association 42
 perception 378

442 Index

- prognosis 405–6
- prognostic implications of depressive schizophrenia (*cont.*)
 - symptoms 225–7, 228, 229–30
 - social course 227, 228, 229–30
 - social disability 226–7
 - society perceptions 378–89
 - symptom development 220, 221, 222, 223
- school-based programs 136–7
- schools 133
 - children with mental disorders 330–1
 - children's services 345–6
 - dropping out 32
 - failure at 139
 - population-focused interventions 339
 - referral of children 335
- screening, high-risk groups 167
- Second Step Violence Prevention Curriculum 136–7
- self-help
 - anxiety disorders 257
 - manuals for anxiety disorders 272
 - programs 163
 - public knowledge/attitudes 405, 408
- self-limiting condition 257, 268–9
- self-medication, substance abuse 73
- self-perception of need, anxiety disorders 258
- self-reliance 113
- self-reporting 65
- service availability, public
 - knowledge/attitudes 403
- service delivery, perceived need 397
- service provision
 - actual/predicted levels 212, 213
 - levels 207, 208
 - local modifying factors 208, 210
 - public knowledge/attitudes 410–11
- service use
 - anxiety disorders 284–6
 - cross-national comparisons 98
 - disability 284–6
 - epidemiology 512
 - patterns 205–7
 - sociodemographic factors 211
- services
 - Aboriginal peoples 327–8
 - decision-making 9
 - need 157
 - public provision 127
- severe mental illness 49, 50, 512
 - distinction from minor disorders 59–60
- severity, mental disorder classification 48
- sexually transmitted disease, low-income countries 369
- SF-12 Mental Health Summary Scale 17, 19, 20, 284, 285
- SF-36 Mental Health Summary Scale 285
- Shona models of illness 374
- Shona Symptom Questionnaire 367, 373
- sick role, healing procedure 187
- sickness 5
- smoking, depression 238–9
- Social Class and Mental Illness* study 55
- social deprivation, service needs weighting 2078
- social disability in schizophrenia 226–7
 - early course 218
- social disadvantage 133
- social disapproval
 - opinions 380
 - psychotic illness 385, 386
- social distance measurement 407
- social engineering 419
- social environment, causes of mental illness 406–7
- social equity 55
- social factors 133
- social functioning impairment 346
- social inequity 42
- social justice 328, 3567
 - Canadian Aboriginal peoples 360–1
- social perception, gender roles 138
- social phobia 264, 265, 267, 269, 277–9
 - cognitive-behavioral treatments 277–8
 - disability 283–6
 - past-therapy outcome 278
 - treatment drop-out 279–83
- social problems, medicalization 359
- social welfare benefits 52, 250, 251, 382
- social workers 110, 167
- society
 - reactions to psychotic illness 378–89
 - restrictions on people with psychotic illness 385
- sociodemographic factors, service utilization 211
- socioeconomic status 42
 - deprivation 89
- somatic symptoms 187–8
 - communication of emotional distress 315
 - reporting 312
- somatization 187
 - dimensional view 311
 - economic factors 316
- Somatization Symptom Index (SSI) 311
- somatoform disorders 187, 196, 309
 - anxiety overlap 318
 - classification 310–11
 - depression overlap 318
 - epidemiology 310–11
 - integrated model 312–15

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)

- medical assessment processes 309–10
- medical training 319
- need for treatment 319–20
- neurobiological factors 313–14
- prevalence 310, 319
- risk factors 313–14
- treatment approaches 317–18
- vulnerability 313–14
- specialty care 111
- stigma
 - opinions 380
 - programs for reduction 166–7
 - psychiatric disorders 73
 - psychotic illness 388
 - public knowledge/attitudes 407–8, 411
- Stirling County study 195, 237
- stress management, public
 - knowledge/attitudes 405
- stressful life experiences 140–1
 - vulnerable children 142
- substance abuse disorders 73–4
 - depression 239
 - prevalence 17, 18
 - public order 31
 - self-medication 73
 - see also* alcohol-use disorders; drug use/abuse
- suicide
 - Aboriginal peoples 360
 - adolescents 239
 - depression 250
 - rate 210
 - young men 239
- suicide, attempted 2457
- antisocial behavior 251
- Canterbury Suicide Project 247–52
- care 252
- criminality 250–1
- mortality 248–9, 251
- prevention 246
- psychiatric disorders 246
- psychiatric morbidity 249–50
- psychosocial outcome 250
- public health 246
- repeat attempts 249
- schizophrenia 224
- single-vehicle accidents 249
- subsequent monitoring 250
- supernatural illness 374
- supply, care 199
- supply-side controls 9
- survival analysis, discrete-time 69
- Sweden, community-based surveys 422
- symptom relationship with need for care 423
- system-accreditation processes 75
- teachers
 - children with mental disorders 330–1, 335
 - children's mental health needs 346–7
- tests, medical 310
- therapeutic alliance 109
- third-party payers 9, 163–4
- Torres Strait Islanders 356, 357, 358
- traditional beliefs 367
- traditional medical practitioners 372
- training
 - primary care 166
 - US 44
- trauma interventions 24
- treatment
 - appropriate 212, 32
 - clinical 13
 - complexity 13
 - compliance 279–83
 - coordination for children 337–8
 - demand 12
 - economic outcome 124
 - guidelines 77
 - inadequate 26870
 - inappropriate 21, 623, 272–3, 334
 - children 336–7
 - ineffective 334
 - need
 - anxiety 90–2, 93
 - depression 90–2
 - perceived 52
 - prevalence 53
 - service distribution 399
 - unmet need 25
 - see also* outcome
- triage 30
- trials *see* clinical trials
- tuberculosis 369, 375
- underprovision 199
- undertreatment *see* treatment, inadequate
- underutilization measure 120
- unemployment 32, 139, 210
- United Kingdom
 - Defeat Depression Campaign 409–10, 411, 417
 - epidemiological survey 9, 141–5
 - National Health Service 9
 - practice guidelines 154
 - prioritizing of health care 29–30
- United States
 - access to treatment 110–11
 - community-based surveys 422
 - Depression Awareness Recognition and Treatment (DART) 409
 - epidemiological survey 9, 14–15
 - evidence-based approaches 146–7

Cambridge University Press

052166229X - Unmet Need in Psychiatry: Problems, Resources, Responses

Edited by Gavin Andrews and Scott Henderson

Index

[More information](#)**444** **Index**

- evidence-based medicine 13
- health insurance 42, 43, 55, 100
- United States (*cont.*)
 - parity 128–9
 - ICPE 97, 98
 - Indigenous mental health 358
 - mental health service 412, 100
 - national health insurance 8
 - public health 13
 - public mental health budget 124–5
 - school role 331
 - service system 101, 1045
 - service use 105–7
 - specialty care 111
 - training of mental health specialists 44
- universalist approach in low-income countries 365–6
- universality, cross-cultural investigations 365
- unmet need 12, 92
 - epidemiological survey 14, 15
 - treatment 25
- utilization
 - care 199
 - comorbidity 94
 - managed care 128–9
 - national patterns 205–7
 - review firms 164
 - treatment/services 86
- validity 51, 56
 - cross-cultural investigations 365
- vascular depression hypothesis 236
- victims, bullying interventions 136–7
- violence
 - low-income countries 375
 - marital 64
 - prevention interventions 136–7
 - public knowledge/attitudes 407–8
- vulnerability, inherent 236
- vulnerability-focused interventions for children 339
- welfare benefits 52, 250, 251, 382
- well-being
 - indirect cost 64
 - perceived need assessment 391–2
 - see also* National Survey of Mental Health and Well-Being (Australia)
- ‘Whole-of-School’ interventions 339
- willingness to pay 121
- witchcraft 374
- withdrawing syndromal patterns, prevention 138–42
- work
 - self-reported loss 65
 - see also* employment; unemployment
- workplace cost approach 66, 67
- World Bank Burden of Disease (GBD) Study 258, 29, 160, 165
 - alcohol-use disorders 294
- World Health Organization (WHO)
 - complementary medicine 179
 - global burden of mental disorders 160, 162
 - ICPE 97
 - primary care education program 163
 - Study of Psychological Problems in General Health Care 157–68
 - Traditional Medicine Unit 179
- yoga, public knowledge/attitudes 405
- youth workers 335
- Zimbabwe 328
 - health system 369
 - medical school curriculum 374
 - Primary Mental Health Care Project 364, 373–4