

1

A brief history of cosmology

MALCOLM S. LONGAIR
Cavendish Laboratory, Cambridge, UK

Abstract

Some highlights of the history of modern cosmology and the lessons to be learned from the successes and blind alleys of the past are described. This heritage forms the background to the lectures and discussions at this Second Carnegie Centennial Symposium, which celebrates the remarkable contributions of the Carnegie Institution in the support of astronomical and cosmological research.

1.1 Introduction

It is a great honor to be invited to give this introductory address at the Second Carnegie Centennial Symposium to celebrate the outstanding achievements of the Observatories of the Carnegie Institution of Washington. I assume that the point of opening this meeting with a survey of the history of cosmology is not only to celebrate the remarkable achievements of modern observational and theoretical cosmology, but also to provide lessons for our time, which may enable us all to avoid some of the errors that we now recognize were made in the past. I am bound to say that I am not at all optimistic that this second aim will be achieved. I recall that, when I gave a similar talk many years ago with the same intention, Giancarlo Setti made the percipient remark:

Cosmology is like love; everyone likes to make their own mistakes.

By its very nature, the subject involves the confrontation of theoretical speculation with cosmological observations, the scepticism of the hardened observer about taking anything a theorist says seriously, the problems of pushing observations to the very limits of technological capability, and sometimes beyond these, resulting in dubious data, and so on. These confrontations have happened many times in the past. My suspicion is that the community of astronomers and cosmologists is now sufficiently large for false dogma and insecure observations to have only limited shelf-lives, but we must remain vigilant. Nonetheless, it is intriguing to survey the present state of cosmology, with its extraordinary successes and challenges, and recognize the many similarities to those that faced the great scientists of the past. I leave it to readers to draw their own preferred analogies.

The history of cosmology is a vast and fascinating subject, and I will only touch on some of the highlights of that story. I have given a more detailed account of that history elsewhere (Longair 1995), and it is a subject that repays careful study. To my regret, there will be little space to do justice to the technological achievements that have made modern cosmology a

2 *M. S. Longair*

rather exact science (see Longair 2001). Without these developments, none of us would be celebrating the achievements of modern cosmology at this symposium.

Before getting down to the history, let me contribute a personal appreciation of Andrew Carnegie's philanthropy. He gave away \$350M of his fortune of \$400M to charitable causes. Among the more remarkable of these was the founding of about 3000 libraries worldwide, including five in my home town of Dundee in Scotland, only about 30 miles from Dunfermline, Carnegie's birthplace. Here is a quotation from the Carnegie Libraries of Scotland web-site:

When the library was officially opened on 22 October 1908, [Charles] Barrie [a former Lord Provost] was asked to perform the opening ceremony ... There was a banquet afterwards in the Victoria Art Galleries hosted by Lord Provost Longair.

Lord Provost Longair was my great, great uncle. I remember as a small boy going regularly to the Dundee Public Libraries to learn about rockets and space flight. Little did I realise then that, more than 50 years later, I would be participating in the celebrations of the centenary of the founding of the Carnegie Institution.

1.2 **Observational Cosmology to 1929**

The earliest cosmologies of the modern era were speculative cosmologies. The "island universe" model of Descartes, published in *The World* of 1636, involved an interlocking jigsaw puzzle of solar systems. Wright's *An Original Theory of the Universe* of 1750 involved spheres of stars and solar systems, while Kant in 1755 and Lambert in 1761 developed the first hierarchical, or fractal, pictures of the Universe (see Harrison 2001). The problem with these early cosmologies was that they lacked observational validation. When these ideas were put forward, the only star whose distance was known was the Sun. The first parallax measurements of stars were only made in the 1830s by Friedrich Bessel, Friedrich Georg Wilhelm Struve and Thomas Henderson.

The first quantitative estimates of the scale and structure of the Universe were made by William Herschel in the late 18th century. Herschel's model of the large-scale structure of the Universe was based upon star counts and provided the first quantitative evidence for the "island universe" picture of Wright, Kant, Swedenborg and Laplace. In deriving his famous model for our Galaxy, Herschel assumed that all stars have the same absolute luminosities. The importance of interstellar extinction in restricting the number counts of stars to a relatively local region of our Galaxy was only fully appreciated in the early 20th century.

John Michell had already warned Herschel that the assumption that the stars have a fixed luminosity was incorrect. This is the same John Michell who was Woodwardian Professor of Geology at Queen's College, Cambridge, before becoming the rector of Thornhill in Yorkshire in 1767. He designed and built what we now know as the Cavendish experiment to measure the mean density of the Earth. Nowadays, he is rightly remembered as the inventor of black holes. In 1767, he showed that there must be a dispersion in the absolute luminosities of the stars from observations of bright star clusters. Despite this warning, Herschel ignored the problem and proceeded to produce a number of different versions of the structure of our Galaxy. In 1802, Herschel measured the magnitudes of visual binary stars and was forced to agree with Michell's conclusion. Equally troubling was the fact that observations with his magnificent 40-foot telescope showed there was no edge to the

A brief history of cosmology

3

Galaxy. He continued to find stars the fainter he looked—evidently, the stellar system was unbounded. Eventually, Herschel lost faith in his model of the Galaxy.

The desire to observe the Universe with telescopes of greater and greater aperture continued throughout the nineteenth century. The largest reflecting telescope constructed during that century was the great 72-inch reflector at Birr Castle in Ireland by William Parsons, the 3rd Earl of Rosse. This “Great Leviathan” was moved by ropes and astronomical objects could be tracked by moving the barrel of the telescope between the two large walls, which also accommodated a movable observing platform at the Newtonian focus of the telescope. Observations were made by eye and so the “length of the exposure” was limited to about a tenth of a second. Despite the difficulties of making observations and the inclement weather in central Ireland, Lord Rosse was able to resolve nebulae into stars and, perhaps most important of all, discovered the spiral structure of galaxies, the most famous drawing being his sketch of M51.

The revolutions that led to the discipline of extragalactic astronomy as we know it today were the use of photography to record astronomical images and the shift from refracting to reflecting telescope designs. The Yerkes 40-inch refractor was the end of the line so far as refracting telescopes were concerned. The much more compact reflecting design had the advantage of greater collecting area, but was much more sensitive to tracking and guiding errors. Many key technologies were developed during the latter half of the nineteenth century, thanks to pioneers such as Lewis Morris Rutherfurd, John Draper, Andrew Common and George Carver. These pioneers solved the problems of the tracking and pointing of reflecting telescopes, an invention of particular importance being the adjustable plate holder, which enables the observer to maintain the pointing of the telescope with high precision.

The resulting technical advances contributed to the remarkable achievement of James Keeler and his colleagues at the Lick Observatory in enhancing the performance of the 91-cm Crossley reflector to become the premier instrument for astronomical imaging. During the commissioning of the Crossley reflector in 1900, Keeler obtained spectacular images of spiral nebulae, including his famous image of M51. Not only were the details of its spiral structure observed in unprecedented detail, but there were also large numbers of fainter spiral nebulae of smaller angular size. If these were objects similar to the Andromeda Nebula M31, they must lie at very great distances from our Solar System. Tragically, just as this new era of astronomy was dawning, Keeler died of a stroke later that same year at the early age of only 42 (Osterbrock 1984).

George Ellery Hale plays a central role in the celebrations of the centenary of the Carnegie Institution. He is rightly regarded as the most successful astronomical entrepreneur of the modern era. He maintained an unswerving determination to construct successively larger and larger telescopes from the time of his directorship of the Yerkes Observatory in the 1890s through the period when he became Director of the Mount Wilson Observatory in 1903 until his death in 1938. In 1895, he had persuaded his father to buy the 1.5 meter blank for a 60-inch reflecting telescope. The design was to be an enlarged version of the Calver-Common design for the 91-cm Crossley reflector at the Lick Observatory. Before the 60-inch telescope was completed, however, he persuaded J. D. Hooker to fund an even bigger telescope, the 100-inch telescope to be built on Mount Wilson. The technological challenges were proportionally greater, the mass of the telescope being 100 tons, but the basic Calver-Common design was retained. The optics were the responsibility of George Ritchey, an optical designer of genius, who was to come up with the ingenious optical configuration

4 *M. S. Longair*

known as the Ritchey-Chrétien design, which enabled excellent imaging to be achieved over a wide field of view. The 60- and 100-inch telescopes became the prime telescopes for the study of the spiral nebulae, but these accomplishments were not achieved without an enormous effort on Hale's part.

The story of Hale's construction of these telescopes is a heroic tale. Equally impressive is Andrew Carnegie's generosity in enabling Hale to realise his vision. Carnegie's fateful visit to the Mount Wilson Observatory in 1906 has been recorded in the volume *The Legacy of George Ellery Hale* (Wright, Warnow, & Weiner 1972). Carnegie was clearly impressed by what he saw during his visit. As recorded in the local Pasadena newspaper, he remarked:

"We do not know what may be discovered here," he said. Franklin had little idea what would be the result of flying his kite. But we do know that this will mean the increase of our knowledge in regard to this great system of which we are part.

Mr. Hale has discovered here 1600 worlds about one of the stars which were not known before. We have found helium in the Sun, and after finding it there, we find it in the Earth. It all goes to show that all things are of a common origin.

To anyone who has had the fortune to be responsible for the operation of a large observatory facility, these remarks have heartening resonances. Carnegie did not quite get the science right so far as the stars were concerned, but he got it absolutely right so far as helium is concerned. Helium was discovered astronomically long before it was identified in the laboratory and is but one of many examples of how astronomical observations can provide key insights into the behavior of matter under circumstances which are only later reproduced in the laboratory. Plainly, Hale had carried out a very successful campaign in enthusing Carnegie about the importance of progress in astronomy.

Following his visit, Carnegie pledged an additional \$10M to the endowment of the Carnegie Institution, specifically requesting that the benefaction be used to enable the work of the Observatory to proceed as rapidly as possible. This is the purist music to the ears of any Observatory Director, who knows that, while it is usually possible in the end to find the capital resources for ambitious projects, these cannot succeed without matching funds for operations in the long term. Carnegie's vision and understanding are models for benefactors of astronomy.

The construction of the 60- and 100-inch telescopes were stressful and Hale suffered a nervous breakdown in 1910. It is touching to read Carnegie's letter to Hale of 1911 (with the original spelling), urging him to take care of his health:

November 27, 1911

Mr dear Friend,-

Delited to read your long note this morning; not too long—every word tells, but pray show your good sense by keeping in check your passion for work, so that you may be spared to put the capstone upon your career, which should be one of the most remarkable ever lived.

Ever yours,

Andrew Carnegie.

Carnegie's benefaction was crucial for the completion of the 100-inch Hooker Telescope at Mount Wilson. The telescope was by far the largest in the world and incorporated all the lessons learned from the works of earlier telescope builders. Completed in 1918, this

A brief history of cosmology

5

instrument was to dominate observational cosmology for the next 30 years until the commissioning of the Palomar 200-inch telescope in 1948.

The Hooker 100-inch telescope played a central role in the resolution of what became known as the “Great Debate,” which concerned the related issues of the size of our own Galaxy and the nature of the white, or spiral, nebulae. This confrontation between Harlow Shapley and Heber D. Curtis is too well known to need much amplification here (see Christianson, this volume). In 1899, Scheiner had obtained a spectrogram of M31 and stated that the spectrum suggested a “cluster of Sun-like stars.” In 1922, Öpik estimated the distance of M31 by comparing the mass-to-light ratio of the central region of M31 with that of our own Galaxy and found a distance of 440 kpc, suggesting that it lay well outside the confines of our Galaxy. The discovery of variable stars in spiral nebulae by Duncan in 1922 led to a flurry of activity and Hubble’s famous discovery of Cepheid variables in M31.

Central to Hubble’s use of Cepheid variable stars in M31 to measure its distance was the discovery of the period-luminosity relation for Cepheids in the Magellanic Clouds by Henrietta Leavitt (Leavitt 1912). Leavitt, like Annie Cannon, was profoundly deaf. While she is best remembered for her work on the Cepheid variables, her main work was the establishment of the North Polar Sequence, the accurate determination of the magnitude scale for stars in a region of sky which would always be accessible to observers in the Northern Hemisphere. By the time of her death in 1921, she had extended the North Polar Sequence from 2.7 to 21st magnitude, with errors less than 0.1 magnitudes. To achieve this, she used observations from 13 telescopes ranging from 0.5 to 60 inches in diameter and compared her scale using 5 different photographic photometric techniques. Without this fundamental work, the magnitude scale for galaxies could not have been established.

It is intriguing that by far the most stubborn pieces of observational evidence against what might be termed the long distance scale were van Maanen’s measurements of the proper motions of spiral arms. It is now well understood how difficult it is to measure tiny displacements of any diffuse object—van Maanen’s evidence was only refuted in 1933 by Edwin Hubble after a considerable observational effort.

Hubble’s paper of 1925 establishing the extragalactic nature of the spiral nebulae is impressive enough (Hubble 1925), but to my mind his paper of the following year entitled *Extragalactic Nebulae* is even more compelling (Hubble 1926). In this paper, he provided the first more or less complete description of galaxies as extragalactic systems. The paper includes a morphological classification of galaxies into the classic Hubble types, estimates of the relative numbers of different types, estimates of mass-to-luminosity ratios for different types of galaxies and their average number densities. Finally, the mean mass density in galaxies in the Universe as a whole was derived for the first time. Adopting Einstein’s static model for the Universe, the radius of curvature of the spherical geometry was $\mathcal{R} = 27,000$ Mpc and the total number of galaxies 3.5×10^{15} . Thus, by 1926, the first application of the ideas of relativistic cosmology to the Universe of galaxies had been made. Hubble concluded that the observations already extended to about 1/300 of the radius of the closed Einstein universe. The prophetic last sentence of his great paper reads

... with reasonable increases in the speed of the plates and size of telescopes, it may become possible to observe an appreciable fraction of the Einstein Universe.

It is no surprise that Hale began his campaign to raise funds for the 200-inch telescope in

6 *M. S. Longair*

1928. By the end of the year, he had received the promise of a grant of \$6M from the Rockefeller foundation.

Hubble's important insights were soon followed by an even more remarkable discovery. In 1917, Vesto M. Slipher had published his heroic pioneering spectroscopic observations of 25 spiral nebulae (Slipher 1917). He realised that, for the spectroscopy of low surface brightness objects such as the spiral nebulae, the crucial factor was the f -ratio of the spectrograph camera, not the size of the telescope. The observations involved very long integrations of 20, 40 or even 80 hours with small telescopes.

The velocities of the galaxies inferred from the Doppler shifts of their absorption lines were typically about 570 km s^{-1} , far in excess of the velocity of any known object in our Galaxy. Furthermore, most of the velocities corresponded to the galaxies moving away from the Solar System, that is, the lines were *redshifted* to longer (red) wavelengths. In 1921, Carl Wilhelm Wirtz concluded that, when the data were averaged in a suitable way, "an approximate linear dependence of velocity upon apparent magnitude is visible" (Wirtz 1921). By 1929, Hubble had assembled approximate distances of 24 galaxies for which velocities had been measured, mostly by Slipher, all within 2 Mpc of our Galaxy. I have always been impressed that Hubble was able to find the law which bears his name from the very crude distance indicators which he had available. The first seven objects within 500 kpc had Cepheid distances; the distances of the next 13 were found assuming the brightest stars all had the same absolute magnitude; the last four, in the Virgo cluster, were estimated on the basis of the mean luminosities of nebulae in the cluster. From these meager data, Hubble derived his famous redshift-distance relation (Hubble 1929). If the redshifts z are interpreted as the Doppler shifts of galaxies due to their recession velocities v , the relation can be written $v = H_0 r$, where H_0 is *Hubble's constant*.

Milton Humason had by then mastered the use of the 100-inch telescope for obtaining the spectra of faint galaxies and by 1934 Humason and Hubble had extended the velocity-distance relation to 7% of the speed of light (Humason & Hubble 1934). Furthermore, Hubble realised that he could test for the isotropy and homogeneity of the Universe by counting the numbers of faint galaxies. Hubble established that the numbers of galaxies increased with increasing apparent magnitude in almost exactly the fashion expected if they were uniformly distributed in space.

Even before 1929, however, it was appreciated that Hubble's law was expected according to world models based upon the general theory of relativity.

1.3 Theoretical Cosmology to 1939

Let us turn to theoretical cosmology and the history of Einstein's static model for the Universe. Working independently, Lobachevsky in Kazan in Russia and Bolyai in Hungary solved the problem of the existence of geometries that violated Euclid's fifth axiom in 1825. These were the first self-consistent hyperbolic (non-Euclidean) geometries. In his great text *On the Principles of Geometry* (1825), Lobachevsky worked out the minimum parallax of any star in hyperbolic geometry

$$\theta = \arctan\left(\frac{a}{\mathcal{R}}\right) \quad (1.1)$$

where a is the radius of the Earth's orbit and \mathcal{R} the radius of curvature of the geometry. In his textbook, he found a minimum value of $\mathcal{R} \geq 1.66 \times 10^5 \text{ AU}$. What is intriguing is that this

A brief history of cosmology

7

estimate was made 8 years before Bessel's announcement of the first successful parallax measurement of 61 Cygni. In making his estimate, Lobachevsky used the observational upper limit to the parallax of bright stars. In a statement which will warm the heart of observational astronomers, and which is particularly apposite in the light of what we will hear at this meeting, he remarked

There is no means other than astronomical observations for judging the exactness which attaches to the calculations of ordinary geometry.

The pioneering works of Lobachevsky and Bolyai led to Riemann's introduction of quadratic differential forms, his generalization of their results to non-Euclidean geometries, and his discovery of spaces of positive curvature—that is, spherical non-Euclidean geometries.

Unlike his other great discoveries, Einstein's route to general relativity was long and tortuous. Four ideas were important in his search for a self-consistent relativistic theory of gravity:

- The influence of gravity on light
- The principle of equivalence
- Riemannian spacetime
- The principle of equivalence

Toward the end of 1912, he realised that what was needed was non-Euclidean geometry. Einstein consulted his old school friend, Marcel Grossmann, about the most general forms of transformation between frames of reference for metrics of the form

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu. \quad (1.2)$$

Grossmann soon came back with the answer that the most general transformation formulae were the Riemannian geometries, but that they had the “bad feature” that they are nonlinear. Einstein instantly recognized that, on the contrary, this was a great advantage since any satisfactory theory of relativistic gravity must be nonlinear.

After further years of struggle, during which he and Grossmann were ploughing very much a lone furrow, general relativity was formulated in its definitive form in 1915 (Einstein 1915). In 1916, Willem de Sitter and Paul Ehrenfest suggested in correspondence that a spherical 4-dimensional spacetime would eliminate the problems of the boundary conditions at infinity, which pose insuperable problems for Newtonian cosmological models. In 1917, Einstein realised that, in general relativity, he had for the first time a theory which could be used to construct fully self-consistent models for the Universe as a whole (Einstein 1917). At that time, the expansion of the Universe had not been discovered.

One of objectives of Einstein's program was to incorporate into the structure of general relativity what he called *Mach's Principle*, meaning that the local inertial frame of reference should be determined by the large-scale distribution of matter in the Universe. There was, however, a further problem, first noted by Newton, that static model universes are unstable under gravity. Einstein proposed to solve both problems by introducing an additional term into the field equations, the *cosmological constant* Λ . In Newtonian terms, the cosmological constant corresponds to a repulsive force \vec{f} acting on a test particle at distance \vec{r} , $\vec{f} = \frac{1}{3}\Lambda\vec{r}$. Unlike gravity, this force is independent of the density of matter. The Λ -term has negligible influence on the scale of the Solar System and is only appreciable on cosmological scales.

The equation that describes the expansion becomes

8 *M. S. Longair*

$$\frac{d^2R}{dt^2} = -\frac{4\pi G\rho_0}{3R^2} + \frac{1}{3}\Lambda R \quad (1.3)$$

The first term on the right-hand side describes the deceleration due to gravity and the second what Zel'dovich referred to as the “repulsive effect of the vacuum” (Zel'dovich 1968). At that time, the physical significance of the Λ term was not understood.

Einstein believed that he had incorporated Mach's Principle into general relativity. In his words,

The inertial structure of spacetime was to be “exhaustively conditioned and determined” by the distribution of material throughout the Universe.

Further, he stated that the extension of the field equations was “not justified by our actual knowledge of gravitation,” but was “logically consistent.” Furthermore, the cosmological term was “necessary only for the purpose of making possible a quasi-static distribution of matter, as required by the fact of the small velocities of stars.”

From Einstein's field equations of general relativity, it followed that the geometry of Einstein's static universe is closed and the radius of curvature of the geometrical sections is $\mathcal{R} = c/(4\pi G\rho_0)^{1/2}$, where ρ_0 is the mean density of the static Universe. The value of Λ was directly related to the mean density of the Universe, $\Lambda = 4\pi G\rho_0$. Einstein believed that he had incorporated Mach's Principle into general relativity, in that static solutions of the field equations did not exist in the absence of matter.

Almost immediately, de Sitter (1917) showed that one of Einstein's objectives had not been achieved. He found solutions of Einstein's field equations in the absence of matter, $\rho = p = 0$. The metric he derived had the form

$$ds^2 = dr^2 - R^2 \sin^2\left(\frac{r}{R}\right) (d\phi^2 + \cos^2\phi d\theta^2) + \cos^2\left(\frac{r}{R}\right) c^2 dt^2 \quad (1.4)$$

Although there is no matter present in the Universe, a test particle still moves along a perfectly well-defined path through spacetime. As de Sitter remarked, “If no matter exists apart from the test body, has this inertia?” One prediction of de Sitter's paper was the fact that distant galaxies would be observed with a redshift, although in his solution the metric was stationary—this phenomenon became known as the *de Sitter effect*.

In 1922, Kornel Lanczos showed that by a simple change of coordinates, the de Sitter solution could be interpreted as an expansion of the system of coordinates in hyperbolic space (Lanczos 1922).

$$ds^2 = -dt^2 + \cosh^2 t [d\phi^2 + \cos^2\phi (d\psi^2 + \cos^2\psi d\chi^2)] \quad (1.5)$$

Lanczos wrote that:

It is interesting to observe how one and the same geometry can appear with quite different physical interpretations according to the interpretations placed upon the particular coordinates.

At almost exactly the same time, the Soviet meteorologist and theoretical physicist Alexander Alexandrovich Friedman published the first of his two classic papers on relativistic cosmology (Friedman 1922, 1924). His key realization was that isotropic world models had to have isotropic curvature everywhere. In the paper of 1922, Friedman found solutions for expanding world models with closed spatial geometries, including those that expand to a maximum radius and then collapse to a singularity. In the paper of 1924, he showed that

A brief history of cosmology

9

there exist expanding solutions that are unbounded with hyperbolic geometry. The differential equations that he derived were:

$$(1922) \quad \left(\frac{\dot{R}}{R}\right)^2 + \left(\frac{2R\ddot{R}}{R^2}\right) + \frac{c^2}{R^2} - \Lambda = 0 \quad (1.6)$$

$$(1924) \quad \left(\frac{\dot{R}}{R}\right)^2 + \left(\frac{2R\ddot{R}}{R^2}\right) - \frac{c^2}{R^2} - \Lambda = 0 \quad (1.7)$$

In both cases,

$$\frac{3\dot{R}^2}{R^2} + \frac{3c^2}{R^2} - \Lambda = \kappa c^2 \rho \quad (1.8)$$

The solutions of these equations correspond exactly to the standard world models of general relativity and are appropriately known as the *Friedman world models*. The history of general relativity in the Soviet Union is a remarkable story and Friedman's role in introducing Soviet scientists to the theory and the subsequent difficult development of these studies in the USSR needs to be better known. It has been carefully described by Zelmanov (1967) in a review that has not been translated into English.

It has always been considered somewhat surprising that it was some years before Friedman's important papers were given the recognition they deserve. In 1923, Einstein believed he had found an error in the first of Friedman's papers and published his concern in *Zeitschrift für Physik*. Friedman showed that Einstein was incorrect and Einstein subsequently published his withdrawal of his objection in the same journal. My guess is that Einstein's concern was remembered, but not his acknowledgment of his error.

In 1927, Georges Lemaître independently discovered the Friedman solutions and only then became aware of Friedman's pioneering contributions (Lemaître 1927). Both Lemaître and Howard P. Robertson (1928) were aware of the fact that the Friedman solutions result locally in a velocity-distance relation. Lemaître derived what he termed the "apparent Doppler effect," in which "the receding velocities of extragalactic nebulae are a cosmical effect of the expansion of the Universe" with $v \propto r$. Robertson found a similar result stating that "we should expect . . . a correlation $v \approx (cl/R)$," where l is distance and v the recession velocity. From nearby galaxies, he found a value for Hubble's constant of $500 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

The discovery of the velocity-distance relation for galaxies was interpreted as evidence for the expansion of the Universe as a whole. There remained problems of interpretation of the notions of time and distance in cosmology because the field equations could be set up in any frame of reference. By 1935, the problem had been solved independently by Robertson and George Walker (Robertson 1935; Walker 1935). For isotropic, homogeneous world models, they showed that the metric of spacetime had to have the form

$$ds^2 = dt^2 - \frac{R^2(t)}{c^2} \left[\frac{dr^2}{(1 + \kappa r^2)} + r^2(d\theta^2 + \sin^2 \theta d\phi^2) \right] \quad (1.9)$$

where κ is the curvature of space at the present epoch, r is a comoving radial distance coordinate and $R(t)$ is the scale factor which describes how the distance between any two world lines change with cosmic time t . The *Robertson-Walker metric* contains all the geometries consistent with the assumptions of isotropy and homogeneity of the Universe; the curvature $\kappa = \mathcal{R}^{-2}$, where \mathcal{R} , the radius of curvature of the spatial sections of the isotropic curved

10 *M. S. Longair*

space, can be positive, negative or zero. The physics of the expansion is absorbed into the scale factor $R(t)$.

With the discovery of the velocity-distance relation, Einstein regretted the inclusion of the cosmological constant into the field equations. According to George Gamow, Einstein stated that the introduction of the cosmological constant was “the biggest blunder of my life” (Gamow 1970). In 1932, Einstein and de Sitter showed that there is one special solution of the equations with $\Lambda = 0$ and $\kappa = 0$, corresponding to Euclidean space sections (Einstein & de Sitter 1932). This *Einstein-de Sitter model* has density at the present epoch $\rho_0 = 3H_0^2/8\pi G$. This density is often referred to as the *critical density* and the Einstein-de Sitter model as the *critical model*, because it separates the ever-expanding models with open, hyperbolic geometries from those that will eventually collapse to a singularity and that have closed, spherical geometry. When Einstein and de Sitter inserted $H_0 = 500 \text{ km s}^{-1} \text{ Mpc}^{-1}$ into the expression for ρ_0 , they found $\rho_0 = 4 \times 10^{-25} \text{ kg m}^{-3}$. Although this value was somewhat greater than the mean density in galaxies derived by Hubble, they argued that it was of the correct order of magnitude and that there might well be a considerable amounts of “dark matter” present in the Universe.

1.4 Astrophysical Cosmology up to 1939

1.4.1 *Dark Matter*

Astrophysical evidence for dark matter was not long in coming. In 1933, Fritz Zwicky made the first dynamical studies of rich clusters of galaxies, in particular, of the Coma cluster (Zwicky 1933, 1937). The method Zwicky used to estimate the total mass of the cluster involved the *virial theorem*, which had been derived by Arthur Eddington in 1916 to estimate the masses of star clusters (Eddington 1916). The theorem relates the total internal kinetic energy $T = \frac{1}{2}M\langle v^2 \rangle$ of the galaxies in a cluster to its gravitational potential energy, $|U| = GM^2/2R_{cl}$ in statistical equilibrium under gravity. Eddington showed that $T = \frac{1}{2}|U|$ and so the mass of the cluster can be found, $M \approx 2R_{cl}\langle v^2 \rangle/G$.

Zwicky measured the velocity dispersion of the galaxies in the Coma cluster and found that there was much more mass in the cluster than could be attributed to the visible parts of galaxies. In solar units, the ratio of mass to optical luminosity of a galaxy such as our own is about 3, whereas for the Coma cluster the ratio was found to be about 500—there must be about 100 times more dark or hidden matter as compared with visible matter in the cluster. Zwicky’s pioneering studies have been confirmed by all subsequent studies of rich clusters of galaxies.

1.4.2 *The Age of the Universe and Eddington-Lemaître Models*

Despite Einstein’s renunciation of the cosmological constant Λ , there remained a very grave problem for those models in which Λ is set equal to zero. In all world models with $\Lambda = 0$, the age of the Universe is less than H_0^{-1} . Using Hubble’s estimate of $H_0 = 500 \text{ km s}^{-1} \text{ Mpc}^{-1}$, the age of the Universe must be less than 2×10^9 years old, a figure in conflict with the age of the Earth derived from studies of the ratios of abundances of long-lived radioactive species, which gave ages significantly greater than this value.

Eddington and Lemaître recognized that this problem could be eliminated if Λ were positive (Eddington 1930; Lemaître 1931a). The effect of a positive cosmological constant is to counteract the attractive force of gravity when the Universe has grown to a large enough