

Index

Page numbers in *italics* indicate tables

- abscess, tretment-related, 310
- access sites, endovascular, 121
- acid-base balance, hypothermia and, 141–142
- activities of daily living
 - Barthel ADL Index, 74
 - Modified Rankin Scale, 73
- adenosine cardiac standstill, 444
- airway
 - compromised, intubation for, 288
 - difficult, 289
- Alberta stroke program early CT score (ASPECTS), 104
- anemia
 - cerebral blood flow affected by, 21
 - after subarachnoid hemorrhage, 471–472, 481
- anesthesia for intracranial angioplasty, 121
- aneurysm
 - clipping vs. coiling, 437, 459–461
 - craniotomy for
 - aneurysm rupture during, 443–444
 - indications, 437–438
 - microscopic approaches, 440–442
 - minimally invasive, 440
 - proximal control, 442–443
 - repair techniques, 443–444
 - surgical techniques, 439–440, 445
 - timing and preparation, 438–439
 - development, 424
- embolization
 - coil, 111–112, 437–438, 448–450
 - development of, 447
- endovascular therapy
 - clinical trials, 459–461
 - complications, 456–457
 - periprocedural care, 455–456
 - site of aneurysm and, 453–455
 - techniques and devices, 447–453

- incomplete treatment, 458–459
- interventional treatment
 - indications for, 111
 - options and procedures, 111–112
- location, 423
- rupture, 423–424
 - during endovascular therapy, 456–457
 - and hydrocephalus, 146, 159
 - intraoperative, 439, 443–444
- surgery for
 - decompressive, 96
 - Hunt and Hess scale of risk, 72
 - lumbar drainage in, 160
 - monitoring in, 40
- angiography
 - of aneurysm, 447–448
 - for craniotomy, 438–439
 - postoperative, 458
 - in arterial malformations, 114
 - in cavernoma, 406
 - in cerebellar hemorrhage, 366–367
 - in cerebellar infarction, 212
 - in cerebral hemorrhage, 392
 - in cerebral venous thrombosis, 502–505
 - in cervical artery dissections, 227
 - four-vessel as baseline, 122
 - in intracerebral hemorrhage, 338
 - in intracranial stenosis, 122
 - pre- and intraoperative, in excision of AVMs, 389
 - in subarachnoid hemorrhage, 427–428
 - in vasospasm, 109
- angiomas
 - classification, 114
 - endovascular therapy, 114
 - venous, 401
- angioplasty
 - after thrombolysis in basilar artery occlusion, 201
 - balloon, 109
 - balloon selection, 123
 - complications, 125
 - postprocedural care, 125
 - prophylactic, 432
 - vs. stent-assisted, 122–123
 - techniques, 123–125
 - in vasospasm, 457, 472
 - for spasmolysis, 109–111
 - prophylactic, 467
 - in intracranial stenosis, 120
- anoxic depolarization, 130
- antibiotic-impregnated catheters, 151
- antibiotics
 - prophylactic
 - in cerebrospinal fluid leak, 531
 - in ventilator-associated pneumonia, 307
 - for ventricular drainage, 150–151, 357
 - selective decontamination by, 433
 - strategies for use of, 311, 311
- anticoagulants
 - after basilar artery occlusion, 202
 - after cerebral venous thrombosis, 509–510
 - in cerebral venous thrombosis, 234–235
 - in cervical artery dissections, 227
 - endovascular therapy and, 121
 - in endovascular therapy, 452, 455
 - and hematoma expansion, 265–268
 - novel direct oral, 267–268
 - in pregnancy, 236
 - reversal, 265–266, 266, 268
- antiedema therapy, 82
 - barbiturates, 84–85
 - elevated head position, 85–86
 - hyperventilation, 85, 316
 - in intracranial hemorrhage, 86–87
 - osmotherapy, 82–84, 316–317
 - steroids as, 86
 - tromethamine, 85
- antihypertensives, 248, 249–251, 279
 - in acute hypertensive response, 278–280
 - in intracerebral hemorrhage, 276–277
 - hematoma expansion and, 262–263, 264, 268
 - immediately post-stroke, 280–282
 - stop vs. continue, 245–246
- anti-inflammatory treatment
 - in posthemorrhagic edema, 86
 - prophylactic for delayed ischemia, 467
- antiplatelet medication
 - after endovascular therapy, 125
 - in endovascular therapy, 121, 455
 - and hematoma expansion, 265
 - in stenosis, 106, 107
 - after stenting, 107, 109
 - in stenting, 107
- APACHE Scores, 68–69
- arterial malformations, endovascular therapy for, 112–114
 - embolization techniques, 114–117
- arteriovenous malformations, 394
 - radiosurgery for, 394–396
 - surgical excision, 387–389

- arteriovenous malformations (cont.)
 - decision-making, 390–392
 - with hemorrhage, 392–393
 - unexpected, in craniotomy, 324, 392–393
 - arteritis
 - giant cell, 229
 - Takayasu, 229–230
 - ataxia
 - in cerebellar hemorrhage, 365
 - in cerebellar infarction, 209
 - autoregulation, cerebral, 4–5, 20–22, 243
 - head injury and, 530
 - in hypertension, 274
 - in intracerebral hemorrhage, 275
 - bacterial infection as cause of stroke, 228–229
 - balloon
 - angioplasty, 109
 - balloon selection, 123
 - complications, 125
 - postprocedural care, 125
 - prophylactic, 432
 - vs. stent-assisted, 122–123
 - techniques, 123–125
 - in vasospasm, 457, 472
 - embolization, 447
 - occlusion of parent artery in aneurysm, 448
- barbiturates, 84–85
- Barthel ADL Index, 74
- basilar artery occlusion, 194–195
 - brainstem hemorrhage in, 378
 - clinical trials of thrombolysis, 198–200
 - diagnostics and imaging, 195–196
 - differential diagnoses, 196
 - intensive care after, 202–203
 - symptoms, 195
- Bedside Shivering Assessment Scale, 137, 137
- Behcet’s disease, 230
- blood pressure, systemic
 - management
 - guidelines, 247, 277–278, 282
 - in cerebellar hemorrhage, 370
 - in intracerebral hemorrhage, 262–263, 264, 276–277, 280–282, 338
 - in subarachnoid hemorrhage, 430
 - stroke and, 243–245
 - target levels, 246
 - See also hypertension, hypotension- blood–brain barrier
 - in edema formation, 81, 82
 - hypothermia and, 130
 - metalloproteinases and, 130
 - osmotherapy and, 82
- brain damage
 - ischemic, 518–519
 - primary, 519–520
 - prevention, 526
 - secondary, 521
 - ischemic, 520–521
- brain symmetry index, 61
- brain tissue oxygenation
 - cerebral perfusion pressure and, 39
 - and hyperventilation, 39
 - monitoring, 37–39
 - in intracranial hemorrhage, 41
 - in ischemic stroke, 40
 - by near-infrared spectroscopy, 63
 - neurochemical monitoring and, 39
 - in subarachnoid hemorrhage, 40, 468, 485
 - thresholds, 38–39
- brainstem auditory evoked potentials, 212
 - for monitoring, 61
- brainstem cavernoma, 382–383
- brainstem compression, 367, 371
- brainstem hematoma, 378
 - surgery in, 381–383
- brainstem hemorrhage, 378–379
 - prognosis, 379–380
 - respiratory effects, 288
 - surgery in, 381
- CADASIL, 232
- calcium influx, hypothermia and, 130
- calcium-channel blockers
 - in delayed ischemia, 465, 487
 - in vasospasm, 109, 110, 433, 465
- Call–Fleming syndrome, 231–232
- Canadian Neurological Scale, 71
- capillary telangiectases, 400–401
- carbon dioxide, cerebral blood flow modulated by, 21, 287
- cardiac assist devices, 495
- cardiac investigations after basilar artery occlusion, 203
- cardiac rhythm
 - hypothermia and, 140, 141
 - subarachnoid hemorrhage and, 492
 - vasopressors and, 473
- cardiopulmonary dysfunction, CNS-mediated, 490
- cardiomyopathies, 492–493, 495

- clinical features and diagnosis, 492–494
- pathophysiology, 490–492
- catecholamine surge, 491, 495
- catheters
 - balloon, for intracranial angioplasty, 123
 - brain tissue oxygenation, 38
 - Camino, 8
 - cerebral microdialysis, 44, 46
 - cooling, 132
 - for intracranial angioplasty, 122
 - and hemorrhage risk, 8
 - and infection risk, 8, 308
- intraventricular, 8
 - antibiotic-impregnated, 151, 357
 - placement, 148–149, 355
- lumbar
 - complications, 163
 - placement, 159
 - in pressure monitoring, 162
- micro-, for intracranial angioplasty, 124
- urinary, and infection, 307–308
- cavernoma, 400–402
 - clinical presentation, 408, 409–410
 - diagnostic imaging, 406–408
 - epidemiology, 402–404
 - histopathology, pathobiology and pathogenesis, 404–406
 - management, 410–413
 - surgical, 382–383, 413–415
 - related intracranial hemorrhage, 408–410
- central volume principle, 27
- cerebellar hemorrhage, 363–364
 - complications, 367
 - diagnosis, 365–367
 - prognosis and outcome, 372–374, 373
 - signs and symptoms, 364–365, 364
 - treatment, 367–370
 - surgical, 370–371
 - ventricular drainage, 371
- cerebellar infarction
 - combination therapy in, 183
 - course and outcome, 213–214
 - decompressive surgery in, 95, 182–183
 - diagnosis, 210–213
 - epidemiology and risk factors, 207
 - mortality, 219
 - pathogenesis, 207–208
 - prognosis and outcome, 218–219, 220
 - signs and symptoms, 208–210, 215
 - treatment strategies, 214–218, 216, 219–223
- cerebral autoregulation, 4–5, 20–22, 243
 - head injury and, 530
 - in hypertension, 274
 - in intracerebral hemorrhage, 275
- cerebral blood flow, 20
 - in acute ischemic stroke, 25
 - hyperventilation and, 85
 - in intracerebral hemorrhage, 275–276
 - in intracranial hemorrhage, 25
 - as marker of ischemia, 22–24
 - monitoring, 26–33, 27
 - comparison of methods, 29
 - indications for, 33
 - in subarachnoid hemorrhage, 25–26
 - in subarachnoid hemorrhage, 468
 - thresholds for ischemia, 22
- cerebral cavernous malformation. *See* cavernoma
- cerebral infarction, delayed, 429
- cerebral ischemia, 22
 - cerebral blood flow as marker of, 22–24
 - delayed, 464–465
 - adjunctive therapies, 474
 - detection, 468–469
 - hemodynamic goals, 471
 - hemodynamic interventions, 469–470, 472, 473–474
 - indications for treatment, 469
 - prevention, 464–465
 - microdialysis to detect, 44, 47, 48–50, 468
- cerebral perfusion pressure, 5, 20
 - brain tissue oxygenation and, 39
 - vs. intracranial pressure, 9
 - reduced, compensatory mechanisms, 22
- cerebral vasospasm. *See* vasospasm
- cerebral venous thrombosis, 233–234
 - causes, 506–507
 - clinical features, 502–505
 - diagnosis, 502–505
 - differential, 505–506
 - epidemiology, 501
 - intracranial pressure monitoring in, 12
 - malignant, decompressive surgery in, 96–97
 - outcome and prognosis, 507–508
 - recurrent, 509–510
 - risk factors, 506
 - treatment, 234–235, 508–512
- cerebrospinal fluid
 - altered dynamics after craniectomy, 98, 100
 - blockage of, and hydrocephalus, 146, 158
 - leaks, 530–531

- cerebrospinal fluid (cont.)
 - role in regulation of ICP, 3
 - shunt, 152
 - surveillance, 151
 - antibiotic-impregnated catheter and, 152
 - See also* lumbar drainage, ventricular drainage
- cerebrovascular malformations, 400–402
- cerebrovascular reserve, 21
- cervical artery dissection, 226–227
- clipping of aneurysms, 443–444
 - aneurysm rupture during, 443–444
 - vs. coiling, 437, 459–461
 - galvanic corrosion of clips, 445
- clot burden
 - and choice of treatment, 103
 - high, recanalization in, 106
- CMRO₂. *See* metabolic demand
- coagulopathy, 237–238
 - hematoma expansion and, 265–268
 - hypothermia and, 142
 - intracerebral hemorrhage and, 323, 338
 - and ventricular drainage, 153
- coil embolization
 - development of, 447
 - in aneurysm, 111–112, 437–438
 - vs. clipping, 437, 459–461
 - devices, 448–450
 - of fistulas, 116
 - stent-assisted, 450–453
- coma
 - airway compromise in, 288
 - in basilar artery occlusion, 200
 - causes, 287
 - in cerebellar hemorrhage, 365
 - in cerebral venous thrombosis, 501
 - Glasgow Coma Scale, 66–68
 - in cerebellar infarction, 210
 - as predictor of extubation, 293
 - microdialysis in, 48, 50–51
- combination therapy
 - decompressive surgery and hypothermia, 185
 - drugs and hypothermia, 135–136
 - hypothermia and drainage, 183
 - intraventricular fibrinolysis and drainage, 162
 - thrombolysis and interventional therapies, 175–177
 - thrombolysis and minimally invasive surgery, 315–318
- complement in edema formation, 86
- consciousness
 - brainstem hemorrhage and, 379
 - in cerebellar hemorrhage, 364, 367
 - in cerebellar infarction, 210
 - Glasgow Coma Scale, 66–68
 - in cerebellar infarction, 209
 - as predictor of extubation, 293
 - respiratory drive and, 287
- contra coup lesion, 524
- contusions, cerebral, 520, 523–524
 - treatment, 528–529
- cooling
 - invasive methods, 132–133, 186
 - non-invasive methods, 132, 185–186
 - optimal temperature, 186
 - rewarming, 134
- core temperature
 - monitoring in cooling, 132
 - regulation, 131
 - and shivering, 136
 - target, in hypothermia, 133
- craniectomy, decompressive, 10, 90–92
 - in cerebral contusions, 528–529
 - clinical trials, 530, 532–533
 - complications, 97–100
 - converting craniotomy to, 324
 - indications, 92–97
 - suboccipital, 217, 371
 - in traumatic brain injury, 14
- cranioplasty, 91, 100
- craniotomy
 - in aneurysm
 - aneurysm rupture during, 443–444
 - indications, 437–438
 - microscopic approaches, 440–442
 - minimally invasive, 440
 - proximal control, 442–443
 - repair techniques, 443–444
 - surgical techniques, 439–440, 445
 - timing and preparation, 438–439
 - for arteriovenous malformations, 387–389
 - decision-making, 390–392
 - with hemorrhage, 392–393
 - decompressive
 - in extradural hematoma, 527
 - in intracerebral hematoma, 528
 - in subdural hematoma, 527–528, 532
 - in intracerebral hemorrhage, 323–326
 - in prevention of hematoma expansion, 260–262
 - suboccipital, 371
- CSF. *See* cerebrospinal fluid

CT	
in basilar artery occlusion, 195–196	
in cavernoma, 407	
in cerebellar hemorrhage, 366	
in cerebellar infarction, 210–211	
in cerebral venous thrombosis, 502–503	
in cervical artery dissections, 227	
in head injury, 524	
Marshall classification, 69	
in intracerebral hemorrhage, 338–339	
of malignant MCA infarction, 182	
to measure cerebral blood flow, 28, 30	
in subarachnoid hemorrhage, 426–428	
in subdural hematoma, 523	
CT perfusion to measure cerebral blood flow, 27, 28–31, 30, 468–469	
decompressive surgery, 90–92, 180	
in cerebellar infarction, 182–183	
in cerebral contusions, 528–529	
in cerebral venous thrombosis, 509–510	
clinical trials, 530, 532–533	
complications, 97–100	
in extradural hematoma, 527	
indications, 92–97	
in intracerebral hematoma, 528	
clinical trials, 528	
for ischemic stroke, 179	
in malignant MCA infarction	
clinical trials, 180–181, 180	
complications, 182	
in subdural hematoma, 527–528	
suboccipital, 217, 371	
diaschisis, 23	
diffuse axonal injury, 519–520	
in hematoma, 522	
disability	
acceptability, 95	
Modified Rankin Scale, 73	
studies of, after decompressive surgery	
in cerebellar infarction, 95	
in hemispheric infarction, 92–95	
dislocation, third ventricle, 59	
dissection, cervical artery, 226–227, 521	
dizziness, diagnostic value of, 208	
drainage	
lumbar, 163–164	
complications, 162–163	
intraoperative, 439	
in intraventricular hemorrhage, 356–357	
in subarachnoid hemorrhage, 159–161	
in traumatic brain injury, 14	
ventricular, 8, 145–147	
in brainstem hemorrhage, 381	
in cerebellar hemorrhage, 371	
in cerebellar infarction, 183, 217	
complications, 149–150	
in intracranial hemorrhage, 11–12	
in intraventricular hemorrhage, 353	
insertion, 147–149	
management, 150–152, 357	
-related infections, 357	
removal, 152	
and risk of hemorrhage, 8	
risk of infection, 309	
thromboprophylaxis, 152–153	
drugs	
for control of shivering, 137	
hypothermia alters pharmacokinetics, 139–140, 140	
dural opening in decompressive craniectomy, 91, 528	
dysphagia, 299–300, 306	
ECG	
hypothermia and, 140	
in subarachnoid hemorrhage, 492, 494	
eclampsia, 236	
edema	
cytotoxic, 520	
ischemic, 90	
pathophysiology, 81	
in malignant MCA infarction, 190–191	
predictors of, 192	
treatment, 192–193	
perihemorrhagic	
management, 315–318	
pathophysiology, 81–82, 315	
post-trauma, 520–521	
pulmonary, 433	
space-occupying	
in cerebellar infarction, 207, 211, 213–214, 218–223	
diagnosis, 210–213	
evoked potentials in, 61	
predictors of, 192	
signs and symptoms, 208–210	
treatment, 192–193	
See also antiedema therapy	
EKOS ultrasound device, 175	
electroencephalography, continuous, for monitoring, 60–61	

- electrolytes
 - hypothermia and, 141–142
 - subarachnoid hemorrhage and, 429, 433, 482
- embolism, postoperative, 235
- embolization
 - balloon, 447
 - coil
 - in aneurysm, 111–112, 437–438
 - vs. clipping, 437, 459–461
 - development of, 447
 - devices, 448–450
 - of fistulas, 116
 - stent-assisted, 450–453
 - of arterial malformations, 112–114
 - preoperative, 388
 - before radiosurgery, 396
 - techniques, 114–117
- empyema, treatment-related, 310
- endoscopic techniques for clot evacuation, 331, 332, 337
 - in intraventricular hemorrhage, 356
- endothelin antagonists in subarachnoid hemorrhage, 434, 466
- endovascular therapy
 - complications, 125
 - in aneurysm
 - clinical trials, 459–461
 - complications, 456–457
 - periprocedural care, 455–456
 - site of aneurysm and, 453–455
 - techniques and devices, 447–453
 - in arterial malformations, 112–114
 - embolization techniques, 114–117
 - in basilar artery occlusion, 202
 - in cerebral venous thrombosis, 509
 - in intracranial stenosis
 - antiplatelet therapy before, 121
 - indications for, 120
 - vs. medical management, 120
 - postprocedural care, 125
 - procedural considerations, 121–122
 - techniques, 122–125
 - in vasospasm, 457–458, 472–473
 - See also* interventional therapy
- energy demand, 298
- enteral nutrition, 299–301
 - vs. parenteral, 301–302
- epilepsy
 - after cerebral venous thrombosis, 511
 - cavernoma and, 408, 410
 - surgery for, 413–414
 - See also* seizures
- European Stroke Scale, 73
- evoked potentials, for monitoring, 61
- external lumbar drainage, 163–164
 - complications, 162–163
 - intraoperative, 439
 - in intraventricular hemorrhage, 356–357
 - in subarachnoid hemorrhage, 159–161
 - in traumatic brain injury, 14
- external ventricular drainage, 8, 145–147
 - in brainstem hemorrhage, 381
 - in cerebellar hemorrhage, 371
 - in cerebellar infarction, 183, 217
 - complications, 149–150
 - insertion, 147–149
 - in intracranial hemorrhage, 11–12
 - management, 150–152, 357
 - related infections, 357
 - removal, 152
 - and risk of hemorrhage, 8
 - risk of infection, 309
 - thromboprophylaxis, 152–153
 - in ventricular hemorrhage, 353
- Fabry disease, 232–233
- feeding. *See* nutrition
- fever
 - and intracerebral hemorrhage, 263–264
 - after subarachnoid hemorrhage, 474, 481
- fibromuscular dysplasia, 230
- Fick principle, 20
 - in monitoring of cerebral blood flow, 26–27
- Fisher Grade Scale, 72
 - Modified, 427, 465
- fistulas
 - classification of Cognard, 116
 - embolization, 114, 116–117
- focal syndrome, 501
- FOUR Score, 68
- gait instability, diagnostic value of, 209
- Glasgow Coma Scale, 66–68
 - in cerebellar infarction, 210
 - as predictor of extubation, 293
- Glasgow Outcome Scale, 74–75
- glucose
 - microdialysis of, 50
 - monitoring after subarachnoid hemorrhage, 484

glutamate release	intracerebral, 523–524
hypothermia and, 130	treatment, 528
and ischemia, 485	subdural, 522–523
glycemic control	chronic, 522–523, 532
hypothermia and, 138	treatment, 527–528
and intracerebral hemorrhage, 264, 302–303	hemicraniectomy, decompressive, 90–92, 180
after subarachnoid hemorrhage, 433, 482	complications, 97–100
in traumatic brain injury, 51	in hemorrhagic stroke, 317
glycerol, 84	indications, 92–97
Graeb Score, 147, 351, 352	in malignant MCA infarction, 94
guidelines	clinical trials, 180–181
blood pressure management, 277–278, 277, 282	complications, 182
head injury, 517, 533–534	in subarachnoid hemorrhage, 96
insertion of ICP monitors, 13	hemispheric infarction
guidewires for intracranial angioplasty, 123–124	decompressive craniectomy in, 92–95
handicap, Modified Rankin Scale, 73	<i>See also</i> malignant MCA infarction
head elevation as antiedema therapy, 85–86	hemispheric syndrome, 81
head injury	hemodilution, 471
APACHE Scores, 68–69	hemorrhagic transformation, detection, 59
clinical trials, 532–533	hemostasis, surgical, methods for, 324, 342, 388
epidemiology, 517	in intraoperative aneurysm rupture, 444
FOUR Score, 68	heparins, 508–509
Glasgow Coma Scale, 66–68	herniation, 529
in cerebellar infarction, 210	brainstem, FOUR Score and, 68
as predictor of extubation, 293	as cause of death in stroke, 90
guidelines, 517, 533–534	in cerebral venous thrombosis, 96
Marshall CT classification, 69	after craniectomy, 98
primary brain damage in, 519–520	levels of, in space-occupying infarction, 191
prognosis and outcome, 533	lumbar drainage and, 161, 162–163
treatment, 527–532	paradoxical, 99
<i>See also</i> traumatic brain injury	prevention, 530
headache	herpes simplex encephalitis, 229
in cerebral venous thrombosis, 501, 505	hibernation, 275
diagnostic value of, 209	HIV, 309
differential diagnosis, 511	hormonal therapy, cerebral venous thrombosis and, 512
in subarachnoid hemorrhage, 426	Hunt and Hess scale, 72–73
hematological disorders, 237–238	hydrocephalus
and cerebral venous thrombosis, 506–507	aneurysm rupture and, 146, 159
and infection, 309	in cerebellar hemorrhage, 367
hematoma	in cerebellar infarction, 182
expansion, 257–259, 276	drainage in, 13
future strategies, 268–269	posthemorrhagic, 158, 161–162
pathophysiology, 259	drainage, 146–147, 159–161, 161–162, 353
predictors, 259–260	hyperbaric oxygen, 532
and prognosis, 350	hypertension
restricting, 260–268, 276–277	acute hypertensive response, 274, 275
extradural, 521–522	antihypertensives in, 278–280
treatment, 527	induced, 469, 470
	in intracerebral hemorrhage, 274

- hypertension (cont.)
 - hematoma expansion and, 262–263, 276
 - immediately post-stroke, 280–282
 - management, 262–263, 264, 276–277
 - management, 245–251
 - guidelines, 277–278, 282
 - syndromes, in cerebral venous thrombosis, 501
- hypertonic saline, 14, 83–84, 316–317, 433, 470
- hyperventilation
 - brain tissue oxygenation and, 39
 - in control of intracranial pressure, 85
 - in regulation of intracranial pressure, 4, 21, 287
- hypotension, 251
- hypothermia, therapeutic, 129, 185
 - clinical application
 - hemorrhagic stroke, 135
 - ischemic stroke, 133–134
 - in combination therapy, 135–136
 - future directions, 134–135
 - to lower intracranial pressure, 15
 - mechanisms of action, 129–131, 185
 - modes of cooling
 - invasive, 186
 - non-invasive, 185–186
 - onset and duration, 186–187
 - physiology and physics of, 131
 - side effects, 136–139
 - during thrombolytic therapy, 134
- imaging
 - in aneurysm, 447–448
 - for craniotomy, 438–439
 - of arterial malformations, 114
 - after basilar artery occlusion, 203
 - in basilar artery occlusion, 195–196
 - in cavernoma, 406–408
 - intraoperative, 414
 - in cerebellar hemorrhage, 366–367
 - in cerebellar infarction, 210–212
 - in cerebral venous thrombosis, 502–505
 - in head injury, 524
 - in ischemic stroke, 103
 - perfusion, after subarachnoid hemorrhage, 468–469
 - pre-/intraoperative, 388
 - in cavernoma resection, 414
 - in excision of arteriovenous malformations, 389
 - in intracerebral hemorrhage, 338–339
- immune-modulating nutrition, 302
- immunity in stroke patients, 306
- immunosuppression
 - for cervical artery dissections, 227
 - hypothermia and, 139
- infection
 - antibiotic strategies, 311
 - as cause of stroke, 228–229
 - blood-borne, 308
 - catheter-related, 8, 308
 - urinary, 307–308
 - hospital-acquired, modifiable risk factors, 310, 310
 - hypothermia and, 139
 - intracerebral hemorrhage-related, 308–309, 308
 - lumbar catheter, 163
 - management, 310–312
 - postoperative, in decompressive surgery, 97
 - treatment-related, 309–310
 - urinary tract, 307–308
 - ventricular drain-related, 357
 - ventriculostomy-related, 150–152
 - wound, hypothermia and, 139
- inflammation
 - in edema formation, 81, 82, 86, 315
 - effects after traumatic brain injury, 15
 - hypothermia and, 130
 - systemic inflammatory response syndrome, 308
 - and vasospasm, 467
- interventional treatment
 - aneurysm, options and procedures, 111–112
 - arterial malformations, 112–114
 - embolization techniques, 114–117
 - in combination with thrombolysis, 175–177
 - ischemic stroke, 103
 - indications, 103–104, 106–107
 - options and procedures, 104–106, 107–109
 - vasospasm
 - indications, 109
 - options and procedures, 109–111
 - See also* endovascular therapy
- intracerebral hemorrhage, 257, 335
 - acute hypertensive response and, 274, 275
 - anticoagulants and, 265–268
 - coagulopathy and, 323
 - decompressive hemicraniectomy in, 96
 - extending to ventricles, 145–146, 161–162, 259, 337
 - hematoma expansion, 257–259, 276
 - pathophysiology, 259
 - predictors, 259–260
 - restricting, 260–268, 276–277

- future strategies, 268–269
- hypertension and, 274
 - pathophysiology, 274–275
- preoperative management, 338–339
- related infection, 308–309, 308
- surgical removal, 336
 - craniotomy for, 323–326
 - effects of subgroup on, 321–323, 336
 - effects of timing on, 322, 325, 339–340
 - minimally invasive, 329–332, 336–337, 340–343
 - selection of patients, 320–321, 329
- intracranial atherosclerotic disease.
 - as cause of stroke, 120
 - See also under* stenosis
- intracranial compliance, 4, 5
- intracranial dynamics, 3–5
- intracranial hemorrhage
 - antiedema therapy in, 86–87
 - arterial malformations and, 112
 - brain tissue oxygenation monitoring in, 41
 - cavernoma-related, 408–410
 - cerebral blood flow in, 25
 - intracranial pressure monitoring in, 11–12
- intracranial pressure, 3
 - barbiturates and, 84–85
 - vs. cerebral perfusion pressure, 9
 - decompressive craniectomy in high, 91, 180
 - hypertonic saline and, 83–84
 - hyperventilation and, 85
 - hypothermia and, 129
 - mechanical ventilation in high, 287
 - monitoring, 3–8
 - in aneurysmal subarachnoid hemorrhage, 12–13
 - Brain Trauma Foundation guidelines, 13
 - in cerebral vein thrombosis, 12
 - complications, 8
 - during craniectomy, 92
 - in intracranial hemorrhage, 11–12
 - in ischemic stroke, 9–11
 - by lumbar catheter, 162
 - in traumatic brain injury, 13–15
 - treatment of raised, 529–530
 - tromethamine and, 85
 - ventricular drainage in high, 146, 150, 353
 - waveform, 5
- intraventricular hemorrhage, 145–146, 348–349
 - clearance, 146–147, 161–162, 353–357
 - endoscopic techniques, 337
 - conventional therapies, 353
 - grading, 351
 - investigational therapies, 353–354
 - pathophysiology, 349
 - prognosis, 349–351
 - score, 350, 351, 352
- intubation
 - in cerebellar hemorrhage, 370
 - indications for, in intracerebral hemorrhage, 286–288
 - options for, 288–290
- ischemia
 - cerebral, 22
 - delayed, 464–465
 - microdialysis to detect, 44, 47, 48–50, 468
 - cerebral blood flow as marker of, 22–24
 - continuous EEG to detect, 61
 - delayed cerebral
 - adjunctive therapies, 474
 - detection, 468–469
 - Fisher Grade Scale predicts, 72
 - hemodynamic goals, 471
 - hemodynamic interventions, 469–470, 472, 473–474
 - indications for treatment, 469
 - prevention, 464–465
 - edema in, 90
 - pathophysiology, 81
 - excitotoxic, 130
 - hypothermia and, 130, 186
 - traumatic brain injury and, 518–519
- ischemic brain damage, 520–521
 - prevention, 526
- Kernohan's phenomenon, 523
- lactate/pyruvate ratio, 46
 - as biomarker of ischemia, 48, 468, 485
- LeRoux Scale, 351, 352
- leucocyte activity, hypothermia and, 130, 139
- locked-in syndrome, 195
- lucid interval, in hematoma, 521, 522
- lumbar drainage, 158, 163–164
 - complications, 162–163
 - in intracranial hypertension, 161
 - intraoperative, 439
 - in intraventricular hemorrhage, 356–357
 - placement, 159
 - in subarachnoid hemorrhage, 159–161
 - in traumatic brain injury, 14
- lumbar puncture, in subarachnoid hemorrhage, 428

- malignant cerebral venous thrombosis, decompressive surgery
 - in, 96–97
 - malignant MCA infarction, 90, 179, 190
 - clinical features and prognosis, 190–191
 - decompressive surgery in, 92–95, 94
 - clinical trials, 180–181, 180
 - complications, 182
 - edema in
 - predictors of, 192
 - treatment, 192–193
 - herniation in, 191
 - hypothermia in, 185
 - imaging and prediction of, 182
 - signs and symptoms, 191
 - Marshall CT classification for head injury, 69
 - matrix metalloproteinases
 - in hematoma expansion, 269
 - hypothermia and, 130
 - mean transit time, 27
 - MELAS, 233
 - meningitis, 229, 309–310
 - antibiotic strategy, 311
 - ventriculostomy -related, 357
 - metabolic crisis
 - biomarkers, 46
 - glycemic control and, 51
 - metabolic demand
 - cerebral blood flow related to, 20
 - in intracranial hemorrhage, 25, 275–276
 - in subarachnoid hemorrhage, 25
 - measurement, 27
 - metabolic derangements after subarachnoid hemorrhage
 - cerebral, 483–485, 486, 496
 - systemic, 481–482, 486, 496
 - metabolic effects of hypothermia, 139, 141
 - microdialysis, 39, 44–47
 - in subarachnoid hemorrhage, 48–50, 483–484
 - in traumatic brain injury, 47–48
 - microemboli detection, transcranial Doppler for, 56
 - midline shift
 - in head injury, 69
 - in malignant MCA infarction, 179, 191
 - ultrasonography to monitor, 59
 - mismatch, PI/DWI, 172–173
 - Modified Fisher Grade Scale, 427, 465
 - Modified Rankin Scale, 73, 94
 - monitoring
 - brain tissue oxygenation, 37–39
 - in intracranial hemorrhage, 41
 - in ischemic stroke, 40
 - by near-infrared spectroscopy, 63
 - neurochemical, 39, 44–47
 - in subarachnoid hemorrhage, 40
 - by continuous EEG, 60–61
 - cerebral blood flow, 26–33, 27
 - comparison of methods, 29
 - indications for, 33
 - continuous non-invasive, 54
 - evoked potentials for, 61
 - in subarachnoid hemorrhage, 470–471, 494
 - intracranial pressure, 3–8
 - in aneurysmal subarachnoid hemorrhage, 12–13
 - during BP reduction, 278
 - Brain Trauma Foundation guidelines, 13
 - vs. cerebral perfusion pressure, 9
 - in cerebral vein thrombosis, 12
 - complications, 8
 - during craniectomy, 92
 - in intracranial hemorrhage, 11–12
 - in ischemic stroke, 9–11
 - by lumbar catheter, 162
 - in traumatic brain injury, 13–15
 - neuro-, in subarachnoid hemorrhage, 431
 - neurochemical, 39, 44–47
 - ultrasonography for
 - transcranial color-coded duplex, 58–60
 - transcranial Doppler, 6, 55–57
- Monro–Kellie doctrine, 3, 180, 529
- mortality
 - APACHE Score, 68
 - in cerebellar hemorrhage, 372, 372
 - in excision of arteriovenous malformation, 390
 - in subarachnoid hemorrhage, 425
- motor effects of cerebellar infarction, 208
- motor evoked potentials, 212
- Moya Moya syndrome, 230
- MR perfusion to measure cerebral blood flow, 27, 30, 31, 468
- MRI
 - of arterial malformations, 114
 - in basilar artery occlusion, 195–196
 - in brainstem hemorrhage, 379
 - in cavernoma, 407–408
 - of brainstem, 382
 - in cerebellar hemorrhage, 366
 - in cerebellar infarction, 211–212
 - in cerebral venous thrombosis, 503–504
 - in cervical artery dissections, 226
 - in evaluation for thrombolysis, 172–174

- in head injury, 524
- in prediction of edema, 192
- in subarachnoid hemorrhage, 428
- myocardial injury in subarachnoid hemorrhage, 493
- myocardial performance, hypothermia and, 140–141
- nausea, diagnostic value of, 208
- near-infrared spectroscopy for monitoring of cerebral blood flow, 32, 63
- neurogenic cardiopulmonary injury, 490
 - clinical features and diagnosis, 492–494
 - pathophysiology, 490–492
- neurological deficits
 - basilar artery occlusion and, 194, 195
 - Canadian Stroke Scale, 71
 - in cavernoma, 408
 - in cerebellar hemorrhage, 365, 365
 - in cerebellar infarction, 208
 - in cerebral venous thrombosis, 501, 507
 - delayed, 429, 464, 468
 - lumbar drainage and, 160
 - European Stroke Scale, 73
 - NIH Stroke Score, 70
 - Scandinavian Stroke Score, 70
- neurological examination in cerebellar infarction, 210
- neuromonitoring, 5–8, 44–47
- neuromuscular blocking agents in control of shivering, 186
- neurosonology. *See* ultrasound
- NIH Stroke Score, 70–71
 - in cerebellar infarction, 210
- nutrition
 - effect on prognosis, 297–298
 - enteral, 299–301
 - vs parenteral, 301–302
 - immune-modulating, 302
 - oral supplementation, 298–299
- nystagmus in cerebellar infarction, 209
- oligemia, 22, 26
- Onyx embolic agent, 114, 453
- opioids
 - to control shivering, 137
 - pharmacokinetics, hypothermia and, 140
- oral contraception, cerebral venous thrombosis and, 512
- osmotherapy, 82–84, 316–317
 - in brainstem hemorrhage, 381
- outcome
 - after decompressive surgery, 181–182
 - in cerebellar infarction, 95, 183
 - in hemispheric infarction, 92–95
 - edema and, 81
 - in endovascular therapy, 114
 - Glasgow Outcome Scale, 74–75
 - Modified Rankin Scale, 73
 - nutrition and, 297–298
 - of thrombolysis, time window and, 170
- overdrainage, herniation due to, 162–163, 164
- Oxfordshire Community Stroke Project classification, 69–70
- oximetry, brain tissue, 15
- oxygen
 - consumption, cerebral metabolic
 - effects of hypothermia, 129
 - See also* metabolic demand
 - extraction fraction (OEF), 20
 - in cerebral autoregulation, 22
 - in cerebral vasospasm, 25
 - in ischemic stroke, 24
 - saturation, cerebral blood flow and, 21
- oxygenation, brain tissue. *See* brain tissue oxygenation
- paralytic agents in rapid sequence intubation, 290
- parenchymal lesions, 505–506
- parenteral nutrition, 301–302
- pbrO₂. *See* brain tissue oxygenation
- penumbra, PI/DWI-mismatch and, 172–173
- perfusion studies
 - CT to measure cerebral blood flow, 27, 28–31, 30, 468–469
 - MR to measure cerebral blood flow, 27, 30, 31, 468
 - transcranial sonography, 60
- perioperative stroke, 235–236
- PET to monitor cerebral blood flow, 27, 30
- pharmacokinetics, hypothermia alters, 139–140, 140
- pneumonia
 - antibiotic strategies, 311
 - aspiration, dysphagia and, 300
 - hospital-acquired, 306–307
 - ventilator-associated, 307
 - hypothermia and, 139
- positive pressure ventilation, 291–292
 - in brainstem hemorrhage, 380
- pregnancy
 - cerebral venous thrombosis in, 512
 - ruptured aneurysm in, 455
 - stroke in, 236–237
- primary pontine hemorrhage, 378, 379
 - prognosis, 380
- procoagulatory states, 237, 506

- prognosis
 - after decompressive surgery, 92–95, 181–182
 - enteral nutrition and, 299–301
 - Scandinavian Stroke Score for, 70
- prophylaxis
 - angioplasty as, 432, 467
 - antibiotic
 - in cerebrospinal fluid leak, 531
 - in ventilator-associated pneumonia, 307
 - for ventricular drainage, 150–151, 357
 - thrombo-, for ventricular drainage, 152–153
- prothrombotic conditions, 237, 506
- proximal control, 442–443
- pulmonary complications, 493–494
 - See also* cardiopulmonary dysfunction
- pulmonary edema, 433, 492, 493–494
- radiosurgery
 - in arteriovenous malformations, 394–396
 - for cavernoma, 415
- radiotherapy for arteriovenous malformations, 396
- re-bleeding
 - in subarachnoid hemorrhage, 428
 - See also* hematoma expansion
- recanalization
 - in basilar artery occlusion, 196–202
 - in cerebral venous thrombosis, 504
 - indications, 103
 - interventional treatment, 103
 - indications, 103–104, 106–107
 - options and procedures, 104–106, 107–109
 - middle cerebral artery, timing and prognosis, 56
 - outcome and, in basilar artery occlusion, 201
 - rates, 56
- renal replacement therapy in ischemic stroke, 10–11
- reocclusion, basilar artery, 202
- respiratory drive
 - brainstem hemorrhage and, 288
 - intracerebral hemorrhage and, 286–287
 - and mode of ventilation, 291
- respiratory failure in intracerebral hemorrhage, 286
- rewarming, 134
 - electrolytes in, 141
- risks of treatment, balancing, 390
- saline, hypertonic, 14, 83–84, 316–317, 433, 470
- salt-wasting syndrome, 433, 482
- scales, 67
 - APACHE Score, 68–69
 - ASPECTS, 104
 - Barthel ADL Index, 74
 - Bedside Shivering Assessment, 137, 137
 - Canadian Neurological, 71
 - CVT risk score, 508
 - European Stroke, 73
 - Fisher Grade, 72
 - Modified, 427, 465
 - FOUR Score, 68
 - Glasgow Coma, 66–68
 - Glasgow Outcome Scale, 74–75
 - Graeb Score, 147, 351, 352
 - Hunt and Hess, 72–73
 - IVH Score, 350, 351, 352
 - LeRoux Scale, 351, 352
 - Modified Rankin, 73, 94
 - NIH Stroke Score, 70–71
 - Oxfordshire Community Stroke Project classification, 69–70
 - Scandinavian Stroke, 70
 - Spetzler–Martin Grading System, 390, 395, 395
 - for subarachnoid hemorrhage, 427
 - Therapy Intensity Level, 530
 - WFNS Grading for SAH, 71–72
- Scandinavian Stroke Scale, 70
- sedatives
 - in brainstem hemorrhage, 380
 - to control shivering, 132, 137, 186
 - pharmacokinetics, hypothermia and, 140
 - for rapid sequence intubation, 289–290
- seizures
 - in cavernoma, 408, 412
 - surgery for, 413–414
 - in cerebral venous thrombosis, 505, 511
 - differential diagnosis, 505
 - after subarachnoid hemorrhage, 434
- sepsis, 308
 - antibiotic strategy, 311
- shivering, 133, 136–138, 186
 - Bedside Shivering Assessment Scale, 137, 137
- shunt
 - infections, antibiotic strategy, 312
 - venous, in regulation of ICP, 4
 - ventricular, 13, 152
- SIADH, 482
- sickle cell disease
 - and infection, 309
 - ischemic stroke risk, 57
- sinking skin syndrome, 100, 182
- skull fracture, depressed, 531–532

- somatosensory evoked potentials, 212
 - for monitoring, 61
- spasmolysis, 458, 472
 - indications, 109
 - options and procedures, 109–111
- SPECT in monitoring of cerebral blood flow, 27
- Spetzler–Martin Grading System, 114, 390, 395, 395
- spinal cord infarction, 237
- statin therapy in subarachnoid hemorrhage, 434
- stenosis, arterial
 - intracranial
 - balloon vs. stent-assisted angioplasty in, 122
 - stenting vs. medical management in, 120
 - stenting in, 106
 - indications for, 106–107
 - options and procedures, 107–109
- stent, 106
 - in aneurysm, 450–453
 - assisted coiling, 112
 - devices, 107
 - in failed thrombectomy, 106
 - flow-diverting, 112, 453
 - indications for, 106–107
 - intracranial, 120
 - placement, 125
 - techniques, 123
 - vs. medical management, 120
 - options and procedures, 107–109
 - retriever, use for thrombectomy, 104
 - for thrombectomy, 202
 - Wingspan, 120
- stereotactic aspiration
 - in hematoma expansion, 261, 268
 - of intracerebral hemorrhage, 331
 - with fibrinolysis, 317, 331, 332
- stereotactic evacuation of intracerebral hemorrhage, 329–332, 336–337, 340–343
- steroids as antiedema therapy, 86
- strokeectomy, 217
- subarachnoid hemorrhage, 111, 146, 423, 425–426
 - brain tissue oxygenation and, 40
 - cardiopulmonary dysfunction in, 490
 - cardiomyopathies, 492–493, 495
 - features and diagnosis, 492–494
 - pathophysiology, 490–492
 - teatment, 495–496
 - cerebral blood flow in, 25–26
 - clinical presentation, 426
 - complications of, 428–429
 - decompressive craniectomy in, 96
 - diagnosis, 426–428
 - Fisher Grade scale, 72
 - hydrocephalus after, 146, 159–160
 - intracranial pressure monitoring in, 12–13
 - lumbar drainage in, 159–161
 - management, 494–495
 - emergency department, 429–431
 - intensive care, 431–434
 - monitoring, 470–471
 - novel strategies, 434–435
 - metabolic derangements after
 - cerebral, 483–485
 - systemic, 481–482
 - microdialysis in, 48–50, 483–484
 - monitoring in, 494
 - Moya Moya syndrome and, 230
 - pathogenesis, 423–425
 - pathophysiology of secondary injury, 480–481
 - peri-mesencephalic, 424
 - rare causes, 424
 - recurrent, 428
 - WFNS Grading System, 71–72
- surgery
 - in aneurysm
 - aneurysm rupture during, 443–444
 - indications, 437–438
 - microscopic approaches, 440–442
 - minimally invasive, 440
 - proximal control, 442–443
 - repair techniques, 443–444
 - surgical techniques, 439–440, 445
 - timing and preparation, 438–439
 - for arteriovenous malformations
 - decision-making, 390–392
 - with hemorrhage, 392–393
 - radiosurgery, 394–396
 - techniques, 387–389
 - in brainstem hematoma, 382–383
 - in brainstem hemorrhage, 381
 - for cavernoma, 413–415
 - radiosurgery, 415
 - decompressive, 10, 90–92, 180
 - in cerebellar infarction, 182–183
 - in cerebral contusions, 528–529
 - in cerebral venous thrombosis, 509–510
 - clinical trials, 530, 532–533
 - complications, 97–100

- surgery (cont.)
 - converting craniotomy to, 324
 - in extradural hematoma, 527
 - in hemorrhagic stroke, 317
 - indications, 92–97
 - in intracerebral hematoma, 528
 - for ischemic stroke, 179
 - in malignant MCA infarction, 94, 182
 - in malignant MCA infarction, clinical trials, 180–181, 180
 - in subarachnoid hemorrhage, 96
 - in subdural hematoma, 527–528
 - suboccipital, 217, 371
 - in traumatic brain injury, 14
 - for intracerebral hemorrhage, 336
 - craniotomy, 323–326
 - effects of subgroup on, 321–323, 336
 - effects of timing on, 322, 325, 339–340
 - minimally invasive, 329–332, 336–337, 340–343
 - selection of patients, 320–321, 329
 - in intraventricular hemorrhage
 - minimally invasive, 356
 - in prevention of hematoma expansion, 260–262
 - sympathetic storm, 490
 - systemic inflammatory response syndrome, 308
- Therapy Intensity Level score, 530
- thermal diffusion flowmetry to measure cerebral blood flow, 30, 31–32
- thermoregulation, 131, 186
 - sedatives and, 137
- thrombectomy
 - devices, 106
 - mechanical, 103, 104–106
 - in basilar artery occlusion, 201–202
- thromboembolism
 - during endovascular therapy, 456
 - posthemorrhagic, 266
- thrombolysis, 103
 - in basilar artery occlusion, 198–201
 - contraindications, 200
 - blood pressure after, 246
 - Bridging Concept, 175
 - in cerebral venous thrombosis, 509
 - in cervical artery dissections, 227
 - clinical trials, 169–171, 174
 - based on MRI, 172–174
 - in combination therapies, 175–177
 - in combination therapy, 136
 - in hematoma expansion, 268
 - hypothermia and, 134
 - ongoing trials, 134–135
 - before interventional treatment, 104
 - intraarterial, 174–175, 198
 - clinical trials, 175
 - intraventricular, 147, 161
 - intraventricular combined with lumbar drainage, 162
 - intraventricular combined with ventricular drainage, 353–356
 - local intraarterial, 104
 - in failed thrombectomy, 106
 - and minimally invasive surgery, 317, 331, 332
 - in pregnancy, 237- thrombophilia, 237–238
- thromboprophylaxis
 - for ventricular drain, 152–153
 - See also* anticoagulants
- time window
 - and choice of treatment, 103, 104
 - for decompressive surgery, 182
 - for surgery in hematoma, 521, 522, 527
 - in therapeutic hypothermia, 133
 - for thrombolysis
 - in basilar artery occlusion, 200–201
 - hypothermia and, 134
 - outcome and, 170
- tracheotomy, 294
 - in brainstem hemorrhage, 380
- transfusion in subarachnoid hemorrhage, 471–472
- trauma, as cause of dissection, 226, 521
- traumatic brain injury
 - brain tissue oxygenation monitoring in, 37–39
 - decompressive surgery in, clinical trials of, 533
 - epidemiology, 517
 - hemicraniectomy in, 98
 - intracranial pressure monitoring in, 13–15
 - ischemia resulting from, 518–519
 - management, 517
 - Marshall CT classification, 69
 - microdialysis in, 47–48
- triple-H therapy, 109, 432–433, 469, 471, 487
 - adverse effects, 473
- ultrasonography
 - EKOS device, 175
 - intraoperative, 389
 - transcranial color-coded duplex, 55
 - for monitoring, 6, 55–57
 - transcranial Doppler, 54
 - in cerebellar infarction, 212

- to measure cerebral blood flow, 32, 109, 467–468
 - for monitoring, 58–60
 - in vasospasm, 432
- urinary tract infections, 307–308
- antibiotic strategies, 312
- urokinase, 104
- vasculitis as cause of stroke, 227–230
- vasospasm, 480–481
- and brain tissue oxygenation, 39, 40
 - continuous EEG to detect, 61
 - and delayed cerebral ischemia, 464–465
 - detection of, 467–468
 - diagnosis, 428
 - endovascular therapy in, 457–458, 472–473
 - Fisher Grade scale predicts, 72
 - grading criteria, 57
 - interventional therapy, 109
 - indications, 109
 - options and procedures, 109–111
 - microdialysis in, 48
 - prevention, 464–465
 - by lumbar drainage, 160–161
 - primary, 231–232
 - in subarachnoid hemorrhage, 25, 160–161, 431–432
 - novel strategies for, 434–435
 - transcranial Doppler to detect, 57, 109, 432
- vegetative state, 533
- venography in cerebral venous thrombosis, 502–505
- venous angioma, 401
- venous blood shunting in regulation of ICP, 4
- ventilation, mechanical
- in brainstem hemorrhage, 380
 - in cardiopulmonary complications, 496
 - indications for in intracerebral hemorrhage, 286–288
 - lung protective, 292
 - methods of, 288–290
 - specialized, 292
 - modes of, 291
 - parameters, 291
- pneumonia and, 139, 307
 - positive end-expiratory pressure, 291–292
 - weaning, 292–294
- ventricular drainage, 8, 145–147
- in brainstem hemorrhage, 381
 - in cerebellar hemorrhage, 371
 - in cerebellar infarction, 183, 217
 - complications, 149–150
 - insertion, 147–149
 - in intracranial hemorrhage, 11–12
 - management, 150–152, 357
 - related infections, 357
 - removal, 152
 - and risk of hemorrhage, 8
 - risk of infection, 309
 - thromboprophylaxis, 152–153
 - in ventricular hemorrhage, 353
- ventriculitis, 309–310
- ventriculostomy-related, 150–152, 357
- ventriculostomy
- endoscopic third, in cerebellar infarction, 218
 - See also* ventricular drainage
- vertigo, diagnostic value of, 208
- viral infection as cause of stroke, 229
- vitamin-K-antagonists and hematoma expansion, 265–266
- volume status
- after subarachnoid hemorrhage, 494–495
 - and triple-H therapy, 432
 - hypotension, 251
 - hypothermia and, 141
 - subarachnoid hemorrhage and, 433, 466, 469
- warming in therapeutic hypothermia
- rewarming, 134
 - skin-surface, 133, 137
- weaning
- from external ventricular drain, 152
 - from mechanical ventilation, 292–294
- WFNS Grading System for SAH, 71–72
- wound infection, hypothermia and, 139