

# Index

---

## A

- Accept–reject simulator (AR), 115–20
  - importance sampling and, 210–12
  - mixed logit and, 145
  - problems with, 118–20
  - smoothed, 120–2
  - truncated densities, 209–10
- Aggregation, 29–32, 54
- Akaike Information Criterion (AIC), 367
- Allocation parameter, 91
- Alternative-specific constants, 20–1, 33, 62, 67, 73, 153–5, 167, 308
- Anticipatory behavior, 113, 146
- Antithetic draws, 216–18, 221, 236
- AR. *See* Accept–reject simulator
- Asymptotic distribution, 219, 245–6, 249, 255–6, 264, 288, 290, 293
- Attraction variables, 54
- Automobile demand models, 134
- Average conditional distribution, 269–70

## B

- BART. *See* Bay Area Rapid Transit (BART)
- Base-year shares, 33
- Bay Area Rapid Transit (BART), 71
- Bayes' rule, 262, 285–6
- Bayesian Information Criterion (BIC), 367
- Bayesian methods, 282–314
  - hierarchical, 283, 299–305
  - mixed logit and, 299–305
  - overview of, 284–91
  - probit and, 313–14
- Berndt–Hall–Hall–Hausman method (BHHH), 192–8, 258
- Bernstein-von Mises theorem, 266–7, 288–92
- BFGS. *See* Broyden–Fletcher–Goldfarb–Shanno method
- BHHH. *See* Berndt–Hall–Hall–Hausman method
- Binary logit, 4, 35, 39–40, 162
- Binary probit, 7, 14, 97, 115, 317
- BLP, 8, 317–28, 334–5, 339, 342, 346
- Bootstrap, 201–2, 355, 370
- Broyden–Fletcher–Goldfarb–Shanno method (BFGS), 197

## C

- California Energy Commission (CEC), 48
- Car–bus example, 17, 20
- Causal perspective, 3
- Central limit theorem, 245–6, 249, 252, 292
- Chatter, in simulations, 250
- Choice probabilities, 18–21. *See specific models*
- Choice sets, 11–14, 37, 57, 73, 157, 282
- Choleski factor, 116–17, 123–4, 129–30, 208–9, 265, 297, 299, 302, 307–8, 338, 364
- Clark algorithm, 115
- College/work model, 169–82
- Concavity, 191–2
- Conditional distribution, 114, 260, 262–7, 269–70, 274–8, 280, 335–42, 349, 358, 362
  - average, 269–70
  - derivation of, 262–4
- Conditional likelihood function, 65
- Constants, recalibration of, 33
- Consumer surplus, 55–7
- Contingent valuation, 164–6
- Contraction, 317, 322–4, 335, 343
- Control function, 8, 317–18, 334–40
- Convenient error partitioning, 6, 113, 173, 176
- Convergence, criterion for, 198–9
- Cross-nested logits, 89, 91
- Cross-subject variance, 112

## D

- Davidon–Fletcher–Powell method (DFP), 197–8
- Densities, drawing from, 205–36
- Derivatives, 29–31, 39, 47, 57–61, 63, 73, 93–5, 119–20, 123, 186–7, 191, 193, 197, 202–4, 243–4, 249–50, 253–4, 287, 289, 334, 351, 354–5, 365
- DFP. *See* Davidon–Fletcher–Powell method
- Diffuse prior, 297–8, 301
- Dimensionality, curse of, 178
- Discrete mixing distribution, 355–61
- Double-bounded methods, 165
- Dynamic choice models, 50–2, 114, 169–82, 314
- Dynamic optimization, 169–82

## E

- Elasticities, 29–31, 45, 47–8, 57–60, 141, 144, 286–7, 329, 332–3, 339–40

## 386 Index

- Electric cars, 48  
 Elimination by aspects model, 81  
 EM algorithm, 347–70  
 Endogeneity, 315–46  
 Energy efficient vehicles, 48  
 Energy supplier, choice of, 152–3, 270–80, 305–13  
 Exclusivity conditions, 12–13  
 Exhaustiveness, 12–13  
 Exogenous sample, 60–6  
 Expectation-maximization algorithm. *See* EM algorithm  
 Explanatory variables, 8, 14, 29, 31–2, 37, 50, 60, 62–3, 70–1, 81, 85, 88, 103, 134, 136, 138, 140, 157, 162, 203, 240–2, 270, 285, 309, 315, 317, 319–21, 324, 327, 331, 336, 339, 343, 348  
 Exploded logit, 157, 321, 342–3, 367  
 Exponential transformation, 81  
 Extreme value terms, 16, 18–19, 34–5, 39, 40, 43, 51, 55–6, 74, 76, 79–80, 89, 92, 137, 139, 142–3, 145, 154, 156–7, 162, 169, 173–4, 176, 181, 207, 262, 267, 300, 304, 308, 319, 321, 337–9, 341, 356
- F**  
 Factor-analytic structure, 103  
 Fishing site model, 147–50, 259  
 Fixed coefficients, 139–40, 308–10, 317–18  
 Forecasting, 11, 32–3, 36, 44, 71–4, 144, 156, 280, 328–9  
 Full information maximum likelihood, 8, 335
- G**  
 Generalized autoregressive structure, 103  
 Generalized extreme value models (GEV), 11, 17–18, 50, 76–96  
   heteroskedastic logit, 92  
   nested logit and, 76–88, 91–2  
   ordered, 89  
   PCL model and, 88, 90–1, 94, 96  
   substitution patterns, 77–8  
 Generalized least squares (GLS), 325  
 Generalized method of moments (GMM), 326–8, 331, 333–4  
 Geographic aggregation, 54  
 GEV. *See* Generalized extreme value models  
 Geweke simulator. *See* GHK simulator  
 GHK simulator, 115, 122–33  
   as importance sampler, 131–3  
 Gibbs sampling, 212, 214, 284, 293–5, 299, 300–4, 306, 309–11, 313  
   MH algorithm and, 213–14, 293–4, 300–3, 309  
 Global maximum, 199, 270, 283  
 Goodness of fit, 68–9  
 Goods-leisure tradeoff, 53  
 Gouieroux–Monfort model, 7  
 Gumbel distribution, 34
- H**  
 Hajivassiliou simulator. *See* GHK simulator  
 Halton sequences, 224–38  
   multidimensional, 226–7  
   randomized, 231–3  
   scrambled, 233–5  
 Heating system example, 38–40, 66  
 Hessian matrix, 186  
   expected, 194, 200–2  
   information identity, 202–4  
   information matrix, 200–4  
 Heteroskedastic extreme value models (HEV), 92  
 Heteroskedasticity, 18–19, 168–9  
 HEV models, 92  
 HEV. *See* Heteroskedastic extreme value models  
 Hierarchical Bayesian procedures, 283, 299–305  
   for probit, 313–14  
 Hydrogen vehicles, 365–6  
 Hyper-parameters, 305  
 Hypothesis testing, 70–1, 81, 280, 288, 292
- I**  
 Identification, 19–29, 100–6  
 IIA, 45–50  
   advantages of, 48–9  
   nested models, 78, 88  
   probit model, 108–10  
   tests of, 49–50  
 Importance sampling, 131, 210–12, 293  
 Inclusive value, 83, 85, 87–8, 139–40  
 Independence assumption, 18, 35  
 Independence from irrelevant alternatives. *See* IIA  
 Individual-level parameters, 259–81  
 Information identity, 194–6, 200–4, 248, 250, 255, 257  
 Information matrix, 200, 204, 248, 289  
 Initial conditions problem, 114  
 Instrumental variables estimation (IV), 317, 330  
 Integral expressions, 3–6  
 Inverted gamma, 297–9, 307  
 Inverted Wishart, 298–301, 308, 312
- K**  
 Keane simulator. *See* GHK simulator
- L**  
 Lagged response, 51, 113, 146  
 Latent class model, 135, 261, 357, 359  
 Likelihood function. *See* Log-likelihood  
 Likelihood ratio index, 68–9, 72, 149  
 Limiting distribution, 245–6, 249, 255, 257, 292  
 Linear regression models, 29, 52, 62, 261, 317, 321, 324–5  
 Liu–Mahmassani method, 199–200  
 Local maximum, 199  
 Log-likelihood function, 7, 37, 52, 63, 65, 67–8, 70–1, 84, 118–19, 129–30, 144, 162, 166, 185, 187, 189, 191–5, 197–9, 215, 225, 237–9, 248, 324, 341  
   curvature of, 193–5, 197  
   globally concave, 37, 191  
   maximization of, 185–204  
   simulated, 8, 118, 144, 185, 237, 264–5, 282–3, 324–6  
 Logistic distribution, 35, 162  
 Logit model, 4, 18, 24–5, 29, 34–75  
   applicability of, 42

BART ridership, 71–4  
 choice-based samples, 66–7  
 consumer surplus, 55–7  
 cross-elasticities, 48, 60  
 decomposition into, 81–4  
 derivation of, 74–5  
 exploded logit, 157  
 extreme value distribution, 34–5, 56  
 heteroskedastic logit, 92, 140  
 IIA, 45–50  
 limitations of, 42–52, 97  
 logit-kernel probit, 121  
 log-sum term, 56  
 maximum-likelihood estimator, 61–3  
 ordered logit, 162–3, 165, 167  
 power of, 42–52  
 probability, 37  
 smoother, 121–2  
 substitution patterns, 45–50  
 taste variation, 42–5  
 tests of IIA, 49–50  
 Lognormal distributions, 148, 282, 310–12  
 Log-sum term, 56, 83, 181–2

**M**

Marginal cost, 328–34, 339–40  
 Marginal cost pricing, 330–1, 339–40  
   with control function, 339–40  
   fixed markup over, 331  
   with GMM, 331  
   with MSL/IV, 330–1  
 Marketing efforts, 315–16  
 Market-share data, 33  
 Markov chain Monte Carlo methods (MCMC), 8, 214, 293  
 Maximization methods, 185–204  
 Maximum-likelihood estimation, 61, 118, 129, 323, 327, 343  
 Maximum simulated likelihood (MSL), 8, 118, 144, 237–9, 255–6, 264, 283, 291, 305, 324–6  
 MCMC. *See* Markov chain Monte Carlo methods (MCMC)  
 MC-squared methods. *See* Markov chain Monte Carlo methods (MCMC)  
 Method of moments (MOM), 237–8, 240–2, 247–8, 250, 256, 326, 328  
 Method of simulated moments (MSM), 8, 237–8, 240–3, 256–7, 326, 328  
 Method of simulated scores (MSS), 8, 238, 243–5, 257  
 Metropolis–Hastings algorithm (MH), 205, 213–14, 284, 293–4, 300–3, 309–10, 312  
   Bayesian procedures and, 293–4, 300–3, 309–10, 312  
   Gibbs sampling and, 213–14, 293–4, 300–3  
 MH. *See* Metropolis–Hastings algorithm (MH)  
 Mixed logit, 2, 6–7, 16–17, 19, 29, 37, 45, 48, 50, 52, 122, 137–9, 141–5, 156–9, 167–8, 299–305  
   accept–reject simulator, 145  
   Bayesian procedures and, 299–305  
   defined, 134  
   error components, 139–41

  mixed nested logit, 167–8  
   mixing distribution, 141–3  
   random coefficients, 137–9  
   ranked data, 156–9  
   relation to probit model, 134  
   RUMs and, 141–3  
   simulation, 144–5  
   substitution patterns, 141  
 Mixed models, 166–9  
 Mixed nested logit, 167–8  
 Mixed probit, 19, 168–9, 314, 337  
 Mixing distribution, 50, 135, 138, 141, 143, 167, 355–65  
 ML. *See* Maximum-likelihood method  
 MOM. *See* Method-of-moments  
 Monopoly pricing, 332–4, 339  
 MSL. *See* Maximum simulated likelihood  
 MSM. *See* Method of simulated moments  
 MSS. *See* Method of simulated scores  
 Multi-level nested logits, 182  
 Multiple ordered responses, 163–4  
 Multiresponse probit, 164, 166  
 Multivariate ordered probit, 164

**N**

Nash equilibrium, 332–4, 339  
 Nested logit, 16, 29, 37, 48, 76–7  
   estimation and, 84–5  
   generalized, 91–2  
   GEV models and, 76–7, 94–5  
   IIA, 77–8, 88  
   overlapping nests, 89–92  
   PCL model, 90–1  
   software, 84  
   substitution patterns, 77–8  
   three-level, 86–8  
   two-level, 81–4  
 Newton–Raphson method, 187–92  
 Nonlinear representative utility, 52–4  
 Nonparametric estimation, 356, 361, 365  
 Non-rational model, 171  
 Nonsimulation procedures, for integrals, 114  
 Non-uniform conditioning, 66  
 Normal densities, 111–13, 116–17, 120, 124–5, 131–2, 136, 205–6, 208–9, 265, 296, 300, 302, 362  
   Choleski factor and, 208–9  
   multivariate, 208–9  
   transformations of, 206  
 Normalization, 11, 16, 21, 23–9, 91, 100–8, 110, 163  
 Numerical maximization, 37, 84, 185–204, 311

**O**

Ordered logit, 162–3, 165–7  
 Ordered probit, 162–4, 166  
 Ordered responses, 159–64  
 Outside good, 319, 342  
 Overlapping nests, 89–92, 139–40, 167

**P**

Paired combinatorial logit (PCL) model, 77, 89–91, 93, 96

## 388 Index

- Panel data, 50–2, 97, 110–15, 134, 145–7, 158  
 PCL. *See* Paired combinatorial logit model  
 Percent correctly predicted statistic, 69  
 Posterior distribution, 265–7, 288–99  
   Bernstein–von Mises theorem, 266, 267, 288–91  
   drawing from, 293–4  
   mean of, 291–3, 295–9  
 Prime numbers, Halton draws and, 221–35  
 Probit model, 7, 14, 23, 100–6, 108–10, 115–20, 122–33, 142, 218, 313–14  
   accept–reject simulator for, 115–20  
   antithetic draws and, 218  
   approximation by mixed logit, 142  
   Bayesian procedures and, 313–14  
   GHK simulator, 122–33, 168  
   identification, 23, 100–6  
   IIA, 108–10  
   mixed, 19, 168–9, 314, 337  
   normalization, 100–6  
   random coefficients, 106  
   ranked data, 158–9  
   route choice, 103  
   simulation, 115  
   substitution patterns, 108–10  
   unobserved utility, 97  
 Proportionate substitution, 48  
 Proposal density, 211, 393  
 Pseudo-observations, 157–8
- Q**
- Quadratic functions, 188–90, 197, 199, 347  
 Quadrature methods, 92, 115, 134  
 Quasi-random methods, 7, 230, 236. *See also* Halton sequences
- R**
- Random coefficients, 103, 106, 137–40, 147, 168, 259–60, 270, 308–9, 317, 342–4, 348, 356, 362  
 Random draws, 8, 149, 205–15, 218–19, 224, 228–31, 236, 311, 314  
 Random utility models (RUMs), 14, 141–3, 145. *See specific models*  
 Ranked data, 156–9  
 Recalibration, of constants, 33  
 Regime-switching models, 261  
 Representative utility, 15, 29–31, 35–8, 40, 42–3, 45–7, 51–62, 65, 67, 73, 76, 81, 106, 113  
 Respondent fatigue, 26  
 Revealed-preference data, 152–6  
 Robust covariance matrix, 201  
 RUMs. *See* Random utility models
- S**
- Sampling, 60–7, 131–3, 200–2, 210–12, 218–21, 238, 245–6, 264–5, 274–8, 280, 284, 286, 288–91, 293–5, 300–1, 304, 306–7, 309–11, 313, 323  
 of alternatives, 64–6  
 choice-based, 66–7  
 exogenous, 60–6  
 Sandwich estimator, 201  
 Scale parameter, 26, 40–2, 155  
 Segmentation, 31–2, 219  
 Sigmoid shape, 38  
 Simulated log-likelihood function. *See* Maximum simulated likelihood  
 Simulated mean of the posterior (SMP), 292–3, 301, 305  
 Simulated moments. *See* Method of simulated moments  
 Simulated scores. *See* Method of simulated scores  
 Site choice model, 148  
 Smoothing, 121–2, 144–5, 274  
 Stated-preference experiments, 152–6, 228, 270, 365  
 Steepest ascent method, 196–8  
 Step function, 119, 121, 145  
 Step size, 189–91  
 Student choice model, 171–82  
 Substitution patterns, 36, 42, 45–50, 77–8, 108–10, 141  
   logit model, 42, 45–50  
   mixed logit, 141  
   nested logit models, 77–8  
   probit model, 108–10  
   proportionate, 47–8  
   *See also specific models*  
 Systematic sampling, 205, 218–21
- T**
- Target density, 131, 211  
 Taste variation, 42–5, 97, 106–7, 123, 134, 138, 146, 282  
 Taylor expansion, 187, 191, 196, 249, 253–4  
 Triangular density, 312  
 t-statistic, 70, 72, 85, 199, 344  
 Type I extreme value distribution, 34
- U**
- Uncertainty, 58, 171, 178–82, 297, 368  
 Uniform conditioning property, 65  
 Univariate densities, drawing from, 208–11  
 Unobserved attributes, 89, 315–16, 319–20, 325–6, 329, 339, 345–6
- V**
- Variance reduction, 214–36  
 Vehicle choice, 322, 325, 343–4, 349–55
- W**
- Weibull distribution, 34  
 White noise, 35–6, 76  
 Work/college model, 169–82