

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

Condensed Matter Field Theory

Second edition

Modern experimental developments in condensed matter and ultracold atom physics present formidable challenges to theorists. This book provides a pedagogical introduction to quantum field theory in many particle physics, emphasizing the applicability of the formalism to concrete problems.

This second edition contains two new chapters developing path integral approaches to classical and quantum nonequilibrium phenomena. Other chapters cover a range of topics, from the introduction of many-body techniques and functional integration, to renormalization group methods, the theory of response functions, and topology. Conceptual aspects and formal methodology are emphasized, but the discussion focuses on practical experimental applications drawn largely from condensed matter physics and neighboring fields.

Extended and challenging problems with fully-worked solutions provide a bridge between formal manipulations and research-oriented thinking. Aimed at elevating graduate students to a level where they can engage in independent research, this book complements graduate level courses on many particle theory.

ALEXANDER ALTLAND is Professor of Theoretical Condensed Matter Physics at the Institute of Theoretical Physics, University of Köln. His main areas of research include mesoscopic physics, the physics of interacting many particle systems, and quantum nonlinear dynamics.

BENJAMIN D. SIMONS is Professor of Theoretical Condensed Matter Physics at the Cavendish Laboratory, University of Cambridge. His main areas of research include strongly correlated condensed matter systems, mesoscopic and ultracold atom physics.

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

Condensed Matter Field Theory

Second edition

Alexander Altland and Ben Simons



Cambridge University Press
978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition
Alexander Altland and Ben Simons
Frontmatter
[More information](#)

CAMBRIDGE UNIVERSITY PRESS
Cambridge, New York, Melbourne, Madrid, Cape Town, Singapore,
São Paulo, Delhi, Dubai, Tokyo

Cambridge University Press
The Edinburgh Building, Cambridge CB2 8RU, UK

Published in the United States of America by Cambridge University Press, New York

www.cambridge.org
Information on this title: www.cambridge.org/9780521769754

© A. Altland and B. Simons 2010

This publication is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without the written
permission of Cambridge University Press.

First published 2006
Reprinted 2007, 2008, 2009
Second edition published 2010

Printed in the United Kingdom at the University Press, Cambridge

A catalogue record for this publication is available from the British Library

ISBN 978-0-521-76975-4 Hardback

Cambridge University Press has no responsibility for the persistence or
accuracy of URLs for external or third-party Internet websites referred to in
this publication, and does not guarantee that any content on such websites is,
or will remain, accurate or appropriate.

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

Contents

<i>Preface</i>	<i>page</i> ix
1 From particles to fields	1
1.1 Classical harmonic chain: phonons	3
1.2 Functional analysis and variational principles	11
1.3 Maxwell's equations as a variational principle	15
1.4 Quantum chain	19
1.5 Quantum electrodynamics	24
1.6 Noether's theorem	30
1.7 Summary and outlook	34
1.8 Problems	35
2 Second quantization	39
2.1 Introduction to second quantization	40
2.2 Applications of second quantization	50
2.3 Summary and outlook	83
2.4 Problems	83
3 Feynman path integral	95
3.1 The path integral: general formalism	95
3.2 Construction of the path integral	97
3.3 Applications of the Feynman path integral	112
3.4 Problems	146
3.5 Problems	146
4 Functional field integral	156
4.1 Construction of the many-body path integral	158
4.2 Field integral for the quantum partition function	165
4.3 Field theoretical bosonization: a case study	173
4.4 Summary and outlook	181
4.5 Problems	181
5 Perturbation theory	193

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

vi

5.1	General structures and low-order expansions	194
5.2	Ground state energy of the interacting electron gas	208
5.3	Infinite-order expansions	223
5.4	Summary and outlook	232
5.5	Problems	233
6	Broken symmetry and collective phenomena	242
6.1	Mean-field theory	243
6.2	Plasma theory of the interacting electron gas	243
6.3	Bose–Einstein condensation and superfluidity	251
6.4	Superconductivity	265
6.5	Field theory of the disordered electron gas	301
6.6	Summary and outlook	329
6.7	Problems	331
7	Response functions	360
7.1	Crash course in modern experimental techniques	360
7.2	Linear response theory	368
7.3	Analytic structure of correlation functions	372
7.4	Electromagnetic linear response	389
7.5	Summary and outlook	399
7.6	Problems	400
8	The renormalization group	409
8.1	The one-dimensional Ising model	412
8.2	Dissipative quantum tunneling	422
8.3	Renormalization group: general theory	429
8.4	RG analysis of the ferromagnetic transition	444
8.5	RG analysis of the nonlinear σ -model	456
8.6	Berezinskii–Kosterlitz–Thouless transition	463
8.7	Summary and outlook	474
8.8	Problems	475
9	Topology	496
9.1	Example: particle on a ring	497
9.2	Homotopy	502
9.3	θ -0terms	505
9.4	Wess–Zumino terms	536
9.5	Chern–Simons terms	569
9.6	Summary and outlook	588
9.7	Problems	588
10	Nonequilibrium (classical)	602
10.1	Fundamental questions of (nonequilibrium) statistical mechanics	607
10.2	Langevin theory	609

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

10.3 Boltzmann kinetic theory	623
10.4 Stochastic processes	632
10.5 Field theory I: zero dimensional theories	643
10.6 Field theory II: higher dimensions	654
10.7 Field theory III: applications	665
10.8 Summary and Outlook	684
10.9 Problems	684
11 Nonequilibrium (quantum)	693
11.1 Prelude: Quantum master equation	695
11.2 Keldysh formalism: basics	700
11.3 Particle coupled to an environment	716
11.4 Fermion Keldysh theory (a list of changes)	720
11.5 Kinetic equation	723
11.6 A mesoscopic application	729
11.7 Full counting statistics	745
11.8 Summary and outlook	753
11.9 Problems	753
<i>Index</i>	766

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

Preface

In the past few decades, the field of quantum condensed matter physics has seen rapid and, at times, almost revolutionary development. Undoubtedly, the success of the field owes much to ground-breaking advances in experiment: already the controlled fabrication of phase coherent electron devices on the nanoscale is commonplace (if not yet routine), while the realization of ultra-cold atomic gases presents a new arena in which to explore strong interaction and condensation phenomena in Fermi and Bose systems. These, along with many other examples, have opened entirely new perspectives on the quantum physics of many-particle systems. Yet, important as it is, experimental progress alone does not, perhaps, fully explain the appeal of modern condensed matter physics. Indeed, in concert with these experimental developments, there has been a “quiet revolution” in condensed matter theory, which has seen phenomena in seemingly quite different systems united by common physical mechanisms. This relentless “unification” of condensed matter theory, which has drawn increasingly on the language of low-energy quantum field theory, betrays the astonishing degree of *universality*, not fully appreciated in the early literature.

On a truly microscopic level, all forms of quantum matter can be formulated as a many-body Hamiltonian encoding the fundamental interactions of the constituent particles. However, in contrast with many other areas of physics, in practically all cases of interest in condensed matter the structure of this operator conveys as much information about the properties of the system as, say, the knowledge of the basic chemical constituents tells us about the behavior of a living organism! Rather, in the condensed matter environment, it has been a long-standing tenet that the degrees of freedom relevant to the low-energy properties of a system are very often not the microscopic. Although, in earlier times, the passage between the microscopic degrees of freedom and the relevant low-energy degrees of freedom has remained more or less transparent, in recent years this situation has changed profoundly. It is a hallmark of many “deep” problems of modern condensed matter physics that the connection between the two levels involves complex and, at times, even controversial mappings. To understand why, it is helpful to place these ideas on a firmer footing.

Historically, the development of modern condensed matter physics has, to a large extent, hinged on the “unreasonable” success and “notorious” failures of *non-interacting* theories. The apparent impotency of interactions observed in a wide range of physical systems can be attributed to a deep and far-reaching principle of *adiabatic continuity*: the

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

quantum numbers that characterize a many-body system are determined by fundamental symmetries (translation, rotation, particle exchange, etc.). Providing that the integrity of the symmetries is maintained, the elementary “quasi-particle” excitations of an interacting system can be usually traced back “adiabatically” to those of the bare particle excitations present in the non-interacting system. Formally, one can say that the radius of convergence of perturbation theory extends beyond the region in which the perturbation is small. For example, this *quasi-particle correspondence*, embodied in Landau’s Fermi-liquid theory, has provided a reliable platform for the investigation of the wide range of Fermi systems from conventional metals to $^3\text{helium}$ fluids and cold atomic Fermi gases.

However, being contingent on symmetry, the principle of adiabatic continuity and, with it, the quasi-particle correspondence, must be abandoned at a *phase transition*. Here, interactions typically effect a substantial rearrangement of the many-body ground state. In the symmetry-broken phase, a system may – and frequently does – exhibit elementary excitations very different from those of the parent non-interacting phase. These elementary excitations may be classified as new species of quasi-particle with their own characteristic quantum numbers, or they may represent a new kind of excitation – a *collective mode* – engaging the cooperative motion of many bare particles. Many familiar examples fall into this category: when ions or electrons condense from a liquid into a solid phase, translational symmetry is broken and the elementary excitations – phonons – involve the motion of many individual bare particles. Less mundane, at certain field strengths, the effective low-energy degrees of freedom of a two-dimensional electron gas subject to a magnetic field (the quantum Hall system) appear as quasi-particles carrying a rational *fraction* (!) of the elementary electron charge – an effect manifestly non-perturbative in character.

This reorganization lends itself to a hierarchical perspective of condensed matter already familiar in the realm of particle physics. Each phase of matter is associated with a unique “non-interacting” reference state with its own characteristic quasi-particle excitations – a product only of the fundamental symmetries that classify the phase. While one stays within a given phase, one may draw on the principle of continuity to infer the influence of interactions. Yet this hierarchical picture delivers two profound implications. Firstly, within the quasi-particle framework, the underlying “bare” or elementary particles remain invisible (witness the fractionally charged quasi-particle excitations of the fractional quantum Hall fluid!). (To quote from P. W. Anderson’s now famous article “More is different,” (*Science* **177** (1972), 393–6), “the ability to reduce everything to simple fundamental laws does not imply the ability to start from those laws and reconstruct the universe.”) Secondly, while the capacity to conceive of new types of interaction is almost unbounded (arguably the most attractive feature of the condensed matter environment!), the freedom to identify non-interacting or free theories is strongly limited, constrained by the space of fundamental symmetries. When this is combined with the principle of continuity, the origin of the observed “universality” in condensed matter is revealed. Although the principles of adiabatic continuity, universality, and the importance of symmetries have been anticipated and emphasized long ago by visionary theorists, it is perhaps not until relatively recently that their mainstream consequences have become visible.

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

How can these concepts be embedded into a theoretical framework? At first sight, the many-body problem seems overwhelmingly daunting. In a typical system, there exist some 10^{23} particles interacting strongly with their neighbors. Monitoring the collective dynamics, even in a classical system, is evidently a hopeless enterprise. Yet, from our discussion above, it is clear that, by focussing on the coordinates of the collective degrees of freedom, one may develop a manageable theory involving only a restricted set of excitations. The success of quantum field theory in describing low-energy theories of particle physics as a successive hierarchy of broken symmetries makes its application in the present context quite natural. As well as presenting a convenient and efficient microscopic formulation of the many-body problem, the quantum field theory description provides a vehicle to systematically identify, isolate, and develop a low-energy theory of the collective field. Moreover, when cast as a field integral, the quantum field theory affords a classification of interacting systems into a small number of universality classes defined by their fundamental symmetries (a phenomenon not confined by the boundaries of condensed matter – many concepts originally developed in medium- or high-energy physics afford a seamless application in condensed matter). This phenomenon has triggered a massive trend of unification in modern theoretical physics. Indeed, by now, several sub-fields of theoretical physics have emerged (such as conformal field theory, random matrix theory, etc.) that define themselves not so much through any specific application as by a certain conceptual or methodological framework.

In deference to the importance attached to the subject, in recent years a number of texts have been written on the subject of quantum field theory within condensed matter. It is, therefore, pertinent for a reader to question the motivation for the present text. Firstly, the principal role of this text is as a primer aimed at elevating graduate students to a level where they can engage in independent research. Secondly, while the discussion of conceptual aspects takes priority over the exposure to the gamut of condensed matter applications, we have endeavored to keep the text firmly rooted in practical experimental application. Thirdly, as well as routine exercises, the present text includes extended problems which are designed to provide a bridge from formal manipulations to research-oriented thinking. Indeed, in this context, readers may note that some of the “answered” problems are deliberately designed to challenge: it is, after all, important to develop a certain degree of *intuitive* understanding of formal structures and, sadly, this can be acquired only by persistent and, at times, even frustrating training!

With this background, let us now discuss in more detail the organization of the text. To prepare for the discussion of field theory and functional integral techniques we begin in Chapter 1 by introducing the notion of a classical and a quantum field. Here we focus on the problem of lattice vibrations in the discrete harmonic chain, and its “ancestor” in the problem of classical and quantum electrodynamics. The development of field integral methods for the many-body system relies on the formulation of quantum mechanical theories in the framework of the second quantization. In Chapter 2 we present a formal and detailed introduction to the general methodology. To assimilate this technique, and motivate some of the examples discussed later in the text, a number of separate and substantial applications are explored in this chapter. In the first of these, we present (in second-quantized form) a somewhat cursory survey of the classification of metals and insulators, identifying a

canonical set of model Hamiltonians, some of which form source material for later chapters. In the case of the one-dimensional system, we will show how the spectrum of elementary collective excitations can be inferred using purely operator methods within the framework of the bosonization scheme. Finally, to close the chapter, we will discuss the application of the second quantization to the low-energy dynamics of quantum mechanical spin systems. As a final basic ingredient in the development of the quantum field theory, in Chapter 3 we introduce the Feynman path integral for the single-particle system. As well as representing a prototype for higher-dimensional field theories, the path integral method provides a valuable and recurring computational tool. This being so, we have included in this chapter a pedagogical discussion of a number of rich and instructive applications which range from the canonical example of a particle confined to a single or double quantum well, to the tunneling of extended objects (quantum fields), quantum dissipation, and the path integral formulation of spin.

Having accumulated all of the necessary background, in Chapter 4 we turn to the formulation and development of the field integral of the quantum many-particle system. Beginning with a discussion of coherent states for Fermi and Bose systems, we develop the many-body path integral from first principles. Although the emphasis in the present text is on the field integral formulation, the majority of early and seminal works in the many-body literature were developed in the framework of diagrammatic perturbation theory. To make contact with this important class of approximation schemes, in Chapter 5 we explore the way diagrammatic perturbation series expansions can be developed systematically from the field integral. Employing the ϕ^4 -theory as a canonical example, we describe how to explore the properties of a system in a high order of perturbation theory around a known reference state. To cement these ideas, we apply these techniques to the problem of the weakly interacting electron gas.

Although the field integral formulation provides a convenient means to organize perturbative approximation schemes as a diagrammatic series expansion, its real power lies in its ability to identify non-trivial reference ground states, or “mean-fields,” and to provide a framework in which low-energy theories of collective excitations can be developed. In Chapter 6, a fusion of perturbative and mean-field methods is used to develop analytical machinery powerful enough to address a spectrum of rich applications ranging from metallic magnetism and superconductivity to superfluidity. To bridge the gap between the (often abstract) formalism of the field integral, and the arena of practical application, it is necessary to infer the behavior of correlation functions. Beginning with a brief survey of concepts and techniques of experimental condensed matter physics, in Chapter 7 we highlight the importance of correlation functions and explore their connection with the theoretical formalism developed in previous chapters. In particular, we discuss how the response of many-body systems to various types of electromagnetic perturbation can be described in terms of correlation functions and how these functions can be computed by field theoretical means.

Although the field integral is usually simple to formulate, its properties are not always easy to uncover. Amongst the armory of tools available to the theorist, perhaps the most adaptable and versatile is the method of the renormalization group. Motivating

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

our discussion with two introductory examples drawn from a classical and a quantum theory, in Chapter 8 we become acquainted with the renormalization group method as a concept whereby nonlinear theories can be analyzed beyond the level of plain perturbation theory. With this background, we then proceed to discuss renormalization methods in more rigorous and general terms, introducing the notion of scaling, dimensional analysis, and the connection to the general theory of phase transitions and critical phenomena. To conclude this chapter, we visit a number of concrete implementations of the renormalization group scheme introduced and exemplified on a number of canonical applications.

In Chapter 9, we turn our attention to low-energy theories with non-trivial forms of long-range order. Specifically, we will learn how to detect and classify topologically non-trivial structures, and to understand their physical consequences. Specifically, we explore the impact of topological terms (i.e. θ -terms, Wess–Zumino terms, and Chern–Simons terms) on the behavior of low-energy field theories solely through the topology of the underlying field configurations. Applications discussed in this chapter include persistent currents, ’t Hooft’s θ -vacua, quantum spin chains, and the quantum Hall effects.

So far, our development of field theoretic methodologies has been tailored to the consideration of single-particle quantum systems, or many-body systems in thermal equilibrium. However, studies of classical nonequilibrium systems have a long and illustrious history, dating back to the earliest studies of thermodynamics, and these days include a range of applications from soft matter physics to population dynamics and ecology. At the same time, the control afforded by modern mesoscopic semiconducting and metallic devices, quantum optics, as well as ultracold atom physics now allow controlled access to quantum systems driven far from equilibrium. For such systems, traditional quantum field theoretical methodologies are inappropriate.

Starting with the foundations of non-equilibrium statistical mechanics, from simple one-step processes, to reaction–diffusion type systems, in Chapter 10 we begin by developing Langevin and Fokker–Planck theory, from which we establish classical Boltzmann transport equations. We then show how these techniques can be formulated in the language of the functional integral developing the Doi–Peliti and Martin–Siggia–Rose techniques. We conclude our discussion with applications to nonequilibrium phase transitions and driven lattice gases. These studies of the classical nonequilibrium system provide a platform to explore the quantum system. In Chapter 11, we develop the Keldysh approach to quantum non-equilibrium systems based, again, on the functional integral technique. In particular, we emphasize and exploit the close connections to classical nonequilibrium field theory, and present applications to problems from the arena of quantum transport.

To focus and limit our discussion, we have endeavored to distill material considered “essential” from the “merely interesting” or “background.” To formally acknowledge and identify this classification, we have frequently included reference to material which we believe may be of interest to the reader in placing the discussion in context, but which can be skipped without losing the essential thread of the text. These intermissions are signaled in the text as “Info” blocks.

At the end of each chapter, we have collected a number of pedagogical and instructive problems. In some cases, the problems expand on some aspect of the main text requiring only

Cambridge University Press

978-0-521-76975-4 - Condensed Matter Field Theory, Second Edition

Alexander Altland and Ben Simons

Frontmatter

[More information](#)

an extension, or straightforward generalization, of a concept raised in the chapter. In other cases, the problems rather complement the main text, visiting fresh applications of the same qualitative material. Such problems take the form of case studies in which both the theory and the setting chart new territory. The latter provide a vehicle to introduce some core areas of physics not encountered in the main text, and allow the reader to assess the degree to which the ideas in the chapter have been assimilated. With both types of questions to make the problems more inclusive and useful as a reference, we have included (sometimes abridged, and sometimes lengthy) answers. In this context, Section 6.5 assumes a somewhat special role: the problem of phase coherent electron transport in weakly disordered media provides a number of profoundly important problems of great theoretical and practical significance. In preparing this section, it became apparent that the quantum disorder problem presents an ideal environment in which many of the theoretical concepts introduced in the previous chapters can be practiced and applied – to wit diagrammatic perturbation theory and series expansions, mean-field theory and collective mode expansions, correlation functions and linear response, and topology. We have therefore organized this material in the form of an extended problem set in Chapter 6.

This concludes our introduction to the text. Throughout, we have tried to limit the range of physical applications to examples which are rooted in experimental fact. We have resisted the temptation to venture into more speculative areas of theoretical condensed matter at the expense of excluding many modern and more-circumspect ideas which pervade the condensed matter literature. Moreover, since the applications are intended to help motivate and support the field theoretical techniques, their discussion is, at times, necessarily superficial. (For example, the hundreds pages of text in this volume could have been invested in their entirety in the subject of superconductivity!) Therefore, where appropriate, we have tried to direct interested readers to the more specialist literature.

In closing, we would like to express our gratitude to Jakob Müller-Hill, Tobias Micklitz, Jan Müller, Natalja Strelkova, Franjo-Frankopan Velic, Andrea Wolff, and Markus Zowislok for their invaluable assistance in the proofreading of the text. Moreover, we would also like to thank Julia Meyer for her help in drafting problems. Finally we would like to acknowledge Sasha Abanov for his advice and guidance in the drafting of the chapter on Topology.

As well as including additional material on the formulation of functional field integral methods to classical and quantum nonequilibrium physics in Chapters 10 and 11, in preparing the second edition of the text, we have endeavored to remove some of the typographical errors that crept into the first edition. Although it seems inevitable that some errors will still have escaped identification, it is clear that many many more would have been missed were it not for the vigilance of many friends and colleagues. In this context, we would particularly like to acknowledge the input of Piet Brouwer, Christoph Bruder, Chung-Pin Chou, Jan von Delft, Karin Everschor, Andrej Fischer, Alex Gezerlis, Sven Gnutzmann, Colin Kiegel, Tobias Lück, Patrick Neven, Achim Rosch, Max Schäfer, Matthias Sitte, Nobuhiko Taniguchi, and Matthias Vojta.