

ENCYCLOPEDIA OF MATHEMATICS AND ITS APPLICATIONS

Asymptotics and Mellin-Barnes Integrals

R. B. PARIS

D. KAMINSKI



CAMBRIDGE
UNIVERSITY PRESS

PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS
The Edinburgh Building, Cambridge CB2 2RU, UK
40 West 20th Street, New York, NY 10011-4211, USA
10 Stamford Road, Oakleigh, VIC 3166, Australia
Ruiz de Alarcón 13, 28014 Madrid, Spain
Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

© Cambridge University Press 2001

This book is in copyright. Subject to statutory exception
and to the provisions of relevant collective licensing agreements,
no reproduction of any part may take place without
the written permission of Cambridge University Press.

First published 2001

Printed in the United Kingdom at the University Press, Cambridge

Typeface Times Roman 10/13 pt. *System* L^AT_EX 2_ε [KW]

A catalogue record for this book is available from the British Library.

Library of Congress Cataloging in Publication Data

Paris, R.B.

Asymptotics and Mellin-Barnes integrals/R. B. Paris, D. Kaminski.

p. cm. – (Encyclopedia of mathematics and its applications; v. 85)

Includes bibliographical references and index

ISBN 0-521-79001-8

1. Mellin transform. 2. Asymptotic expansions. I. Kaminski, D. (David), 1960– II. Title
III. Series.

QA432.P34 2001

515'.723 – dc21

00-067492

ISBN 0 521 79001 8 hardback

Contents

Preface		<i>page</i> xiii
1	Introduction	1
1.1	Introduction to Asymptotics	1
1.1.1	Order Relations	1
1.1.2	Asymptotic Expansions	4
1.1.3	Other Expansions	15
1.2	Biographies of Mellin and Barnes	25
2	Fundamental Results	29
2.1	The Gamma Function $\Gamma(z)$	29
2.1.1	The Asymptotic Expansion of $\Gamma(z)$	29
2.1.2	The Stirling Coefficients	32
2.1.3	Bounds for $\Gamma(z)$	33
2.2	Expansion of Quotients of Gamma Functions	35
2.2.1	Inverse Factorial Expansions	36
2.2.2	A Recursion Formula when $\alpha_r = \beta_r = 1$	40
2.2.3	Examples	44
2.2.4	An Algebraic Method for the Determination of the A_j	46
2.2.5	Special Cases	49
2.3	The Asymptotic Expansion of Integral Functions	55
2.3.1	An Example	58
2.4	Convergence of Mellin-Barnes Integrals	63
2.5	Order Estimates for Remainder Integrals	69
2.5.1	An Example	69
2.5.2	Lemmas	71

3	Properties of Mellin Transforms	79
3.1	Basic Properties	79
3.1.1	Definition	79
3.1.2	Translational and Differential Properties	81
3.1.3	The Parseval Formula	82
3.2	Analytic Properties	85
3.3	Inverse Mellin Transforms	89
3.3.1	Integrals Connected with e^{-z}	89
3.3.2	Some Standard Integrals	91
3.3.3	Discontinuous Integrals	93
3.3.4	Gamma-Function Integrals	96
3.3.5	Ramanujan-Type Integrals	99
3.3.6	Barnes' Lemmas	103
3.4	Mellin-Barnes Integral Representations	106
3.4.1	The Confluent Hypergeometric Functions	107
3.4.2	The Gauss Hypergeometric Function	110
3.4.3	Some Special Functions	112
4	Applications of Mellin Transforms	117
4.1	Transformation of Series	117
4.1.1	The Mellin Transform Method	117
4.1.2	The Poisson-Jacobi Formula	120
4.2	Examples	122
4.2.1	An Infinite Series	122
4.2.2	A Smoothed Dirichlet Series	125
4.2.3	A Finite Sum	128
4.3	Number-Theoretic Examples	133
4.3.1	A Harmonic Sum	133
4.3.2	Euler's Product	136
4.3.3	Ramanujan's Function	137
4.3.4	Some Other Number-Theoretic Sums	141
4.4	Solution of Differential Equations	146
4.4.1	Potential Problems in Wedge-Shaped Regions	146
4.4.2	Ordinary Differential Equations	149
4.4.3	Inverse Mellin Transform Solutions	152
4.5	Solution of Integral Equations	156
4.5.1	Kernels of the Form $k(xt)$	157
4.5.2	Kernels of the Form $k(x/t)$	160
4.6	Solution of Difference Equations	164
4.6.1	Solution by Mellin Transforms	166
4.6.2	The Hypergeometric Difference Equation	167
4.6.3	Solution of the Inhomogeneous First-Order Equation	172

4.7	Convergent Inverse Factorial Series	174
5	Asymptotic Expansions	179
5.1	Algebraic Asymptotic Expansions	179
5.1.1	The Exponential Integral $E_1(z)$	180
5.1.2	The Parabolic Cylinder Function $D_\nu(z)$	182
5.1.3	A Bessel Function Integral	184
5.1.4	The Mittag-Leffler Function $\mathcal{E}_a(z)$	186
5.2	Remainder Integrals	190
5.2.1	Error Bounds	190
5.2.2	Numerical Evaluation	195
5.3	Saddle-Point Approximation of Integrals	197
5.3.1	An Integral Due to Heading and Whipple	198
5.3.2	The Bessel Function $J_n(nx)$	200
5.3.3	A Gauss Hypergeometric Function	202
5.4	Exponential Asymptotic Expansions	205
5.4.1	The Exponential Integral $E_1(z)$	206
5.4.2	The Bessel Function $J_\nu(z)$	208
5.4.3	The Parabolic Cylinder Function $D_\nu(z)$	210
5.4.4	An Infinite Sum	213
5.5	Faxén's Integral	216
5.6	Integrals with a 'Contour Barrier'	220
5.6.1	An Illustrative Example	221
5.6.2	An Integral Involving a Bessel Function	225
5.6.3	Example in §4.2.3 Revisited	229
6	The Stokes Phenomenon and Hyperasymptotics	234
6.1	The Stokes Phenomenon	234
6.1.1	A Qualitative Description	234
6.1.2	The Modified Bessel Function $K_\nu(z)$	236
6.2	Mellin-Barnes Theory	240
6.2.1	Exponentially-Improved Expansion	241
6.2.2	Estimates for $ R_{N,M}(z) $	244
6.2.3	The Stokes Multiplier	246
6.2.4	The Stokes Multiplier for a High-Order Differential Equation	249
6.2.5	Numerical Examples	256
6.2.6	Asymptotics of the Terminant $T_\nu(z)$	259
6.3	Hyperasymptotics	265
6.3.1	Mellin-Barnes Theory of Hyperasymptotics	266
6.3.2	Optimal Truncation Schemes	271

6.3.3	A Numerical Example	277
6.4	Exponentially-Improved Asymptotics for $\Gamma(z)$	279
6.4.1	Origin of the Exponentially Small Terms	280
6.4.2	The Expansion of $\Omega(z)$	282
6.4.3	A Numerical Example	286
7	Multiple Mellin-Barnes Integrals	289
7.1	Some Double Integrals	289
7.2	Residues and Double Integrals	295
7.3	Laplace-Type Double Integrals	299
7.3.1	Mellin-Barnes Integral Representation	303
7.3.2	The Newton Diagram	304
7.4	Asymptotics of $I(\lambda)$	305
7.4.1	Two Internal Points	306
7.4.2	Three and More Internal Points	311
7.4.3	Other Double Integrals	322
7.5	Geometric Content	323
7.5.1	Remoteness	324
7.5.2	Asymptotic Scales	326
7.6	Laplace-Type Integrals of Higher Dimension	328
7.6.1	Representation of Treble Integrals	328
7.6.2	Asymptotics with One Internal Point	329
7.6.3	Asymptotics with Two Internal Points	332
7.6.4	Other Considerations for Treble Integrals	339
7.7	Numerical Examples	340
8	Application to Some Special Functions	352
8.1	Asymptotics of the Euler-Jacobi Series	353
8.1.1	Introduction	353
8.1.2	The Saddle Point Approach	356
8.1.3	The Asymptotics of $S_p(a)$	361
8.1.4	Mellin-Barnes Integral Approach	365
8.1.5	The Asymptotic Expansion of $S_p(a)$ for $a \rightarrow 0+$	368
8.1.6	The Coefficients A_k	371
8.1.7	The Stokes Phenomenon for $p \simeq 2$	374
8.2	An Asymptotic Formula for $\zeta(\frac{1}{2} + it)$	380
8.2.1	Introduction	380
8.2.2	An Expansion for $\zeta(s)$	382
8.2.3	An Exponentially-Smoothed Gram-Type Expansion	384
8.2.4	A Riemann-Siegel-Type Expansion	385
8.3	The Asymptotics of Pearcey's Integral	389
8.3.1	Introduction	389

8.3.2	A Mellin-Barnes Integral Representation	391
8.3.3	Asymptotics of $\hat{P}(x, y)$ for $ x \rightarrow \infty$	393
8.3.4	Asymptotics of $\hat{P}(x, y)$ for $ y \rightarrow \infty$	398
8.3.5	Asymptotics of $P(X, Y)$ for Real X, Y	400
8.3.6	Generalisations of $P(X, Y)$	401
Appendix		404
References		409
Index		419

Introduction

1.1 Introduction to Asymptotics

Before venturing into our examination of Mellin-Barnes integrals, we present an overview of some of the basic definitions and ideas found in asymptotic analysis. The treatment provided here is not intended to be comprehensive, and several high quality references exist which can provide a more complete treatment than is given here: in particular, we recommend the tracts by Olver (1974), Bleistein & Handelsman (1975) and Wong (1989) as particularly good treatments of asymptotic analysis, each with their own strengths.[†]

1.1.1 Order Relations

Let us begin our survey by defining the *Landau symbols* O and o and the notion of asymptotic equality.

Let f and g be two functions defined in a neighbourhood of x_0 . We say that $f(x) = O(g(x))$ as $x \rightarrow x_0$ if there is a constant M for which

$$|f(x)| \leq M |g(x)|$$

for x sufficiently close to x_0 . The constant M depends only on how close to x_0 we wish the bound to hold. The notation $O(g)$ is read as ‘big-oh of g ’, and the constant M , which is often not explicitly calculated, is termed the *implied constant*.

In a similar fashion, we define $f(x) = o(g(x))$ as $x \rightarrow x_0$ to mean that

$$|f(x)/g(x)| \rightarrow 0$$

[†] Olver provides a good balance between techniques used in both integrals and differential equations; Bleistein & Handelsman present a relatively unified treatment of integrals through the use of Mellin convolutions; and Wong develops the theory and application of (Schwartz) distributions in the setting of developing expansions of integrals.

as $x \rightarrow x_0$, subject to the proviso that $g(x)$ be nonzero in a neighbourhood of x_0 . The expression $o(g)$ is read as ‘little-oh of g ’, and from the preceding definition, it is immediate that $f = o(g)$ implies that $f = O(g)$ (merely take the implied constant to be any (arbitrarily small) positive number).

The last primitive asymptotic notion required is that of asymptotic equality. We write

$$f(x) \sim g(x)$$

as $x \rightarrow x_0$ to mean that

$$\lim_{x \rightarrow x_0} \frac{f(x)}{g(x)} = 1,$$

provided, of course, that g is nonzero sufficiently close to x_0 . The tilde here is read ‘is asymptotically equal to’. An equivalent formulation of asymptotic equality is readily available: for $x \rightarrow x_0$,

$$f(x) \sim g(x) \quad \text{iff} \quad f(x) = g(x)\{1 + o(1)\}.$$

Example 1. The function $\log x$ satisfies the order relation $\log x = O(x - 1)$ as $x \rightarrow \infty$, since the ratio $(\log x)/(x - 1)$ is bounded for all large x . In fact, it is also true that $\log x = o(x - 1)$ for large x , and for $x \rightarrow 1$, $\log x \sim x - 1$.

Example 2. Stirling’s formula is a well-known asymptotic equality. For large n , we have

$$n! \sim (2\pi)^{\frac{1}{2}} e^{-n} n^{n+\frac{1}{2}}.$$

This result follows from the asymptotic expansion of the gamma function, a result carefully developed in §2.1.

Example 3. The celebrated Prime Number Theorem is an asymptotic equality. If we denote by $\pi(x)$ the number of primes less than or equal to x , then for large positive x we have the well-known result

$$\pi(x) \sim \frac{x}{\log x}.$$

With the aid of Gauss’ logarithmic integral,[†]

$$\text{li}(x) = \int_2^x \frac{dt}{\log t}$$

we also have the somewhat more accurate form

$$\pi(x) \sim \text{li}(x) \quad (x \rightarrow \infty).$$

[†] We note here that $\text{li}(x)$ is also used to denote the same integral, but taken over the interval $(0, x)$, with $x > 1$. With this larger interval, the integral is a Cauchy principal value integral. The notation in this example appears to be in use by some number theorists, and is also sometimes written $\text{Li}(x)$.

That both forms hold can be seen from a simple integration by parts:

$$\text{li}(x) = \frac{x}{\log x} - \frac{2}{\log 2} + \int_2^x \frac{dt}{(\log t)^2}.$$

An application of l'Hôpital's rule reveals that the resulting integral on the right-hand side is $o(x/\log x)$, from which the $x/\log x$ form of the Prime Number Theorem follows.

A number of useful relationships exist for manipulating the Landau symbols. The following selections are all easily obtained from the above definitions, and are not established here:

$$\begin{array}{ll} \text{(a)} O(O(f)) = O(f) & \text{(e)} O(f) + O(f) = O(f) \\ \text{(b)} o(o(f)) = o(f) & \text{(f)} o(f) + o(f) = o(f) \\ \text{(c)} O(fg) = O(f) \cdot O(g) & \text{(g)} o(f) + O(f) = O(f) \\ \text{(d)} O(f) \cdot o(g) = o(fg) & \text{(h)} O(o(f)) = o(O(f)) = o(f). \end{array} \quad (1.1.1)$$

It is easy to deduce linearity of Landau symbols using these properties, and it is a simple matter to establish asymptotic equality as an equivalence relation. In the transition to calculus, however, some difficulties surface.

A moment's consideration reveals that differentiation is, in general, often badly behaved in the sense that if $f = O(g)$, then it does not necessarily follow that $f' = O(g')$, as the example $f(x) = x + \sin e^x$ aptly illustrates: for large, real x , we have $f = O(x)$, but the derivative of f is not bounded (i.e., not $O(1)$).

The situation for integration is a good deal better. It is possible to formulate many results concerning integrals of order estimates, but we content ourselves with just two.

Example 4. For functions f and g of a real variable x satisfying $f = O(g)$ as $x \rightarrow x_0$ on the real line, we have

$$\int_{x_0}^x f(t) dt = O\left(\int_{x_0}^x |g(t)| dt\right) \quad (x \rightarrow x_0).$$

A proof can be fashioned along the following lines: for $f(t) = O(g(t))$, let M be the implied constant so that $|f(t)| \leq M |g(t)|$ for t sufficiently close to x_0 , say $|t - x_0| \leq \eta$. (For $x_0 = \infty$, a suitable interval would be $t \geq N$ for some large positive N .) Then

$$-M |g(t)| \leq f(t) \leq M |g(t)| \quad (|t - x_0| \leq \eta),$$

whence the result follows upon integration.

Example 5. If f is an integrable function of a real variable x , and $f(x) \sim x^\nu$, $\text{Re}(\nu) < -1$ as $x \rightarrow \infty$, then

$$\int_x^\infty f(t) dt \sim -\frac{x^{\nu+1}}{\nu+1} \quad (x \rightarrow \infty).$$

A proof of this claim follows from $f(x) = x^\nu\{1 + \psi(x)\}$ where $\psi(x) = o(1)$ as $x \rightarrow \infty$, for then

$$\int_x^\infty f(t) dt = -\frac{x^{\nu+1}}{\nu+1} + \int_x^\infty t^\nu \psi(t) dt.$$

But $\psi(t) = o(1)$ implies that for $\epsilon > 0$ arbitrarily small, there is an $x_0 > 0$ for which $|\psi(t)| < \epsilon$ whenever $t > x_0$. Thus, the remaining integral may be bounded as

$$\left| \int_x^\infty t^\nu \psi(t) dt \right| < \epsilon \int_x^\infty |t^\nu| dt \quad (x > x_0).$$

Accordingly, we find

$$\int_x^\infty f(t) dt = -\frac{x^{\nu+1}}{\nu+1} + o\left(\frac{x^{\nu+1}}{\nu+1}\right) = -\frac{x^{\nu+1}}{\nu+1}\{1 + o(1)\},$$

from which the asymptotic equality is immediate. $\square$

It is in the complex plane that we find differentiation of order estimates becomes better behaved. This is due, in part, to the fact that the Cauchy integral theorem allows us to represent holomorphic functions as integrals which, as we have noted, are better behaved in the setting of Landau symbols. A standard result in this direction is the following:

Lemma 1.1. *Let f be holomorphic in a region containing the closed annular sector $\mathcal{S} = \{z : \alpha \leq \arg(z - z_0) \leq \beta, |z - z_0| \geq R \geq 0\}$, and suppose $f(z) = O(z^\nu)$ (resp. $f(z) = o(z^\nu)$) as $z \rightarrow \infty$ in the sector, for fixed real ν . Then $f^{(n)}(z) = O(z^{\nu-n})$ (resp. $f^{(n)} = o(z^{\nu-n})$) as $z \rightarrow \infty$ in any closed annular sector properly interior to $\mathcal{S}$ with common vertex z_0 .*

The proof of this result follows from the Cauchy integral formula for $f^{(n)}$, and is available in Olver (1974, p. 9).

1.1.2 Asymptotic Expansions

Let a sequence of continuous functions $\{\phi_n\}$, $n = 0, 1, 2, \dots$, be defined on some domain, and let x_0 be a (possibly infinite) limit point of this domain. The sequence $\{\phi_n\}$ is termed an *asymptotic scale* if it happens that $\phi_{n+1}(x) = o(\phi_n(x))$ as $x \rightarrow x_0$, for every n . If f is some continuous function on the common domain of the asymptotic scale, then by an (infinite) *asymptotic expansion* of f with respect to the asymptotic scale $\{\phi_n\}$ is meant the formal series $\sum_{n=0}^\infty a_n \phi_n(x)$, provided the coefficients a_n , independent of x , are chosen so that for any nonnegative integer N ,

$$f(x) = \sum_{n=0}^N a_n \phi_n(x) + O(\phi_{N+1}(x)) \quad (x \rightarrow x_0). \quad (1.1.2)$$

In this case we write

$$f(x) \sim \sum_{n=0}^{\infty} a_n \phi_n(x) \quad (x \rightarrow x_0).$$

Such a formal series is uniquely determined in view of the fact that the coefficients a_n can be computed from

$$a_N = \lim_{x \rightarrow x_0} \frac{1}{\phi_N(x)} \left\{ f(x) - \sum_{n=0}^{N-1} a_n \phi_n(x) \right\} \quad (N = 0, 1, 2, \dots).$$

The formal series so obtained is also referred to as an asymptotic expansion of *Poincaré type*, or an asymptotic expansion in the sense of Poincaré or, more simply, a Poincaré expansion. Examples of asymptotic scales and asymptotic expansions built with them are easy to come by. The most commonplace is the asymptotic power series: an *asymptotic power series* is a formal series

$$\sum_{n=0}^{\infty} a_n (x - x_0)^{v_n},$$

where the appropriate asymptotic scale is the sequence $\{(x - x_0)^{v_n}\}$, $n = 0, 1, 2, \dots$, and the v_n are constants for which $(x - x_0)^{v_{n+1}} = o((x - x_0)^{v_n})$ as $x \rightarrow x_0$. Any convergent Taylor series expansion of an analytic function f serves as an example of an asymptotic power series, with x_0 a point in the domain of analyticity of f , $v_n = n$ for any nonnegative integer n , and the coefficients in the expansion are the familiar Taylor coefficients $a_n = f^{(n)}(x_0)/n!$.

Asymptotic expansions, however, need not be convergent, as the next two examples illustrate.

Example 1. WATSON'S LEMMA. A well-known result of Laplace transform theory is that the Laplace transform of a piecewise continuous function on the interval $[0, +\infty)$ is $o(1)$ as the transform variable grows without bound. By imposing more structure on the small parameter behaviour of the function being transformed, a good deal more can be said about the growth at infinity of the transform.

Lemma 1.2. Let $g(t)$ be an integrable function of the variable $t > 0$ with asymptotic expansion

$$g(t) \sim \sum_{n=0}^{\infty} a_n t^{(n+\lambda-\mu)/\mu} \quad (t \rightarrow 0+)$$

for some constants $\lambda > 0$, $\mu > 0$. Then, provided the integral converges for all sufficiently large x , the Laplace transform of g , $\mathcal{L}[g; x]$, has the asymptotic behaviour

$$\mathcal{L}[g; x] \equiv \int_0^{\infty} e^{-xt} g(t) dt \sim \sum_{n=0}^{\infty} \Gamma\left(\frac{n+\lambda}{\mu}\right) \frac{a_n}{x^{(n+\lambda)/\mu}} \quad (x \rightarrow \infty).$$

Proof. To see this, let us put, for positive integer N and $t > 0$,

$$g_N(t) = g(t) - \sum_{n=0}^{N-1} a_n t^{(n+\lambda-\mu)/\mu}$$

so that the Laplace transform has a finite expansion with remainder given by

$$\mathcal{L}[g; x] = \sum_{n=0}^{N-1} \Gamma\left(\frac{n+\lambda}{\mu}\right) \frac{a_n}{x^{(n+\lambda)/\mu}} + \int_0^\infty e^{-xt} g_N(t) dt. \quad (1.1.3)$$

Since $g_N(t) = O(t^{(N+\lambda-\mu)/\mu})$, there are constants K_N and t_N for which

$$|g_N(t)| \leq K_N t^{(N+\lambda-\mu)/\mu} \quad (0 < t \leq t_N).$$

Use of this in the remainder term in our finite expansion (1.1.3) allows us to write

$$\begin{aligned} \left| \int_0^{t_N} e^{-xt} g_N(t) dt \right| &\leq K_N \int_0^{t_N} e^{-xt} t^{(N+\lambda-\mu)/\mu} dt \\ &< \Gamma\left(\frac{N+\lambda}{\mu}\right) \frac{K_N}{x^{(N+\lambda)/\mu}}. \end{aligned} \quad (1.1.4)$$

By hypothesis, $\mathcal{L}[g; x]$ exists for all sufficiently large x , so the Laplace transform of g_N must also exist for all sufficiently large x , by virtue of (1.1.3). Let X be such that $\mathcal{L}[g_N; x]$ exists for all $x \geq X$, and put

$$G_N(t) = \int_{t_N}^t e^{-Xv} g_N(v) dv.$$

The function G_N so defined is a bounded continuous function on $[t_N, \infty)$, whence the bound

$$L_N = \sup_{[t_N, \infty)} |G_N(t)|$$

exists. Then for $x > X$, we have

$$\begin{aligned} \int_{t_N}^\infty e^{-xt} g_N(t) dt &= \int_{t_N}^\infty e^{-(x-X)t} e^{-Xt} g_N(t) dt \\ &= (x - X) \int_{t_N}^\infty e^{-(x-X)t} G_N(t) dt \end{aligned}$$

after one integration by parts. After applying the uniform bound L_N to the integral that remains, we arrive at

$$\left| \int_{t_N}^\infty e^{-xt} g_N(t) dt \right| \leq (x - X) L_N \int_{t_N}^\infty e^{-(x-X)t} dt = L_N e^{-(x-X)t_N} \quad (1.1.5)$$

for $x > X$.

Together, (1.1.4) and (1.1.5) yield

$$\left| \int_0^\infty e^{-xt} g_N(t) dt \right| < \Gamma\left(\frac{N+\lambda}{\mu}\right) \frac{K_N}{x^{(N+\lambda)/\mu}} + L_N e^{-(x-X)t_N}$$

which, since $L_N e^{-(x-X)t_N}$ is $o(x^{-\nu})$ for any positive ν , establishes the asymptotic expansion for $\mathcal{L}[g; x]$. $\square$

As a simple illustration of the use of Watson's lemma, consider the Laplace transform of $(1+t)^{\frac{1}{2}}$. From the binomial theorem, we have the convergent expansion as $t \rightarrow 0$

$$(1+t)^{\frac{1}{2}} = 1 + \frac{1}{2}t - \frac{1}{8}t^2 + \sum_{n=3}^{\infty} (-)^{n-1} \frac{1 \cdot 3 \cdot 5 \cdots (2n-3)}{2^n n!} t^n.$$

Since $(1+t)^{\frac{1}{2}}$ is of algebraic growth, its Laplace transform clearly exists for $x > 0$, and Watson's lemma produces the asymptotic expansion

$$\mathcal{L}[(1+t)^{\frac{1}{2}}; x] \sim \frac{1}{x} + \frac{1}{2x^2} - \frac{1}{4x^3} + \sum_{n=3}^{\infty} (-)^{n-1} \frac{1 \cdot 3 \cdot 5 \cdots (2n-3)}{2^n x^{n+1}}$$

as $x \rightarrow \infty$. The resulting asymptotic series is divergent, since the ratio of the $(n+1)$ th to n th terms in absolute value is $(2n-1)/(2x)$ which, for fixed x , tends to ∞ with n . The reason for this divergence is a simple consequence of our applying the binomial expansion for $(1+t)^{\frac{1}{2}}$ (valid in $0 \leq t \leq 1$) in the Laplace integral beyond its interval of convergence.

Example 2. The confluent hypergeometric function† $U(1; 1; z)$ (which equals the exponential integral $e^z E_1(z)$) has the integral representation

$$U(1; 1; z) = \int_0^\infty \frac{e^{-t} dt}{t+z} \quad (1.1.6)$$

for z not a negative number or zero. In fact, it is relatively easy to show that this integral representation converges uniformly in the closed annular sector $\mathcal{S}_{\epsilon, \delta} = \{z : |z| \geq \epsilon, |\arg z| \leq \pi - \delta\}$ for every positive ϵ and every positive $\delta < \pi$. Such a demonstration can proceed along the following lines.

Put $\theta = \arg z$ for $z \in \mathcal{S}_{\epsilon, \delta}$ and observe that for any nonnegative t , $|t+z|^2 = t^2 + |z|^2 + 2|z|t \cos \theta \geq t^2 + |z|^2 - 2|z|t \cos \delta \geq |z|^2 \sin^2 \delta$. Thus, the integrand of (1.1.6) admits the simple bound

$$e^{-t}|t+z|^{-1} \leq e^{-t}|z|^{-1} \operatorname{cosec} \delta$$

whence we have, upon integrating the bound,

$$|U(1; 1; z)| \leq |z|^{-1} \operatorname{cosec} \delta$$

† An alternative notation for this function is $\Psi(1; 1; z)$.

for $z \in \mathcal{S}_{\epsilon, \delta}$. The uniform convergence of the integral follows, from which we see that $U(1; 1; z)$ is holomorphic in the z plane cut along the negative real axis.

Through repeated integration by parts, differentiating in each case the factor $(t + z)^{-k}$ appearing at each step, we arrive at

$$U(1; 1; z) = \sum_{k=1}^n (-)^{k-1} (k-1)! z^{-k} + R_n(z), \quad (1.1.7)$$

where the remainder term $R_n(z)$ is

$$R_n(z) = (-)^n n! \int_0^\infty \frac{e^{-t}}{(t+z)^{n+1}} dt. \quad (1.1.8)$$

Evidently, each term produced in the series in (1.1.7) is a term from the asymptotic scale $\{z^{-j}\}$, $j = 1, 2, \dots$, so that if we can show that for any n , $R_n(z) = O(z^{-n-1})$, we will have established the asymptotic expansion

$$U(1; 1; z) \sim \sum_{k=1}^{\infty} (-)^{k-1} (k-1)! z^{-k}, \quad (1.1.9)$$

for $z \rightarrow \infty$ in the sector $|\arg z| \leq \pi - \delta < \pi$.

To this end, we observe that the bound used in establishing the uniform convergence of the integral (1.1.6), namely $1/|t+z| \leq 1/|z| \sin \delta$, can be brought to bear on (1.1.8) to yield

$$|R_n(z)| \leq \frac{n!}{(|z| \sin \delta)^{n+1}}.$$

The expansion (1.1.9) is therefore an asymptotic expansion in the sense of Poincaré. It is, however, quite clearly a divergent series, as ratios of consecutive terms in the asymptotic series diverge to ∞ as $(n!/|z|^{n+1})/((n-1)!/|z|^n) = n/|z|$, as $n \rightarrow \infty$, irrespective of the value of z . Nevertheless, the divergent character of this asymptotic series does not detract from its computational utility. $\square$

In Tables† 1.1 and 1.2, we have gathered together computed and approximate values of $U(1; 1; z)$, with approximate values derived from the finite series approximation

$$S_n(z) = \sum_{k=1}^n (-)^{k-1} (k-1)! z^{-k},$$

obtained by truncating the asymptotic expansion (1.1.9) after n terms. It is apparent from the tables that the calibre of even modest approximations to $U(1; 1; z)$ becomes quite good once $|z|$ is of the order of 100, and is good to two or more significant digits for values of $|z|$ as small as 10. This naturally leads one to

† In Tables 1.1 and 1.2 we have adopted the convention of writing $x(y)$ in lieu of the more cumbersome $x \times 10^y$.

Table 1.1. *Computed and approximate values of $U(1; 1; z)$ for real values of z*

z	$U(1; 1; z)$	$S_5(z)$	$S_{10}(z)$
10	0.915633(−1)	0.916400(−1)	0.915456(−1)
50	0.196151(−1)	0.196151(−1)	0.196151(−1)
100	0.990194(−2)	0.990194(−2)	0.990194(−2)

Table 1.2. *Computed and approximate values of $U(1; 1; z)$ for imaginary values of z*

z	$U(1; 1; z)$
10 <i>i</i>	0.948854(−2) − 0.981910(−1) <i>i</i>
50 <i>i</i>	0.399048(−3) − 0.199841(−1) <i>i</i>
100 <i>i</i>	0.999401(−4) − 0.999800(−2) <i>i</i>
z	$S_5(z)$
10 <i>i</i>	0.940000(−2) − 0.982400(−1) <i>i</i>
50 <i>i</i>	0.399040(−3) − 0.199841(−1) <i>i</i>
100 <i>i</i>	0.999400(−4) − 0.999800(−2) <i>i</i>
z	$S_{10}(z)$
10 <i>i</i>	0.950589(−2) − 0.982083(−1) <i>i</i>
50 <i>i</i>	0.399048(−3) − 0.199841(−1) <i>i</i>
100 <i>i</i>	0.999401(−4) − 0.999800(−2) <i>i</i>

wonder how the best approximation can be obtained, in view of the utility of these finite approximations and the divergence of the full asymptotic expansion: how can we select n so that the approximation furnished by $S_n(z)$ is the best possible?

The strategy we detail here, called *optimal truncation*, is easily stated: for a fixed z , the successive terms in the asymptotic expansion will reach a minimum in absolute value, after which the terms must necessarily increase without bound given the divergent character of the full expansion; see Fig. 1.1. It is readily shown that the terms in $S_n(z)$ attain their smallest absolute value when $k \sim |z|$ (except when $|z|$ is an integer, in which case there are two equally small terms corresponding to $k = |z| - 1$ and $k = |z|$). If the full series is truncated just before this minimum modulus term is reached, then the finite series that results is the optimally truncated series, and will yield the best approximation to the original function, in the present case, $U(1; 1; z)$.

To see that this is so, observe for $U(1; 1; z)$ that for $z > 0$ the remainder in the approximation after n terms of the asymptotic series,

$$R_n(z) = U(1; 1; z) - S_n(z) = (-)^n n! \int_0^\infty \frac{e^{-t} dt}{(t + z)^{n+1}},$$




Fig. 1.1. Magnitude of the terms $a_k = (-)^{k-1} \Gamma(k) z^{-k}$ in the expansion $S_n(z)$ against ordinal number k when $z = 10$.

is of the sign opposite to that in the last term in $S_n(z)$ and further, is of the same sign as the first term left in the full asymptotic series after excising $S_n(z)$. In absolute value, we also have

$$|R_n(z)| = \frac{n!}{z^{n+1}} \int_0^\infty \frac{e^{-t} dt}{(1 + t/z)^{n+1}} < \frac{n!}{z^{n+1}},$$

so the remainder term is numerically smaller in absolute value than the modulus of the first neglected term. Since the series $S_n(z)$ is an alternating series, it follows that $S_n(z)$ is alternately bigger than $U(1; 1; z)$ and less than $U(1; 1; z)$ as n increases. The sum $S_n(z)$ will therefore be closest in value to $U(1; 1; z)$ precisely when we truncate the full expansion just before the numerically smallest term (in absolute value) in the full expansion. From the preceding inequality, it is easy to note that the remainder term will then be bounded by this minimal term.

To see the order of the remainder term at optimal truncation, we substitute $n \sim z$ ($\gg 1$) in the above bound for $R_n(z)$, and employ Stirling's formula to approximate the factorial, to find

$$|R_n(z)| < \frac{n!}{z^{n+1}} \simeq (2\pi)^{\frac{1}{2}} \frac{e^{-n} n^{n+\frac{1}{2}}}{z^{n+1}} \simeq \left(\frac{2\pi}{z}\right)^{\frac{1}{2}} e^{-z}.$$

This shows that at optimal truncation the remainder term for $U(1; 1; z)$ is of order $z^{-\frac{1}{2}} e^{-z}$ as $z \rightarrow +\infty$ and consequently that evaluation of the function by this scheme will result in an error that is *exponentially small* in z ; these results can be extended to deal with complex values of z – see Olver (1974, p. 523) for a more detailed treatment. We remark that this principle is found to apply to a wide range of asymptotic series yielding in each case an error term at optimal truncation that is typically exponentially small in the asymptotic variable.

We observe that not all asymptotic series present the regular behaviour of the coefficients depicted in Fig. 1.1. In certain compound expansions, with coefficients

containing gamma functions in the numerator, it is possible to find situations where some of the arguments of the gamma functions approach a nonpositive integer value. This gives rise to a series of ‘peaks’ superimposed on the basic structure of Fig. 1.1. A specific example is provided by the compound expansion

$$z^{-2/\mu}(I_1 + I_2), \quad (1.1.10)$$

where $I_r = \sum_{k=0}^{\infty} a_k^{(r)}$ ($r = 1, 2$) and, for positive parameters m_1, m_2 and μ ,

$$a_k^{(1)} = \frac{(-)^k}{k!} \Gamma\left(\frac{1+\mu k}{m_1}\right) \Gamma\left(\frac{m_1 - m_2(1+\mu k)}{m_1 \mu}\right) z^{-(1+\mu k)/m_1}$$

with a similar expression for $a_k^{(2)}$ with m_1 and m_2 interchanged. Expansions of this type arise in the treatment of certain Laplace-type integrals discussed in Chapter 7. If the parameters m_1, m_2 and μ are chosen such that the arguments of the second gamma function in $a_k^{(1)}$ and $a_k^{(2)}$ are not close to zero or a negative integer, then the variation of the modulus of the coefficients with ordinal number k will be similar to that shown in Fig. 1.1. If, however, the parameter values are chosen so that these arguments become close to a nonpositive integer† for subsets of k values, then we find that the variation of the coefficients becomes irregular with a sequence of peaks of variable height. Such a situation for the coefficients $a_k^{(1)}$ is shown in Fig. 1.2 for two sets of parameter values. The truncation of such series has been investigated in Liakhovetski & Paris (1998), where it is found that even if the series I_1 is truncated at a peak (provided that the corresponding peak associated with the coefficients $a_k^{(2)}$ is included) increasingly accurate asymptotic approximations are obtained by steadily increasing the truncation indices in the series I_1 and I_2 until they correspond roughly to the global minimum of each curve. An inspection of Fig. 1.2, however, would indicate that these optimal points are not as easily distinguished as in the case of Fig. 1.1.

The notion of optimal truncation will surface in a significant way in the subject matter of the Stokes phenomenon and hyperasymptotics, and so we defer further discussion of it until Chapter 6, where a detailed analysis of remainder terms is undertaken. We do mention, however, that apart from optimally truncating an asymptotic series, one can sometimes obtain dramatic improvements in the numerical utility of an asymptotic expansion if one is able to extract exponentially small (measured against the scale being used) terms prior to developing an asymptotic expansion. This particular situation can be seen in the following example.

† If the parameter values are such that the second gamma-function argument equals a nonpositive integer for a subset of k values, then the expansion (1.1.10) becomes nugatory. In the derivation of (1.1.10) by a Mellin-Barnes approach this would result in a sequence of double poles and the formation of logarithmic terms.




Fig. 1.2. Magnitude of the coefficients $a_k^{(1)}$ against ordinal number k for $\mu = 3$, $m_1 = 1.5$ when (a) $m_2 = 1.2$, $z = 3.0$ and (b) $m_2 = 1.049$, $z = 3.6$. For clarity the points have been joined.

Example 3. Let us consider the finite Fourier integral

$$J(\lambda) = \int_{-1}^1 e^{i\lambda(x^3/3+x)} dx$$

with λ large and positive. Introduce the change of variable $u = \frac{1}{3}x^3 + x$ and observe that over the interval of integration, the change of variable is one-to-one, fixes the origin and maps ± 1 to $\pm \frac{4}{3}$ respectively, resulting in

$$J(\lambda) = \int_{-4/3}^{4/3} e^{i\lambda u} x'(u) du,$$

where $x(u)$ is the function inverse to the $x \mapsto u$ change of variable. An explicit formula for $x(u)$ is available to us from the classical theory of equations, resulting from the trigonometric solution to the cubic equation, and takes the form

$$x = 2 \sinh \theta, \quad \text{where } 3\theta = \operatorname{arcsinh}\left(\frac{3}{2}u\right),$$

or

$$x = \left(\frac{3}{2}u + \sqrt{\frac{9}{4}u^2 + 1}\right)^{1/3} - \left(\frac{3}{2}u - \sqrt{\frac{9}{4}u^2 + 1}\right)^{-1/3}.$$

It is a straightforward matter to deduce that $x^{(k)}(-u) = (-)^{k-1}x^{(k)}(u)$, where $x^{(n)}(u)$ as usual indicates the n th derivative of the inverse function.

By repeatedly applying integration by parts, the latter representation for $J(\lambda)$ can be seen to yield a finite asymptotic expansion with remainder,

$$J(\lambda) = \sum_{n=1}^N \left\{ e^{4i\lambda/3} x^{(n)}\left(\frac{4}{3}\right) - e^{-4i\lambda/3} x^{(n)}\left(-\frac{4}{3}\right) \right\} \frac{(-)^{n-1}}{(i\lambda)^n} \\ + \frac{(-)^N}{(i\lambda)^N} \int_{-4/3}^{4/3} e^{i\lambda u} x^{(N+1)}(u) du. \quad (1.1.11)$$

In view of the Riemann-Lebesgue lemma, the remainder term is seen to be $o(\lambda^{-N})$, so the finite expansion (1.1.11) leads, after exploiting $x^{(k)}(-\frac{4}{3}) = (-)^{k-1} x^{(k)}(\frac{4}{3})$, to the large- λ expansion†

$$J(\lambda) \sim 2 \sin\left(\frac{4}{3}\lambda\right) \sum_{n=0}^{\infty} \frac{(-)^n}{\lambda^{2n+1}} x^{(2n+1)}\left(\frac{4}{3}\right) - 2 \cos\left(\frac{4}{3}\lambda\right) \sum_{n=1}^{\infty} \frac{(-)^n}{\lambda^{2n}} x^{(2n)}\left(\frac{4}{3}\right).$$

If we evaluate the first few derivatives $x^{(n)}(\frac{4}{3})$ and employ optimal truncation for modest values of λ , say $\lambda = 4, 5, 6, 7$, we obtain the approximate values shown in the fourth column of Table 1.3. The columns labelled N_s and N_c show respectively, for each value of λ , the number of terms of the sine and cosine series in the expansion of $J(\lambda)$ retained after optimally truncating each series. As comparison with the last column of Table 1.3 reveals, the asymptotic approximations obtained for these modest values of λ are of poor calibre.

However, an improvement in the numerical utility of the expansion can be obtained by rewriting the integral representation of $J(\lambda)$ in the following manner. Because of the exponential decay in the integrand, we can, by Cauchy's theorem, write

$$J(\lambda) = \left\{ - \int_1^{\infty e^{\pi i/6}} + \int_{-1}^{\infty e^{5\pi i/6}} + \int_{\infty e^{5\pi i/6}}^{\infty e^{\pi i/6}} \right\} e^{i\lambda(x^3/3+x)} dx. \quad (1.1.12)$$

The third integral in this sum can be expressed in terms of the Airy function

$$\text{Ai}(z) = \frac{1}{2\pi i} \int_{\infty e^{-\pi i/3}}^{\infty e^{\pi i/3}} \exp\left(\frac{1}{3}t^3 - zt\right) dt,$$

namely,

$$2\pi\lambda^{-1/3} \text{Ai}(\lambda^{2/3}) = \int_{\infty e^{5\pi i/6}}^{\infty e^{\pi i/6}} e^{i\lambda(x^3/3+x)} dx$$

upon making the substitution $x = i\lambda^{-1/3}$. From this, and integration by parts applied to each of the remaining integrals in (1.1.12), we arrive at the same expansion and approximation for $J(\lambda)$ that we found earlier, only now the expansion

† This expansion does not fit the form of a Poincaré-type expansion as we have defined it previously, but rather is an example (after separating sine and cosine terms) of a compound asymptotic expansion, discussed in the next section.

Table 1.3. *Comparison of optimally truncated asymptotic approximation, asymptotic approximation and exponentially decaying correction and computed values of the Fourier integral $J(\lambda)$*

λ	N_s	N_c	Optimally truncated series	Optimally truncated series with Airy term	$J(\lambda)$
4	2	2	-0.213739	-0.153525	-0.154260
5	3	2	0.055788	0.083551	0.083545
6	6	5	0.164661	0.177709	0.177703
7	6	5	0.022816	0.029031	0.029034

includes the term involving the Airy function:

$$J(\lambda) \sim \frac{2\pi}{\lambda^{1/3}} \text{Ai}(\lambda^{2/3}) + 2 \sin\left(\frac{4}{3}\lambda\right) \sum_{n=0}^{\infty} \frac{(-)^n}{\lambda^{2n+1}} x^{(2n+1)} \left(\frac{4}{3}\right) \\ - 2 \cos\left(\frac{4}{3}\lambda\right) \sum_{n=1}^{\infty} \frac{(-)^n}{\lambda^{2n}} x^{(2n)} \left(\frac{4}{3}\right).$$

The Airy function of positive argument can be shown to exhibit exponential decay as the argument increases, so the additional Airy function term in the above expression is $o(\lambda^{-k})$ for any nonnegative integer k and can be eliminated entirely from the asymptotic expansion in view of the definition of asymptotic expansions of Poincaré type. If it is instead retained, the resulting approximations for the same modest values of λ used in Table 1.3 show dramatic improvement, giving several significant figures of the computed values of $J(\lambda)$ as a comparison of the last two columns of Table 1.3 reveals. $\square$

Another interesting fact concerning asymptotic power series stems from the observation that given an arbitrary sequence of complex numbers $\{a_n\}_{n=0}^{\infty}$, there is a function $f(z)$ holomorphic in a region containing a closed annular sector which has the formal series $\sum_{n=0}^{\infty} a_n z^{-n}$ as its asymptotic expansion.

One such construction[†] proceeds by taking the closed annular sector to be $\mathcal{S} = \{z : |\arg z| \leq \theta, |z| \geq R > 0\}$ – other sectors can be used by translating and rotating this initial choice. Then set

$$f(z) = \sum_{n=0}^{\infty} \frac{a_n e_n(z)}{z^n}$$

where for nonzero a_n ,

$$e_n(z) = 1 - \exp(-z^\phi r^n / |a_n|),$$

[†] This account is drawn from Olver (1974, § 1.9). Other examples along this line are also to be found there.

for numbers ϕ and r chosen to satisfy $0 < \phi < \pi/(2\theta)$ and $0 < r < R$. Should an a_n vanish, the corresponding e_n is taken to be the zero function, so that the corresponding term in the sum defining f is effectively excised.

With these terms so defined, in the sector of interest we have $|\arg(z^\phi)| < \frac{1}{2}\pi$ and

$$\left| \frac{a_n e_n(z)}{z^n} \right| \leq r^n |z|^{\phi-n} \leq |z|^\phi \left(\frac{r}{R} \right)^n, \quad (1.1.13)$$

since $|1 - e^{-\zeta}| \leq |\zeta|$ when $|\arg \zeta| \leq \frac{1}{2}\pi$. The series defining f therefore converges uniformly on compact subsets of our sector, and so defines a holomorphic function there.

That f has the desired asymptotic expansion can be seen from

$$f(z) - \sum_{n=0}^{N-1} \frac{a_n}{z^n} = - \sum_{n=0}^{N-1} \frac{a_n}{z^n} \exp\left(-\frac{z^\phi r^n}{|a_n|}\right) + \sum_{n=N}^{\infty} \frac{a_n e_n(z)}{z^n}, \quad (1.1.14)$$

where it bears noting that the infinite series here is uniformly convergent. Because of the exponential decay of each term in the finite sum on the right, the entire sum is $o(z^{-n})$ for any n as $z \rightarrow \infty$ in our sector. The remaining series on the right-hand side is easily bounded using (1.1.13) to give

$$\left| \sum_{n=N}^{\infty} \frac{a_n e_n(z)}{z^n} \right| \leq |z|^\phi \sum_{n=N}^{\infty} \left(\frac{r}{|z|} \right)^n = |z|^\phi \left(\frac{r}{|z|} \right)^N \frac{|z|}{|z| - r} = O(z^{\phi-N}).$$

Upon replacing N by $N + \lfloor \phi \rfloor + 1$, we obtain a similar expression to that in (1.1.14), for which the right-hand side is $O(z^{-N})$ but for which there are “extra” terms on the left-hand side. These additional terms, $a_n z^{-n}$ for $n \geq N$, are also $O(z^{-N})$ and so can be absorbed into the order estimate that results on the right-hand side.

1.1.3 Other Expansions

Expansions other than Poincaré-type also have currency in asymptotic analysis. Here, we mention but three types.

To begin, let $\{\phi_n\}$ be an asymptotic scale as $x \rightarrow x_0$. A formal series $\sum f_n(x)$ is a *generalised asymptotic expansion* of a function $f(x)$ with respect to the asymptotic scale $\{\phi_n\}$ if

$$f(x) = \sum_{n=0}^N f_n(x) + o(\phi_N(x)) \quad (x \rightarrow x_0, N = 0, 1, 2, \dots).$$

In this event, we write, as we have for Poincaré-type expansions,

$$f(x) \sim \sum_{n=0}^{\infty} f_n(x) \quad (x \rightarrow x_0, \{\phi_n\}),$$

indicating with the formal series the asymptotic scale used to define the expansion.

The important difference between Poincaré and generalised asymptotic expansions is that the functions f_n appearing in the formal series expansion for f need not, themselves, form an asymptotic scale.

Example 1. Define the sequence of functions $\{f_n\}$, for nonnegative integer n and nonzero x , by

$$f_n(x) = \frac{\cos nx}{x^n}.$$

For $x \rightarrow \infty$, it is apparent that each $f_n(x) = O(x^{-n})$, and that $\{\phi_n(x)\} = \{x^{-n}\}$ is an asymptotic scale. However, the sequence $\{f_n(x)\}$ fails to be an asymptotic scale, as a ratio of consecutive elements in the sequence gives

$$\frac{f_{n+1}(x)}{f_n(x)} = \frac{\cos(n+1)x}{x \cos nx},$$

which fails to be $o(1)$ for all x sufficiently large.

Generalised asymptotic expansions are less commonplace than expansions of Poincaré type, and are not used in our development of asymptotic expansions of Mellin-Barnes integrals.

A different mechanism for extending Poincaré-type expansions presents itself naturally in the setting of the method of stationary phase or steepest descent, and in the domain of expansions of solutions of differential equations. The idea here is to replace the series expansion of a function, as in (1.1.2), by several different series, each with different scales.

Put more precisely, by a *compound asymptotic expansion* of a function f , we mean a finite sum of Poincaré-type series expansions

$$\begin{aligned} f(x) \sim & A_1(x) \sum_{n=0}^{\infty} a_{1n} \phi_{1n}(x) + A_2(x) \sum_{n=0}^{\infty} a_{2n} \phi_{2n}(x) \\ & + \cdots + A_k(x) \sum_{n=0}^{\infty} a_{kn} \phi_{kn}(x) \quad (x \rightarrow x_0), \end{aligned}$$

where, for $1 \leq m \leq k$, the sequences $\{\phi_{mn}\}$ are asymptotic scales, the coefficient functions $A_m(x)$ are continuous, and for $N_1, N_2, \dots, N_k \geq 0$, we have

$$\begin{aligned} f(x) = & A_1(x) \left\{ \sum_{n=0}^{N_1} a_{1n} \phi_{1n}(x) + O(\phi_{1, N_1+1}(x)) \right\} \\ & + A_2(x) \left\{ \sum_{n=0}^{N_2} a_{2n} \phi_{2n}(x) + O(\phi_{2, N_2+1}(x)) \right\} \\ & + \cdots + A_k(x) \left\{ \sum_{n=0}^{N_k} a_{kn} \phi_{kn}(x) + O(\phi_{k, N_k+1}(x)) \right\} \quad (x \rightarrow x_0). \end{aligned}$$

It is entirely possible that some of the series $A_j(x) \sum a_{jn} \phi_{jn}(x)$ could, by virtue of the coefficient function $A_j(x)$, or choice of scale $\{\phi_{jn}\}$, be $o(\phi_{mn})$ for some $m \neq j$, and so be absorbed into the error terms implied in other series in the compound expansion. However, in some numerical work, the retention of such negligible terms, when measured against the other scales in the expansion, can add to the numerical accuracy of asymptotic approximations of f , especially for values of x that are at some distance from x_0 . This, in turn, extends the utility of such expansions.

In some circumstances, it may be possible to embed the scales $\{\phi_{mn}\}$ in a larger scale, say $\{\psi_v\}$, and so collapse the sum of Poincaré expansions into a single-series expansion involving this larger scale $\{\psi_v\}$. Success in this direction depends in part on the coefficient functions $A_j(x)$.

Example 2. STEEPEST DESCENT METHOD. An integral of the form

$$I(\lambda) = \int_C g(z) e^{\lambda f(z)} dz,$$

is said to be of *Laplace type* if the functions f and g are holomorphic in a region containing the contour C , and the integral converges for some λ . In the most common setting, C is an infinite contour, and the parameter λ is large in modulus. Thus, we require that the integral $I(\lambda)$ exist for all λ sufficiently large in some sector.

The idea behind the steepest descent method is deceptively simple: deform the integration contour C into a sum of contours, $C_1, C_2, \dots, C_k$, so that along each of the contours C_n , the *phase* function $f(z)$ has a single point z_n – a *saddle point*[†] – at which $f'(z_n)$ vanishes, and as z varies along the contour C_n , $\lambda[f(z) - f(z_n)] \leq 0$, with this difference tending to $-\infty$ as $|z| \rightarrow \infty$ along the contour. If this deformation is possible, the contours $C_1, C_2, \dots, C_k$ are termed *steepest descent contours*, and the integral can be recast as

$$I(\lambda) = \sum_{n=1}^k e^{\lambda f(z_n)} \int_{C_n} g(z) e^{\lambda[f(z) - f(z_n)]} dz.$$

In the case where $f''(z_n) \neq 0$ for all saddle points z_n , each integral in the sum can be represented as a Gaussian integral, namely

$$e^{\lambda f(z_n)} \int_{C_n} g(z) e^{\lambda[f(z) - f(z_n)]} dz = e^{\lambda f(z_n)} \int_{-\infty}^{\infty} g(z(t)) e^{-|\lambda| t^2} z'(t) dt.$$

The transformation $z \mapsto t$ will map one branch of the steepest descent curve from z_n to ∞ into the positive real t axis, and the remainder of the steepest descent curve will be mapped into the negative real t axis. By splitting the integral into integrals

[†] Saddle points of Fourier-type integrals are often referred to as stationary points.

taken over negative and positive real t axes separately, a further reduction to a sum of two Laplace transforms can be achieved, to each of which Watson's lemma can then be applied.

For a concrete example, we consider the Pearcey integral

$$P(x, y) = \int_{-\infty}^{\infty} \exp\{i(t^4 + xt^2 + yt)\} dt,$$

where, for the purpose of illustration, we will assume $|x|$ and $|y|$ are both large, with $x < 0$ and $y > 0$. We will also replace x by $-x$ and take $x > 0$. Thus, we consider

$$P(-x, y) = x^{\frac{1}{2}} \int_{-\infty}^{\infty} \exp\{ix^2(u^4 - u^2 + yx^{-3/2}u)\} du, \quad (1.1.15)$$

where we have applied the simple change of variable $t = x^{\frac{1}{2}}u$. Denoting the phase function of this integral by

$$\psi(u) = u^4 - u^2 + yx^{-3/2}u,$$

we have

$$\begin{aligned} \psi'(u) &= 4(u^3 - \tfrac{1}{2}u + \tfrac{1}{4}yx^{-3/2}) \\ &= 4(u^3 - (u_1 + u_2 + u_3)u^2 + (u_1u_2 + u_1u_3 + u_2u_3)u - u_1u_2u_3), \end{aligned}$$

where the roots of $\psi'(u) = 0$ are indicated by u_1, u_2 and u_3 . Because $\psi'(u)$ is a real cubic polynomial, we always have one real zero. If x is sufficiently large compared to y , we can ensure that the other two zeros of $\psi'(u)$ are also real, and that all three are distinct. Additionally, the elementary theory of equations furnishes us with

$$\sum u_i = 0, \quad \sum_{i < j} u_i u_j = -\tfrac{1}{2}, \quad u_1 u_2 u_3 = -\tfrac{1}{4}yx^{-3/2},$$

from which we deduce that one $u_i < 0$, and the other two are positive. Let us label these so that $u_1 < 0 < u_2 < u_3$.

We mention here that the theory of equations also provides a trigonometric form for the roots u_i , namely,

$$\begin{aligned} u_1 &= -\sqrt{2/3} \cdot \sin(\phi + \tfrac{1}{3}\pi), \\ u_2 &= \sqrt{2/3} \cdot \sin \phi, \\ u_3 &= \sqrt{2/3} \cdot \sin(\tfrac{1}{3}\pi - \phi), \end{aligned} \quad (1.1.16)$$

where the angle ϕ is given by

$$\sin(3\phi) = y(\tfrac{2}{3}x)^{3/2} \quad (1.1.17)$$

which, under the hypothesis of $y(\tfrac{2}{3}x)^{-3/2} < 1$, can be guaranteed to be real. The zeros displayed in (1.1.16) undergo a confluence when the angle ϕ tends to $\tfrac{1}{6}\pi$. The curve this value of ϕ defines is the so-called caustic in the real

plane: $y = (\frac{2}{3}x)^{3/2}$. The saddles u_i are therefore, successively, the locations of a local minimum, a local maximum and a local minimum of $\psi(u)$.

For real x and y , we may rotate the contour of integration in (1.1.15) onto the line from $\infty e^{9\pi i/8}$ to $\infty e^{\pi i/8}$ through an application of Jordan's lemma. Since there are three real saddle points for $(-x, y)$ satisfying $\phi < \frac{1}{6}\pi$, we may further represent $P(-x, y)$ as a sum of three contour integrals,

$$P(-x, y) = x^{\frac{1}{2}} \sum_{j=1}^3 \int_{\Gamma_j} e^{ix^2\psi(u)} du, \quad (1.1.18)$$

where the contours Γ_j are the steepest descent curves: Γ_1 , beginning at $\infty e^{9\pi i/8}$, ending at $\infty e^{5\pi i/8}$ and passing through $u_1 < 0$; Γ_2 , beginning at $\infty e^{5\pi i/8}$, ending at $\infty e^{-3\pi i/8}$ and passing through $u_2 > 0$; and Γ_3 , beginning at $\infty e^{-3\pi i/8}$, ending at $\infty e^{\pi i/8}$ and passing through $u_3 > u_2$. Along these contours, the phase $i\psi(u)$ is real and decreases to $-\infty$ as we move along the Γ_j away from the saddle points so that each integral is effectively a Gaussian integral. The general situation is depicted in Fig. 1.3.

Let us set

$$d_j = \{(-)^j(1 - 6u_j^2)\}^{\frac{1}{2}} \quad (j = 1, 2, 3).$$

In accordance with the steepest descent methodology mentioned previously, we set $\psi(u) - \psi(u_j) = (-)^{j+1}d_j^2v^2$, to find at each saddle point u_j ,

$$v = (u - u_j) \left\{ 1 + \frac{4u_j(u - u_j)}{6u_j^2 - 1} + \frac{(u - u_j)^2}{6u_j^2 - 1} \right\}^{1/2}$$

whence reversion yields the expansion, for each j ,

$$u - u_j = \sum_{k=1}^{\infty} b_{k,j} v^k,$$




Fig. 1.3. Steepest descent curves through the saddles u_1 , u_2 and u_3 .

convergent in a neighbourhood of $v = 0$. We observe that $b_{1,j} = 1$ for each $j = 1, 2, 3$. Substitution into each term in (1.1.18) followed by termwise integration will furnish

$$\int_{\Gamma_j} e^{ix^2\psi(u)} du \sim e^{ix^2\psi(u_j)+(-)^{j+1}\pi i/4} \frac{\pi^{\frac{1}{2}}}{x d_j} S_j(x, \phi),$$

where $S_j(x, \phi)$ denotes the formal asymptotic sum

$$S_j(x, \phi) = \sum_{k=0}^{\infty} (2k+1) b_{2k+1,j} \frac{\Gamma(k+\frac{1}{2})}{\Gamma(\frac{1}{2})} ((-)^{j+1}i)^k (d_j x)^{-2k}.$$

It then follows that

$$P(-x, y) \sim \sqrt{\frac{\pi}{x}} \sum_{j=1}^3 \frac{e^{ix^2\psi(u_j)+(-)^{j+1}\pi i/4}}{d_j} S_j(x, \phi)$$

for large x . This is evidently a compound asymptotic expansion with each constituent asymptotic series corresponding to a single saddle point of $P(-x, y)$. We shall meet the Pearcey integral again in Chapter 8, in a less restricted setting. $\square$

There also arise situations in which functions depending on parameters other than the asymptotic one may possess asymptotic expansions which not only depend on such auxiliary parameters, but may also undergo discontinuous changes of scale as these parameters vary. Such a discontinuity in the scale can occur, even if the function involved is holomorphic in the control parameter. In more specific terms, let us suppose that a function $F(\lambda; \mu)$ has asymptotic parameter λ and control parameter μ . For $\lambda \rightarrow \lambda_0$, and $\mu < \mu_0$, say, one might have an asymptotic form

$$F(\lambda; \mu) \sim A_1(\lambda; \mu) \sum_{n=0}^{\infty} a_{1n}(\mu) \phi_{1n}^- + \cdots + A_k(\lambda; \mu) \sum_{n=0}^{\infty} a_{kn}(\mu) \phi_{kn}^-,$$

where $\{\phi_{jn}^-\}$ ($1 \leq j \leq k$) are asymptotic scales in the variable λ , while for $\mu > \mu_0$, a different expansion might hold, say

$$F(\lambda; \mu) \sim B_1(\lambda; \mu) \sum_{n=0}^{\infty} b_{1n}(\mu) \phi_{1n}^+ + \cdots + B_r(\lambda; \mu) \sum_{n=0}^{\infty} b_{rn}(\mu) \phi_{rn}^+,$$

for different scales $\{\phi_{jn}^+\}$ ($1 \leq j \leq r$) in λ . For the value $\mu = \mu_0$, a third expansion may hold, involving yet another scale $\{\phi_{jn}\}$ ($1 \leq j \leq s$),

$$F(\lambda; \mu_0) \sim C_1(\lambda) \sum_{n=0}^{\infty} c_{1n} \phi_{1n} + \cdots + C_s(\lambda) \sum_{n=0}^{\infty} c_{sn} \phi_{sn} \quad (\lambda \rightarrow \lambda_0).$$

Distinct forms such as these may apply, even if F is analytic in a neighbourhood of μ_0 , and the limiting forms of the expansions may not exist as $\mu \rightarrow \mu_0^{\pm}$,