

# The Prime Number Theorem

G. J. O. JAMESON

*Lancaster University*



PUBLISHED BY THE PRESS SYNDICATE OF THE UNIVERSITY OF CAMBRIDGE  
The Pitt Building, Trumpington Street, Cambridge, United Kingdom

CAMBRIDGE UNIVERSITY PRESS  
The Edinburgh Building, Cambridge CB2 2RU, UK  
40 West 20th Street, New York, NY 10011-4211, USA  
477 Williamstown Road, Port Melbourne, VIC 3207, Australia  
Ruiz de Alarcón 13, 28014 Madrid, Spain  
Dock House, The Waterfront, Cape Town 8001, South Africa

<http://www.cambridge.org>

© Graham Jameson 2003

This book is in copyright. Subject to statutory exception  
and to the provisions of relevant collective licensing agreements,  
no reproduction of any part may take place without the written  
permission of Cambridge University Press.

First published 2003

Printed in the United Kingdom at the University Press, Cambridge

*A catalog record for this book is available from the British Library.*

*Library of Congress Cataloging in Publication data*

Jameson, G. J. O. (Graham James Oscar)  
The prime number theorem / G.J.O. Jameson.  
p. cm. – (London Mathematical Society student texts ; 53)  
Includes bibliographical references and index.  
ISBN 0-521-81411-1 – ISBN 0-521-89110-8 (pb.)  
I. Numbers, Prime. I. Title. II. Series

QA246 .J36 2003  
512'.72-dc21 2002074199

ISBN 0 521 81411 1 hardback  
ISBN 0 521 89110 8 paperback

# Contents

<i>Preface</i>	<i>page vii</i>
<b>1 Foundations</b>	<b>1</b>
1.1 Counting prime numbers	2
1.2 Arithmetic functions	6
1.3 Abel summation	9
1.4 Estimation of sums by integrals; Euler's summation formula	18
1.5 The function $\text{li}(x)$	28
1.6 Chebyshev's theta function	33
1.7 Dirichlet series and the zeta function	38
1.8 Convolutions	50
<b>2 Some important Dirichlet series and arithmetic functions</b>	<b>56</b>
2.1 The Euler product	57
2.2 The Möbius function	61
2.3 The series for $\log \zeta(s)$ and $\zeta'(s)/\zeta(s)$	70
2.4 Chebyshev's psi function and powers of primes	75
2.5 Estimates of some summation functions	82
2.6 Mertens's estimates	89
<b>3 The basic theorems</b>	<b>98</b>
3.1 Extension of the definition of the zeta function	99
3.2 Inversion of Dirichlet series; the integral version of the fundamental theorem	114
3.3 An alternative method: Newman's proof	124
3.4 The limit and series versions of the fundamental theorem; the prime number theorem	130
3.5 Some applications of the prime number theorem	138
<b>4 Prime numbers in residue classes: Dirichlet's theorem</b>	<b>147</b>
4.1 Characters of finite abelian groups	148

4.2	Dirichlet characters	153
4.3	Dirichlet $L$ -functions	160
4.4	Prime numbers in residue classes	173
<b>5</b>	<b>Error estimates and the Riemann hypothesis</b>	<b>178</b>
5.1	Error estimates	179
5.2	Connections with the Riemann hypothesis	192
5.3	The zero-free region of the zeta function	195
<b>6</b>	<b>An “elementary” proof of the prime number theorem</b>	<b>206</b>
6.1	Framework of the proof	207
6.2	Selberg’s formulae and completion of the proof	212
 <i>Appendices</i>		
A	Complex functions of a real variable	223
B	Double series and multiplication of series	225
C	Infinite products	228
D	Differentiation under the integral sign	230
E	The $O, o$ notation	232
F	Computing values of $\pi(x)$	234
G	Table of primes	238
H	Biographical notes	240
 <i>Bibliography</i>		
<i>Index</i>		

# 1

## Foundations

In this chapter, we assemble a number of ideas and techniques that will eventually be fitted together to achieve our aim. Their only common feature is that they are needed to prove the prime number theorem, so the chapter has no single unifying theme. However, each *section* of the chapter is devoted to a very clearly defined topic. Some of these ideas are analytic, others number-theoretic, but there would be no advantage in trying to keep the two strands apart: they reinforce each other in a fruitful partnership.

Our objective is to find a formula that approximates  $\pi(x)$ , the number of primes not greater than  $x$ . We start, in section 1.1, by identifying some candidates as suggested by numerical evidence. We also give a brief account of the long history leading to the successful proof of the prime number theorem.

The term “arithmetic function” is used for a sequence defined using number-theoretic properties in some way. A great deal of number theory consists of the study of such functions. Now we can express  $\pi(x)$  as the partial sum  $\sum_{n \leq x} u_P(n)$ , where  $u_P$  is the arithmetic function defined as follows:

$$u_P(n) = \begin{cases} 1 & \text{if } n \text{ is prime,} \\ 0 & \text{otherwise.} \end{cases}$$

Typically, arithmetic functions appear to be very irregular, but this is smoothed out by addition, and one can hope to find an estimate for their partial sums. This identifies our problem as one of a certain type.

We go on to describe two essential techniques for rewriting and estimating discrete sums, Abel summation and integral estimation. Both are used constantly in all that follows.

After this, we are in a position to describe the first real progress towards the prime number theorem, achieved by Chebyshev in 1850. Chebyshev recognised that an estimation of  $\pi(x)$  can be deduced from an estimation of  $\theta(x) = \sum_{p \in P[x]} \log p$  (where  $P[x]$  denotes the set of primes not greater than

$x$ ), and showed that the latter sum can be estimated by a comparatively short (but ingenious) argument. In this way, he demonstrated that  $\pi(x)$  lies between  $cx/\log x$  and  $Cx/\log x$  for two constants  $c, C$ .

Finally, we introduce a concept that will permeate the rest of our study to the extent that it could serve as a subtitle for this book. A *Dirichlet series* is a series of the form  $\sum_{n=1}^{\infty} a(n)/n^s$ , in which  $s$  is a complex variable. The case  $a(n) = 1$  defines the *Riemann zeta function*. Every arithmetic function has a corresponding Dirichlet series; multiplication of the series corresponds to “convolution” of the arithmetic functions. Our “fundamental theorems” in chapter 3 will derive information about the partial sums of  $a(n)$  from the nature of the function defined by the series, with the prime number theorem appearing as a special case.

### 1.1 Counting prime numbers

As the reader surely knows, prime numbers are those that have no positive divisors except 1 and the number itself. The special significance of prime numbers is due to the following fact, which we will assume known:

*Every positive integer is expressible as a product of primes. The expression is unique if the primes are listed in increasing order.*

Effectively, this means that the primes are the basic “atoms” in the multiplicative system of integers. (If  $n$  is itself prime, we are regarding it as a “product” of one prime, itself.)

The first result on the number of primes was already known to Euclid. Here it is, with Euclid’s beautiful proof.

**Proposition 1.1.1** *There are infinitely many prime numbers.*

*Proof* Choose finitely many primes  $p_1, p_2, \dots, p_n$ . We will show that they cannot constitute the total set of primes. Consider the number

$$N = p_1 p_2 \dots p_n + 1.$$

Then  $N$  is not a multiple of any  $p_j$ , because it clearly leaves remainder 1 when divided by  $p_j$ . However, by the above statement,  $N$  is expressible as a product of primes. Let  $q$  be any one of these. Then  $q$  is a further prime, different from all the  $p_j$ , which therefore indeed fail to constitute the total set of primes.  $\square$

**Note** This reasoning actually shows a bit more: if the primes are listed as  $p_1, p_2, \dots$  in increasing order, then  $p_{n+1} \leq p_1 p_2 \dots p_n + 1$ .

With this settled, it is natural to ask how many prime numbers there are up to any given number. This is the topic of our study. Let us give it some notation:

$$\pi(x) = \text{the number of primes not greater than } x.$$

This notation is standard in number theory; there is no real danger of confusion with the number  $\pi$ . It will suit our purposes to regard  $\pi(x)$  as a function of a real variable  $x$ . As such a function, it is, of course, constant between primes and jumps by 1 at each prime.

The first impression given by the sequence of primes

$$2, 3, 5, 7, 11, \dots, 101, 103, 107, 109, 113, 127, \dots, 163, 167, 173, 179, \dots$$

is one of extreme irregularity. There are bunches, gaps and relatively uniform stretches. It would appear to be a daunting task to find a simple expression that approximates to  $\pi(x)$  for all large enough  $x$ . The only simple observation is that the primes tend to become more sparse as one goes on. However, an examination of the numerical values of  $\pi(x)$  suggests that a reasonable approximation is given by  $x/(\log x)$ , a considerably better one by

$$\frac{x}{\log x - 1},$$

and a still better one by the “logarithmic integral”, defined as follows:

$$\text{li}(x) = \int_2^x \frac{1}{\log t} dt.$$

Some of these numerical values are as follows (given to the nearest integer):

$n$	$\pi(n)$	$\frac{n}{\log n}$	$\frac{n}{\log n - 1}$	$\text{li}(n)$
1,000	168	145	169	177
10,000	1,229	1,086	1,218	1,246
50,000	5,133	4,621	5,092	5,166
100,000	9,592	8,686	9,512	9,630
500,000	41,538	38,103	41,246	41,607
1,000,000	78,498	72,382	78,031	78,628
10,000,000	664,579	620,421	661,459	664,918

By the year 1800, long before the age of computers, mathematicians had performed the remarkable feat of calculating these figures by hand up to

$n = 400,000$ . In the age of computers, it has of course become much easier to calculate values of  $\pi(x)$ . Some readers will be interested in doing so on their own computer: various methods for this are discussed in appendix F.

Let us formulate precisely the conjecture suggested by these figures. Given two functions  $f(x)$ ,  $g(x)$ , both tending to infinity as  $x \rightarrow \infty$ , we write

$$f(x) \sim g(x) \quad \text{as } x \rightarrow \infty$$

to mean that

$$\frac{f(x)}{g(x)} \rightarrow 1 \quad \text{as } x \rightarrow \infty.$$

Our conjecture is the statement

$$\pi(x) \sim \text{li}(x) \quad \text{as } x \rightarrow \infty.$$

In fact, as we will show in section 1.5,

$$\frac{x}{\log x} \sim \frac{x}{\log x - 1} \sim \text{li}(x) \quad \text{as } x \rightarrow \infty,$$

so at this level it is equivalent to state the conjecture using any of the three functions.

The conjecture is in fact true. The statement  $\pi(x) \sim \text{li}(x)$  is called the *prime number theorem*. It is indisputably one of the most celebrated theorems in mathematics. Ways of proving it, together with related results, more precise versions and generalizations, form the subject of this book. Of course, numerical evidence of the above type can never constitute a proof of the general statement.

An informal interpretation of the theorem is that the “average density” of primes around a large number  $x$  approximates to  $1/(\log x)$ , or that the “probability” of  $n$  being prime is (in some sense)  $1/(\log n)$ .

Let us return to the historical trail (for a much more detailed historical account, see [Nar]). Legendre, in 1798, postulated the approximations  $x/(\log x)$  and  $x/(\log x - 1)$ . He suggested (wrongly) that an even better approximation would be given by  $x/(\log x - A)$ , with  $A = 1.0836$ . Meanwhile, Gauss proposed  $\text{li}(x)$ . It seems that Gauss recorded his conjecture around 1793 (at the age of 14!) but did not communicate it to anyone until 1849.

The search for a proof remained one of the main areas of mathematical endeavour during the rest of the nineteenth century. In 1850, a giant stride was made by Chebyshev, who showed, by essentially number-theoretic methods, that there are constants  $c$  and  $C$  (not very far from 1) such that

$$c \text{li}(x) \leq \pi(x) \leq C \text{li}(x)$$



for all large enough  $x$ . However, no refinement of his methods seemed to offer any hope of proving the desired limit.

A completely different approach was proposed by Riemann in 1859. His starting point was a remarkable identity already discovered by Euler in 1737, expressing the “zeta function”

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$$

as an infinite *product* involving the primes. Riemann considered this as a function of a *complex* variable  $s$ , defined by the above formula for  $\text{Re } s > 1$ . He showed how to extend the definition of the function to the rest of the complex plane and outlined a programme showing how, if certain properties of the extended zeta function could be established, the prime number theorem would follow. His paper was a bold imaginative leap; it was hardly an obvious idea to use the theorems of complex analysis to count prime numbers! However, Riemann was not able to justify all his steps, and one of them, the “Riemann hypothesis”, has remained unsolved to this day, regarded by many as the most important unsolved problem in mathematics.

It was not until 1896 that Riemann’s programme was successfully completed. It was then done so independently by the French mathematician Jacques Hadamard and the Belgian Charles-Jean de la Vallée Poussin. They were able to bypass the Riemann hypothesis and establish other properties of the zeta function that were sufficient for the purpose. Hadamard lived until 1963 (aged 97) and de la Vallée Poussin until 1962 (aged 95): their mathematical labours cannot have done any harm to their health!

Further variations and modifications of their methods were developed by Mertens, Landau and others, but to this day the simplest, and most powerful, proofs of the prime number theorem rely on the zeta function and complex analysis, as suggested by Riemann. In chapters 1–3, we present a version (and a variant) that benefits from a century of “tidying up”, but which still recognisably owes its existence to Riemann.

After the successful outcome of Riemann’s programme, it remained a matter of great interest to ask whether the theorem could after all be proved by number-theoretic methods, without complex analysis. This was eventually achieved in 1949, again independently by two people, A. Selberg and P. Erdős. Proofs of this sort are called “elementary” as opposed to “analytic”. However, “elementary” does not mean “simple”! Half a century later, known proofs of this sort are still more complicated than analytic ones and are less successful in providing error estimates or in delivering other theorems of the same sort. A version is presented in chapter 6.

## Exercises

- 1 Let  $n \geq 1$  and let  $E = \{30n + r : 0 \leq r \leq 29\}$ . For which values of  $r$  is  $30n + r$  not a multiple of 2, 3 or 5? By considering the possible positions of multiples of 7, show that  $E$  contains at most seven primes (seven cases, no short cuts!).
- 2 Show that, for any  $n \geq 2$ , there is eventually a gap of length at least  $n$  between successive primes. [Hint: Consider  $n! + 2$  or  $p_1 p_2 \dots p_n + 2$ .]
- 3 Let the primes be listed, in order, as  $p_1, p_2, \dots$ . Use Euclid's proof to show by induction that  $p_n \leq 2^{2^{n-1}}$  for each  $n$ . Deduce that

$$\pi(x) \geq \frac{\log \log x}{\log 2}.$$

## 1.2 Arithmetic functions

Formally, an arithmetic function is simply a sequence, with real or complex values. A sequence is, of course, a *function* on the set  $\mathbb{N}$  of positive integers. To emphasize that we are thinking of them as functions, we shall usually use notation like  $a(n)$ , rather than  $a_n$ , for the value corresponding to the integer  $n$ . The term "arithmetic function" is used especially when  $a(n)$  is defined using number-theoretic properties in some way. A large part of number theory consists, in one way or another, of the study of these functions.

We list some examples. First, two very simple ones, mainly to establish the notation we will use:

$$u(n) = 1 \quad \text{for all } n \quad (\text{the "unit function"});$$

$$e_j(n) = \begin{cases} 1 & \text{if } n = j, \\ 0 & \text{if } n \neq j. \end{cases}$$

Next, given any subset  $E$  of  $\mathbb{N}$  (for example, the set  $P$  of primes), define

$$u_E(n) = \begin{cases} 1 & \text{if } n \in E, \\ 0 & \text{if } n \notin E. \end{cases}$$

Clearly,  $u$  itself is the case  $E = \mathbb{N}$ . Third, three more obviously number-theoretic examples:

$\tau(n)$  = the number of (positive) divisors of  $n$ , including 1 and  $n$ ;

$\omega(n)$  = the number of prime divisors of  $n$ ;

$\Omega(n)$  = the number of prime factors of  $n$ , counted with repetitions.

(This notation is more or less standard, though some writers use  $d$  instead of  $\tau$ .) Note that  $\tau(1) = 1$  and  $\omega(1) = \Omega(1) = 0$ . For  $n > 1$ , these functions can easily be described in terms of the prime factorization of  $n$ , as follows.

**Proposition 1.2.1** *Suppose that  $n > 1$ , with prime factorization*

$$n = \prod_{j=1}^m p_j^{k_j}.$$

Then

$$\tau(n) = \prod_{j=1}^m (k_j + 1), \quad \omega(n) = m, \quad \Omega(n) = \sum_{j=1}^m k_j.$$

*Proof* The expressions for  $\omega(n)$  and  $\Omega(n)$  are just the definition. Divisors of  $n$  are of the form  $\prod_{j=1}^m p_j^{r_j}$ , where, for each  $j$ , the possible values of  $r_j$  are  $0, 1, \dots, k_j$ . This gives the expression for  $\tau(n)$ .  $\square$

In particular, if  $p$  is prime, then  $\tau(p^k) = k + 1$ ,  $\omega(p^k) = 1$  and  $\Omega(p^k) = k$ . To give another example, since  $72 = 2^3 \cdot 3^2$ , we have  $\tau(72) = 12$ ,  $\omega(72) = 2$  and  $\Omega(72) = 5$ .

Given arithmetic functions  $a, b$ , we denote the pointwise product by  $ab$ , so that  $(ab)(n) = a(n)b(n)$ . Obviously,  $au = a$  for any  $a$ .

**Summation functions.** Given an arithmetic function  $a(n)$ , its *summation function*  $A(x)$  is defined by

$$A(x) = \sum_{n \leq x} a(n).$$

It is useful to regard  $A(x)$  as a function of a *real* variable  $x$ . As such a function, it is, of course, constant between integers and has a jump discontinuity at each integer where  $a(n) \neq 0$ . Clearly,  $\pi(x)$  is the summation function of  $u_P(n)$  (note: in particular cases, the established notation will not usually allow a notational device like the substitution of  $A$  for  $a$ ).

Individual values of arithmetic functions may fluctuate wildly – as in most of the examples just given. However, in many cases summation smooths out the fluctuation, and it may be possible to find an asymptotic expression for the summation function for large  $x$ . In the case of  $\tau(n)$  and  $\omega(n)$ , the first step is to apply a bit of “lateral thinking” to obtain alternative expressions for the summation functions, as in the next result. The notation  $[x]$  means the largest integer not greater than  $x$ . We shall use the notation  $P[x]$  to mean the set of primes not greater than  $x$  (but there is no generally agreed notation for this). Also,  $j|n$  means that  $j$  divides into  $n$ .

**Proposition 1.2.2** Write  $S_\tau(x) = \sum_{n \leq x} \tau(n)$  and  $S_\omega(x) = \sum_{n \leq x} \omega(n)$ . Then

$$S_\tau(x) = \sum_{j \leq x} \left[ \frac{x}{j} \right], \quad S_\omega(x) = \sum_{p \in P[x]} \left[ \frac{x}{p} \right].$$

*Proof* Clearly,  $S_\tau(x)$  is the number of ordered pairs  $(j, n)$  such that  $j|n$  and  $n \leq x$ . For fixed  $j$  (instead of fixed  $n$ ), the number of such pairs is the number of multiples  $rj$  not greater than  $x$ . This number is obviously  $[x/j]$ . The stated expression follows.

The argument for  $S_\omega(x)$  is similar, counting the number of pairs  $(p, n)$  as above, but with  $p$  prime.  $\square$

The “double counting” principle seen in this proof is often useful. We shall see later how the identities in 1.2.2 can be used to derive asymptotic expressions for  $S_\tau(x)$  and  $S_\omega(x)$ , and similar results will be obtained for some other arithmetic functions. Since  $\pi(x)$  is itself a summation function, our main objective, the prime number theorem, is a result of exactly this type. But, as the reader may suspect, this case will cost us a lot more effort than most of the others.

**Multiplicative functions.** We denote by  $(m, n)$  the greatest common divisor of  $m$  and  $n$ . An arithmetic function  $a$  is said to be

*completely multiplicative* if  $a(mn) = a(m)a(n)$  for all  $m, n$ ;

*multiplicative* if  $a(mn) = a(m)a(n)$  whenever  $(m, n) = 1$ .

Clearly, if  $a$  is multiplicative and not identically zero, then  $a(1) = 1$ . Also,  $a$  is fully determined by the values  $a(p^k)$  for prime  $p$ , since if the prime factorization of  $n$  is  $\prod_{j=1}^r p_j^{k_j}$ , then  $a(n) = \prod_{j=1}^r a(p_j^{k_j})$ . Of course, if  $a$  is completely multiplicative, then  $a(p^k) = a(p)^k$ , and the function is already fully determined by the values  $a(p)$ .

We list some examples.

- (i) For any  $s$ , let  $a(n) = n^s$ . Then  $a$  is completely multiplicative.
- (ii)  $e_1$  is completely multiplicative, but  $e_j$  is not multiplicative for  $j \geq 2$ , since  $e_j(1) = 0$ .
- (iii)  $\tau$  is multiplicative. This follows from 1.2.1, since if  $(m, n) = 1$ , then  $m$  and  $n$  have different prime divisors. It is not completely multiplicative, since  $\tau(2) = 2$ ,  $\tau(4) = 3$ .
- (iv)  $u_P$  is not multiplicative, since  $u_P(1) = 0$ .
- (v) Neither  $\omega$  nor  $\Omega$  is multiplicative; however, we have  $\Omega(mn) = \Omega(m) + \Omega(n)$ , and similarly for  $\omega$  when  $(m, n) = 1$ .

- (vi) *Liouville's function* is defined by  $\lambda(n) = (-1)^{\Omega(n)}$ . It is completely multiplicative, by the statement in (v).
- (vii) Let

$$\chi(n) = \begin{cases} 0 & \text{if } n \text{ is even} \\ 1 & \text{if } n \equiv 1 \pmod{4} \\ -1 & \text{if } n \equiv -1 \pmod{4}. \end{cases}$$

By considering the different cases, one checks easily that  $\chi$  is completely multiplicative.

As the reader may already know, there are many further interesting arithmetic functions. Some will make their appearance in later sections.

### Exercises

- 1 Find the smallest  $n$  such that: (i)  $\Omega(n) = 4$ , (ii)  $\omega(n) = 4$ , (iii)  $\tau(n) = 4$ .
- 2 Show that  $\sum_{n=1}^{30} \tau(n) = 111$  without calculating individual values of  $\tau(n)$ .
- 3 Calculate  $\sum_{n=1}^{100} \omega(n)$  without calculating individual values of  $\omega(n)$ .
- 4 Show that  $\tau(n)$  is odd if and only if  $n$  is a square.
- 5 Show that, for any  $n \geq 2$ ,

$$2^{\omega(n)} \leq \tau(n) \leq 2^{\Omega(n)} \leq n.$$

[You may assume that  $k+1 \leq 2^k$  for all  $k \geq 0$ .]

- 6 Let  $S$  be the set of squares. Show that  $u_S$  is multiplicative.
- 7 Let  $a(n) = (-1)^{n-1}$  for  $n \geq 1$ . Show that  $a$  is multiplicative.
- 8 Prove that, for any  $\varepsilon > 0$ , we have  $\tau(n)/n^\varepsilon \rightarrow 0$  as  $n \rightarrow \infty$ . [Again use  $k+1 \leq 2^k$ . For each prime  $p < 2^{1/\varepsilon}$ , show that there is a constant  $C_p$  such that  $k+1 \leq C_p p^{\varepsilon k}$  for all  $k$ .]

## 1.3 Abel summation

### *Discrete version*

Abel summation, in its various forms, will be a very basic tool in all that follows, so we will describe it rather thoroughly. It is the process of expressing a sum of products  $\sum a(r)f(r)$  in terms of *partial sums* of the  $a(r)$ 's and *differences* of the  $f(r)$ 's. Our choice of notation reflects the different roles played by  $a(r)$  and  $f(r)$ . The process is exactly analogous to integration by

parts for functions, and indeed it is sometimes called “summation by parts” or “partial summation”. As already mentioned, a central theme in analytic number theory is the estimation of partial sums (rather than individual values) of arithmetic functions. This explains why Abel summation is so often appropriate. Throughout the following, we assume that  $a(r)$ ,  $f(r)$  are given numbers (real or complex) for  $r \geq 1$ , and write  $A(n) = \sum_{r=1}^n a(r)$  for  $n \geq 1$  (also  $A(0) = 0$ ). If we have another sequence  $b(r)$ , then  $B(n)$  is defined similarly. The basic result is very simple, as follows.

**Proposition 1.3.1** *For integers  $n > m \geq 0$ ,*

$$\sum_{r=m+1}^n a(r)f(r) = \sum_{r=m}^{n-1} A(r)[f(r) - f(r+1)] + A(n)f(n) - A(m)f(m).$$

*In particular,*

$$\sum_{r=1}^n a(r)f(r) = \sum_{r=1}^{n-1} A(r)[f(r) - f(r+1)] + A(n)f(n).$$

*Proof* The proof looks nicer if we write  $A_r$  and  $f_r$  instead of  $A(r)$  and  $f(r)$ ! For all  $r \geq 1$ , we have  $a_r = A_r - A_{r-1}$  (recall  $A_0 = 0$ ). Hence

$$\begin{aligned} \sum_{r=m+1}^n a_r f_r &= (A_{m+1} - A_m)f_{m+1} + (A_{m+2} - A_{m+1})f_{m+2} + \cdots \\ &\quad \cdots + (A_n - A_{n-1})f_n \\ &= -A_m f_{m+1} + \sum_{r=m+1}^{n-1} A_r (f_r - f_{r+1}) + A_n f_n \end{aligned} \tag{1.1}$$

$$= -A_m f_m + \sum_{r=m}^{n-1} A_r (f_r - f_{r+1}) + A_n f_n. \tag{1.2}$$

The second statement is the case  $m = 0$ . □

This simple identity has numerous corollaries and applications.

**Corollary 1.3.2** *Suppose that  $f(r)$  is real and non-negative, and decreases with  $r$ . Suppose that  $a(r)$ ,  $b(r)$  are such that  $A(r) \leq C B(r)$  for all  $r$ . Then*

$$\sum_{r=1}^n a(r)f(r) \leq C \sum_{r=1}^n b(r)f(r).$$

*Proof* This follows at once from 1.3.1, because  $f(r) - f(r+1)$  and  $f(n)$  are non-negative. □

**Note** Taking  $a(r) = 1$  in 1.3.1, we obtain an expression for  $\sum_{r=1}^n f(r)$  itself:

$$\sum_{r=1}^n f(r) = \sum_{r=1}^{n-1} r[f(r) - f(r+1)] + nf(n).$$

Next, we describe some applications to infinite series.

**Proposition 1.3.3** *Suppose that  $A(n)f(n) \rightarrow 0$  as  $n \rightarrow \infty$ . Then if one of the series*

$$\sum_{r=1}^{\infty} a(r)f(r) \quad \text{and} \quad \sum_{r=1}^{\infty} A(r)[f(r) - f(r+1)]$$

*converges, then so does the other, to the same sum.*

*Proof* Recall that the sum of a series means the limit as  $n \rightarrow \infty$  of the sum of  $n$  terms. Given this, the statement follows at once from 1.3.1.  $\square$

Note that by taking limits in the first statement in 1.3.1, we have also

$$\sum_{r=m+1}^{\infty} a(r)f(r) = \sum_{r=m}^{\infty} A(r)[f(r) - f(r+1)] - A(m)f(m).$$

**Proposition 1.3.4 (Dirichlet's test for convergence)** *Suppose that:*

- (i)  $|A(r)| \leq C$  for all  $r$ ,
- (ii)  $f(r) \rightarrow 0$  as  $r \rightarrow \infty$ ,
- (iii)  $\sum_{r=1}^{\infty} |f(r) - f(r+1)|$  is convergent.

Then  $\sum_{r=1}^{\infty} a(r)f(r)$  converges, say to  $S$ , where

$$|S| \leq C \sum_{r=1}^{\infty} |f(r) - f(r+1)|.$$

Condition (iii) can be replaced by: (iiia)  $f(r)$  is non-negative and decreasing. We then have  $|S| \leq Cf(1)$ .

*Proof* By the comparison test for series,  $\sum_{r=1}^{\infty} A(r)[f(r) - f(r+1)]$  is convergent, with the sum  $S$  satisfying the stated inequality. Also, clearly,  $A(r)f(r) \rightarrow 0$  as  $r \rightarrow \infty$ . The statement follows, by 1.3.3. Under condition (iiia), we have

$$\sum_{r=1}^n |f(r) - f(r+1)| = \sum_{r=1}^n [f(r) - f(r+1)] = f(1) - f(n+1),$$

so that  $\sum_{r=1}^{\infty} |f(r) - f(r+1)| = f(1)$ .  $\square$

Note that, under the conditions of this result,  $\sum_{r=1}^{\infty} A(r) [f(r) - f(r+1)]$  is *absolutely* convergent, while  $\sum_{r=1}^{\infty} a(r)f(r)$  may well not be.

In particular, condition (i) implies that, for all real  $s > 0$ , the series  $\sum a(r)/r^s$  converges, say to  $S$ , where  $|S| \leq C$ . Later, we shall be considering series of this type for *complex*  $s$ ; these are more easily handled by the continuous version of Abel summation, discussed below.

Abel summation is often used to give estimations of partial sums, given information about other partial sums. The next example illustrates this.

**Example 1** Given that  $|\sum_{r=1}^n a(r)/r| \leq C$  for all  $n$ , to show that  $|A(n)| \leq (2n-1)C$  for all  $n$ .

Write  $a(r)/r = b(r)$ . Then we have  $|B(n)| \leq C$  for all  $n$  and, by 1.3.1,

$$A(n) = \sum_{r=1}^n rb(r) = -B(1) - B(2) - \dots - B(n-1) + nB(n).$$

The stated inequality is now clear.

### *Continuous version*

Given an arithmetic function  $a(r)$ , its summation function  $A(x)$  is defined for all positive, real  $x$ , not just integers. Meanwhile, in products of the type we are considering, the numbers  $f(r)$  (as our notation already suggests) are often the values at integers of a function  $f(x)$  of a real variable. When this happens, we can replace the discrete sums in the above expressions by integrals, using the elementary fact that  $f(r+1) - f(r) = \int_r^{r+1} f'(t) dt$  (this remains valid when  $f$  is complex-valued: for clarification, see appendix A). The resulting identity, in its various forms, is sometimes called *Abel's summation formula*. It is very useful, because integrals are often easier to evaluate than discrete sums.

Throughout the following, the numbers  $a(r)$  (for integers  $r \geq 1$ ) are assumed given, and  $A(x) = \sum_{r \leq x} a(r)$ . We start by giving the continuous equivalent of the first expression in 1.3.1. The similarity with integration by parts is even more apparent.

**Theorem 1.3.5** *Let  $y < x$ , and let  $f$  be a function (with real or complex values) having a continuous derivative on  $[y, x]$ . Then*

$$\sum_{y < r \leq x} a(r)f(r) = A(x)f(x) - A(y)f(y) - \int_y^x A(t)f'(t) dt.$$



*Proof* Let  $m, n$  be integers such that  $n \leq x < n + 1$  and  $m \leq y < m + 1$ . Then (treating the symbols  $<$  and  $\leq$  with due respect) we have

$$\sum_{y < r \leq x} a(r)f(r) = \sum_{r=m+1}^n a(r)f(r).$$

We use identity (1.1) from the proof of 1.3.1. Since  $A(t) = A(r)$  for  $r \leq t < r + 1$ , we have

$$\begin{aligned} \sum_{r=m+1}^{n-1} A(r)[f(r) - f(r+1)] &= - \sum_{r=m+1}^{n-1} A(r) \int_r^{r+1} f'(t) dt \\ &= - \sum_{r=m+1}^{n-1} \int_r^{r+1} A(t)f'(t) dt \\ &= - \int_{m+1}^n A(t)f'(t) dt. \end{aligned}$$

(Note that the integral of  $A(t)f'(t)$  on  $[r, r+1]$  is not affected by the possibly new value of  $A(t)$  at the single point  $r+1$ .) Also,  $A(t) = A(n)$  for  $n \leq t \leq x$ , so

$$A(x)f(x) - A(n)f(n) = A(n)[f(x) - f(n)] = \int_n^x A(t)f'(t) dt,$$

and similarly

$$A(m)f(m+1) - A(y)f(y) = A(m)[f(m+1) - f(y)] = \int_y^{m+1} A(t)f'(t) dt.$$

Using these identities to substitute for  $A(m)f(m+1)$  and  $A(n)f(n)$  in (1.1), we obtain the stated identity.  $\square$

Most of the time we shall apply the formula to sums starting at  $r = 1$ . Two variant forms for this case are given in the next result.

**Proposition 1.3.6** *Let  $f$  have a continuous derivative on  $[1, x]$ . Then:*

- (i)  $\sum_{r \leq x} a(r)f(r) = A(x)f(x) - \int_1^x A(t)f'(t) dt,$
- (ii)  $\sum_{r \leq x} a(r)[f(x) - f(r)] = \int_1^x A(t)f'(t) dt.$

*Proof* (i) Take  $y = 1$  in 1.3.5. Then  $A(y)f(y) = a(1)f(1)$ , while the sum on the left is  $\sum_{2 \leq r \leq x} a(r)f(r)$ . When combined, this gives the required sum.

(ii) This follows, because  $\sum_{r \leq x} a(r)f(x) = A(x)f(x)$ .  $\square$

Sometimes we will be interested in a function  $f$  that is undefined at 1 (typically  $f(t) = 1/\log t$ ). The following variant of the formula deals with this situation.

**Proposition 1.3.7** *If  $f$  has a continuous derivative on  $[2, x]$  and  $a(1) = 0$ , then*

$$\sum_{2 \leq r \leq x} a(r)f(r) = A(x)f(x) - \int_2^x A(t)f'(t) dt.$$

*Proof* Taking  $y = 2$  in 1.3.5, we have

$$\begin{aligned} \sum_{2 \leq r \leq x} a(r)f(r) &= a(2)f(2) + \sum_{2 < r \leq x} a(r)f(r) \\ &= a(2)f(2) + A(x)f(x) - A(2)f(2) - \int_2^x A(t)f'(t) dt. \end{aligned}$$

Since  $a(1) = 0$ , we have  $A(2) = a(2)$ , so the statement follows.  $\square$

To assist in the process of getting used to these formulae, we list a number of particular cases (with general  $a(r)$  but specific  $f(t)$ ). We remark that when similar expressions appear in later sections, the notation will often be  $n$  instead of  $r$ . The reader may like to write out and compare some of the corresponding discrete expressions.

- (i)  $\sum_{r \leq x} \frac{a(r)}{r} = \frac{A(x)}{x} + \int_1^x \frac{A(t)}{t^2} dt;$
- (ii)  $\sum_{y < r \leq x} a(r) \log r = A(x) \log x - A(y) \log y - \int_y^x \frac{A(t)}{t} dt;$
- (iii)  $\sum_{r \leq x} ra(r) = xA(x) - \int_1^x A(t) dt;$
- (iv)  $\sum_{r \leq x} (x-r)a(r) = \int_1^x A(t) dt;$
- (v) if  $a(1) = 0$ , then  $\sum_{2 \leq r \leq x} \frac{a(r)}{\log r} = \frac{A(x)}{\log x} + \int_2^x \frac{A(t)}{t(\log t)^2} dt.$

**Example 2** To estimate  $S(x) = \sum_{r \leq x} a(r)/r$ , given that  $|A(x)| \leq Cx$  for all  $x$ . By (i), we have

$$|S(x)| \leq C + \int_1^x \frac{C}{t} dt = C(\log x + 1).$$

Continuous Abel summation has applications to infinite series exactly analogous to those of discrete Abel summation, as follows.

**Proposition 1.3.8** *Suppose that  $f$  has a continuous derivative on  $[1, \infty)$ , and that  $A(x)f(x) \rightarrow 0$  as  $x \rightarrow \infty$ . Then*

$$\sum_{r=1}^{\infty} a(r)f(r) = - \int_1^{\infty} A(t)f'(t) dt,$$

*in the sense that if either side converges, then so does the other, to the same value. Further, we then have*

$$\sum_{r>x} a(r)f(r) = -A(x)f(x) - \int_x^{\infty} A(t)f'(t) dt.$$

*Proof* The first statement follows from 1.3.6(i) on considering limits as  $x$  tends to  $\infty$ . The second statement follows similarly from 1.3.5, or by subtracting the identity in 1.3.6(i).  $\square$

Usually we will use this in the direction of deducing convergence of the series from that of the integral, as in the following corollary.

**Corollary 1.3.9** *Suppose that:*

- (i)  $|A(t)| \leq C$  for all  $t$ ,
- (ii)  $f(t) \rightarrow 0$  as  $t \rightarrow \infty$ ,
- (iii)  $\int_1^{\infty} |f'(t)| dt$  is convergent (say to  $I$ ).

*Then  $\sum_{r=1}^{\infty} a(r)f(r)$  is convergent (say to  $S$ ), and  $|S| \leq CI$ .*

*Proof* This follows from 1.3.8 and the general fact that if  $\int_1^{\infty} |g(t)| dt$  converges (say to  $J$ ), then so does  $\int_1^{\infty} g(t) dt$ , with value not greater than  $J$  in modulus. (Again, this remains valid in the complex case.)  $\square$

So we have, for example (given condition (i)):

$$\sum_{n=1}^{\infty} \frac{a(n)}{n} = \int_1^{\infty} \frac{A(t)}{t^2} dt.$$

Except when  $f(t)$  is real and decreasing, the integral estimate for  $|S|$  in 1.3.9 is usually more useful than the discrete one of 1.3.4, because it is easier to evaluate the integral than the discrete sum.

Finally, we apply Abel's formula to some specific choices of  $a(r)$ . As yet, our repertoire of arithmetic functions is very limited, but two cases are well worth mentioning. First, take  $a(r) = 1$  for all  $r$ , so that the sum

$\sum_{r \leq x} a(r)f(r)$  is simply  $\sum_{r \leq x} f(r)$ . Then  $A(x) = [x]$ , and the two identities in 1.3.6 become

$$\sum_{r \leq x} f(r) = [x]f(x) - \int_1^x [t]f'(t) dt,$$

$$\sum_{r \leq x} [f(x) - f(r)] = \int_1^x [t]f'(t) dt.$$

Second, consider sums of the form  $\sum_{p \in P[x]} f(p)$ . Recall that  $u_P(r)$  is 1 for  $r$  prime, 0 otherwise. With this notation, the stated sum can be rewritten  $\sum_{r \leq x} u_P(r)f(r)$ . Clearly,  $\sum_{r \leq x} u_P(r) = \pi(x)$ , so by 1.3.6(i),

$$\sum_{p \in P[x]} f(p) = \sum_{r \leq x} u_P(r)f(r) = \pi(x)f(x) - \int_1^x \pi(t)f'(t) dt.$$

Since  $\pi(t) = 0$  for  $t < 2$ , we can amend the interval of integration to  $[2, x]$ .

**Example 3** To estimate  $S(x) = \sum_{r \leq x} \log(x/r)$ .

By the second of the formulae just given for the case  $a(r) = 1$ ,

$$S(x) = \sum_{r \leq x} (\log x - \log r) = \int_1^x \frac{[t]}{t} dt.$$

To estimate this, write  $t = [t] + \{t\}$ , where  $0 \leq \{t\} < 1$ . Then

$$S(x) = \int_1^x \frac{t - \{t\}}{t} dt = (x - 1) - q(x),$$

where

$$q(x) = \int_1^x \frac{\{t\}}{t} dt,$$

so that

$$0 \leq q(x) \leq \int_1^x \frac{1}{t} dt = \log x.$$

### Exercises

- Express  $\sum_{p \in P[x]} (1/p)$  in terms of  $\pi(x)$ .
- Let  $S(x) = \sum_{r \leq x} a(r) \log r$ . Show that if  $0 \leq A(x) \leq C$  for all  $x \geq 1$ , then  $|S(x)| \leq C \log x$  for all  $x \geq 1$ . (Use either discrete or continuous Abel summation, or both!)

- 3 Suppose that  $|a(r)| \leq 1$  and  $|A(n)| \leq C$  for all  $n$ . Using the appropriate integral expression, together with the fact that  $|A(t)| \leq t$  for  $1 \leq t \leq C$ , show that

$$\left| \sum_{r=1}^{\infty} \frac{a(r)}{r} \right| \leq \log C + 1.$$

- 4 Assume (or prove) that if  $B(n) \rightarrow B$  as  $n \rightarrow \infty$ , then

$$\frac{1}{n}[B(1) + \cdots + B(n)] \rightarrow B \quad \text{as } n \rightarrow \infty.$$

Now suppose that  $\sum_{r=1}^{\infty} a(r)/r$  is convergent. Write  $a(r)/r = b(r)$ . Use discrete Abel summation to give an expression for  $A(n)$  in terms of  $B(n)$ , and deduce that  $A(n)/n \rightarrow 0$  as  $n \rightarrow \infty$ .

- 5 Suppose that  $|A(t)| \leq C$  for all  $t$ , and that  $f(t)$  is differentiable and decreasing and tends to 0 as  $t \rightarrow \infty$ . Prove that, for all  $x \geq 1$ ,

$$\left| \sum_{r>x} a(r)f(r) \right| \leq 2Cf(x).$$

- 6 Prove that:

(a) if  $|A(x)| \leq Cx/(\log x)^2$  for all  $x \geq 2$ , then  $\sum_{r=1}^{\infty} a(r)/r$  is convergent;

(b) if  $a(r) \geq 0$  for all  $r$  and  $A(x) \geq Cx/\log x$  for all  $x \geq 2$ , then  $\sum_{r=1}^{\infty} a(r)/r$  is divergent.

- 7 Let  $b(r) = a(r) \log r$ . Show that:

(a) if  $|A(x)| \leq Cx$  for all  $x \geq 1$ , then  $|B(x)| \leq Cx(\log x + 1)$  for all  $x \geq 1$ ;

(b) if  $a(1) = 0$  and  $|B(x)| \leq Cx \log x$  for all  $x \geq 2$ , then  $|A(x)| \leq C(x + \text{li}(x))$  for all  $x \geq 2$ .

- 8 Prove that, for any real  $x > 1$ ,

$$\sum_{r>x} \frac{1}{r^2} \leq \frac{1}{x} + \frac{1}{x^2}.$$

- 9 Show that for any real  $x$ , the partial sums  $\sum_{r=1}^n \sin rx$  are bounded for all  $n$  (use an identity for  $\cos(r - \frac{1}{2})x - \cos(r + \frac{1}{2})x$ ). Deduce that the series  $\sum_{r=1}^{\infty} \frac{1}{r} \sin rx$  is convergent.

## 1.4 Estimation of sums by integrals; Euler's summation formula

### *Basic integral estimation*

The use of an integral to estimate a discrete sum is a very useful technique, which we shall apply repeatedly. In its simplest form, the relationship is described in the following result.

**Proposition 1.4.1** *Suppose that  $f$  is a decreasing function on  $[m, n]$ , where  $m, n$  are integers. Then*


$$f(m+1) + \cdots + f(n) \leq \int_m^n f(t) dt \leq f(m) + \cdots + f(n-1).$$

*The opposite inequality applies if  $f$  is increasing.*

*Proof* Suppose that  $f(t)$  is decreasing. For  $r-1 \leq t \leq r$ , we have  $f(r) \leq f(t) \leq f(r-1)$ . So, by integration on this interval of length 1,

$$f(r) \leq \int_{r-1}^r f(t) dt \leq f(r-1).$$

By adding these inequalities for  $r = m+1, \dots, n$ , we obtain the statement.



(This statement is well illustrated by the picture. The integral is the area under the curve, while the two sums are the total area of the small and big rectangles respectively.)  $\square$

For  $x \geq 1$  (not necessarily an integer), write

$$S(x) = \sum_{1 \leq r \leq x} f(r), \quad I(x) = \int_1^x f(t) dt.$$

Taking  $m = 1$  in 1.4.1, we obtain, in the case when  $f(t)$  is decreasing,

$$f(2) + \cdots + f(n) \leq \int_1^n f(t) dt \leq f(1) + \cdots + f(n-1),$$

in other words,

$$S(n) - f(1) \leq I(n) \leq S(n - 1), \quad (1.3)$$

or, equivalently,

$$I(n + 1) \leq S(n) \leq I(n) + f(1), \quad (1.4)$$

with opposite inequalities when  $f(t)$  is increasing. This shows how  $I(n)$  can be used to give bounds for  $S(n)$  when  $n$  is an integer. We shall often require estimates for  $S(x)$  in terms of  $I(x)$  when  $x$  is not an integer. Slightly different statements are appropriate for decreasing and increasing functions, as follows.

**Proposition 1.4.2** *Suppose that  $f(t)$  is non-negative and decreasing for all  $t \geq 1$ , and define  $S(x)$ ,  $I(x)$  as above. Then for all  $x \geq 1$ ,*

$$I(x) \leq S(x) \leq I(x) + f(1).$$

*Proof* Suppose that  $n \leq x < n + 1$ , where  $n$  is an integer. Then  $S(x) = S(n)$  and  $I(n) \leq I(x) \leq I(n + 1)$ . The statement now follows from (1.4).  $\square$

**Example 1** Let  $S(x) = \sum_{r \leq x} \frac{1}{r}$ . Since

$$\int_1^x \frac{1}{t} dt = \log x,$$

we have  $\log x \leq S(x) \leq \log x + 1$ .

**Proposition 1.4.3** *Suppose that  $f(t)$  is non-negative and increasing for  $t \geq 1$ . Define  $S(x)$ ,  $I(x)$  as above. Then for all  $x \geq 1$ , we have  $S(x) = I(x) + r(x)$ , where  $|r(x)| \leq f(x)$ .*

*Proof* We now have  $S(n) - f(n) \leq I(n) \leq S(n) - f(1)$  (the reverse of (1.3)). Let  $n \leq x < n + 1$ . Then

$$S(x) = S(n) \leq I(n) + f(n) \leq I(x) + f(x).$$

Since  $f(t)$  is increasing and  $x - n < 1$ , we have  $\int_n^x f(t) dt \leq f(x)$ , and hence

$$I(x) \leq I(n) + f(x) \leq S(n) - f(1) + f(x) \leq S(x) + f(x). \quad \square$$

**Note** Of course, similar results apply if the sum and integral both start at 0 instead of 1 (in 1.4.2,  $f(1)$  is then replaced by  $f(0)$ ).

**Example 2** Let  $S(x) = \sum_{r \leq x} \log r$ . Then

$$I(x) = \int_1^x \log t \, dt = x \log x - x + 1,$$

and  $|S(x) - I(x)| \leq \log x$ .

Even when we know  $S(n)$  explicitly, as we do when  $f(t) = t$ , 1.4.3 gives a quick route to the estimation of  $S(x)$  for non-integral  $x$ .

Returning to the case where  $f(t)$  is decreasing, we derive the “integral test” for convergence of an infinite series.

**Proposition 1.4.4** *Let  $f(t)$  be decreasing and non-negative for all  $t \geq 1$ . Then the series  $\sum_{r=1}^{\infty} f(r)$  is convergent if and only if the integral  $\int_1^{\infty} f(t) dt$  is convergent. If the sum of the series is  $S$  and the value of the integral is  $I$ , then  $I \leq S \leq I + f(1)$ .*

*Further, if we write*

$$S^*(x) = \sum_{r > x} f(r), \quad I^*(x) = \int_x^{\infty} f(t) \, dt,$$

*then for integers  $n$ ,*

$$I^*(n+1) \leq S^*(n) \leq I^*(n),$$

*and for general  $x$ ,*

$$I^*(x) - f(x) \leq S^*(x) \leq I^*(x) + f(x).$$

*Proof* Note that convergence of the series (or integral) means that  $S(n)$  (or  $I(n)$ ) tends to a limit as  $n \rightarrow \infty$ . Both  $S(n)$  and  $I(n)$  increase with  $n$ , so they tend to a limit if and only if they are bounded above. By (1.3),

$$S(n) - f(1) \leq I(n) \leq S(n - 1).$$

If the integral converges to  $I$ , then  $I(n) \leq I$  for all  $n$ , hence  $S(n) \leq I + f(1)$ . So  $S(n) \rightarrow S$ , where  $S \leq I + f(1)$ . Conversely, if the series converges to  $S$ , then  $I(n) \leq S(n) \leq S$  for all  $n$ , so  $I(n) \rightarrow I$ , where  $I \leq S$ .

By 1.4.1 again, for integers  $n, p$  with  $n < p$ ,

$$\int_{n+1}^{p+1} f(t) \, dt \leq f(n+1) + \cdots + f(p) \leq \int_n^p f(t) \, dt.$$

By letting  $p \rightarrow \infty$ , we obtain the stated inequalities for  $S^*(n)$ . Now let  $n \leq x < n+1$ . Then  $S^*(x) = S^*(n) \geq I^*(n+1)$ . Now

$$I^*(n+1) = I^*(x) - \int_x^{n+1} f(t) \, dt,$$