

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)

## Index

---

*No references are given here to the mention of a topic in the résumés at the end of each chapter.*

- A $\beta$  cutaneous afferents
  - conduction velocity, 77, 400, 418
  - electrical threshold, 400, 418
  - RII reflex, 414
  - skin stimulation mimicking mixed nerve stimulation, 77, 85, 204, 253, 299
  - tactile sensation, 391, 400
- abdominal skin reflexes
  - afferent pathway, 400
  - central pathway, 395, 400
  - convergence of tactile and noxious afferents, 402
  - functional organisation, 402
  - habituation, 394, 395
  - methodology, 394
  - normal descending facilitation, 433
  - protective function, 399, 401–2
  - reciprocal organisation, 402
  - spatial/temporal summation, 394
  - voluntary contraction, 412
  - upper motoneurone lesions, 433
- acetylcholine
  - neuromuscular junction, 151
  - recurrent collaterals, 151, 152
- activity-dependent hyperpolarisation of Ia
  - afferents, 12–13, 341
- A $\delta$  cutaneous afferents
  - conduction velocity, 400
  - electrical threshold, 400
  - pain sensation, 391, 399–400
  - RIII reflex, 399–400
- afferent conduction times used to suggest
  - heteronymous Ia excitation, 70–2
  - Ib inhibition, 253–5

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**602 Index**

- afferent feedback accompanying a movement, 515–16
  - cutaneous, 388, 516
  - joint, 272, 516
  - Ia–group II, 133–5, 136, 515–16
  - Ib, 267–8, 516
  - state-dependent modulation of sensory feedback, 530
- afferent fibres, *see* cutaneous, FRA, group Ia, group Ib
  - group II, joint afferents
- afterhyperpolarisation (AHP) of motoneurones
  - on-going EMG, 28
  - paired H reflex technique, 156, 157–9, 161, 178
  - PSTHs, 31, 32, 34, 37, 257
  - spatial facilitation, 47
  - unitary H reflex, 39
- alpha ( $\alpha$ ) motoneurone
  - $\alpha$  axon, motor threshold ( $1 \times MT$ ), 6, 7
  - final common path, 1
- alpha motoneurone excitability
  - methodology, 561–3
  - studies in patients
    - Parkinson's disease, 584–6
    - spasticity, 560–3
    - spinal spasticity, 580
    - stroke, 576
- alpha motoneurone, excitation through
  - cervical propriospinal neurones, 460–3
  - early cutaneous reflexes, 427–30
  - group II interneurones: heteronymous, 304–5
  - homonymous, 304
  - lumbar propriospinal neurones, 494–6
  - monosynaptic Ia connections: heteronymous, 81–6;
    - homonymous, 79
  - Ib excitatory interneurones, 258
  - transcortical cutaneous pathways, 421–4, 430–2
  - transcortical Ia pathways, 90–2, 548–9
  - withdrawal reflexes, 399–414
- alpha motoneurone, inhibition through
  - cutaneous non-noxious spinal reflexes, 429
  - non-reciprocal group I inhibition (wrist), 211–14, 522–4
  - Ib inhibition (hinge joint), 256–8
  - reciprocal Ia inhibition, 211
  - recurrent inhibition, 169–71: heteronymous, 169–70;
    - homonymous, 169
  - withdrawal reflexes, 405
- alpha motoneurones, recruitment order
  - cervical propriospinal excitation, 471
  - corticospinal excitation through cervical propriospinal neurones, 471, 519
  - cutaneomuscular responses, 424–7, 519
- F wave, 22
- H reflex, 4
- MEP: at rest, 45; during contraction, 44–5
- monosynaptic Ia excitation, 3–4, 79–81
  - exceptions, 79, 81
  - reversed by presynaptic inhibition, 347–8; functional implications, 348
- most reflex actions, 4
- slowing eccentric contractions, 519
- voluntary contraction (EMG), 25, 518
- alpha-gamma co-activation, *see* servo-assistance
- alpha-gamma linkage in reciprocal Ia inhibition
  - postulated from connections, 201
  - voluntary contraction, 198, 219, 223
- amyotrophic lateral sclerosis, changes in spinal pathways
  - presynaptic inhibition of Ia terminals, 369
  - recurrent inhibition, 184, 186, 581
- anaesthetic blockade
  - A $\delta$  afferents, responsible for the RIII reflex, 393, 400
  - cutaneous afferents, group II excitation, 317, 318
  - $\gamma$  efferents, 119; principle, 118–19
  - group I afferents
    - natural reciprocal inhibition, 217
    - non-reciprocal group I inhibition (wrist), 525
    - presynaptic inhibition, 220, 361
    - reciprocal Ia inhibition, 220
    - unloading during walking, 315, 316
- ankle and hip strategies in bipedal stance, 542
- ascending projections
  - A $\beta$  cutaneous afferents, 421, 549
  - A $\delta$  cutaneous afferents, 411
  - cervical propriospinal neurones, 454, 479, 531
  - feedback inhibitory interneurones to cervical propriospinal neurones, 454
  - FRA, 388
  - group Ia afferents, 90–2, 548
  - motor learning, 530–1
  - Ib interneurones, 246–7
  - Renshaw cells, 154
- Ashworth score, 573
- autogenetic inhibition, 244, *see* group Ib inhibition
  - from homonymous muscle, 244; from extensors, 246
  - not responsible for the clasp-knife phenomenon, 244–5
- Babinski response, 433–4
  - dysfunction of the upper motor neurone, 434
  - extensor hallucis longus, 402, 403, 434
  - late Babinski response after spinal transection, 433
  - methodology, 392, 402, 434

- normal plantar response, 402–3, 404
- pathophysiology, 434
  - disinhibition of withdrawal reflex, 434
  - suppression of normal response, 433, 434
- background from animal experiments
  - cervical propriospinal pathways, 452–5
  - cutaneous reflexes, 385–91
    - FRA reflex pathways, 388–91
    - low-threshold-induced cutaneous reflexes, 385–7
    - withdrawal reflexes, 387
  - group Ib pathways, 244–8
  - group II pathways, 288–93
  - lumbar propriospinal pathways, 490–1
  - monosynaptic Ia excitation, 64–6
  - muscle spindles and fusimotor drive, 113–17
  - post-activation depression, 96
  - presynaptic inhibition of Ia terminals, 337–40
  - reciprocal Ia inhibition, 197–201
  - recurrent inhibition, 151–4
- bag intrafusal fibres, 114
  - bag<sub>1</sub> fibres, 114, 115, 117
  - bag<sub>2</sub> fibres, 114, 115, 289
- ball joint, *see* organisation of spinal circuitry at wrist level
- ballistic contraction, recurrent inhibition, 176
- beta ( $\beta$ , skeleto-fusimotor) efferents, 117
  - background from animal experiments, 115, 117: dynamic, 117; static, 117
  - evidence in humans, 117
- beta( $\beta$ ) motoneurons, 117
- bidirectional connections
  - monosynaptic Ia excitation, 73, 74
  - non-reciprocal group I inhibition (wrist), 205, 206
  - Ib excitation, 258
  - reciprocal Ia inhibition, 205
- bi-triphasic effects
  - cutaneous responses to non-noxious stimuli
    - bi-triphasic effects in the lower limb, 415
    - triphasic effects in the upper limb, 415
  - disynaptic reciprocal inhibition-presynaptic inhibition, 343–4
  - group I-group II excitations, 293, 297, 301
  - non-reciprocal group I inhibition-Ib excitation (wrist), 258
  - Ia excitation-Ib inhibition, 249, 253, 255–6, 257
  - Ia excitation-recurrent inhibition, 35, 163, 164, 166–7, 168–9, 170
  - reciprocal Ia inhibition-Ib excitation, 255, 258
- blockade of one mechanism controlling the stretch reflex, 560
- botulinum toxin, 556, 559
- braking an eccentric contraction, 521
- central delay in spinal pathways
  - cervical propriospinal excitation, 457, 458, 459
  - cutaneomuscular spinal responses, 419
  - feedback inhibition of cervical propriospinal neurones, 458–9, 464
  - group II excitation, 303
  - lumbar propriospinal excitation, 491–2, 493
  - monosynaptic Ia excitation: homonymous, 66–7, 70; heteronymous, 73
  - non-reciprocal group I inhibition (wrist), 205, 206
  - Ib excitation, 255
  - Ib inhibition, 253–5
  - presynaptic inhibition, 339; D1 inhibition, 343, 344
  - reciprocal Ia inhibition, 205
  - recurrent inhibition, 166, 171, 206
  - transcortical cutaneous responses, 421–3
  - withdrawal (early) responses, 400–1
  - withdrawal (late) responses in
    - normal subjects, 410
    - patients with spinal transection, 408
- central delay inferred from
  - afferent conduction times
    - cutaneomuscular spinal responses, 419
    - heteronymous monosynaptic Ia excitation, 70–2
    - Ib inhibition, 253–5
    - reciprocal Ia inhibition, 205
- bidirectional connections
  - heteronymous monosynaptic Ia excitation, 73
  - non-reciprocal group I inhibition (wrist), 205, 206
  - reciprocal Ia inhibition, 205
- delay with respect to monosynaptic Ia excitation
  - cervical propriospinal excitation, 459
  - lumbar propriospinal excitation, 493
  - group II excitation, 303
  - Ib inhibition, 253–5
  - recurrent inhibition, 166
- delay with respect to reciprocal Ia inhibition
  - Ib excitation, 255
  - presynaptic inhibition of Ia terminals, 344
- cerebral palsy
  - contribution of spasticity to motor impairment, 559, 574
  - reciprocal Ia excitation, 233, 579
  - reciprocal Ia inhibition, 233, 579
  - recurrent inhibition, 187
- cervical propriospinal relay, background from animal experiments, 452–5
- conflicting results in the monkey, 454–5
  - apparent weakness, 454–5

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**604 Index**

- cervical propriospinal relay (*cont.*)
  - disclosure with strychnine, 455
  - evidence for a functional propriospinal system, 455
- connections in the cat, 453, 452–3, 454
  - ascending projections, 454; efference copy, 454
  - excitatory convergence, 452: from descending tracts, 453; from peripheral afferents, 453
  - diverging projections, 453: on motoneurons, 453; Ia interneurons, 453
  - inhibitory connections, 453–4: feedback, 454; feedforward, 453; inhibitory propriospinal neurons, 453
- function, 454
  - co-ordinated synergies, 453, 529
  - target-reaching, 454
  - updating, 454, 529
- phylogenetic differences, 455, 479
- cervical propriospinal relay, methodology, 455–60
  - critique of the tests, 457–8, 459
  - cutaneous suppression, 458–9
    - evidence for propriospinally mediated effect
    - normative values: MEP, 459, 474; ongoing EMG, 459
    - stimulation and recording, 458, 481: single shocks and trains, 481; stimulus graded with respect to motor response, 458, 481; window of analysis, 458–9
    - suppression of the MEP, 459, 473
    - suppression of the on-going EMG, 458–9, 471–2, 473
  - underlying principles, 458, 471
- evidence for propriospinally mediated effect
  - cutaneous suppression: central delay, 458, 471, 472; disfacilitation, 459, 471–3
  - group I excitation, 455, 474: central delay, 457, 459; diffuse pattern of input, 457, 460; disappearance when stimulation is increased, 457, 464, 467; low threshold, 457, 460
  - lesion at C6–C7 spinal level, 479–81
  - rostral location of the relevant neurons, 459–60: corticospinal excitation, 463; corticospinal inhibition, 467; cutaneous suppression, 459, 464, 473; group I excitation, 458, 459
  - group I excitation, facilitation of, 455–8: H reflex, 457, 472; MEP, 456, 457; on-going EMG, 456, 457; PSTHs, 456, 457
  - how to disclose propriospinally mediated corticospinal excitation, 468
- cervical propriospinal relay, organisation, 460–71
  - afferent excitatory input, 460–1
    - afferents responsible: group I, 460; including Ia, 460; cutaneous, 460
    - low threshold, 460
    - weakness, 460–1
    - widespread sources, 457, 460, 461
- corticospinal excitation, 461–2, 463
  - initial sparing, 462, 463
  - peripheral inhibition, 463
  - rostral location, 463
  - spatial facilitation: PSTH, 461–3; reflex, 463
- divergence, 469, 476, 529
- excitatory convergence, 469
  - corticospinal and afferent inputs, 460, 461, 469
  - different afferent inputs, 469
- feedback inhibition of propriospinal neurons, 463–7
  - afferent projections, 464: group I, 464, 465; cutaneous, 464, 465
  - corticospinal projections, 464–6, 467
- interactions between excitatory and inhibitory inputs, 467–8
  - corticospinal, 467; peripheral, 461, 467
  - explanation for the conflicting conclusions in primates, 467–8
  - natural vs. artificial stimulation of the system, 468
- lack of projections to motoneurons of intrinsic hand muscles, 460
- organisation in subsets, 468–9
- projections to motoneurons of different
  - muscles: distal, 460, 522; proximal, 460, 516–17, 521–2
  - types, 469–71: corticospinal excitation, 469–71; propriospinal neurons, 470, 471
- cervical propriospinal relay, physiological implications,
  - transmission of the descending command, 471–9
- cutaneous disfacilitation of the descending command, 471–4, 472
  - evidence for: ECR, 471–3; other motor nuclei, 473–4
  - quantitative contribution, 474
- facilitation of propriospinal neurons, 474–6
  - evidence for, 475
  - focused descending drive, 476
  - onset of contraction, 472, 474–5
  - tonic contraction, 476–7
- cervical propriospinal relay, possible function, 476–9
  - integration of peripheral and descending inputs, 477–8
    - control of force and speed of movement, 477, 518
  - integration at premotoneuronal level, 471
  - safety factor to transmission of command, 477
  - servo-assistance and diffuse distribution, 477, 516–17
  - synergies through the propriospinal system, 476, 529
- motor learning, 479, 531
- movements in which the system might be involved, 479
  - co-contraction of antagonists, 533

- expanded repertoire, 479
- multi-joint movements, 476, 478, 529
- rapid movements, 477, 479
- reaching, 529–30
- selectivity of movement
  - focus of descending drive, 476
  - lateral inhibition, 477, 517
- termination of movement, 475, 478, 518
- tonic contraction, 476–7
- cervical propriospinal relay, studies in patients, 479–85
  - C6–C7 lesion, 479–81: peripheral modulation of MEPs, 479–80, 481; recovery, 481
- Parkinson's disease, 484–5
  - asymmetry of cutaneous suppression of on-going EMG, 484
  - compensatory mechanism, 485, 592
  - increased transmission of the descending command, 485, 591
  - methodology, 484
- stroke, 481–4, 579
  - asymmetry of cutaneous suppression of on-going EMG, 481–3; disfacilitation, 483
  - changes throughout recovery, 482, 484
  - increased transmission of the descending command, 483, 579
  - methodology, 481
  - spasticity, 579
  - synkinetic contractions, 484, 579
  - underlying mechanisms, 483–4, 579
- chain intrafusal fibres, 114, 115, 117, 289
- changes in the recruitment gain of the H reflex, *see* recruitment gain
- clasp-knife phenomenon
  - characteristics, 558
  - differences from the lengthening reaction, 244
- responsible afferents
  - absence of involvement of muscle group II pathways, 326
  - FRA, 244, 245, 326
- clenching, synergy between ECR and finger flexors, 211–12, 525
- clonidine, *see* group II excitation in spinal spasticity
- clonus
  - $\alpha$ - $\gamma$  co-activation, 139–40
  - characteristics, 558
- co-contraction, *see* voluntary co-contraction
- coherence analysis of EMG/EMG or EEG/EMG signals, 48–9
  - coherence techniques, 48–9
  - cross-correlation, 48
  - use during movement, 512
- concentric contractions
  - muscle spindle afferent discharge, 135
- Ib afferent discharge, 267–8
- reciprocal inhibition, 222–3, 519
- servo-assistance through
  - $\alpha$ - $\gamma$  coactivation, 516
  - cervical propriospinal neurones, 477, 516–17
- conduction velocity
  - A $\beta$  cutaneous afferents, 77, 400, 418
  - A $\delta$  cutaneous afferents, 400
  - group Ia afferents, 36, 64, 72, 118, 302–3
  - group Ib afferents, 245
  - group II afferents, 302–3
  - group II/Ia ratio, 303
- contamination of the H reflex by Ib inhibition, 14–16, 67, 81
  - curtailment of Ia EPSPs, 14
  - evidence for Ib inhibition in quadriceps, 14–15
  - limitation of the H reflex, 14–16, 301–2, 310–12, 493, 500
- convergence
  - cervical propriospinal neurones
    - corticospinal and group I inputs, 461–3
    - various peripheral afferents, 469
  - feedback inhibitory interneurones to cervical propriospinal neurones
    - corticospinal and cutaneous inputs, 478, 483
    - corticospinal and group I inputs, 464–7
    - cutaneous and group I afferents, 469
  - feedback inhibitory interneurones to lumbar propriospinal neurones
    - corticospinal and group I inputs, 310, 500
    - corticospinal and group II inputs, 310
  - group I inhibitory interneurones (wrist), 212
    - group I inputs from antagonist and homonymous muscles, 212
    - group I inputs from elbow and antagonist muscles, 212
  - group II interneurones
    - corticospinal and group II inputs, 46–7, 307–9
    - group I and group II inputs, 305–6
    - other FRA afferents, 388, 514
- interneurones mediating cutaneous reflexes
  - A $\beta$  and A $\delta$  cutaneous afferents, 402, 405
- lumbar propriospinal neurones
  - corticospinal and peripheral inputs, 46–7, 307–9, 498
  - group I and group II inputs, 305–6
  - group I from different muscles, 47, 496
- Ia interneurones
  - excitation: Ia and corticospinal, 216; Ia and cutaneous, 214–15; Ia and vestibulospinal inputs, 216–17
  - inhibition from Renshaw cells, 171–3, 205–8
- Ib interneurones

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**606 Index**

- convergence (*cont.*),
  - first- and last-order interneurons, 267
  - multiple inputs, 265–7: from different muscles, 257–8; Ib and corticospinal, 265; cutaneous, 261–3; joint, 263–5; nociceptive, 265; Ia, 260–1; vestibulospinal, 265
  - various states: rest, 267; strong contractions, 265; weak contractions, 265–7
- PAD interneurons on Ia terminals
  - depression: corticospinal, 353; cutaneous, 350
  - facilitation: corticospinal, 353; group I, 348–50; vestibulospinal, 353
- Renshaw cells
  - excitation: reticulospinal, 186
  - inhibition: corticospinal, 172, 173
- spinal interneurons, 511
- cooling
  - evidence for group II excitation, 294, 297–9, 316, 317, 318
  - evidence for transcortical Ia pathway, 91
  - underlying principle, 297
- co-ordinated activation of various synergies
  - cervical propriospinal neurones
    - multi-joint movements, 476, 478, 529
    - reaching, 529–30
  - FRA hypothesis, 389, 513, 514, 529
  - hierarchical control schema, 527, 529–30
  - motor learning, 530–1
    - efference copy, 479, 531
    - fusimotor drive, 137, 138, 530
    - projections to ascending tracts, 530–1
  - spinal heteronymous connections, 527, 528–9
    - group II, 304–5, 512, 529, 538; flexible synergies, 529
    - Ia, 63, 93–4, 512, 528, 538, 550; flexible synergies, 94, 529; relaxation of antagonists, 528
  - state-dependent modulation of sensory feedback, 530
- cortical stimulation, general features
  - advantages, 45
  - comparison of responses to electrical and magnetic stimulation, 44, 424, 430–2
  - initial studies, 40
  - input–output relationship, 44
  - limitations, 44–5
  - mono- and non-monosynaptic transmission, 40
  - multiple descending volleys, 40–1, 309: D wave, 40, 43, 498; I waves, 40–1, 43, 44, 498
  - projection to motoneurons of different type: at rest, 45, 471; during contraction, 44–5
  - TMS-induced responses: lower limb, 43–4; upper limb, 43
  - trans-synaptic activation of pyramidal tract neurones by TMS, 43
- cortical stimulation, methodology
  - coils, 42–3
  - cross-talk, 39–40, 44
  - electrical stimulation, 40
  - modulation of PSTHs, 40, 41, 42
  - motor evoked potential (MEP), 39, 41, 42
  - recording, 39–40
  - stability of the coil, 44
  - transcranial magnetic stimulation (TMS), 42–3, 44
- cortical stimulation used for
  - demonstration of
    - non-monosynaptic transmission of the cortical command, 44, 307–9, 461–3, 498–500
    - presynaptic inhibition of Ia terminals, 44, 342, 343–4
    - transcortical cutaneous pathways, 424, 430–2
  - investigation of
    - corticospinal projections on spinal pathways, 44
    - excitability of corticospinal neurones, 44
- corticofugal syndrome, 557
- corticospinal control, background from animal experiments
  - $\alpha$  motoneurons (monosynaptic), 452, 454–5
  - cervical propriospinal neurones, 452, 454–5
  - cutaneous (private) pathways, 388
  - feedback inhibitory interneurons to cervical propriospinal neurones, 454, 455
  - $\gamma$  motoneurons, 114
  - group II interneurons, 292
  - lumbar propriospinal neurones, 491
  - Ia interneurons, 200
  - Ib interneurons, 248
  - PAD interneurons on Ia terminals, 339
  - PAD interneurons on Ib terminals, 248
  - Renshaw cells, 153
- corticospinal projections
  - $\alpha$  motoneurons
    - contralateral (monosynaptic), 40, 42, 463; possible function, 479
    - ipsilateral, 579
  - cervical propriospinal neurones, 461–3
  - feedback inhibitory interneurons to cervical propriospinal neurones, 464–7
  - feedback inhibitory interneurons to lumbar propriospinal neurones, 310, 500
  - $\gamma$  and  $\beta$  motoneurons, 130
  - group II interneurons, 46–7, 307–9
  - Ia interneurons, 216
  - Ib interneurons, 265; tonic, 577

- lumbar propriospinal neurones, 46–7, 307–9, 498–500
- non-reciprocal group I interneurons (wrist), 524
- PAD interneurons on Ia terminals, 350–3; tonic, 370, 576
- PAD interneurons on Ib terminals, 270
- Renshaw cells, 173
- cremasteric reflexes, 433
- critical firing stimulus, *see* unitary H reflex, 38–9
- critique of tests
  - $\alpha$  motoneurone excitability, 563
- cervical propriospinal excitation, 457–8, 459
- cutaneous reflexes, 396–8
- fusimotor drive, 118, 119, 126
- group II excitation, 299–302
- lumbar propriospinal excitation, 492–3, 500
- monosynaptic Ia excitation: heteronymous, 70, 72–3; homonymous, 70
- Ib inhibition, 255–6
- presynaptic inhibition of Ia terminals, 340, 344–5, 346; vibration of the homonymous tendon, 341, 368, 371, 565
- reciprocal Ia inhibition, 208–9
- recurrent inhibition: heteronymous, 168–9; homonymous, 160–1
- cross-talk
  - H reflex, 6
  - MEP, 39–40
- cumulative sum (CUSUM), 33–4
- cutaneomuscular responses, *see* cutaneous non-noxious spinal reflexes, transcortical cutaneous pathways
- cutaneous control of proprioceptive pathways
  - activation of cutaneous receptors during movement, 388, 516
- cervical propriospinal neurones
  - excitation, 460, 469
  - inhibition, 478, 518
- $\gamma$  motoneurons, 127–30, 138
- group II interneurons, 291, 306–7
- lumbar propriospinal neurones, 496
- Ia interneurons, 214–15
- Ib inhibitory interneurons
  - nociceptive, 265
  - tactile, 261–3: facilitation, 262; inhibition, 261–2
- PAD interneurons
  - late FRA, 408
  - tactile afferents, 350
- spinal pathways, 384, 385, 391
- cutaneous non-noxious spinal reflexes
  - afferent pathway, 418
  - central pathway, 418–21
    - oligosynaptic, 419, 427
    - private, 419–21: cutaneomuscular responses, 420; RII, 419
    - spinal, 418–19: cutaneomuscular responses, 419; monosynaptic reflex modulation, 419; RII, 418–19
  - E1 cutaneomuscular excitation in the upper limb, 417
    - distribution, 415, 427
    - functional implications, 428, 514
    - latency, 415
    - maturation, 423, 428
    - task-related changes, 427, 428
    - threshold, 415
  - early cutaneomuscular effects in the lower limb, 429
    - distribution, 415: E1 excitation, 415, 416, 429; inhibition, 416, 429
    - functional implications, 430, 514
    - task-related changes, 429–30
    - threshold, 415
  - monosynaptic reflex modulation, 415–18
    - FCR, 415–18: pattern, 415–16; underlying mechanisms, 416–18
    - lower limb, 397, 415, 419
  - projections to motoneurons of different type, 424–7
    - change in the recruitment gain, 425
    - effects of natural stimuli, 425, 426
    - effects on: H reflex, 425, 426; single units, 424–5, 426
    - functional implications, 425–7: grip, 425–7, 519, 534–5; locomotion, 427
  - RII reflexes at rest, 414
    - biceps femoris, 414
    - other muscles, 396, 417
- cutaneous reflexes at rest, *see* placing reaction, RII, RIII reflexes
- cutaneous reflexes, background from animal experiments, 385–91
  - FRA pathways, 388–91: early, 389–90; late, 390–1
  - initial findings, 385
  - modulation of the descending command, 388
  - private pathways of non-noxious reflexes, 385–7
    - corticospinal facilitation, 388
    - descending control, 388
    - forelimb, 387
    - stumbling reaction in locomotion, 387
    - toe extensor reflex, 385–7
  - projections to motoneurons of different type, 387
    - descending facilitation, 388
  - withdrawal reflexes, 387

## 608 Index

- cutaneous reflexes, methodology, 391–9
  - critique of tests, 396–8
    - encroachment upon other afferents, 398, 435
    - recordings: at rest, 398, 414; monosynaptic reflexes, 398, 418; on-going EMG, 398, 415; PSTHs, 398
    - stimuli: electrical, 398; mechanical, 396; radiant heat, 396–8
  - evidence for spinal reflexes
    - non-noxious reflexes, 418–19
    - withdrawal reflexes: early, 400–1; late, 407
  - habituation, 391: abdominal skin reflexes, 394; RII reflex, 394; withdrawal reflexes, 394
  - modulation of motoneurone excitability, 396
    - monosynaptic reflexes, 396
    - on-going EMG, 396, 404, 414–15
    - PSTHs, 396, 417
    - spike-triggered averaging of EMG, 396, 417
  - responses at rest, 394–6
    - RII, 394
    - RIII, 394; other withdrawal reflexes, 394
  - spatial and/or temporal summation required, 391, 392, 394, 396
  - stimuli, 391–4
    - cutaneomuscular responses, 392, 415
    - electrical, 391–2: on: nerves, 391–2; skin, 392; single shocks vs. trains, 392
    - mechanical, 392–4: abdominal reflexes, 394; plantar responses, 392; probe to indent the skin, 392
    - withdrawal reflexes, 392
  - underlying principles, 391
- cutaneous reflexes, studies in patients, 432–8
  - complete spinal transection, 433
    - Babinski response, 433
    - from spinal shock to spasticity, 433
    - late withdrawal reflexes, 407–10
  - grasp reflex, 436
  - Parkinson's disease, 436–7
    - transcortical inhibition, 437, 589
    - withdrawal reflexes, 436–7, 589
  - peripheral neuropathies, 437
  - upper motoneurone lesions
    - abolition of normal reflexes, 433
    - Babinski response, 433–4
    - cutaneomuscular responses, 423, 436
    - withdrawal reflexes: lower limb, 434–5; upper limb, 435
- cutaneous silent period
  - non-noxious stimuli
    - lower limb, 415, 416; spinal origin, 429
    - upper limb (II), 415, 417; transcortical origin, 423
  - noxious stimuli
    - afferent pathway, 401
    - central delay, 401
    - evidence for spinal pathway, 400–1
    - lower limb, 397, 404–5
    - Parkinson's disease, 436–7, 589
    - upper limb, 405, 406; evidence for motoneurone inhibition, 405, 406
    - upper motoneurone lesions, 435
- cutaneous suppression of the on-going EMG through
  - activation of
    - cutaneous non-noxious spinal pathways, 415, 429
    - cutaneous transcortical pathways, 415, 423
  - feedback inhibitory interneurons to cervical propriospinal neurones, 464
    - evidence for disfacilitation, 459, 471–3
    - methodology, 458, 481; window of analysis, 458–9
    - normative values, 459
    - pattern, 478
  - Ib interneurons, 271–2, 518
  - withdrawal reflex pathways
    - lower limb, 404–5
    - upper limb, 405
- D wave after cortical stimulation, 40, 41, 43
- D1 inhibition
  - critique, 344–5, 364, 365–6
  - early part, assessment of presynaptic inhibition of Ia terminals, 343–4, 367, 565–6
  - late part: lower limb, 344; upper limb, 344, 372
- D2 inhibition, 344
- decerebrate rigidity
  - common features with spasticity, 560
  - differences from spasticity, 560
  - underlying mechanisms, 560
- descending control of spinal pathways
  - normal, *see* cortico-, reticulo-, vestibulo-spinal control
  - spasticity, 571
- differences between spinal pathways in cervical and lumbar enlargements
  - cutaneomuscular responses: cervical, 415, 427; lumbar, 415, 429
  - function of  $\gamma$  drive: cervical, 137–8; lumbar, 137
  - heteronymous Ia excitation: cervical, 83–6; lumbar, 81–3
  - heteronymous Ib inhibition: cervical, 257; lumbar, 257
  - heteronymous recurrent inhibition: cervical, 167, 170–1; lumbar, 166, 169–70
- PAD interneurons
  - corticospinal control: cervical, 353; lumbar, 350–3



- focus drive, 351, 359–60
- stroke: cervical, 369; lumbar, 368–9
- voluntary contraction, 363
- propriospinal neurones
  - cutaneous inhibition: cervical, 458–9, 471–4; lumbar, 496
  - peripheral excitation: cervical, 460–1; lumbar, 494–6
  - voluntary contraction, cervical, 471–9; lumbar, 500–2
  - uncertainty about extrapolations, 511
- differences between spinal pathways to proximal and distal muscles, 521–2
- distal muscles, 522
- proximal muscles, 521–2
- disfacilitation of motoneurones
- disfacilitation vs. inhibition, 473
- feedback inhibition of cervical propriospinal neurones
  - differences with inhibition, 416–18, 473
  - evidence for disfacilitation, 459, 471–3, 483
  - site of disfacilitation, 473
- feedback inhibition of group II interneurones, 307, 310
- feedback inhibition of lumbar propriospinal neurones, 307, 310, 496, 500, 502
- relaxation of antagonists, 527, 528
- spasticity, inhibitory pathways, 571: Ia interneurones, 232, 570; Ib interneurones, 568
- suppression of on-going EMG, 25
- disinhibition of motoneurones
  - non-reciprocal group I inhibition, 214
  - reciprocal Ia inhibition, 222, 223, 225, 520
- disynaptic effects
  - excitation
    - cervical propriospinal excitation, 457
    - lumbar propriospinal excitation, 491–2
    - group II excitation, 302–4
  - inhibition
    - Ib inhibition, 253–5
    - non-reciprocal group I inhibition (wrist), 205, 206
    - reciprocal Ia inhibition, 205
    - recurrent inhibition, 166
- DOPA
  - disclosure of late FRA pathways in the spinal cat, 390
  - effect on group II excitation in spinal spasticity, 322
  - see also* Parkinson's disease, 582–92
- duration
  - group II excitation, 295, 300
  - Ib inhibition, 155, 255
  - monosynaptic Ia excitation, 261
  - presynaptic inhibition, 338
  - reciprocal Ia inhibition, 26–7, 201, 203–4
  - recurrent inhibition, 152, 159, 166, 171, 206
- dystonia
  - effects of rTMS, 372
  - non-reciprocal group I inhibition (wrist), 234, 372
  - presynaptic inhibition of Ia terminals, 371–2
    - correlation with dystonia, 372
    - D1 inhibition, 371–2: early phase, 371–2; late phase, 372
    - group III-induced presynaptic inhibition, 372
    - specific descending control, 372
    - unaffected side, 372
- E1, early cutaneomuscular excitation, *see* cutaneous non-noxious spinal reflexes
- E2, late cutaneomuscular excitation, *see* transcortical cutaneous pathways
- early discharge, 121, 298
- eccentric contractions
  - fusimotor drive, 135; walking, 544
  - reciprocal inhibition, 519
- servo-assistance through
  - $\alpha$ - $\gamma$  coactivation, 516
  - cervical propriospinal neurones, 516–17
- slowing contractions, 521
  - fusimotor drive, 521
  - recruitment order of motoneurones, 519
- walking, 544
- efference copy, 454, 479, 531
- electrical stimulation
  - afferent fibres, *see* threshold to electrical stimulation
  - cutaneous reflexes, 391–2
  - motor cortex, 40
- electrodes for recording
  - microneurography, 119
  - needle: F wave, 21; PSTH, 30, 31, 37; unitary H reflex, 37
  - surface: F wave, 23; H reflex, 5: bipolar, 5; monopolar, 6; MEP, 39; on-going EMG, 28; PSTH:30
  - surface differential, PSTH, 31
  - wire, PSTH:31, 37
- endings
  - cutaneous mechanoreceptors, 113; activation by vibration, 131
  - free nerve endings, 113
  - see also* Golgi tendon organs, primary endings, secondary endings

**610 Index**

- EPSP (excitatory post-synaptic potential)  
 curtailment of the compound Ia EPSP of the H reflex by  
 oligosynaptic IPSPs: 14–15  
 Ia EPSPs in motoneurons, 3; evoked by H and tendon reflex  
 afferent volleys, 118  
 rise time of the test Ia EPSP, 9  
 excitation, *see*  $\alpha$  motoneurone excitation; disinhibition of  
 motoneurons
- F wave, general features, 21–2, 24  
 all muscles, 22  
 amplitude, 23  
 chronodispersion, 23  
 interest in distal muscles, 22  
 latency, 23  
 motoneurons involved in the F wave, 22  
 variability and persistence, 22–3
- F wave, methodology  
 basic methodology, underlying principles, 21–2  
 used as a measure of motoneurone excitability,  
 23–4
- F wave, studies in patients  
 Parkinson's disease, 585  
 peripheral neuropathies, 24  
 proximal nerve lesions, 24  
 spasticity, 24, 562–3
- fatigue, compensation for muscle fatigue  
 recurrent inhibition, 174, 179, 515  
 spindle activation, 135, 512
- feedback  
 need for afferent feedback, 136–7  
 negative feedback to motoneurons, via Renshaw cells, 151  
 positive feedback to group II interneurons through the  $\gamma$   
 loop, 291, 323, 325, 567–8, 581
- feedback inhibition of cervical propriospinal neurones,  
 background from animal experiments  
 cat, 454: ascending projections, 454; corticospinal control,  
 454; function, 454  
 monkey, 455  
 phylogenetic differences, 455
- feedback inhibition of cervical propriospinal neurones,  
 methodology  
 cutaneous suppression of on-going EMG and MEP, 458–9  
 evidence for: disfacilitation of motoneurons, 459, 471–3;  
 inhibition of propriospinal neurones, 458–9
- feedback inhibition of cervical propriospinal neurones,  
 organisation  
 afferent input  
 group I, 464: heteronymous, 464; homonymous, 464, 465  
 cutaneous, 464: not involved in cutaneomuscular  
 responses, 420; pattern, 465, 478  
 convergence: corticospinal and cutaneous inputs, 478;  
 corticospinal and group I inputs, 464–7; cutaneous and  
 group I inputs, 469  
 corticospinal excitation, 464–6, 467  
 initial sparing, 466  
 reversal when increasing the input, 464–5; underlying  
 mechanism, 466  
 disfacilitation  
 disfacilitation vs. inhibition, 416–18, 473  
 evidence for disfacilitation: cutaneous, 459, 464, 471–3;  
 group I, 464  
 site of disfacilitation, 473  
 various motoneurone pools, 473–4  
 explanation for conflicting conclusions in primates, 467–8  
 interactions between excitatory and inhibitory inputs to  
 propriospinal neurones, 467–8: corticospinal, 467;  
 peripheral, 461, 467  
 projections to propriospinal neurones, 464, 467  
 feedback inhibition of cervical propriospinal neurones,  
 possible function  
 control of force and speed, 477, 518  
 natural vs. artificial stimulation of the system, 468  
 selectivity of movement, 477, 517  
 termination of movement, 478, 518
- feedback inhibition of lumbar propriospinal neurones  
 corticospinal excitation, 310, 500; disclosure, of connections  
 with corticospinal facilitation, 500  
 group I excitation, 307, 496–7: heteronymous, 496–7;  
 homonymous, 496  
 focus of the descending motor command, 502–3
- feedforward inhibition of cervical propriospinal neurones  
 cat, 453  
 humans, 467, 477  
 monkey, 455
- flexible synergies through  
 fusimotor drive, 135, 517  
 inhibition of group II excitation, 514, 538  
 lateral inhibition of cervical propriospinal neurones, 477,  
 517  
 Ib inhibition, 273, 517  
 presynaptic inhibition of Ia terminals, 94, 359–60, 517, 529  
 recurrent inhibition, 94, 183–4, 529, 538
- flexion reflexes, *see* withdrawal reflexes
- flexor spasms, 140, 434–5
- follow-up length servo, 114: postulated from connections, 114
- force of movement, spinal mechanisms contributing to,  
 517–18

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)

- feedback inhibition of cervical propriospinal neurones, 477, 518
- fusimotor drive, 135, 518
- non-reciprocal group I inhibition (wrist), 524
- Ib inhibition, 271, 518
  - dynamic control through Ia projections, 272
  - hinge joint movements, 517
  - maintenance of an operating level, 271, 515
  - smooth profile of force, 271, 515
  - tonic contractions, 515
- presynaptic inhibition of Ia terminals, 358, 359, 518
- reciprocal Ia inhibition, 224, 518
- recurrent inhibition, 179, 518
- FRA (flexion reflex afferents)
  - characteristics, 388–9
  - clasp-knife phenomenon, 244
  - critique of the concept, 389
  - reasons to group FRA, 388–9
    - ascending projections, 388
    - descending control, 388–9
    - extensive afferent convergence, 385, 388
- FRA hypothesis in movement, 386, 389, 513, 514, 529
- FRA pathways in the cat (early), 389–90
  - alternative FRA pathways, 388–9
  - possible functional role, 389
  - presynaptic inhibition of FRA, 390
  - projections to spinal interneurons, 390
    - group II interneurons, 292
    - Ia interneurons, 199
    - Ib interneurons, 247–8
    - PAD interneurons, 339
    - Renshaw cells, 153
- FRA pathways in the cat (late), 390–1
  - appearance with DOPA, 390
  - inhibition from early FRA pathways, 386, 390
  - inhibitory effects mediated via Ia interneurons, 390
  - mediated via specific interneurons, 390
  - mutual inhibition between flexors and extensors, 386, 390;
    - possible role in locomotion, 390
  - presynaptic inhibition of Ia afferents, 390
- frontal lesions
  - dishabituation, 436
  - grasp reflex, 436
- functional stretch reflex, 91, 539
- fusimotor drive, background from animal experiments, 113–17
  - dynamic  $\gamma$  drive, 114, 115
  - follow-up servo length, 114
  - initial investigations, 113
  - postulated role, 114
  - servo assistance, 114
  - static  $\gamma$  drive, 114, 115
- fusimotor drive, methodology, 117–26
  - comparison of tendon jerk and H reflex, 9, 117
  - differences in several aspects, 2–3, 117–18
  - critique of tests, 118, 119, 126
  - microneurography, 119–24: basic methodology, 119–20; bias
    - towards large axons, 123; identification of endings, 120–2; microelectrode, 119; uncertainties, 123–4
  - nerve blocks: anaesthetics, 119; pressure, 117, 119, 133
  - thixotropy, 122, 124–6: human studies, 124–5; perspectives, 126; principle, 124
- fusimotor drive, organisation, 127–30
  - background fusimotor drive, 127: dynamic, 127; static, 127, 128
  - corticospinal projections to  $\gamma$  and  $\beta$  motoneurons, 130
  - cutaneous projections to  $\gamma$  motoneurons
    - lower limb in: reclining, 127–8; standing, 128, 129
    - possible involvement in late cutaneomuscular responses, 420
    - upper limb, 129–30
- fusimotor drive, receptor-bearing muscle contraction, 133–6
  - activation of:  $\beta$ , 136;  $\gamma_d$ , 136;  $\gamma_s$ , 136
  - appearance of spindle discharge after the EMG, 125, 133
  - co-contraction of antagonists, 135, 532
  - concentric and eccentric contractions, 135
  - contraction of antagonists, 135
  - contribution to excitation of  $\alpha$  motoneurons in, 133–5
    - hinge joint movements, 135, 521
    - tonic contraction, 117, 122, 133, 512
  - decline of spindle activity during long-lasting contractions, 125, 135
  - excitation of synergist motoneurons through
    - heteronymous Ia and group II connections, 93–4, 512, 528–9, 550
  - focus on contracting muscles, 122, 135, 517
  - force-related, 135, 518
  - FRA hypothesis, 389, 513, 514
  - muscle fatigue, 135, 512
  - mutual reinforcing of Ia and group II excitations, 512–14
  - persistence of spindle discharge after EMG disappearance, 122, 124, 125
  - primary and secondary endings, 135, 512
  - reciprocal Ia inhibition, 221–3, 519
  - spindle activity during standing, 135, 537–8; cutaneous
    - control, 128
  - stretched antagonistic muscles, 135

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**612 Index**

- fusimotor drive, receptor-bearing muscle contraction, possible
  - function, 136–8
  - co-contraction, 138
  - cutaneous control, 138
  - need for afferent feedback, 136–7
  - separate control of  $\gamma_s$  and  $\gamma_d$  motoneurons, 138
  - slowing eccentric contractions, 521
- fusimotor drive, remote contractions (Jendrassik manoeuvre), 131–3
- fusimotor drive, servo-assistance
  - motor learning, 137, 138, 530
  - through
    - $\alpha$ - $\gamma$  coactivation, 516
    - cervical propriospinal neurones, 477, 516–17
  - unstable stance, 537–8
  - voluntary co-contractions of antagonists, 135, 532
  - voluntary movement, 135, 516
    - lower limb, 137
    - upper limb: distal muscles, 137–8, 522; proximal muscles, 521–2
  - voluntary tonic contraction, 512–14
  - walking, 544
- fusimotor drive, studies in patients, 138–41
  - Parkinson's disease, 140–1, 586
    - indirect evidence, 586
    - insufficient data, 586
    - microneurography, 586
    - rigidity, 140–1
    - tremor, 141
  - spasticity, 139–40, 563–5
    - animal models, 564
    - clonus, 139–40
    - flawed arguments for, 139, 564
    - how it could produce spasticity, 563
    - importance of contribution, 564–5
    - microneurography, 139, 564
    - possibility of occurrence, 139, 564–65
    - reciprocal Ia inhibition, 574
  - spinal spasticity, 140, 580
    - positive feedback through group II projections, 140, 325, 580
    - reflex activation of  $\gamma$  motoneurons, 140, 580
  - stroke, 139–40
    - contribution to motor deficit, 139, 140, 565
    - $\gamma$  activity, 139, 576
- GABA, 337
- gain hypothesis, *see* recurrent inhibition, 179
- gait, *see* running, walking
- gamma ( $\gamma$ ) drive, *see* fusimotor drive
- gamma efferents and motoneurons
  - background from animal experiments, 114:  $\gamma_d$ , 114;  $\gamma_s$ , 114
  - background  $\gamma$  drive, 127: dynamic, 127; static, 127
  - corticospinal projections, 130
  - cutaneous projections
    - lower limb: reclining, 127–8; standing, 128
    - upper limb, 129–30
  - possible role of positive  $\gamma$  feedback, 291, 325
- glycine, 198–9, 214, 233
- Golgi tendon organs
  - background from animal experiments, 245
  - origin of Ib afferents, 113, 245
  - sensitivity to vibration: cat, 245; humans, 123, 130–1, 245
  - voluntary contraction, 267–8
- grasp reflex, 436
- group I input to
  - cervical propriospinal neurones, 460
- feedback inhibitory interneurons to cervical propriospinal neurones, 464
- feedback inhibitory interneurons to lumbar propriospinal neurones, 307, 496–7
- group Ib excitatory interneurons, 255, 258
- group Ib inhibitory interneurons, 252–3, 256–8
- group II interneurons, 291, 305–6
- lumbar propriospinal neurones, 305–6, 494
- non-reciprocal group I inhibitory interneurons (wrist), 212
- Ia interneurons, 204–5
- PAD interneurons, 348–50
- group Ia afferents
  - background from animal experiments, 64
  - conduction velocity, 36, 64, 72, 118, 303
  - electrical threshold, 64, 67–9, 75, 204, 303
    - activity-dependent hyperpolarisation, 12–13, 341
    - elevation by tendon vibration, 76, 205, 245, 260–1
  - in the inferior soleus nerve, 69
  - large range in human experiments, 77–8; underestimation of Ia effects, 78
  - with respect to motor threshold, 81
- excitability cycle, 11
- monosynaptic projections on motoneurons, 66–77
  - motoneurons of different type, 79–81
  - presynaptic inhibition, 347–8
- projections on interneurons
  - cervical propriospinal neurones, 460
  - lumbar propriospinal neurones, 494
  - group II interneurons, 305–6
  - Ia interneurons, 204–5
  - Ib inhibitory interneurons, 260–1

- non-reciprocal group I interneurons (wrist), 212
- PAD interneurons, 350
- see also* transcortical Ia pathways
- group Ib afferents
  - conduction velocity, 245
  - electrical threshold
    - close to that of Ia afferents, 245
    - use of vibration to distinguish between Ia and Ib afferents:
      - cat, 245; humans, 251, 260–1
  - projections on interneurons
    - cervical propriospinal neurones, 460
    - lumbar propriospinal neurones, 494
    - non-reciprocal group I (wrist), 212
    - Ib excitatory, 255, 258
    - Ib inhibitory, 252–3, 256–8
    - PAD interneurons, 350
- group Ib excitation
  - background from animal experiments: trisynaptic excitation
    - of flexor motoneurons, 246
  - connections and organisation
    - cutaneous facilitation, 259, 262
    - not involved in the peroneal excitation of quadriceps
      - motoneurons, 258–60, 493
    - only found between strict antagonists, 258: lower limb, 250, 258; upper limb, 251, 258
  - methodology, 255: electrical threshold, 258; latency, 255; tendon tap, 251, 255
  - studies in spastic patients, 277–9, 569
    - contribution to adverse co-contraction of the antagonist, 279, 575
    - contribution to spasticity, 278–9, 569
    - evidence for increased Ib excitation, 230, 277–8
    - spinal cord injury, 277, 581
    - stroke, 277, 577
    - underlying mechanisms, 278, 577, 581
- group Ib inhibition (hinge joint), background from animal experiments, 244–8
- contraction-induced Ib inhibition, 248
- general features of Ib pathways, 245–6
  - absence of inhibition from Renshaw cells, 245–6
  - denomination, 245
  - reciprocal organisation, 245
- Golgi tendon organs and Ib afferents, 245
- initial findings, 244–5
- input, extensive convergence, 247–8: descending tracts, 248; peripheral afferents, 247–8
- post-activation depression, 248
- presynaptic inhibition, 247, 248
- projections to, 246–7
  - $\alpha$  motoneurons, 246, 247: alternative pathways, 246; dominant pattern, 246; extended pattern, 246; inverse myotatic reflex, 246
  - ascending tract neurones, 246–7
  - $\gamma$  motoneurons, 246
  - mutual inhibition of Ib interneurons, 246
  - reflex reversal during fictive locomotion, 248
- group Ib inhibition (hinge joint), methodology
  - critique of tests, 255–6
    - biphasic effects: presynaptic inhibition of Ia afferents, 255–6; underestimation, 257
    - electrical stimulation over tendons, 256
    - weakness of isolated Ib inhibition, 256
- evidence for Ib inhibition, 252–5
  - differences from reciprocal Ia inhibition, 255
  - disynaptic inhibition, 253–5, 261: H reflex, 253; PSTHs, 253–5
  - Ib origin, 252–3: absence of cutaneous effect, 253; ischaemia, 252; low electrical threshold, 250, 252; tendon tap, 251, 252–3
  - short duration, 255
- inhibition
  - monosynaptic reflexes, 249: biphasic effects, 249, 250; isolated inhibition, 249, 250, 251
  - PSTHs, 249–51, 252
- group Ib inhibition (hinge joint), organisation
  - control by joint afferents, 263–4, 265: function of protection, 263–5; heteronymous, 263; homonymous, 263
  - convergence of Ia afferents, 260–1
  - corticospinal control, 265
  - cutaneous control, 261–3
    - facilitation, 259, 262: not involved in cutaneomuscular responses, 420; pattern, 262, 272; rest, 262; reversal during contraction, 262
  - inhibition, 254, 261–2: mutual inhibition of interneurons, 254, 261–2; pattern, 261
- descending facilitation of first-order interneurons: 262, 267, 271
- multiple convergence, 265–6, 267
  - first- and last-order interneurons, 267
  - various states: rest, 267; strong contractions, 265; weak contractions, 265–7
- nociceptive control, 265: cutaneous, 265; muscle, 265
- organisation, pattern and strength, 256–8
  - convergence from different muscles, 257–8: lower limb, 257–8; upper limb, 258
  - divergence, 257, 258
  - heteronymous, 257: lower limb, 257, 260; upper limb, 257

## 614 Index

- group Ib inhibition (hinge joint), organisation (*cont.*)
  - homonymous, 256–7
  - organisation in subsets, 257
  - underestimation, 257
  - vestibulospinal control, 265
- group Ib inhibition (hinge joint), physiological tasks, 267–75
  - contraction of antagonistic muscles, 273, 520
  - contraction of synergists, 272–3; underlying mechanism, 269, 272
  - contraction of the target muscle, 268–71
    - distinct from changes in presynaptic inhibition of Ia terminals, 268–70
    - homonymous Ib inhibition, 268, 269; heteronymous Ib inhibition, 268, 269
    - onset, 269, 270
    - tonic, force-related, 268, 269, 515
    - underlying mechanisms, 268–71: descending, 270; peripheral, 270; presynaptic, 270; postsynaptic, 270
  - walking, 273–5
    - heteronymous Ib pathways to biceps femoris, 274, 275, 546
    - Ib pathways to extensors, 273–4; soleus, 273
- group Ib inhibition (hinge joint), possible function
- group I inhibition of antagonists, 273, 520
- Ib inhibition of active motoneurons, 271; hinge joint
  - movements, 517; tonic contraction, 515
  - contribution to force, 271, 518; dynamic control, 272; maintenance of an operating level, 271, 515; smooth profile of force, 271, 515
  - restoration by other afferents, 271–2
  - termination of movement via facilitation by: cutaneous afferents, 262, 271–2, 518; joint afferents, 263, 272, 518
- Ib inhibition of inactive synergists and selectivity, 273, 517
- walking
  - prevention of excessive reflex activity in soleus, 274, 546–7
  - stability of the knee, 274–5, 547
- group Ib inhibition (hinge joint), studies in patients, 275–7
  - hyperekplexia, 276
  - Ib inhibition vs. propriospinal excitation, 277, 568
  - Parkinson's disease (ankle), 276–7, 588
    - correlation with: rigidity, 276, 588
    - effect of treatment, 588
    - underlying mechanisms, 277, 588
  - spasticity (ankle), 276, 568–9
    - how it could produce spasticity, 568
    - importance of contribution, 569
    - methodology, 275–6, 568
  - spinal spasticity, 276, 277, 581
  - stroke, 276, 577: correlation with spasticity, 276; underlying mechanisms, 277, 577
- group II afferents
  - conduction velocity
    - group I/group II ratio: cat, 289; humans, 303
    - peripheral nerves, 289, 302–3: distal, 302, 303; proximal, 302, 303
    - within the spinal cord: cat, 289; humans, 302
  - electrical threshold
    - cat, 289
    - group II/group I ratio: cat, 289; humans, 303
    - high threshold, 293, 295, 297
  - projections (cat)
    - $\alpha$  motoneurons: monosynaptic, 289; via interneurons, 289–91
    - $\gamma$  motoneurons: monosynaptic, 291; via group II interneurons, 291
    - inhibition of Renshaw cells, 153
    - lumbar propriospinal neurons, 289, 494
    - mutual inhibition of interneurons, 291
- group II excitation, *see also* lumbar propriospinal pathways
- group II excitation, background from animal experiments, 288–93
- excitatory input from, 290, 291–2
  - contralateral group II afferents, 292
  - descending tracts, 292
  - ipsilateral group II afferents, 291; convergence-divergence, 291
  - other peripheral afferents, 291–2
- group II interneurons, 289
  - caudal interneurons, 289
  - rostral propriospinal neurons, 289
  - very effective transmission, 290
- initial findings, 288–9
- monoaminergic inhibitory control, 292
  - locus coeruleus, 292
  - projections to  $\gamma$  motoneurons, 292
  - selective control of group II transmission, 292
  - underlying mechanisms, 290, 292
- mutual inhibition of group II interneurons, 291, 292
- post-activation depression, 292–3
- presynaptic inhibition, 292
  - feedback control, 292
  - feedforward control, 292
- projections on  $\alpha$  motoneurons, 290, 291
  - alternative pathways, 288, 291
  - distribution extensors-flexors, 288, 290, 291
  - muscle origin of inhibition, 291

- projections on  $\gamma$  motoneurons, 290, 291; positive feedback, 291
- secondary endings and group II afferents, 289
- group II excitation, methodology, 293–302
- critique of tests, 299–302
  - contamination by group I effects, 14–15, 299–301
  - interactions at interneuronal level, 301; facilitation, 301; occlusion, 301
  - interactions at motoneuronal level, 301
  - limitation of the H reflex by Ib inhibition, 14–16, 301–2, 310–12, 493, 500
  - overlapping of group I and group II effects, 3, 301
  - stretch-induced responses in upright stance, 301
- evidence against excitation due to stimulation of cutaneous afferents, 299
- fusimotor or motor axons, 297, 298
- joint afferents, 299
- evidence for group II effects, 297–9
  - high threshold, 293, 295, 297, 300
  - pharmacological validation, 294, 299, 300
  - slowly conducting afferents, 297–9: cooling, 294, 297–9; height of subjects, 297, 307
- facilitation of, 293–7
  - H reflexes, 293–5, 296
  - on-going EMG, 296, 297, 300
  - PSTHs of single units, 295–7
- stretch-induced responses, 293, 294
  - medium-latency responses, 293
  - short-latency responses, 293
- group II excitation, organisation, 302–10
  - central pathway, 303–4
    - central latency, 303
    - rostral location, 303–4
  - convergence of group I afferents, 305–6: delay when group I excitation is inhibited, 306; facilitation-occlusion, 296, 306; ischaemic blockade of group I afferents, 306
  - corticospinal facilitation of, 307–10
    - group II interneurons, 307–9
    - feedback inhibitory interneurons, 308, 309, 310
  - disfacilitation of motoneurons, 307; by: group I afferents, 307; group II afferents, 307
  - distribution, 304–5
    - bilateral connections, 305
    - heteronymous connections, 304–5: lower limb, 304–5; upper limb, 305
    - homonymous connections, 304
  - lack of evidence for cutaneous convergence, 306–7
- group II excitation, physiological tasks, 310–20
  - postural tasks, 312–14
    - perturbation to stance, 312, 539: bilateral response, 294, 313; effects of the postural set, 311, 313, 541; responses in the kinematic chain, 542; underlying mechanisms, 314
    - unstable stance, 313–14: focus, 538; leaning backwards, 311, 313–14; leaning forwards, 314; underlying mechanisms, 314, 538
  - voluntary contraction of target muscles, 310–12
    - H reflex, contamination by Ib inhibition, 14–15, 26, 27, 301–2, 310–12
    - quadriceps contraction, 312; underlying mechanism, 312
    - semitendinosus contraction, 312; underlying mechanism, 312
  - walking, 314–20
    - effects of unloading soleus, 315–16
    - M2 stretch responses, 316–18: heteronymous muscles, 316; tibialis anterior, 316, 317, 318, 548; triceps surae, 316–17, 318, 548
    - peroneal group II excitation of quadriceps, 318–19, 546; underlying mechanisms, 318–19
- group II excitation, possible function
  - postural tasks, 314
    - perturbations to stance, 312–14, 541
    - unstable stance, 314, 538; focus, 538
- servo-assistance
  - convergence of Ia and group II afferents, 313, 316, 318, 319–20, 512–14
  - FRA hypothesis, 389, 513, 514, 529
  - lack of post-activation depression, 310, 512–14
  - selectivity, 514, 529
- walking
  - contribution to soleus activation, 319–20, 545–6
  - stabilisation of the ankle, 318, 549
  - stabilisation of the knee, 319, 547
- group II excitation, studies in patients, 320–6
  - Parkinson's disease, 326
    - increased group II excitation and rigidity, 326, 327, 583–4, 588–9
  - perturbations to stance, 326, 589–90: functional implications, 590; influence of the postural set, 326, 327, 589–90; underlying mechanisms, 326, 590
- peripheral neuropathies, 320
  - Charcot-Marie-Tooth disease, 320, 321
  - diabetic neuropathy, 320, 321
- spasticity, pathophysiology, 320–6, 567–8
  - absence of correlations with clinical spasticity, 325
  - actions of monoamines, 321, 324–5, 568: group I excitation, 568; group II excitation, 568; interpretation with caution, 321–2, 325, 568; therapeutic action, 324–5, 568



## 616 Index

- group II excitation, studies in patients (*cont.*)
  - group II pathways and clasp-knife phenomenon, 326
  - how it could produce spasticity, 567
  - importance of contribution, 324–6, 568
  - methodology, 320–1, 567
  - positive feedback via excitation of  $\gamma$  motoneurones, 325, 567–8
  - possible mechanisms underlying increased group I excitation, 324; muscle stretch, 324; propriospinal contribution to tendon jerk, 324
- spinal spasticity, 322, 323, 581
  - changes in group I-group II excitation, 322
  - effect of: clonidine, 322, 323; DOPA, 322, 323
  - positive feedback via excitation of  $\gamma$  motoneurones, 323, 325, 567–8, 581
  - underlying mechanisms, 324, 581
- stroke, 322, 323, 577
  - affected side, 322
  - effect of tizanidine, 322
  - unaffected side, 322
  - underlying mechanisms, 322–4, 577
- group II inhibition of motoneurones
  - cat, 288, 291
  - lack of evidence in humans, 307
  - lack of involvement in clasp-knife phenomenon, 326
- group III afferents
  - clasp-knife phenomenon, 326
  - contribution to FRAs, 388
  - contribution to late withdrawal reflexes, 433
  - high-threshold inhibition of quadriceps after tibial nerve stimulation, 307
  - presynaptic inhibition of Ia terminals, 372
- H reflex, general features
  - advantages, 21
  - comparisons with the tendon jerk
    - differences in the afferent volleys, 2–3, 117–18; at the motoneurone pool, 118
    - $\gamma$  drive, 9, 117
    - Jendrassik manoeuvre, 131–3
  - desynchronisation of the reflex volley, 9–10
  - distorting effects of normalisation, 16, 17, 219
  - during voluntary contraction, 4, 18, 69, 79
  - evidence for a two-neurone arc, 66–7
  - initial studies, 2
  - input–output relationship, 17–18
  - less sensitivity to inhibition than the ongoing EMG, 26–7, 203–4
  - limitations, 11–20
  - mechanisms acting on the afferent volley, 11–16
  - monosynaptic pathway, 2, 66–7
  - motoneurones involved, 4, 79–81
  - necessity of activation of many Ia afferents, 154–5
  - ‘pool problems’: comparison with single unit recordings, 20, 21
  - principles of the method, 3, 4
  - recruitment gain, 18–20
  - size-related sensitivity of the reflex, 16–18
  - variant of the method, 4
- H reflex, limitations due to, 11–20
  - alterations in the excitability of Ia afferents, 12–13
  - changes in the recruitment gain, 18–20, 340, 344, 346, 425
  - consequences of size-related sensitivity of the reflex
    - when using conditioning effects: large, 18;
    - modest, 18
  - contamination by oligosynaptic inhibition, 14–16, 37, 67, 70, 81, 301–2, 310–12, 493, 500
  - curtailment of Ia EPSPs, 14
  - evidence for Ib inhibition in quadriceps, 14–15
  - plateau potentials, 20
  - ‘pools problems’, 16–20, 37
  - poor time resolution, 9–10, 70, 205, 253
  - post-activation depression, 13–14, 97–9
  - presynaptic inhibition of Ia terminals, 13, 81
  - size-related sensitivity of the reflex, 16–17
- H reflex, methodology, 4–11
  - ‘adjusting’ the test stimulus intensity, 18, 219
  - basic methodology, 4–11
  - cross-talk, 6
  - electrical stimulation, 6–7: bipolar, 7; monopolar, 6; percutaneous, 6
  - expression as a percentage of  $M_{\max}$ , 6, 8
  - general experimental arrangement, 4; awareness, 4; posture, 4
  - $M_{\max}$ , 7–8
  - magnetic stimulation, 7, 69
  - measurement, 6
  - monitoring the stability of the stimulation, 8
  - muscles in which it may be recorded at rest, 4, 79
  - random alternation, 9
  - recording, 5–6
  - recovery cycle, 10–11, 12
  - recruitment curve, 5, 7; collision with the antidromic motor volley, 7; descending limb, 8
  - stimulus duration: electrical, 6; magnetic, 7
  - stimulus rate, 97–9; at rest, 7, 13–14, 97–9; during contraction, 98, 99



- threshold tracking, 11, 12: advantages, 11; disadvantages, 11;  
principle, 11
- voluntary contraction, 4, 69, 79, 95
- H reflex, studies in patients
- $H_{\max}/M_{\max}$  ratio
- Parkinson's disease, 584–5
- spasticity, 562
- spinal spasticity, 580
- stroke,  $H_{\max}/M_{\max}$  ratio: FCR, 576; soleus, 576
- normative data, 20–1
- peripheral neuropathies and nerve lesions, 95–6
- recovery cycle
- Parkinson's disease, 585
- spasticity, 562
- reflex threshold
- Parkinson's disease, 585
- spasticity, 562
- H reflex, used to study
- cervical propriospinal pathways, 457, 474–5
- cutaneous pathways, 396
- group II pathways, 293–5
- lumbar propriospinal pathways, 493
- monosynaptic Ia excitation
- heteronymous, 70, 71
- homonymous: at rest, 4, 66–7, 68, 79; during contraction,  
69, 79, 80, 95
- Ib inhibition, 249
- presynaptic inhibition of Ia terminals, 340,  
341–6
- reciprocal Ia inhibition, 201
- recurrent inhibition
- activation of Renshaw cells, 155–7, 162
- heteronymous recurrent inhibition, 161–9
- homonymous recurrent inhibition, 155–61
- H' test reflex, *see* recurrent inhibition (homonymous),  
methodology
- handedness-related symmetry-asymmetry
- cervical propriospinal neurones: cutaneous inhibition, 458;  
peripheral excitation, 476–7
- H reflex, 20
- habituation of cutaneous reflexes
- abdominal skin reflexes, 394
- E1 cutaneomuscular response, 415
- grasp reflex, 436
- RII reflex, 394
- withdrawal reflexes
- normal subjects, 394
- Parkinson's disease, 436
- upper motoneurone lesions, 434
- hemiplegia, *see* stroke
- heteronymous connections, *see* group Ib inhibition, group II  
excitation, monosynaptic Ia excitation, recurrent  
inhibition
- homonymous connections, *see* group Ib inhibition, group II  
excitation, monosynaptic Ia excitation, recurrent  
inhibition
- homosynaptic depression, *see* post-activation depression
- hopping
- cutaneous excitation, 432
- heteronymous Ia connections, 93–4, 550
- short-latency Ia stretch reflex, 87–8
- hyperekplexia, changes in spinal pathways
- Ib inhibition, 276
- reciprocal Ia inhibition, 233: lower limb, 233; upper limb, 214,  
233
- recurrent inhibition, 187
- I waves after cortical stimulation, 40–1, 43, 44
- I1, cutaneomuscular inhibition, *see* transcortical cutaneous  
pathways
- inhibition, *see*  $\alpha$  motoneurone inhibition; disfacilitation of  
motoneurones; presynaptic inhibition
- initial part of monosynaptic excitation
- monosynaptic transmission, 16, 67, 346, 349, 352, 362, 364
- uncontaminated by interneuronal inputs
- cortico-motoneuronal excitation, 46, 47, 308–9, 456, 462,  
463, 466, 472, 473, 498, 499
- Ia excitation, 15, 212, 263, 264, 266, 267
- input–output relationship
- H reflex, 17–18
- MEP, 44
- recurrent inhibition, 152
- integration at premotoneuronal level
- cervical propriospinal neurones, 477–8
- Ib interneurones, 247–8, 265–7
- spinal interneurones
- interneuronal transmission, evidence for, *see* convergence,  
initial part of monosynaptic excitation, spatial facilitation
- intrafusal fibres, *see* bag and chain fibres
- ischaemic blockade
- A $\beta$  cutaneous afferents
- effects on RIII reflexes, 393, 411–12
- responses to perturbations to stance, 540
- group I afferents
- group Ib effects, 252
- Jendrassik manoeuvre, 132
- monosynaptic Ia excitation (homonymous), 69
- 'natural' reciprocal inhibition, 217

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**618 Index**

- ischaemic blockade (*cont.*)
    - perturbations to stance, 89
    - post-activation depression, 97, 98; reciprocal Ia inhibition, 221
    - presynaptic inhibition of Ia terminals during motor tasks, 220, 358, 361
    - projections on group II interneurons during walking, 306, 315–16, 317, 318, 502
    - reciprocal Ia inhibition during motor tasks, 220, 225, 229
    - running, 87
  - inverse myotatic reflex, 246
- Jendrassik manoeuvre
  - central changes in the gain of the reflex, 133, 134
  - fusimotor drive, 131; flawed hypothesis, 131–3
  - presynaptic inhibition of Ia terminals, 133, 361–2
  - reciprocal Ia inhibition, 133, 227
- joint afferents
  - activation in extremes of movement, 272, 516
  - contribution to FRA, 388
  - facilitation of motoneurons, 264
  - facilitation of Ib inhibitory interneurons
    - evidence, 263–5
    - function: protection, 263–5; termination of movement, 272, 518
  - inhibition of PAD interneurons, 339
  - lateral articular nerve of the knee joint, 263
  - projections on group II interneurons, 291
- joint stiffness
  - co-contractions of antagonists, 533–4
  - quiet stance, 536–7
- L-acetylcarnitine (L-Ac)
  - recurrent inhibition of motoneurons, 158, 159–60, 168
  - recurrent inhibition of Ia interneurons, 208
- landing, short-latency Ia stretch reflex, 88–9
  - heteronymous Ia connections, 93–4, 550
  - quadriceps, 88–9
  - soleus, 88
  - triceps brachii, 89, 521
- late spinal withdrawal reflexes, *see* withdrawal reflexes (late),
  - patients with complete spinal transection
- lengthening reaction, 244
- load compensation
  - presynaptic inhibition of Ia terminals, 359
  - stretch reflex during running, 87
- local sign of cutaneous effects
  - facilitation of
    - Ib excitation, 259, 262, 271
    - Ib inhibition, 262, 272
- inhibition of
  - cervical propriospinal neurones, 478
  - PAD interneurons, 350
- initial findings, 385
- organisation of withdrawal reflexes, 402–5, 407
- locomotion
  - possible role of late FRA pathways, 390
  - see* running, walking
- locus coeruleus
  - noradrenergic control of group II pathways in the cat, 292
- parkinsonian rigidity, 326, 590
- perturbations to stance
  - normal subjects, 314
  - Parkinson's disease, 326, 589–90
  - patients with hemiplegia, 322
- spinal spasticity, 324, 581
- lumbar propriospinal pathways, background from animal
  - experiments, 490–1
- initial findings, 490–1
- dorsolateral propriospinal neurones, 491
- ventromedial propriospinal neurones, 289, 491
  - descending control, 491; peripheral input, 491
- lumbar propriospinal pathways, methodology,
  - 491–4
- critique of tests: H reflex, 14–16, 301–2, 310–12, 493, 500;
  - PSTHs, 492–3; rostral location, 494
- evidence for propriospinally mediated effect: central delay,
  - 491, 493; diffuse pattern of input, 492, 494; low threshold, 491, 493
- facilitation (early), 493; of the: H reflex, 492, 493; on-going
  - EMG, 492, 493; PSTHs, 491–2, 493, 497
- rostral location of the interneurons, 493–4
- lumbar propriospinal pathways, organisation, 494–500
- afferent excitatory input, 494–6
  - convergence of: afferents from different muscles, 47, 495, 496
  - convergence of: group I, 305–6, 494; including Ia, 494; group II, 494
- diffuse distribution, 494
- disclosure of connections with corticospinal facilitation, 498
- peroneal-induced excitation of quadriceps, 258–60, 494–5, 496
- strength, 494–5
- corticospinal control, 498–500
  - convergence of corticospinal and peripheral inputs, 498–500; which interneurons?, 498–500

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)

- effect on feedback inhibitory interneurons, 500; dominant effect, 500
- feedback inhibition, 496–7
  - heteronymous inhibition, 496–7
  - homonymous inhibition, 496: disclosed by cortical stimulation, 496; weakness, 496
  - lack of cutaneous inhibition, 496
- inhibitory projections on motoneurons, 497–8
  - from plantar muscles to extensors, 497
  - medium-latency reciprocal inhibition, 497–8
  - which interneurons?, 498
- lumbar propriospinal pathways, physiological implications, 500–3
- unstable stance, 502
- voluntary contraction of antagonist muscles, 217, 219, 503, 519–20
- voluntary contraction of the target muscle, 14, 26, 27, 47, 500–2
  - dominant suppression, 500–2; disfacilitation, 502
  - weak facilitation, 500–2
- walking, 503: quadriceps excitation, 318, 502, 547; soleus inhibition, 547
- lumbar propriospinal pathways, possible function, 502–3
  - focus of the descending motor command to active motoneurons, 502–3
- relaxation of antagonist muscles during voluntary contraction, 217, 219, 503, 519–20
- walking
  - prevention of excessive Ia activity in soleus, 503, 547
  - stabilisation of the knee, 319, 547
- lumbar propriospinal pathways, studies in patients, 503
- Parkinson's disease, 277, 503, 588
- spasticity, 277, 503, 566–7
  - how it could produce spasticity, 566
  - importance of contribution, 567
  - methodology, 567
  - monoaminergic depression of group I excitation, 324
- spinal spasticity, 322, 503, 504, 581
  - quadriceps: effect of: clonidine, 322; DOPA, 322; increased group I excitation, 503; group I–group II excitation, 503; positive feedback via excitation of  $\gamma$  motoneurons, 567–8, 581; underlying mechanisms, 324, 581
  - soleus, 503
- stroke, 322, 503, 504, 577
  - affected side, 322, 503
  - effect of tizanidine, 322
  - unaffected side, 322, 503
  - underlying mechanisms, 322–4, 503, 577
- M1 response to stretch, *see* stretch reflex
- M2 response to stretch, normal subjects
  - hand muscles, 90–2, 583
  - lower limb muscles in stance
    - bilateral response, 313
    - effects of the postural set, 313, 541
    - evoked by perturbations of stance, 293, 539
    - functional role, 314, 541
    - group II in origin, 297–9
    - responses in the kinematic chain, 542; ankle and hip strategies, 542
  - overlap with group Ia excitation, 3, 301
  - proximal arm muscles, 92
  - walking, 316–18
    - depending on the perturbation, 548: translation, 316; vertical, 316–18
    - function, 318, 549
    - heteronymous muscles, 316
    - tibialis anterior, 316, 318, 548
    - triceps surae, 316–18, 548
  - wrist muscles, 583
- M2 response to stretch, studies in patients
  - Parkinson's disease
    - increased response, 583: hand muscles, 583, 584; wrist muscles, 583, 584
    - later part, 584; underlying mechanisms, 584
    - lower limb muscles in stance, 326, 584, 589–90; influence of the postural set, 326, 589–90
  - peripheral neuropathies, 320
  - stroke, 322
- maturation
  - cutaneomuscular responses, 423
  - heteronymous Ia excitation, 86
  - plantar response, 404
- medium-latency response to stretch, *see* M2 response
- MEP (motor evoked potential), *see* cortical stimulation
- methodology, comparison of results from different methods
  - changes in excitability of corticospinal neurones electrical vs. magnetic stimulation of the motor cortex, 44, 424, 430–2
  - contamination of the H reflex by Ib inhibition: modulation of the H reflex vs. on-going EMG, 14–15, 16, 301–2, 310–12
  - non-monosynaptic transmission of the cortical command modulation of the H reflex vs. on-going EMG and MEP, 44, 307–8, 309, 310, 471–3, 500
  - presynaptic inhibition of Ia terminals
    - modulation of the H reflex vs. corticospinal excitation: MEP, 44, 168, 343–4; PSTH, 342
    - modulation of H reflex vs. on-going EMG, 27, 340, 365, 545

## 620 Index

- methodology, comparison of results (*cont.*)  
     monosynaptic excitation vs. D1 inhibition of the H reflex, 347
- methodology used to assess motoneurone excitability  
     coherence analysis, 48–9  
     cortical stimulation, 39–41, 45  
     F wave, 21–2, 24  
     H reflex and tendon jerk, 1–21  
     on-going EMG, 24–6, 28  
     PSTHs of single units, 28–30, 37  
     unitary H reflex, 37–9
- methodology used to study spinal pathways  
      $\alpha$  motoneurone excitability, 561–3  
     cervical propriospinal pathways, 455–60  
     cutaneous reflexes, 391–9  
     fusimotor activation, 117–26  
     group Ib pathways, 249–56: excitation, 255; inhibition, 249–55  
     group II excitation, 293–302  
     lumbar propriospinal pathways, 491–4  
     monosynaptic Ia excitation, 66–77; heteronymous, 70–7; homonymous, 66–70  
     post-activation depression (Ia fibre-motoneurone), 97–9  
     presynaptic inhibition of Ia terminals, 340–7  
     reciprocal Ia inhibition, 201–9  
     recurrent inhibition heteronymous, 161–9; homonymous, 155–61
- microneurography, 119–24  
     basic methodology, 119–20  
     bias towards large fibres, 123  
     identification of axons, 120–2  
     dynamic sensitivity, 121–2  
     twitch test, 120–1: Golgi tendon organs, 123; spindle endings, 116, 121, 122, 125  
     limitations, 126  
     microelectrode, 119, 120  
     responses of various endings  
         Golgi tendon organ, 123  
         primary ending, 116, 121, 125, 128, 132  
         secondary ending, 120, 121, 122, 132  
     uncertainties, 116, 123–4  
     *see also* muscle spindles, Golgi tendon organs
- mid-thoracic nucleus, 494
- mirror movements (patients with bilateral corticospinal projections), used to investigate  
     cutaneomuscular responses, 423  
     long-latency stretch responses, 91
- monoamines
- selective blockade of group II excitation  
     cat, 292  
     humans, 299
- spasticity, 321, 324–5, 568  
     effects on group II excitation, 322, 568  
     effects on group I excitation, 322, 568; underlying mechanisms, 324  
     interpretation with caution, 568  
     therapeutic action, 568  
     *see also* clonidine, DOPA, tizanidine
- monosynaptic Ia excitation, background from animal experiments, 64–6
- heteronymous projections, 63, 65–6
- initial findings, 64
- monosynaptic projections to motoneurons, 65:  
     heteronymous, 65; homonymous, 64
- primary endings and Ia afferents, 64
- projections to motoneurons of different type, 65
- stretch reflex, 66
- monosynaptic Ia excitation, methodology, 66–77  
     critique of the tests: heteronymous, 70, 72–3  
         homonymous, 70  
     evidence for Ia excitation, 67–9, 70, 73–7  
         cutaneous stimuli, 77, 85  
         elevation of the threshold by tendon vibration, 76  
         facilitation by homonymous volley, 69  
         ischaemia, 68, 69  
         low electrical threshold, 64, 67–9, 75  
         tendon taps, 67, 71, 75–6
- H reflex, homonymous  
     at rest, 4, 66–7, 68, 79  
     during contraction, 69, 79, 80, 95  
     initial part of the excitation, 15, 16, 67
- large range of electrical thresholds in humans, 77–8  
     decrease in latency, 77, 78  
     underestimation of: monosynaptic latency, 78; strength of Ia projections, 78
- mode of homonymous conditioning stimulation, 69
- modulation of  
     heteronymous H reflex, 70, 71  
     ongoing EMG, 71, 73  
     PSTHs of single units: heteronymous, 33, 70–1, 72, 73, 85; homonymous, 67, 68, 69–70, 79
- monosynaptic latency  
     heteronymous, 70–3; bidirectional connections, 73, 74, 77  
     homonymous: in soleus, 66–7; other muscles, 70
- monosynaptic Ia excitation, organisation, 79–86  
     heteronymous connections, 81–6

- between close synergists: lower limb, 81–2; upper limb, 83–4
- between ‘non-synergist’ ankle muscles, 82
- pattern in the lower limb, 81–3; comparison with
  - distribution of recurrent inhibition, 170
- pattern in the upper limb, 83–6
- to antagonists at another joint, 83
- transjoint connections: lower limb, 83; upper limb, 84–6
- homonymous excitation, 2, 79–81
  - all motoneurons: H reflex, 79; PSTH, 79
  - orderly recruitment, 3–4, 79–81; with exceptions, 79, 81;
  - presynaptic inhibition, 347–8
- limitation due to
  - contamination by Ib inhibition, 14–15, 67, 70, 81, 301–2, 493, 500
  - presynaptic inhibition, 13, 81
- maturation, 86
- phylogenetic differences, 63; lower limb, 83, 92; upper limb, 84
- reciprocal Ia excitation newborn, 86; adult, 86
- monosynaptic Ia excitation, physiological tasks, 87–94
  - contribution to the short-latency spinal stretch reflex, 87–90, 293, 320
- hopping, 87–8
- landing: quadriceps, 88–9; soleus, 88; triceps brachii, 89, 521
- running: quadriceps, 87; soleus, 87, 88, 367
- standing: not involved in quiet stance, 537; perturbations
  - to stance, 89–90, 313, 539, 540–1; unstable stance, 94, 537–8
- walking, Ia feedback to extensors: ankle, 89, 545; knee, 545
- walking, soleus Ia stretch reflex, 89, 548
- monosynaptic Ia excitation, possible function
  - servo-assistance in movement: co-contractions of
    - antagonists, 135, 532; hinge joint movements, 135, 516;
    - tonic contractions, 122, 133, 512
  - stretch reflex: interaction with motor programs, 87; load
    - compensation, 87
- synergies through heteronymous connections, 512
  - grasping, 94, 528
  - stance and gait, 93–4, 528, 537–8, 550
- walking
  - prevention of excessive Ia activity in soleus, 93, 546
  - stabilisation of the ankle: Ia projections, 93, 546; Ia spinal
    - stretch reflex, 89, 548
- see also* servo-assistance, stretch reflex
- monosynaptic Ia excitation, studies in patients (H reflex), 95–6
  - Parkinson’s disease:  $H_{\max}/M_{\max}$  ratio, 584–5; reflex threshold, 585; recovery cycle, 585
- peripheral neuropathies and nerve lesions, 95–6; location, 95
- reflex irradiation, 96, 131
- spasticity, H reflex, 562; developmental slope, 562;
  - $H_{\max}/M_{\max}$  ratio, 562; recovery cycle, 562; reflex
    - threshold, 562; transmission of the afferent volley, 562
- spinal spasticity,  $H_{\max}/M_{\max}$  ratio, 580
- stroke,  $H_{\max}/M_{\max}$  ratio: FCR, 576; soleus, 576
- monosynaptic Ia facilitation to assess presynaptic inhibition of
  - Ia terminals
    - critique, 346
    - evidence for presynaptic inhibition, 347
    - facilitation of the: H reflex, 345–6; in PSTHs, 346–7
    - initial bins, 15, 16, 67
- monosynaptic reflex
  - apparent simplicity, 1, 11
  - different afferent volleys of H and tendon reflexes, 2–3, 117–18
  - initial studies: cat, 1; humans, 2
  - orderly recruitment of motoneurons, 3–4
  - pathway, 2
  - principles of monosynaptic reflex testing, 3, 4
  - reliability, 1–2
  - see also* H reflex, tendon jerk
- motoneurone, *see*  $\alpha$  motoneurons,  $\gamma$  motoneurons
- motor learning, 530–1
  - efference copy, 531
  - $\gamma$  drive, 137, 138, 530
  - projections to ascending tracts, 530–1
- motor tasks, changes in spinal pathways
  - changes in
    - cervical propriospinal excitation, 474–6
    - cutaneomuscular spinal responses, 427, 429–30
    - fusimotor drive, 131–8
    - group II excitation, 310–20
    - lumbar propriospinal excitation, 500–3
    - non-reciprocal group I inhibition (wrist), 524–6
    - Ib inhibition, 267–75
    - presynaptic inhibition of Ia terminals, 355–67
    - reciprocal Ia inhibition, 217–29
    - recurrent inhibition, 173–84
    - synaptic efficacy: group II excitation, 310; monosynaptic Ia
      - excitation, 97; non-reciprocal group I inhibition, 212–14;
      - reciprocal Ia inhibition, 221
    - withdrawal reflexes, 412–14
  - co-contraction of antagonists, 531–5
  - co-ordinated synergies, 527–31
  - gait, 542–50
  - isometric tonic contraction, 512–15
  - movements at ball joints, 522–6
  - movements at hinge joints, 515–22
  - postural tasks, 535–42

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**622 Index**

- motor threshold ( $1 \times MT$ ), 6, 7
- multiple corticospinal volleys, 40–1
  - D wave, 40, 43
  - I waves, 40–1, 43, 44
  - summation with peripheral volleys, 309, 498
- multiple sclerosis, changes in spinal pathways
  - Ib excitation of antagonists, 277
  - post-activation depression, 99, 100
  - presynaptic inhibition of Ia terminals, 369
  - reciprocal Ia inhibition: at rest, 230, 231, 581; during
    - voluntary contraction, 231–2; plasticity, 233
- muscle fibre properties, changes in
  - parkinsonian rigidity, 582–3
  - spasticity, 572–3
- muscle spindles
  - afferent innervation, *see* group Ia and group II afferents
  - differences in cats and humans, 115–16
  - differences in different muscles, 115
  - efferent innervation, *see*  $\gamma$  and  $\beta$  efferents
  - 'in-parallel' response, 113
  - sensory endings, *see* primary and secondary endings
  - spindle density, 117
  - structure, 114, 115
- mutual inhibition of interneurons
  - non-reciprocal group I interneurons (wrist), 214, 524, 525
- Ib interneurons
  - cat, 246
  - cutaneous control, 254, 261–2
  - reversal from facilitation to suppression of Ib inhibition
    - when increasing the input, 256: group I, 258;
    - corticospinal, 265, 269, 270; vestibulospinal, 265
- opposite Ia interneurons
  - cat, 199
  - patients with spasticity, 232
  - voluntary contraction, 222, 223, 225, 520
  - walking, 229
- Renshaw cells, 154, 176
- myotatic reflex, *see* stretch reflex
- myotatic unit, 65
- natural reciprocal inhibition, 217–19
  - origin: descending, 217–19; group I, 217
  - time course, 217, 218; prior to EMG activity, 217
  - underlying mechanisms, 217–19: post-activation depression,
    - 217; presynaptic inhibition, 217; propriospinally
      - mediated inhibition, 217–19, 519–20; reciprocal Ia
        - inhibition, 217, 519
- nerve blocks, *see* anaesthetics, ischaemic, pressure blocks
- nociceptive afferents
  - control of Ib inhibition, 265: cutaneous afferents, 265; muscle
    - afferents, 265
  - see also*: A $\delta$  afferents, FRA pathways, RIII reflex, withdrawal
    - reflexes
- non-monosynaptic group I excitation, *see* cervical and lumbar
  - propriospinal neurones
- non-reciprocal group I inhibition, 245, *see* group Ib inhibition
- non-reciprocal group I inhibition at wrist level, organisation,
  - 522–4
  - absence of evidence for pure Ia origin, 211
  - absence of post-activation depression, 212–14
  - absence of recurrent inhibition, 171–3, 207, 208
  - convergence of group I afferents from several different
    - muscles, 212, 258, 524
  - corticospinal excitation, 524
  - disynaptic connection, 205, 206
  - mutual inhibition of interneurons, 214, 524, 525
  - not between purely antagonistic muscles: clenching, 211–12,
    - 525; wrist abduction, 211, 522
  - organisation in subsets, 524
  - potency, 524
- non-reciprocal group I inhibition at wrist level, physiological
  - tasks, 524–6
- voluntary co-contraction of antagonists, 532
- voluntary contraction
  - changes in pathways to: active motoneurons, 524;
  - antagonistic motoneurons, 524–5
  - mechanisms involved: corticospinal, 524, 525; group I
    - discharge, 525–6
  - possible function, 525–6
- non-reciprocal group I inhibition at wrist level, studies in
  - patients, 234
  - dystonia, 234, 372
  - hyperekplexia, 214
  - Parkinson's disease, 234, 587; voluntary contraction, 591
  - stroke, 234, 577; correlation with spasticity, 577
- noradrenaline, *see* noradrenergic agonists
- noradrenergic agonists used to depress group II
  - cat, 292
  - normal humans, tizanidine, 299, 318
  - spastic patients, 321, 324–5, 568: clonidine, 322; DOPA, 322;
  - tizanidine, 322
- occlusion
  - cervical propriospinal neurones: not involved in the reversal
    - of corticospinal excitation, 466; during contraction,
      - 476
  - group II interneurons: group I and group II excitations, 301,
    - 306

- lumbar propriospinal neurones: corticospinal excitation, 500
- Ia interneurons: Ia and cutaneous, 214; Ia and corticospinal inputs, 216; tonic contraction of the antagonist, 220
- Ib interneurons: multiple afferent inputs, 256; Ib from different muscles, 258; Ib and corticospinal, 265; vestibulospinal, 265; tonic contraction, 270
- PAD interneurons: D1; inhibition, 344–5, 364
- Renshaw cells, voluntary contraction, 176
- oligosynaptic group I excitation, 245, *see* group Ib excitation
- on-going EMG modulation, general features
  - advantages, 27–8
  - afterhyperpolarisation, 28
  - different sensitivity to inhibition than the H reflex, 25, 26–7, 203–4
  - discrepancies with the modulation of the H reflex, 26–7; gating the afferent volley of the H reflex, 14–15, 26, 27, 340
  - disfacilitation vs. inhibition, 25, 416–18, 473
  - initial studies, 24
  - limitations, 28
  - motoneurons involved, 25
  - underlying principles, 24–5
- on-going EMG modulation, methodology
  - assessment of the central delay, 25–6
  - basic methodology with rectified EMG, 24–5
  - cutaneous reflexes, 396, 404, 414–15
  - unrectified EMG, 25
- on-going EMG modulation used to study
  - cervical propriospinal pathways, 457, 458–9, 471–3
  - cutaneomuscular responses, 414–15
  - cutaneous pathways, 396
  - effects of noxious cutaneous stimuli, 404
  - group II pathways, 297
  - lumbar propriospinal pathways, 493, 502
  - monosynaptic Ia excitation (heteronymous), 73
  - presynaptic inhibition of Ia terminals, 27, 340
  - reciprocal Ia inhibition, 203–4
  - recurrent inhibition (heteronymous), 161–9
- onset of movement, focusing action of, 520–1
  - cervical propriospinal excitation, 476
  - fusimotor drive, 135, 521
  - non-reciprocal group I inhibition (wrist), 525
  - Ib inhibition, 270, 521
  - presynaptic inhibition of Ia terminals, 359–60, 520
  - recurrent inhibition, 517, 520
  - see also* timing of changes in spinal pathways during movement
- orderly recruitment, *see*  $\alpha$  motoneurons, recruitment order
- organisation and pattern of connections
  - cervical propriospinal pathways, 460–71
  - cutaneous reflexes (low-threshold afferents), 414–32
  - fusimotor system, 127–30
  - group II pathways, 302–10
  - lumbar propriospinal pathways, 494–500
  - monosynaptic Ia excitation, 79–86
  - non-reciprocal group I inhibition (wrist), 522–4
  - Ib pathways, 256–67
  - presynaptic inhibition of Ia terminals, 347–55
  - reciprocal Ia inhibition, 209–17
  - recurrent inhibition, 169–73
  - withdrawal reflexes, 399–14
- organisation of spinal circuitry at wrist level, 522–4
  - differences with hinge joints, 522–4
  - non-noxious cutaneomuscular responses, 526
  - non-reciprocal group I inhibition, 211–14, 524
    - absence of evidence for: pure Ia origin, 211; post-activation depression, 212–14; recurrent inhibition, 171–3, 208
    - convergence of: group I afferents, 212, 524; from elbow, 212; homonymous, 212; Ib afferents, 212
    - corticospinal excitation, 524
    - organisation in subsets, 524
    - mutual inhibition, 214, 524
    - potency, 524
    - wrist movements, 524–6
- presynaptic inhibition of Ia terminals
  - corticospinal control: facilitation, 353, 524; depression, 353
  - cutaneous control, 350
  - wrist movements, 362–3, 526; non-specific decrease, 363, 526, 532; possible increase, 526
- recurrent inhibition
  - between FCR and ECR, 171, 522
  - wrist movements, 534
- synergies at wrist level: clenching, 211–12, 525; wrist abduction, 211, 522
- organisation of spinal interneurons in subsets
  - with respect to muscle afferent input: cervical propriospinal neurones, 468–9
  - with respect to target motoneurons
    - interneurons mediating withdrawal reflexes, 407
    - non-reciprocal group I interneurons (wrist), 524, 525
    - Ia interneurons, 199
    - Ib inhibitory interneurons, 257
    - PAD interneurons, 348, 358, 363
- oscillations during movement, prevention of, 512
  - limitation of the stretch reflex during co-contraction of antagonists: ball joints, 534–5; hinge joints, 534
- post-activation depression, 97
- presynaptic inhibition, 359



## 624 Index

- PAD (primary afferent depolarisation), 337
- PAD interneurons, cat
- FRA terminals, 390
  - group II terminals, 292
  - Ia terminals, 337
  - Ib terminals, 248
- pain
- parallel with RIII reflex, 399–400
  - RIII reflex used to monitor medication for pain, 437
- paired H reflex technique, *see* recurrent inhibition (homonymous), methodology, 155–61
- Parkinson's disease, findings at rest, transmission in spinal pathways, 584–9
- absence of consistency, 589
  - $\alpha$  motoneurone excitability, 584–6
    - F waves, 585
    - H reflex:  $H_{\max}/M_{\max}$  ratio, 584–5; reflex threshold, 585; recovery cycle, 585
    - inconsistencies, 585–6
    - MEP, 585
  - fusimotor drive, 140–1, 586
  - indirect evidence, 586
  - insufficient data, 586
  - microneurography, 586
  - group II excitation, 326, 584, 588–9
    - correlation with rigidity, 326, 589
    - underlying mechanisms, 588
  - lumbar propriospinal neurones, 277, 503, 588
  - non-reciprocal group I inhibition (wrist), 234, 587
  - Ib inhibition (ankle), 276–7, 588
    - correlation with rigidity, 276, 588
    - effect of treatment, 588
    - underlying mechanisms, 277, 588
  - post-activation depression, 585
  - presynaptic inhibition of Ia terminals, 371, 586–7
    - homonymous vibratory inhibition, 371, 587
    - lower limb, 371, 586
    - upper limb, 371, 587
  - reciprocal Ia inhibition, 233–4, 587
    - tonic inhibition, 587
    - underlying mechanisms, 587–8
  - recurrent inhibition, 187, 589
  - transcortical inhibition, 437, 589
  - withdrawal reflexes, 436–7, 589
- Parkinson's disease, findings during motor tasks, 589–92
- cervical propriospinal excitation, 484–5
    - compensatory mechanism, 485, 592
    - increased transmission of the descending command, 485, 591
  - group II excitation in perturbations to stance, 326, 589–90
    - backward translation, 590
    - functional implications, 590
    - influence of the postural set, 326, 589–90
    - underlying mechanisms, 326, 590
  - relaxation of the antagonists, 590–1
    - functional implications, 591
    - non-reciprocal group I inhibition (wrist), 591
    - reciprocal Ia inhibition, 234, 590–1
    - underlying mechanisms, 591
- parkinsonian akinesia
- post-activation depression, 585
  - presynaptic inhibition of Ia terminals, 371, 586
- parkinsonian rigidity, 582–4
- changes in muscle fibre properties, 582–3
    - conflicting results, 582–3
  - elbow muscles, 582
  - gait, 582
- general features
- clinical characteristics, 582
  - depending on afferent inflow, 582
  - inability to relax, 582, 583
- involvement of spinal pathways
- $\gamma$  drive, 140–1
  - group II excitation, 326, 583–4, 589
  - Ib inhibition, 276, 588
  - presynaptic inhibition of Ia terminals, 371, 586
  - reciprocal Ia inhibition, 587
  - transcortical Ia pathways, 584
- long-latency stretch reflex (M2), 583–4
- correlation with rigidity, 584
  - increase, 583
  - later part, 583, 584; underlying mechanisms, 584
  - origin, 583–4: distal muscles, 583; wrist muscles, 583, 584
- short-latency stretch reflex (M1), 583
- parkinsonian tremor, 141
- pattern of connections, *see* organisation and pattern of connections
- perception of force, 246–7
- period prior to voluntary EMG, *see* timing of changes in spinal pathways during movement
- peripheral neuropathies and nerve lesions
- F wave, 24
  - group II-mediated excitation, 320
  - H reflex, 95–6
  - location of the lesion
    - cutaneomuscular spinal responses, 437
  - H reflex, 95
  - spinally mediated Ia stretch reflex, 320



- perturbations to stance, 538–42
  - ankle and hip strategies, 542
  - stretch responses in ankle muscles, 539–41
    - conflicting results, 539
    - cutaneous contribution, 540
    - long-latency in the antagonist, 540: function, 541; origin, 540; possibly transcortical, 540
    - long-latency in the stretched muscle, 539–40; functional stretch reflex, 539
    - medium-latency, 293: distribution, 293, 539; function, 314, 541; group II origin, 91, 297–9; influence of the postural set, 313, 541; underlying mechanisms, 314
    - short-latency, 293: distribution, 539; function, 320, 540–1; Ia origin, 89–90, 320
  - stretch responses in the kinematic chain, 541–2
    - group II excitation, 313, 542
    - responses in foot muscles, 312–13, 541–2
- studies in patients
  - Parkinson's disease, 326, 589–90
  - peripheral neuropathies, 320
  - stroke, 322
- pharmacology
  - group II excitation, 292, 299, 321, 324–5, 568
- presynaptic inhibition of Ia terminals, 337
- reciprocal Ia inhibition, 198–9
- recurrent inhibition of: motoneurons, 151, 152, 159–60; Ia interneurons, 208
- phylogenetic differences
  - cervical propriospinal system, 455, 479
  - heteronymous group II excitation, 305
  - heteronymous Ia excitation, 63: lower limb, 83, 92; upper limb, 84
  - heteronymous recurrent inhibition: lower limb, 169–70; upper limb, 170–1
  - inhibition from group II afferents, 307
- physiotherapy
  - action on spasticity, 556
  - effects on reciprocal Ia inhibition, 233
  - plasticity, 572
- placing reaction, 427
  - cat, 385–7
  - humans: biceps femoris, 427; peroneus longus, 427
- plantar responses, 402–4
  - Babinski response, 433–4
  - dysfunction of the upper motor neurone, 434
  - extensor hallucis longus, 402, 403, 434
  - pathophysiology, 434
- maturation, 404
- methodology, 392, 402, 434
  - electrical stimulation, 402, 434
  - extensor hallucis longus, 402, 434
  - mechanical stimulation, 392, 402, 434
- normal functional organisation, 402–4
- protective function, 404
- plasticity
  - basis for physiotherapy, 572
  - changes after spinal cord lesions in animals, 572: sprouting, 572; supersensitivity, 572
  - changes in muscle fibre properties in: Parkinson's disease, 582–3; spasticity, 572–3
  - late withdrawal responses in normal subjects, 410; influence of: continuous stimulation, 410; hypnosis, 410; posture, 410
  - possible cause of  $\alpha$  motoneurone hyperexcitability, 561, 572
  - post-activation depression in: Parkinson's disease, 585; spasticity, 99–100, 572
  - reciprocal Ia inhibition, 232–3
    - normal subjects, 210, 232
  - spasticity: frequent peroneal stimulation, 233, 572; training, 233
- plateau potentials
  - normal subjects, 20
  - spasticity, 563, 580
- post-activation depression at the Ia fibre-motoneurone synapse, 96–100
  - background from animal experiments, 96; effect on motoneurons of different size, 96
  - cause of misinterpretations when using the H reflex
    - circumstances other than changes in the stimulus rate, 13, 14
  - vibration of the homonymous tendon, 341, 368, 371, 565
- methodology, 97–9
  - increasing the stimulus rate for the H reflex, 97–9: at rest, 13–14, 97–8, 99; during contraction, 98, 99
  - stretch of the tested muscle, 97, 98
- motor tasks
  - effect on motoneurons of different type, 99
  - functional implications, 97, 512, 516
- studies in patients
  - Parkinson's disease, 585
  - spasticity, 96, 99–100, 566; plasticity, 99–100, 572
  - spinal cord lesions, 99, 100, 580; multiple sclerosis, 99; trauma, 99
  - stroke: affected side, 99, 100, 577; unaffected side, 99, 100, 579

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**626 Index**

- post-activation depression in other spinal pathways
  - background from animal experiments, 248
  - functional significance, 97
  - group II excitation, 310, 512–14
  - non-reciprocal group I inhibition (wrist), 212–14
  - reciprocal Ia inhibition
    - $\alpha$ - $\gamma$  co-activation during tonic contraction, 14, 221; functional implications, 222, 519
    - stretch of the antagonistic (tested) muscle, 217
  - recurrent pathway, 176
  - withdrawal reflexes, 411
- postural tasks, 535–42; *see*: perturbations to stance, quiet stance, unstable stance
- pressure block of  $\gamma$  efferents, 117, 119, 133
- presynaptic inhibition of cortico-motoneuronal projections, 342–3
- presynaptic inhibition of group I terminals to cervical propriospinal neurones, 475, 476
- presynaptic inhibition of group II terminals, 292
- presynaptic inhibition of non-nociceptive cutaneous afferents, 388
- presynaptic inhibition of nociceptive cutaneous afferents
  - cat, 390
  - tactile depression of withdrawal reflexes, 412
- presynaptic inhibition of Ia terminals, background from animal experiments, 337–40
  - descending input, 338, 339
    - excitatory, 339: corticospinal, 339; tonic level of presynaptic inhibition, 338, 339; vestibulospinal, 339
    - inhibitory, 339: corticospinal, 339; reticulospinal, 339
  - general features, 337–9: electrophysiology, 338–9; location, 337; mechanisms, 337; organisation, 338
  - initial findings, 337
  - PAD interneurones, 337
  - peripheral input, 338, 339: excitatory, 339; inhibitory, 339
  - presynaptic inhibition of Ia terminals to Ia interneurones, 200, 340
  - selectivity of the control of PAD interneurones, 339–40
- presynaptic inhibition of Ia terminals, methodology, 340–7
  - activation of PAD interneurones by a heteronymous group I volley, 341–5
    - comparison of H reflex and corticospinal excitation: MEP, 343–4; PSTH, 342
  - critique, 344–5, 364, 365–6: drawbacks inherent to H reflex technique, 344; occlusion, 344–5, 364; presynaptic inhibition of the conditioning volley, 364; spread of the stimulus, 344; superimposed post-synaptic facilitation, 344, 364
  - D1 inhibition, 343–4: lower limb, 344; upper limb, 343–4
  - inhibition of the H reflex by a brief vibration or tap, 341–2
    - underlying principle, 340–1
  - changes with ageing, 368
  - discrepancy between H reflex and on-going EMG, 340
    - critique, 340
    - gait, 365, 545
    - underlying principle, 27, 340
  - evidence for presynaptic inhibition, 347
    - comparison of results with heteronymous facilitation and D1 inhibition, 347
    - experiments during voluntary contraction, 347
    - investigations of single units, 347
  - homonymous vibratory inhibition of the H reflex, 341
    - critique, 341, 368, 371, 565
    - vibration paradox, 341
  - monosynaptic Ia facilitation of the H reflex, 345–6
    - critique, 346
    - large facilitation required, 346
    - underlying principle, 345
    - validation, 346
  - techniques using single units, 346–7
    - PSTHs, 346–7; initial bins, 16, 67, 346, 349, 352, 362, 364
    - unitary H reflex, 347
- presynaptic inhibition of Ia terminals, organisation, 347–55
  - corticospinal projections, 350–3
    - lower limb: dominant depression, 350–1, 352, 353; convergence with cutaneous input, 353; focused drive, 351; possible facilitation, 353
    - upper limb: dominant facilitation, 352, 353, 524; functional implications, 353; possibly tonic, 370; reversed by cutaneous stimuli, 353
  - cutaneous inhibition, 350
    - lower limb, 350, 420; local sign, 350
    - upper limb, 350; absence of local sign, 350; tonic effect, 350
- group I excitation, 348–50
  - Ia afferents, 350; pattern, 350
  - Ib afferents, 350
- organisation in subsets, 348
  - convergence, 348
  - divergence, 348
  - with respect to target motoneurones, 348, 356, 357, 358, 363
- produced by late FRAs, 408
- projections on motoneurones of different type, 347–8, 349
- sensitivity of the stretch reflex to presynaptic inhibition, 354–5
  - functional implications, 355
  - underlying mechanisms, 354–5
- tonic level, 338, 353–4
- vestibulospinal facilitation, 353

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)

- presynaptic inhibition of Ia terminals, physiological
  - implications, 355–67
- co-contraction of antagonists
  - ankle, 361, 362, 532; underlying mechanisms, 361, 362
  - wrist, 532
- contraction of antagonists, 360–1
  - timing: before, 360; onset, 360; ramp, 360–1
  - tonic contraction, 361
  - underlying mechanisms, 360–1
- contractions of remote muscles, 133, 361–2
- contraction of synergists, 359–60
  - presynaptic inhibition to inactive motoneurons, 357, 359
  - underlying mechanisms, 357, 359
- contraction of the target muscle, 355–6, 357, 358
  - force-related, 356, 358
  - timing: before, 358; onset, 355, 356; ramp, 356, 358; ramp end, 358
  - tonic contraction, 358
  - underlying mechanisms, 356, 357, 358–9
- contractions of wrist muscles, 362–3, 526
  - co-contractions of antagonists, 532
  - non-specific decrease, 363, 526, 532
  - possible increase, 526
  - underlying mechanisms, 363
- quiet stance, 363–5: quadriceps, 363, 364; soleus, 363–4; tibialis anterior, 363
- running, 367
- walking, 365–7
  - Ia feedback to motoneurons: quadriceps, 365, 545; soleus, 365–6, 545
  - Ia feedback to Ia interneurons, 229
- presynaptic inhibition of Ia terminals, possible function
  - active motoneurons involved in a selective contraction
    - Ia feedback increased, 359; control of: force and speed, 359, 517–18; timing beginning, 359; later, 359
    - Ia feedback to inactive synergists decreased: selectivity of movement, 94, 359–60, 517, 529
    - reciprocal Ia inhibition depressed, 225, 361, 520
  - antagonistic motoneurons reciprocal Ia inhibition
    - facilitated, 222, 519; stretch reflex, 361
  - co-contractions: reciprocal Ia inhibition depressed, 226–7, 361, 532; stretch reflex, 361
- quiet stance, 364–5
  - quadriceps, 364
  - soleus, 364–5: reciprocal Ia inhibition, 364–5; stretch reflex, 364
- running, 367
- walking
  - prevention of excessive Ia activity in soleus, 366–7, 546
  - stabilisation of: ankle, 229, 546; knee, 365, 547
  - yield of the knee, 365, 547
- presynaptic inhibition of Ia terminals, studies in patients, 367–72
- dystonia, 371–2
  - D1 inhibition, 371–2: early phase, 371–2; late phase, 372
  - group III-induced presynaptic inhibition, 372
  - underlying mechanisms, 372
- methodology, 367–8
  - D1 inhibition, 367, 565–6
  - heteronymous facilitation, 566
  - heteronymous tendon tap, 368, 566
- Parkinson's disease, 371, 586–7
  - correlations with: akinesia, 371; rigidity, 371
  - homonymous vibratory inhibition, 371, 587
  - lower limb, 371, 586
  - upper limb, 371, 587
- spasticity, 368–70, 565–6
  - animal models, 565
  - how it could produce spasticity, 565
  - importance of contribution, 370, 566
  - overestimation, 368, 566: homonymous vibration, 368, 565; sensitivity of the stretch reflex to presynaptic inhibition, 368
  - variations with the level of the lesion, 566
  - voluntary contraction, 370, 574, 575
  - walking, 368, 370–1, 575
- spinal spasticity, 369, 580; underlying mechanisms, 370, 580
- stroke, 368–9, 576
  - lower limb, 368–9, 576
  - upper limb, 369, 576
  - underlying mechanisms, 369–70, 576
- presynaptic inhibition of Ib terminals
  - cat, 248
  - spasticity, 278, 577
  - voluntary contraction, 270, 272
- presynaptic inhibition of reciprocal Ia inhibition
  - background from animal experiments, 200–1
    - Ia terminals on interneurons, 200
    - terminals of the interneurons, 201
  - reciprocal Ia inhibition
    - depressed during co-contractions of antagonists, 226–7, 361, 532
    - depressed to active motoneurons, 225, 361, 520
    - facilitated to antagonist motoneurons, 222, 519

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**628 Index**

- primary endings (muscle spindle)
  - cat, definition, structure, 114, 115
  - dynamic sensitivity to perturbations of: stance, 89
  - walking, 316
  - dynamic sensitivity used in identification, 120–1
  - origin of Ia afferents, 64, 114
  - recordings from, 116, 121, 125, 128, 132
  - sensitivity to stretch and vibration in the cat, 64, 114
  - sensitivity to vibration in humans, 130, 132
- propriospinal neurones, 452, *see* cervical and lumbar propriospinal neurones, group II interneurones
- propriospinally mediated inhibition of motoneurones
  - from plantar muscles to extensor motoneurones, 497; walking, 503, 547
  - reciprocal inhibition of soleus (tibialis anterior voluntary contraction)
    - normal subjects, 217–19, 497–8, 519–20
    - spastic patients, 575
- PSTHs (post-stimulus time histograms), general features
  - advantages, 37
  - initial studies, 29
  - limitations, 36–7
  - underlying models, 29
  - underlying principles, 29, 30
- PSTHs, methodology
  - afterhyperpolarisation, 31, 32, 37
  - assessment of conduction velocity of Ia afferents, 36
  - basic methodology, 29–32
  - central delay, 34
  - cumulative sum (CUSUM), 33–4
  - initial bins, 15, 16
  - latency measurements, 32–3
  - mono- and non-monosynaptic components, 34
  - multiple peaks, 37
  - normalisation, 35–6
  - recording, from a single motor unit, 30–1: characterisation, 31; one motor unit, 30–1; pairs of units, 31; same unit, 31; stable frequency, 31
  - size and threshold of a change, with stimulation triggered by the previous firing, 33, 34–5, 460
  - randomly, 34
  - stimulation: delivered randomly, 31; triggered by the unit firing, 32
  - stimulus intensity, 32
  - window, 32
- PSTHs, used to investigate projections on motoneurones of
  - different type of
    - cervical propriospinal neurones, 471
    - corticospinal excitation through propriospinal neurones, 469–71
    - low-threshold cutaneous afferents, 425
    - monosynaptic Ia excitation, 79–81
    - presynaptic inhibition of Ia terminals, 347–8
- PSTHs, used to study
  - cervical propriospinal pathways, 457
  - conduction velocity of Ia afferents, 36
  - corticospinal projections on
    - cervical propriospinal neurones, 461–3
    - feedback inhibitory interneurones to cervical propriospinal neurones, 464–7
    - feedback inhibitory interneurones to lumbar propriospinal neurones, 310, 500
    - group II interneurones, 46–7, 308–9
    - lumbar propriospinal neurones, 498
    - PAD interneurones, 351–3
  - cutaneous (non-noxious) pathways, 396, 416, 417, 420
  - group II pathways, 295–7
  - lumbar propriospinal pathways, 491–3
  - monosynaptic Ia excitation: heteronymous, 70–3
    - homonymous, 67, 69–70
  - non-reciprocal group I inhibition (wrist), 212
  - Ib inhibition, 249–52, 253–5
  - oligosynaptic limitation of the H reflex, 14–15, 37
  - 'pool problems', 20, 21, 37
  - presynaptic inhibition of Ia terminals, 342, 346–7
    - co-contractions, 361, 362
    - quiet stance, 363, 364
  - reciprocal Ia inhibition, 204
  - recurrent inhibition (heteronymous), 161–8
  - responses to cortical stimulation, 40
- quantitative assessments, difficulties
  - convergence of multiple inputs on interneurones, 261, 301, 306, 474
  - spatial/temporal summation in motoneurones, 301, 316, 474, 512
- quiet stance, 535–7
  - anticipatory control of body sway, 537
- changes in transmission in spinal pathways
  - presynaptic inhibition of Ia terminals: quadriceps, 363, 364; soleus, 363, 364–5
  - recurrent inhibition, 181
  - spinally mediated Ia excitation, 537
  - withdrawal reflexes, 414
- intrinsic stiffness of the ankle, 536–7
- inverted pendulum, 535
- multi-sensory feedback, 535–6; absence of redundancy, 536

- ramp contractions
  - co-contraction of antagonists: recurrent inhibition, 180
  - contraction of the antagonistic muscle
    - fusimotor drive, 135
    - presynaptic inhibition of Ia terminals, 360–1
    - reciprocal Ia inhibition, 221
    - recurrent inhibition, 180
  - contraction of the target muscle
    - fusimotor drive, 135
    - presynaptic inhibition of Ia terminals, 358
    - recurrent inhibition, 174
- reaching, role of the cervical propriospinal system
  - cat, 454
  - possible role in humans, 529–30
- reciprocal Ia excitation
  - cerebral palsy, 233, 579
  - maturation, 86
  - spread of the mechanical stimulus, 86
- reciprocal Ia inhibition, background from animal experiments, 197–201
  - $\alpha$ - $\gamma$  linkage in reciprocal Ia inhibition, 198, 201
  - 'direct' inhibition, 197
  - general features, 198–9: electrophysiology, 199; identification, 198; mode of action, 199; morphology, 198; pharmacology, 198–9
  - initial findings, 197–8
  - input to Ia interneurons from, 199–200: descending tracts, 200; cortico-, 200; rubro-, 200; vestibulo-spinal, 200; FRA, 199
  - mutual inhibition of opposite Ia interneurons, 198, 199
  - 'Ia interneurons', 197
  - presynaptic inhibition, 200–1: Ia terminals, 198, 200; terminals of the interneurons, 201
  - projections of Ia interneurons to, 199:  $\alpha$  motoneurons, 199; opposite Ia interneurons, 199
  - recurrent inhibition of Ia interneurons, 152, 154, 198, 200
- reciprocal Ia inhibition, methodology, 201–9
  - critique of tests, 208–9
    - central delay, 208
    - encroachment on afferents in other nerves: superficial peroneal, 209; upper limb, 209
    - propriospinally-mediated inhibition, 209
    - stimulation of other fibres, 208–9
  - differences from Ib inhibition, 255
  - evidence for disynaptic inhibition
    - bidirectional connections: H reflex, 205, 206; PSTH, 203, 205
    - poor time resolution of the H reflex, 205
  - evidence for inhibition from Renshaw cells, 205–8
    - absence at wrist, 171–3, 207, 208
    - ankle, 171, 206, 207
    - elbow, 171, 172, 208
    - pharmacological validation, 207, 208
  - evidence for Ia origin, 204–5
    - absence of cutaneous effect, 204
    - elevation of the threshold by tendon vibration, 203, 205
    - low electrical threshold, 204
    - tendon tap, 202, 204
  - inhibition of
    - H reflex, 201, 202; tendon jerk, 201, 203
    - on-going EMG, 25, 26–7, 203–4, 223
    - PSTHs of single units, 204
- reciprocal Ia inhibition, organisation, 209–17
  - corticospinal control, 216
  - cutaneous control, 214–15
  - organisation and strength
    - ankle, 202, 209–10: soleus, 202, 210; individual variations, 202, 210; tibialis anterior, 202, 210
    - elbow, 203, 210, 521
    - knee, 210–11
  - recurrent inhibition of Ia interneurons, 171–3, 205–8
  - vestibulospinal facilitation, 216–17
- wrist
  - absence of true reciprocal Ia inhibition, 211–14
    - see non-reciprocal group I inhibition at wrist level*
- reciprocal Ia inhibition, physiological implications, 217–29
  - co-contraction of antagonists, 225–6, 227
  - modulation of: H reflex, 225, 531; PSTHs, 225
  - onset, 225
  - underlying mechanisms, 225–7: decoupling of
    - motoneurons and Ia interneurons, 225–6, 533;  $\gamma$  drive, 135, 532; presynaptic inhibition, 226–7, 361, 532; recurrent inhibition, 181, 227, 531–2
  - contraction of antagonistic muscles, 217–23
    - $\alpha$ - $\gamma$  linkage in reciprocal Ia inhibition, 198, 219, 223
  - mechanisms obscuring reciprocal Ia inhibition: occlusion, 220; post-activation depression, 221; presynaptic inhibition, 220
  - 'natural' reciprocal inhibition, 217–18, 219
  - ramp contractions, 218, 221; prior to EMG activity, 221
  - tonic contractions, 218, 219–20; blockade of the afferent feedback, 218, 220; conflicting results, 219; individual variations, 219–20; methodological considerations, 219
  - underlying mechanisms, 221–3, 519:  $\alpha$ - $\gamma$  coactivation, 221–3; descending drive, 221; presynaptic inhibition, 222, 519; recurrent inhibition, 222
  - contraction of remote muscles, 133, 227
  - contraction of the target muscle, 223–4, 225
    - force-related, 223

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**630 Index**

- reciprocal Ia inhibition, physiological implications (*cont.*)
  - modulation of: H reflex, 223; ongoing EMG, 223
  - underlying mechanisms, 223–5: mutual inhibition of Ia interneurons, 223–4; presynaptic inhibition, 225, 361, 520; recurrent inhibition, 225
  - plasticity in normal subjects, 202, 210, 232
  - postural activity, 227
  - walking, 227–9, 546
    - modulation of: H reflex, 228, 229; ongoing EMG, 228, 229
    - underlying mechanisms, 229; mutual inhibition of Ia interneurons, 229; presynaptic inhibition, 229
- reciprocal Ia inhibition, possible function
  - co-contraction, decoupling of motoneurons and corresponding Ia interneurons, 227; unstable stance, 227
  - depression of reciprocal Ia inhibition of: active motoneurons, 225, 517; corresponding Ia interneurons, 225, 520; control of force, 224, 518
  - inhibition of antagonistic: motoneurons, 223, 519; corresponding Ia interneurons, 222, 223, 520
  - linkage of descending command to motoneurons and corresponding Ia interneurons, 223, 227, 519
  - walking, stabilisation of the ankle, 229, 546; cutaneous facilitation, 214–15, 549
- reciprocal Ia inhibition, studies in patients, 229–34
  - cerebral palsy, 233, 579
  - hyperekplexia, 233: lower limb, 233; upper limb, 214, 233
  - Parkinson's disease, 233–4
    - findings at rest, 233–4, 587: tonic inhibition, 587; underlying mechanisms, 587–8
    - voluntary movement, 234, 590–1; underlying mechanisms, 591
  - spasticity findings at rest, 229–30, 570
  - distribution extensors–flexors, 232, 570
  - how it could produce spasticity, 570
  - importance of contribution, 570
  - methodology, 229, 570
  - underlying mechanisms, 232, 570
- spasticity, findings during motor tasks
  - cause of motor impairment, 231–2, 574–5
  - underlying mechanisms  $\alpha$ – $\gamma$  linkage, 574; corticospinal drive to Ia interneurons, 574; opposite Ia interneurons, 575; presynaptic inhibition, 574; recurrent inhibition, 575
- spasticity, plasticity: frequent peroneal stimulation, 233
- training, 233
- spinal spasticity
  - correlation with recovery, 232
  - multiple sclerosis, 230, 231, 581
  - spinal cord injury, 230; distribution, 581
  - stroke, 230, 577; correlation with recovery, 232, 577; underlying mechanisms, 577
- reciprocal organisation
  - absent for spinal cutaneomuscular responses: lower limb, 429; upper limb, 427
  - Ia pathways: cat, 197; hinge joints, 211
  - Ib pathways: cat, 245; Ib excitation elbow, 258
  - withdrawal reflexes, 402, 404, 405
- recovery from
  - propriospinal lesion, 481
  - stroke
    - propriospinal transmission, 484
    - reciprocal Ia inhibition, 233, 577
- recruitment gain of the H reflex
  - comparison of results with heteronymous facilitation and D1 inhibition, 347
  - effect of cutaneous stimuli, 20, 425
  - limitation of the H reflex method, 18–20
  - single unit investigations required, 20, 347
- recruitment order, *see*  $\alpha$  motoneurons, recruitment order
- recurrent collaterals from motor axons, 152
- recurrent inhibition, background from animal experiments, 151–4
  - functional hypotheses, 154
  - general features, 151–2: electrophysiology, 152; input-output relationship, 152; morphology, 151; organisation, 151; pharmacology, 151, 152; strength, 152
  - initial findings, 151
  - input to Renshaw cells from, 152, 152–3:  $\alpha$  motoneurons, 152–3; descending tracts, 153; segmental afferents excitatory, 153; inhibitory, 153
  - projections from Renshaw cells to, 153–4:  $\alpha$  motoneurons, 153;  $\gamma$  motoneurons, 153; Ia interneurons, 152, 154, 200; other Renshaw cells, 152, 154; VSCT, 154
- recurrent inhibition, methodology (heteronymous), 161–9
  - critique of the tests, 168–9
  - evidence for recurrent inhibition, 166–7
    - antidromic motor discharge, 168; orthodromic motor discharge, 167–8
    - central delay, 166
    - duration, 166
    - threshold related to conditioning motor discharge, 162, 166
  - inhibition evoked by motor discharge
    - antidromic, 162–4, 166
    - orthodromic, 162: H reflex, 35, 162, 164, 165; tendon jerk, 162, 163
  - inhibition of, 161–2
    - MEPs, 165
    - monosynaptic reflexes, 165

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)

- on-going EMG, 165
- PSTHs of single units, 163, 164, 165
- rest vs. contraction, 169
- routine explorations, 169
- recurrent inhibition, methodology (homonymous), 155–61
  - afterhyperpolarisation and recurrent inhibition, 156, 157–9, 161
  - comparison of H' with a reference H reflex, 160
  - conditioning and test reflexes, 155–6, 157
  - contamination by Ib inhibition, 155, 160–1
  - critique of the method, 160–1
  - evidence for recurrent inhibition, 157, 159
    - inhibition related to conditioning reflex size, 157, 158
    - time course, 158, 159
  - lesser sensitivity of H' than of the reference H to PSPs, 161
  - limitations, 160–1
  - underlying principles, 155
  - use of antidromic motor volleys is invalid, 154–5
  - validation
    - animal experiments, 159
    - pharmacological, 158, 159–60
- recurrent inhibition, organisation, 169–73
  - absence of recurrent collaterals to motoneurons of distal muscles: heteronymous, 170, 171; homonymous, 169
  - comparison with distribution of Ia excitation, 170
  - corticospinal control of recurrent inhibition, 173:
    - heteronymous, 172, 173; homonymous, 172, 173; via reticular formation, 173
  - functional implications of heteronymous distribution: lower limb, 170, 183–4, 529, 538; upper limb, 171
  - inhibition of motoneurons, 169–71
    - heteronymous distribution, 169–71: lower limb, 166, 169–70; upper limb, 167, 170–1
    - homonymous projections to muscles, 169: distal, 169; proximal, 169, 521
  - inhibition of Ia interneurons, 171–3
    - absence at wrist, 171–3, 207, 208
    - ankle, 171, 206, 207
    - elbow, 171, 172, 208
  - mutual, between FCR and ECR, 171, 522; function, 534
  - phylogenetic difference: lower limb, 169–70; upper limb, 170–1
- recurrent inhibition, physiological implications, 173–84
  - co-contraction of the antagonists
    - ankle, 177, 180, 531–2: ramp, 180; tonic, 180
    - wrist, 534
  - contraction of antagonist muscles, 180: ramp, 177, 180; tonic, 180
  - contraction of the target muscle, 173–9, 514–15, 517
  - contractions of different types: ballistic, 176; ramp, 174, 175; tonic, 174, 175
  - heteronymous recurrent inhibition, 178–9
  - underlying mechanisms: afterhyperpolarisation, 178; descending changes in Renshaw cell excitability, 178; effects of the natural motor discharge, 176
  - postural tasks: quiet stance, 181, 182; unstable stance, 182, 183
  - walking, 546
- recurrent inhibition, possible function
  - stance: quiet stance, 181; unstable stance, 183–4, 538
  - voluntary co-contraction of the antagonists: hinge joint,
    - inhibition of motoneurons and Ia interneurons, 181, 532; specific control, 181; wrist, 534
  - voluntary contraction of the antagonist
    - inhibition of motoneurons, 180, 520
    - prevention of stretch reflex, 520
  - voluntary contraction of the target muscle, 179
    - compensation for muscle fatigue, 174
    - control of force and speed, 179, 518
    - control of reciprocal Ia inhibition, 179, 222; mutual inhibition of Ia interneurons, 225
    - gain hypothesis, 179, 514–15
    - selectivity: homonymous, 517; heteronymous, 183–4
  - walking, 547
- recurrent inhibition, studies in patients, 184–7
  - cerebral palsy, 187
  - hyperekplexia, 187
  - Parkinson's disease, 187, 589
- spasticity, findings at rest, 184, 185
  - amyotrophic lateral sclerosis, 184, 581
  - animal models, 569
  - how it could produce spasticity, 184, 569
  - importance of contribution, 569
  - methodology, 184, 569
  - underlying mechanisms, 186
- spasticity, findings during motor tasks, 185, 186–7, 575
- spinal spasticity
  - progressive hereditary paraparesis, 185, 186, 581
  - spinal cord injury, 184, 581
- stroke, 184, 185, 577; findings using a flawed technique, 569
- redundancy in the function of different pathways contributing to
  - inhibition of antagonists, 520
  - motor tasks, 511
  - quiet stance, 536
- reference H reflex, *see* recurrent inhibition (homonymous), methodology



## 632 Index

- reflex reversal
  - cutaneous control of Ib inhibition during voluntary contraction, 259, 262, 265–7
  - cutaneous reflexes during walking: cat, 387; humans, 430
  - Ib inhibition during walking: cat, 248; humans, 273–5, 546
  - propriospinally mediated excitation when increasing the input, 464–7
  - state-dependent modulation of sensory feedback, 530
- reinforcement manoeuvre, *see* Jendrassik manoeuvre
- relaxation of antagonists, spinal pathways contributing to, 513, 519–20, 521
- normal subjects
  - absence of redundancy, 520
  - non-reciprocal group I inhibition (wrist), 523, 524–5
  - Ib inhibition, 273, 520
  - post-activation depression (stretched muscle), 217
  - presynaptic inhibition of Ia afferents, 217, 361, 519
  - propriospinally mediated reciprocal inhibition, 217, 503, 519–20
  - reciprocal Ia inhibition, 217, 519
  - recurrent inhibition, 180, 520
  - see also* 'natural reciprocal' inhibition, 217
- spastic patients, 573–5
- Renshaw cells, 151
- Renshaw inhibition, *see* recurrent inhibition
- repetition rate
  - cutaneous reflexes, *see* habituation of cutaneous reflexes, 391
  - H reflex, 97–9: at rest, 7, 13–14, 97–9; during contraction, 98, 99
- reticulospinal control, background from animal experiments
  - cervical propriospinal neurones, 453
  - group II pathways, 292
  - lumbar propriospinal neurones, 491
  - Ib pathways, 248
  - presynaptic inhibition: group II terminals, 292; Ia terminals, 339
- reticulospinal control in spasticity
  - cervical propriospinal neurones, 483–4, 579
  - presynaptic inhibition of Ia terminals, 370
  - Renshaw cells, 185, 186
- rostral location with respect to motoneurones
  - cervical propriospinal neurone, 459–60
  - group II interneurones, 303–4
  - lumbar propriospinal neurones, 493–4
- rTMS, 372
- RII reflex, biceps femoris, 393, 394, 414
  - afferent pathway, 418
  - central pathway, 418–19
  - habituation, 394
  - methodology, 394
  - placing reaction, 427
  - temporal summation, 394, 418
  - threshold, 418
- RIII reflex, other muscles, 396, 417
- RIII reflex, biceps femoris, 393, 394
  - afferent pathway, 399–400: A $\beta$ , 400; A $\delta$ , 393, 400
  - central pathway, 401
  - methodology, 394: single shocks, 400; trains, 393, 400
  - parallel with pain sensation, 393, 399–400
  - threshold, 400
  - use to monitor medication pain, 437
- running, contribution of spinal pathways
  - heteronymous Ia connections, 93–4, 550
  - presynaptic inhibition of Ia terminals, 367
  - short-latency Ia excitation, 87: quadriceps, 87; soleus, 87
- secondary endings (muscle spindle)
  - cat, definition and structure, 114, 115, 289
  - low dynamic sensitivity during perturbations of gait, 316
  - low dynamic sensitivity used for identification, 121–2
  - recordings from, 120, 121, 122, 132
  - sensitivity to stretch and vibration in cats, 114
  - sensitivity to vibration in humans, 130, 132, 289
- selectivity of movement, spinal mechanisms contributing to, 517
- cervical propriospinal system
  - focus of descending drive, 476
  - lateral inhibition, 477, 517
- focus of group II excitation, 514
- fusimotor drive, 122, 135, 517
- lumbar propriospinal neurones (descending inhibition), 502–3
- Ib inhibition, 273, 517
- presynaptic inhibition of Ia terminals, 359–60, 517, 529
- recurrent inhibition: heteronymous, 183–4, 529, 538; homonymous, 517
- sensitivity to facilitation and inhibition
  - H reflexes of different sizes, 16–18
  - H reflex vs. on-going EMG and PSTH, 25, 26–7, 203–4
  - paired H reflex technique: H' and reference H reflexes, 161
- servo-assistance, 114
  - group I effects from tibialis anterior to biceps, stabilisation of the knee in walking, 274–5, 547
  - group Ia projections to  $\alpha$  motoneurones
    - perturbations to stance, 89–90, 539, 540–1
    - unstable stance: heteronymous, 94, 538; homonymous, 538



- voluntary contractions: co-contractions of antagonists, 135, 532; hinge joint movements, 135, 516; tonic contractions, 122, 133, 512
- walking, stabilisation of the: ankle, 89, 93, 546, 548; knee, 545
- group Ia projections to cervical propriospinal neurones, 477, 516–17
- group II projections
  - FRA hypothesis, 389, 513, 514, 529
  - perturbations to stance, 539, 541; influence of the postural set, 313, 541
  - unstable stance, heteronymous, 313–14, 538; homonymous, 538
  - voluntary contractions, 512–14, 529
  - walking, contribution to soleus activation, 319–20, 545–6
- motor learning, 138, 530
- mutual reinforcement of Ia and group II excitations, 512–14
- see also* co-ordinated activation of various synergies through heteronymous Ia and group II connections
- spasticity, clinical features and pathophysiology
  - blockade of one mechanism, 560
  - clinical features, 558
    - clasp-knife phenomenon, 244, 245, 326, 558
    - clonus, 139–40, 558
    - reflex irradiation, 96, 131
    - synkinetic contractions, 557, 579
    - variations with the causative lesion, 557–8
- contribution to motor impairment, 558–60, 574
  - absence of relation with stretch reflex exaggeration, 558, 573–4
  - conflicting results, 559
  - overestimation, 560
  - variations with the underlying cause, 559: cerebral palsy, 559, 574; spinal cord lesion, 559, 574; stroke, 559
- definition, 557
- longitudinal studies, 556–7
- reflex irradiation, 96, 570
  - heteronymous connections, 570
  - transmission of vibration, 570
- relationship to decerebrate rigidity, 560
  - common features, 560
  - differences, 560
  - mechanisms underlying decerebrate rigidity, 560
- stretch reflex exaggeration, 557
  - increase in tone, 558
  - reflex gain, 557, 559
  - reflex threshold, 557, 559
  - tonic stretch reflex, 557
- therapeutic issues, 556
  - blockade of peripheral nerves by alcohol/phenol, 556
  - botulinum toxin, 556, 559
  - oral/intrathecal drugs, 556
  - physiotherapy, 556
- spasticity, findings at rest, transmission in spinal pathways, 560–71
  - absence of correlation with clinical spasticity, 571
  - $\alpha$  motoneurone excitability, 560–3
    - F wave, 562–3
    - H reflex, 96, 562: developmental slope, 562;  $H_{\max}/M_{\max}$  ratio, 562; recovery cycle, 562; reflex threshold, 562; transmission of the afferent volley, 562
  - how it could produce spasticity, 560
  - importance of contribution, 563
  - plateau potentials, 563
- fusimotor drive, 139–40, 563–5
  - animal models, 564
  - clonus, 139–40
  - flawed arguments for, 139, 564
  - how it could produce spasticity, 563
  - importance of contribution, 564, 565
  - microneurography, 139, 564
  - possibility of occurrence, 139, 564–5
- group II excitation, 320–6, 567–8
  - absence of correlations with clinical spasticity, 325
  - actions of monoamines, 321, 324–5, 568: group I excitation, 568; group II excitation, 568; interpretation with caution, 321–2, 325, 568; therapeutic action, 568
  - group II pathways and clasp-knife phenomenon, 326
  - how it could produce spasticity, 567
  - importance of contribution, 324–6, 568
  - methodology, 320–1, 567
  - positive feedback via excitation of  $\gamma$  motoneurons, 325, 567–8
  - possible mechanisms underlying increased group I excitation, 324: muscle stretch, 324; propriospinal contribution to tendon jerk, 324
- non-reciprocal group I inhibition (wrist), 234
- Ib excitation to antagonists, 277–9, 569
  - how it could produce spasticity, 569
  - importance of contribution, 278–9, 569
  - methodology, 569
  - Ib excitation vs. reciprocal Ia inhibition, 230, 277–8; vs. propriospinal excitation, 569
- Ib inhibition (ankle), 276, 568–9
  - how it could produce spasticity, 568
  - importance of contribution, 569
  - methodology, 275–6, 568

## 634 Index

- spasticity, findings at rest, transmission (*cont.*)  
 Ib inhibition vs. propriospinal excitation, 277, 568
- post-activation depression, 96, 99–100, 566  
 how it could produce spasticity, 566  
 importance of contribution, 566  
 methodology, 566
- presynaptic inhibition of Ia terminals, 368–70, 565–6  
 animal models, 565  
 how it could produce spasticity, 565  
 importance of contribution, 370, 566  
 methodology, 367–8: D1 inhibition, 565–6; heteronymous facilitation, 566; heteronymous tendon tap, 566  
 overestimation, 368, 566: homonymous vibration, 368, 565; sensitivity of the stretch reflex to presynaptic inhibition, 368  
 variations with the level of the lesion, 566
- propriospinal excitation, 503, 566–7  
 how it could produce spasticity: cervical, 567; lumbar, 566  
 importance of contribution, 567  
 methodology: cervical, 481, 567; lumbar, 567
- reciprocal Ia inhibition, 229–30, 570  
 distribution extensors–flexors, 232, 570  
 how it could produce spasticity, 570  
 importance of contribution, 570  
 methodology, 229, 570  
 plasticity: frequent peroneal stimulation, 233; training, 233
- recurrent inhibition, 184, 569  
 animal models, 569  
 findings using a flawed technique, 569  
 how it could produce spasticity, 184, 569  
 importance of contribution, 569  
 methodology, 184, 569  
 variations with the level of the lesion, 569
- spasticity, findings at rest, underlying mechanisms, 560–72  
 abnormal descending control, 571  
 $\alpha$  motoneurone excitability, 561  
 $\gamma$  drive, 563–4  
 group II excitation, 322–4, 567  
 Ib excitation to antagonists, 278, 569  
 Ib inhibition, 277  
 presynaptic inhibition of Ia terminals, 369–70  
 propriospinal excitation: cervical, 483–4, 567, 579; lumbar, 567  
 reciprocal Ia inhibition, 232  
 recurrent inhibition in hereditary paraparesis, 186
- plastic changes, 571  
 animal models, 572: sprouting, 572; supersensitivity, 572  
 excitability of  $\alpha$  motoneurons, 561, 572  
 excitability of  $\gamma$  motoneurons, 572  
 post-activation depression, 99–100, 572  
 reciprocal Ia inhibition, 233, 572
- spasticity, findings during motor tasks, 573–5  
 abnormal inhibition of the antagonists, 574, 575; due to  
 Ib facilitation of antagonists, 279, 575  
 presynaptic inhibition of Ia terminals, 575  
 propriospinally mediated inhibition of motoneurons, 575  
 reciprocal Ia inhibition, 231–2, 574–5:  $\alpha$ – $\gamma$  linkage, 574; corticospinal drive to Ia interneurons, 574; opposite Ia interneurons, 575; presynaptic inhibition, 574; recurrent inhibition, 575  
 recurrent inhibition, 575
- presynaptic inhibition of Ia terminals  
 voluntary contraction, 370, 575  
 walking, 368, 370–1, 575
- recurrent inhibition, 185, 186–7, 575
- spindle behaviour, 139, 140, 565
- synkinetic contractions, 484, 579
- spasticity, modification of properties of muscle fibres, 572–3  
 absence of abnormal stretch reflex, 559, 573  
 alterations in properties of muscle fibres, 573  
 changes related to inactivity of  
 motoneurons, 572–3  
 muscles, 572, 573  
 contracture, 557, 559, 560, 572  
 importance in stroke patients, 573
- spatial facilitation, background from animal experiments  
 convergence of excitation and inhibition, 45  
 excitatory convergence in: excitatory interneurons, 45; inhibitory interneurons, 45  
 underlying principles, 45, 46
- spatial facilitation, methodology  
 advantages and limitations, 48  
 initial sparing, 15, 46, 47, 212, 263, 267, 308–9, 456, 462, 463, 466, 472, 473, 498, 499  
 limitations of the H reflex method, 47–8  
 limitations of the PSTH method, 47  
 methodology for PSTHs of single units, summation in, 46–7:  
 excitatory pathways, 46–7; inhibitory pathways, 14–15  
 methodology for the H reflex, 46, 47
- spatial facilitation, used to demonstrate  
 contamination of the H reflex by Ib inhibition, 14–15  
 recurrent inhibition of Ia interneurons, 171–3, 205–8
- spatial facilitation, used to study convergence on  
 cervical propriospinal neurons, 461–3, 469  
 feedback inhibitory interneurons to propriospinal neurons: cervical, 464–7, 469, 478; lumbar, 310, 500  
 group II interneurons, 46–7, 307–9  
 Ia interneurons, 171–3, 202, 205–8, 214–15, 216–17

- Ib inhibitory interneurons, 257–8, 260–7
- lumbar propriospinal neurones, 46–7, 307–9, 496, 498
- non-reciprocal group I inhibitory interneurons (wrist), 212, 258, 524
- PAD interneurons acting on Ia terminals, 350–3
- Renshaw cells, 173
- spatial and/or temporal summation
  - occurring in
    - cervical propriospinal neurones, 460, 461, 469, 474, 476, 477
    - cutaneous reflex pathways, 391, 392, 394, 396
    - feedback inhibitory interneurons to cervical propriospinal neurones, 478, 483
    - group II interneurons, 290, 306, 316
    - Ia interneurons, 204
    - Ib interneurons, 261
  - making difficult quantitative assessment of effects, 261, 316, 474, 512
  - see also* convergence, spatial facilitation
- speed of movement
  - changes in the recruitment order of motoneurons favouring fast units
    - cervical propriospinal neurones, 477, 479
  - presynaptic inhibition of Ia terminals, 348
  - changes in spinal pathways related to the speed of the movement
    - presynaptic inhibition of Ia terminals, 358, 359, 518
    - recurrent inhibition, 174, 518
- spinal cord lesions, clinical features
  - contribution of spasticity to motor impairment, 559, 574, 580
  - from spinal shock to spasticity, 581–2
  - therapeutic issues, 580
- spinal cord lesions, transmission in spinal pathways at rest, 580–1
- $\alpha$  motoneurone excitability, 580
  - F wave, 580
  - H reflex, 580
  - plateau potentials, 580
- cutaneous reflexes
  - abolition of normal responses, 433
  - Babinski response, 433–4
  - cutaneomuscular responses, 423, 436, 437
  - early withdrawal reflexes, 433, 434
  - flexor spasms, 434–5
  - late withdrawal reflexes, 407–10
- fusimotor drive, 140, 580
  - positive feedback through group II projections, 140, 325, 580
  - reflex activation of  $\gamma$  motoneurons, 140, 580
- group II excitation, 322, 581
  - changes in group I-group II excitation, 322
  - effect of: clonidine, 322; DOPA, 322
  - positive feedback via excitation of  $\gamma$  motoneurons, 567–8, 581
  - underlying mechanisms, 324, 581
- lumbar propriospinal pathways, 503, 581; underlying mechanisms, 581
- Ib excitation to antagonists, 277, 581; underlying mechanisms, 278, 581
- Ib inhibition, 276, 277, 581
- post-activation depression, 100, 580
  - multiple sclerosis, 99
  - trauma, 99
  - underlying mechanisms, 99–100, 572
- presynaptic inhibition of Ia terminals, 369, 580; underlying mechanisms, 370, 580
- reciprocal Ia inhibition, 230
  - correlation with recovery, 232
  - focal lesions, distribution, 581
  - multiple sclerosis, 581
- recurrent inhibition, 581
  - progressive paraparesis: amyotrophic lateral sclerosis, 184, 581; hereditary spastic paraparesis, 186, 581
  - spinal cord injury, 184, 581
  - underlying mechanisms, 186, 581
- spinal transection, 433
  - Babinski response, 433
  - flexor spasms, 434–5
  - from spinal shock to spasticity, 433
  - late withdrawal reflexes, 407–10
  - see also* spinal cord lesions
- spino-bulbo-spinal pathways, 424
- standing, 535–42, *see* perturbations to stance, quiet stance, unstable stance
- startle response, 424
- stimulus rate, *see* repetition rate
- strategies for distal and proximal movements, 521–2
- stretch reflex (early spinal response, M1)
  - during co-contraction of antagonists: ball joints, 534–5; hinge joints, 534
  - functional role in, 87–90: hopping, 87–8; landing, 88–9; running, 87; standing, 89–90, 537; walking (ankle), 89, 548, 549
  - Ia origin, 87, 293, 320
  - Ib inhibition reversed during locomotion: cat, 66, 248; humans, 273–5
  - overlap with group II excitation, 3, 301

## 636 Index

- stretch reflex (*cont.*)
- perturbations to stance, 313, 539; function, 320, 540–1
  - quiet stance, 364, 537
  - prevention of stretch reflex during voluntary contraction of the antagonist, 180, 223, 273, 361, 519, 520, 525
  - sensitivity to presynaptic inhibition of Ia fibres, 354–5
  - walking: soleus, depending on stretch velocity, 89, 317, 548; despite presynaptic inhibition, 89, 548; tibialis anterior, 548
  - see also* parkinsonian rigidity, spasticity, stroke
- stroke, clinical features
- changes in properties of muscle fibres, 573
  - contracture, 557, 560, 572
  - contribution to motor impairment, 559, 573
  - incidence of spasticity, 575
  - stretch reflex in: elbow muscles, 559; triceps surae, 559, 573
  - tonic stretch reflex, 557, 575–6
- stroke, transmission in spinal pathways at rest, 576–7
- $\alpha$  motoneurone excitability, 576
  - F wave, 576
  - H reflex: FCR, 576; soleus, 576
  - cervical propriospinal neurones, 481–4, 579
    - asymmetry, 481–3
    - changes throughout recovery, 484
    - increased transmission of the descending command, 483, 579
    - methodology, 481
    - spasticity, 579
    - synkinetic contractions, 484, 579
    - underlying mechanisms, 483–4, 579
  - cutaneous reflexes
    - abolition of normal responses, 433
    - Babinski response, 433–4
    - cutaneomuscular responses, 423, 436, 437
    - withdrawal reflexes, 434
  - fusimotor drive, 139–40
    - contribution to motor deficit, 139, 140, 565
    - $\gamma$  activity, 139, 576
  - group II excitation (affected side), 322, 577
    - effect of tizanidine, 322
    - underlying mechanisms, 322–4, 577
  - lumbar propriospinal neurones, 503, 577; underlying mechanisms, 503, 577
  - non-reciprocal group I inhibition (wrist), 234, 577; correlation with spasticity, 577
  - Ib excitation to antagonists, 277, 577; underlying mechanisms, 278, 577
  - Ib inhibition, 276, 577
    - correlation with spasticity, 276
    - underlying mechanisms, 277, 577
  - post-activation depression (affected side), 99, 100, 577; underlying mechanisms, 99–100, 572
  - presynaptic inhibition of Ia terminals, 368–9, 576
    - lower limb, 368–9, 576
    - upper limb, 369, 576
    - underlying mechanisms, 369–70, 576
  - reciprocal Ia inhibition, 230, 577
    - correlation with recovery, 232, 577
    - underlying mechanisms, 577
  - recurrent inhibition, 184, 577; underlying mechanisms, 186
- stroke, unaffected side, 577–9
- abnormal transmission in spinal pathways, 577–9
    - presynaptic inhibition of Ia terminals, upper limb, 579
    - reciprocal Ia inhibition, 230, 278, 579
    - responses to stretch, 577–9
  - normal transmission in spinal pathways, 579
    - group II excitation, 322
    - lumbar propriospinal excitation, 503, 579
    - Ib excitation, 278, 579
    - Ib inhibition, 276, 579
    - post-activation depression, 99, 579
    - underlying mechanisms, 579
- stumbling reaction during locomotion, 387
- subliminal excitation of motoneurons
- created by a conditioning Ia volley, 154, 257
  - subliminal fringe of excitation of the monosynaptic reflex, 3, 4
- synergies
- see* co-ordinated activation of various synergies
  - synergies at wrist level
    - clenching, 211–12, 525
    - wrist abduction, 211, 522
- synkinetic contractions
- clinical features, 557
  - pathophysiology, 484, 579
- tendon jerk
- comparison with the H reflex, 117–18
    - delay with respect to H reflex, 9
    - differences: in the afferent volleys, 2–3, 117–18; at the motoneurone pool, 118
    - used as a measure of  $\gamma$  drive, 9, 117
  - general features
    - pathway, 2, 64
    - possible propriospinal contribution, 324
    - reflex irradiation, 96, 131
  - use to evoke recurrent inhibition, 162
  - use to study

- effects of noxious stimuli at knee level, 404–5
- spinal pathways at elbow level, 8: cervical propriospinal
  - excitation, 474; Ib excitation, 255; reciprocal Ia inhibition, 201
- tendon tap
  - use to distinguish between Ia and Ib effects, 252–3, 255
  - use to support the Ia origin of
    - group I excitation, 67, 75–6
    - reciprocal Ia inhibition, 204
- termination of movement, spinal mechanisms contributing to, 518
  - cutaneous inhibition of cervical propriospinal neurones, 478, 518
- facilitation of Ib interneurones
  - cutaneous afferents, 262, 271–2, 518
  - joint afferents, 263, 272, 518
  - multiple convergence, 265–7
- recurrent inhibition not involved, 176
- thixotropy, 124–6
  - advantages, limitations, perspectives, 126
  - underlying principles, 124
  - used to study fusimotor activation, 124–5
- threshold
  - cervical propriospinal excitation, 460
  - cutaneomuscular spinal responses, 415
  - feedback inhibition of cervical propriospinal neurones, 464
  - group II excitation, 293, 295, 297
  - lumbar propriospinal excitation, 491
  - monosynaptic Ia excitation, 64, 67–9, 75
  - Ib inhibition, 252
  - presynaptic inhibition (D1), 350
  - reciprocal Ia inhibition, 204
  - recurrent inhibition, 166, 171, 206
  - withdrawal RIII reflex, 400
- threshold to electrical stimulation
  - A $\beta$  cutaneous afferents, 400, 418
  - A $\delta$  cutaneous afferents, 400
  - group Ia afferents, 64, 67–9, 75, 204, 303
    - activity-dependent hyperpolarisation, 341
    - elevation by tendon vibration, 76, 205, 245, 260–1
  - in the inferior soleus nerve, 69
  - large range in human experiments, 77–8
  - with respect to motor threshold, 81
  - group Ib afferents, 245, 252, 258
  - group II afferents, 289, 293, 295, 297
  - group II/Ia ratio, 289, 303
- threshold tracking
  - H reflex, 11, 12
  - unitary H reflex, 37
- timing of changes in spinal pathways during co-contractions of antagonists
  - movement in progress: recurrent inhibition, 180
  - onset: presynaptic inhibition of Ia terminals, 361; reciprocal Ia inhibition, 225; recurrent inhibition, 180
  - period preceding contraction: presynaptic inhibition, 361
- timing of changes in spinal pathways to active motoneurones
  - during movement, 520–1
  - movement in progress
    - development of force, 521
    - fusimotor drive, 135
    - presynaptic inhibition, 358
    - recurrent inhibition, 174
  - onset, 520–1
    - cervical propriospinal pathways, 474–5
    - fusimotor drive, 133, 521
    - Ib inhibition, 270, 521
    - presynaptic inhibition, 355, 520
    - recurrent inhibition, 174, 520
  - period preceding contraction
    - fusimotor drive, 133
    - Ib inhibition, 270
    - presynaptic inhibition of Ia terminals, 358
    - recurrent inhibition, 174
- slowing eccentric contractions, 521
  - fusimotor drive, 521
  - recruitment order of motoneurones, 519
- termination, 518, 521
  - cutaneous inhibition of cervical propriospinal neurones, 478, 518
  - facilitation of Ib inhibition: cutaneous afferents, 272, 518; joint afferents, 272, 518
  - recurrent inhibition not involved, 176
- timing of changes in spinal pathways to antagonistic motoneurones during movement
  - movement in progress
    - presynaptic inhibition, 360–1
    - reciprocal Ia inhibition, 221
    - recurrent inhibition, 180
  - onset
    - presynaptic inhibition, 360
    - reciprocal Ia inhibition, 221
    - recurrent inhibition, 180
  - period preceding contraction
    - presynaptic inhibition of Ia terminals, 360
    - reciprocal Ia inhibition, 221
- see also* relaxation of antagonists

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**638 Index**

- tizanidine used to depress group II excitation  
 cat, 292  
 normal subjects, 294, 299; walking, 317, 318  
 stroke patients, 322, 323
- tonic contractions  
 co-contraction of antagonists  
   fusimotor drive, 135, 532  
   presynaptic inhibition of Ia terminals, 361, 532  
   reciprocal Ia inhibition, 225  
   recurrent inhibition, 180
- contraction of the antagonistic muscle  
 Ib inhibition, 273, 520  
 presynaptic inhibition of Ia terminals, 361  
 reciprocal Ia inhibition, 219–23  
 recurrent inhibition, 180
- contraction of the target muscle, 512–15  
 cervical propriospinal pathways, 476–7  
 cutaneomuscular responses, 514  
 fusimotor drive, 122, 133, 512  
 group II excitation, 310–12  
 lumbar propriospinal neurones, 500–2  
 Ib inhibition, 268, 271, 515  
 post-activation depression, 99  
 presynaptic inhibition of Ia terminals, 358  
 reciprocal Ia inhibition, 223  
 recurrent inhibition, 174
- transcortical cutaneous pathways  
 afferent pathway, 421  
 central pathway, 421–4  
   evidence for transcortical mediation, 422, 424  
   latency, 421–3: calculations, 421–2, 423; rostrocaudal  
     sequence, 421  
   observations in patients, 423  
   possible contribution of other pathways, 424
- E2 excitatory cutaneomuscular responses  
 lower limb, 415, 416  
 upper limb, 415, 417
- I1 inhibitory cutaneomuscular response, 415, 417  
 evidence for transcortical mediation, 423  
 possible function, 428
- maturation, 423, 428
- motor task-related changes  
 hopping, 432  
 voluntary contractions, 427, 428, 429  
 walking, 430–2: evidence for transcortical mediation,  
   430–1, 432; flexor muscles, 430, 431; functional role, 432,  
   549–50; pattern and timing, 430, 549; reflex reversal, 430
- Parkinson's disease, 437
- upper motoneurone lesions, 423, 436, 437
- transcortical group II pathways  
 Parkinson's disease, 584  
 standing, in the antagonist of the stretched muscle, 540;  
   significance, 541
- transcortical Ia pathways  
 hand muscles, 90, 92; functional significance, 92  
 history, 90–1  
 interaction with voluntary intent, 92, 539  
 lower limb, 91–2  
   functional stretch reflex, 91, 539  
   voluntary contraction, 548  
   walking, 548–9: in soleus, 548; in tibialis anterior, 548–9;  
     role, 549–50; upper motoneurone lesions, 548, 549  
 Parkinson's disease, 583, 584
- transcranial magnetic stimulation (TMS), *see* cortical  
 stimulation
- tremor, *see* parkinsonian tremor
- trisinaptic effects  
 disfacilitation of propriospinal neurones, 467  
 Ib excitation, 255
- trunk skin reflexes  
 central delay, 400  
 changes during voluntary contraction, 412  
 reciprocal organisation, 402  
 upper motoneurone diseases, 433  
*see also* abdominal skin reflexes
- unaffected side in hemiplegia, *see* stroke, unaffected side, 577–9
- unitary H reflex, 37–9  
 general features  
   advantages, 39  
   limitations, 39  
   underlying principles, 37, 38  
 methodology  
   basic methodology, 37  
   critical firing stimulus and its significance, 37, 38–9  
   recording, 37  
   stimulation, 37–8
- threshold tracking, 37
- used to study  
 presynaptic inhibition of Ia terminals (heteronymous Ia  
   facilitation), 38–9, 347  
 reciprocal Ia inhibition during contraction of the  
   antagonists, 219  
 single units at rest, 37  
 skewed distribution of corticospinal inputs at rest, 45
- unloading  
 voluntary contraction, 133  
 walking, unloading soleus, 315–16

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)

- EMG suppression, 315–16
  - methodology, 315
- unstable stance
  - co-contraction of synergists, 94, 537–8
  - cutaneomuscular responses, 430, 514
  - fusimotor drive, 135, 538
  - homonymous Ia and group II excitation, 38
  - heteronymous group II excitation, 313–14, 538
    - focus, 538
    - function, 314, 538
    - leaning: backwards, 313–14; forwards, 314
    - underlying mechanisms, 314, 538
  - heteronymous Ia excitation, 94, 538; focus, 183, 538
  - reciprocal Ia inhibition, 227
- updating of the descending command, 454, 477, 529
- upper motoneurone lesion, changes in
  - $\alpha$  motoneurone excitability, 560–3
  - cervical propriospinal pathways, 481–4, 579
  - cutaneous reflexes, 433–6
    - abolition of normal responses, 433
    - Babinski response, 433–4
    - cutaneomuscular responses, 423, 436, 437
    - withdrawal reflexes, 434–5
  - fusimotor drive, 139–40, 563–5
  - group II excitation, 320–6, 567–8
  - lumbar propriospinal pathways, 503, 566–7
  - non-reciprocal group I inhibition (wrist), 234
  - Ib excitation, 277, 569
  - Ib inhibition, 276, 568–9
  - post-activation depression, 99–100, 566
  - presynaptic inhibition of Ia terminals, 368–70, 565–6
  - reciprocal Ia inhibition, 229–33, 570
  - recurrent inhibition, 184–7, 569
  - transcortical group I pathways, 548, 549
  - transcortical cutaneous pathways, 423
  - withdrawal reflexes, 434–5
- upper motoneurone syndrome, 557
  - loss of dexterity, 557
  - negative symptoms, 557
  - positive symptoms, 557; reflex release, 557
  - release of flexor reflexes, 434–5, 557
  - spasticity, 557
  - weakness, 557
    - reciprocal inhibition from the stretched muscle, 574
    - spindle behaviour, 139, 140, 565
    - unaffected side, 579
- vestibulospinal control, background from animal experiments
  - group II interneurons, 292
  - lumbar propriospinal neurones, 491
  - Ia interneurons, 200
  - PAD interneurons, 339
- vestibulospinal projections
  - Ia interneurons, 216–17
  - Ib interneurons, 265
  - PAD interneurons, 353, 364
  - Renshaw cells, 181
- vibration of the homonymous tendon
  - activation of, 130–1
    - cutaneous mechanoreceptors, 131
    - Golgi tendon organs, 123, 130–1
    - primary endings, 130, 132
    - secondary endings, 130, 132
    - spindle endings during contraction, 131
  - activity-dependent hyperpolarisation of Ia afferents, 12–13, 341
  - post-activation depression at the Ia-motoneurone synapse, 341, 368, 371, 565
  - presynaptic inhibition of Ia terminals, 341
- vibration paradox, 341
- vibration, used to produce
  - activation of primary endings in
    - cats, 64, 114
    - humans, 130, 132
  - elevation of the electrical threshold of Ia afferents, 76, 205, 245, 260–1
  - presynaptic inhibition of Ia terminals
    - heteronymous, 341–2
    - homonymous, 341
- voluntary co-contraction of antagonists, 531–5
  - changes in spinal pathways to antagonists, 531–2
    - fusimotor drive, 135, 532
    - non-reciprocal group I inhibition (wrist), 523, 532
    - presynaptic inhibition of Ia terminals: ankle, 361, 532; wrist, 523, 532
    - reciprocal Ia inhibition (ankle), 225–7, 531
    - recurrent inhibition: ankle, 177, 180, 531–2; wrist, 523, 534
  - control of motor output and recurrent inhibition: hinge joints, 532; wrist, 534
  - joint stiffness, 533–4
  - pathways involved in co-contraction of antagonists
    - cervical propriospinal relay, 533
    - corticospinal drive, 225–6, 533
    - spinal pathways, 533
  - stretch reflex: hinge joints, 534; wrist, 534–5



Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**640 Index**

- voluntary contraction, *see*
  - concentric and eccentric contractions;
  - force, speed of movement;
  - onset, termination, timing of movement;
  - ramp and tonic voluntary contractions;
  - relaxation of antagonists;
  - voluntary co-contraction;
  - voluntary contraction of antagonist, remote, synergist, target muscles;
  - voluntary movement
- voluntary contraction of antagonists
  - changes in pathways to inactive antagonists, 519–20
    - non-reciprocal group I inhibition (wrist), 524–5
    - Ib inhibition (ankle), 273, 520
  - presynaptic inhibition of Ia terminals: lower limb, 360–1, 519; wrist, 363, 526
  - propriospinally mediated inhibition of motoneurones, 217–19, 503, 519–20
  - reciprocal Ia inhibition, 217–3, 519
  - recurrent inhibition, 180, 520
  - spindle activation, 135, 221–2
- functional significance of changes in spinal pathways
  - non-reciprocal group I inhibition (wrist), 525–6
  - Ib inhibition (ankle), 273, 520
  - presynaptic inhibition of Ia terminals, 361
  - propriospinally mediated inhibition of motoneurones, 503, 520
  - reciprocal Ia inhibition on: motoneurones, 223, 520;
    - opposite Ia interneurones, 222, 223, 520
  - recurrent inhibition (ankle), 180, 520
  - spindle activation, 137–8
- voluntary contraction of remote muscles (Jendrassik manoeuvre)
  - central changes in the gain of the reflex, 133, 134
  - fusimotor drive, 131; flawed hypothesis, 131–2
  - presynaptic inhibition of Ia terminals, 133, 361–2
  - reciprocal Ia inhibition, 133, 227
- voluntary contraction of synergistic muscles
  - changes in pathways to inactive synergists
    - fusimotor drive, 135
    - Ib inhibition, 273, 517
  - presynaptic inhibition of Ia terminals, 359–60
  - recurrent inhibition, 183
- focus on contracting muscles
  - cervical propriospinal neurones, 476, 477, 517
  - fusimotor drive, 517
  - Ib inhibition, 273, 517
  - presynaptic inhibition of Ia terminals, 359–60, 517, 529
  - recurrent inhibition, 183–4, 529, 538
- voluntary contraction of the target muscle
  - changes in spinal pathways to active muscles, 513, 516–17
    - cervical propriospinal neurones, 471–9
    - cutaneomuscular responses, 427–30, 514
    - fusimotor drive, 133–6, 512–14, 516
    - group II excitation, 310–12
    - lumbar propriospinal neurones, 500–2
    - non-reciprocal group I inhibition (wrist), 523, 524
    - Ib inhibition, 268–71, 515, 517; facilitation by other afferents, 271–2, 515
    - post-activation depression, 99
  - presynaptic inhibition of Ia terminals: lower limb, 355–8;
    - wrist, 362–3, 526
  - reciprocal Ia inhibition, 223–5, 517
  - recurrent inhibition, 173–9, 514–15, 517
  - withdrawal reflexes, 412–14
- functional significance of changes in spinal pathways
  - cervical propriospinal excitation, 477–8, 516–17, 529
  - feedback inhibition of cervical propriospinal neurones:
    - cutaneous, 478, 518; group I, 477, 517, 518
  - fusimotor drive, 136–8, 512; slowing eccentric contractions, 521
  - group II excitation, 312, 514
  - lumbar propriospinal excitation, 502–3
  - non-reciprocal group I inhibition (wrist), 523, 525–6
  - Ib inhibition, 271, 517, 518; facilitation by other afferents, 272, 515
  - presynaptic inhibition of Ia terminals, 359, 517, 518
  - reciprocal Ia inhibition, 225, 518
  - recurrent inhibition, 179, 517, 518
- voluntary movement, control of different features
  - co-ordinated synergies, 527–1
  - force and speed, 359, 517–18
  - inhibition of antagonists: hinge joints, 513, 519–20; wrist, 523, 525–6
  - pathways to active motoneurones, 513, 516–17; excitatory, 516–17; inhibitory, 517
  - recruitment of different motoneurone types, 518–19
  - selectivity, 517
  - strategies for distal and proximal movements, 521–2
  - termination, 518
  - timing, 520–1
  - tonic contraction vs. movement, 522
- walking, characteristics of human gait, 542–4
  - biomechanical characteristics, 542–4
  - different roles of ankle and knee extensors, 273–4
  - lengthening loaded contractions, 544



- pattern of muscle activation, 543, 544
  - use of external energy, 544
- walking, changes in spinal and transcortical pathways
  - fusimotor drive, 137, 544
  - group II excitation, 545–6
    - quadriceps (heteronymous), 318, 546
    - soleus (homonymous), 315–16, 545–6
  - monosynaptic Ia excitation of soleus
- Ib inhibition, 274, 546
  - heteronymous, 93
  - homonymous, 89
  - stretch reflex, 89
- presynaptic inhibition of
  - quadriceps Ia terminals, 365, 545
  - soleus Ia terminals, 365–6, 545
- reciprocal Ia inhibition (ankle), 227–9, 546
- recurrent inhibition (knee), 546
- stretch responses in soleus
  - group II, 316–18, 548
  - monosynaptic Ia, 89, 548
  - transcortical, 548
- stretch responses in tibialis anterior
  - group II, 316, 318, 548
  - monosynaptic Ia, 548
  - transcortical, 548–9
- transcortical cutaneous responses, 430–2, 549
  - evidence for, 430–2
  - pattern and timing, 430, 549
  - reflex reversal, 430
- walking, role of spinal pathways
  - contribution to soleus activation, 319–20, 545–6
  - cutaneous facilitation of reciprocal Ia inhibition, 214–15, 549
  - prevention of excessive Ia activity in soleus, 546–7
    - Ib inhibition, 274, 546–7
  - pattern of Ia projections, 93, 546
  - presynaptic inhibition, 366–7, 546
  - propriospinally mediated inhibition of motoneurons, 503, 547
- stabilisation of the ankle
  - heteronymous Ia projections, 93, 546
  - reciprocal Ia inhibition, 229, 546
  - stretch responses, 318, 546, 549
- stabilisation of the knee
  - cutaneous suppression of Ib inhibition to quadriceps, 274, 547
  - group I-II discharge from tibialis anterior, 319, 547
  - heteronymous Ib effects to biceps, 274–5, 547
  - presynaptic inhibition, 365, 547
  - recurrent inhibition, 547
- stumbling, 94, 548
- transition, 547
- yield of the knee, 365, 547
- walking, role of transcortical pathways
  - cutaneous pathways, 432, 549–50
  - group I pathways, 549–50
  - stretch responses in ankle muscles, 549
- walking, studies in patients with upper motoneurone lesions
  - presynaptic inhibition of Ia terminals, 370–1, 575
  - transcortical group I excitation, 548, 549
  - withdrawal reflexes, 435
- weakness, *see* upper motoneurone syndrome
- withdrawal reflexes, background from animal experiments
  - FRA pathways, 388–91: early, 389–90; late, 390–1
- initial findings, 385
- private pathways of withdrawal reflexes, 387
  - crossed extension reflex, 385
  - extensor activation, 387
  - flexor reflex, 385, 387
  - modular organisation, 386, 387
- withdrawal reflexes, methodology
  - critique of tests, 396–8
  - stimuli
    - effects on late responses of increased stimulus intensity or duration in: normal subjects, 410; patients with spinal transection, 408
    - electrical, 392
    - mechanical, 392, 394
- modulation of motoneurone excitability, 396
  - monosynaptic reflexes, 396; on-going EMG, 396
- reflexes at rest: RIII in the biceps femoris, 394; other
  - withdrawal reflexes, 394, 395
- withdrawal reflexes, organisation, 399–414
  - afferent pathway of the RIII reflex, 399–400
    - afferents: A $\beta$ , 400; A $\delta$ , 400
    - pain sensation, 399–400: nerve, 399; skin, 399–400
- central pathway, 400–1: central delay, 400–1; complete spinal transection, 401
- conditioning by afferents, 411–12
  - nociceptive afferents, 411: early facilitation, 393, 411; late depression, 395, 411; underlying mechanisms, 411
  - tactile afferents, 393, 411–12; underlying mechanisms, 412
  - warmth, 412
- descending effects
  - early FRAs, 412; abdominal skin reflexes, 433; plantar response, 434
  - late FRAs in normal subjects, 411
- functional organisation, 401–7
  - convergence of tactile afferents, 402, 405

Cambridge University Press

0521825814 - The Circuitry of the Human Spinal Cord: Its Role in Motor Control and Movement Disorders

Emmanuel Pierrot-Deseilligny and David Burke

Index

[More information](#)**642 Index**

- withdrawal reflexes, organisation (*cont.*)
  - local sign, 397, 402–3, 404–5, 407
  - modular organisation, 407
  - lower limb responses, 404–5: afferent pathway, 401; central delay, 401; modulation of monosynaptic reflexes, 397, 404–5; receptive fields, 397, 404–5
  - plantar responses, 402–4: extensor hallucis longus, 402; maturation, 404; receptive fields, 402–4
  - trunk muscles, 402
  - upper limb responses, 405–7: central delay, 401, 405; evidence for motoneurone inhibition, 405, 406; responses at rest, 405–6, 407; silent period, 405, 406
- protective function, 399, 401–2, 404, 405, 406, 407
- rIII reflex in the biceps femoris: afferent pathway, 400; central pathway, 401; pain sensation, 399–400; threshold, 400
- reciprocal organisation, 397, 402, 404, 405
- withdrawal reflexes (late) in normal subjects, 397, 410–11
  - characteristics, 410: latency, 410; threshold, 410
  - plasticity, influence of, 410: continuous stimulation, 410; hypnosis, 410; posture, 410
  - supraspinal mediation, 411
- withdrawal reflexes, physiological implications
  - postural tasks, 413, 414
  - voluntary contraction, 412–14
    - leg muscles, 412–13, 414
    - trunk muscles, 412
- withdrawal reflexes, studies in patients
  - Babinski response, 403, 434
  - complete spinal transection, 433
  - Parkinson's disease, 436–7, 589
  - peripheral neuropathies, 437
  - upper motoneurone lesions, lower limb, 434–5
    - alterations of withdrawal reflexes, 434
    - extensor inhibition, 435
    - flexor spasms, 434–5
    - plantar nerve stimulation, 435
    - walking, 435
  - upper motoneurone lesions, upper limb, 435
  - withdrawal reflexes (late), patients with spinal transection, 407–10
    - background from animal experiments, 390–1
  - characteristics
    - effects of increased stimulus intensity or duration, 408, 409
    - inhibition from contralateral FRAs, 408, 409
    - inhibition from early FRAs, 408, 410
    - long-latency ipsilateral flexion, 407–8
  - contralateral effects, 408, 409
    - post-synaptic facilitation of extensors, 408
    - presynaptic inhibition of Ia terminals, 408
  - contribution of high-threshold muscle afferents, 433
  - evidence for spinal reflexes, 407
  - from spinal shock to spasticity, 433
  - larger receptive field, 433
  - methodology, 407, 433
- wrist, *see* organisation of spinal circuitry at wrist level
- yield of the knee, 365, 547