

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)

Index

- actin cytoskeleton
 - role in hyphal extension 95
 - role in polarized growth 103, 108–10
- aflatoxin B1 25–7
 - biosynthesis 30–1, 32
- agriculture *see* crop protection
- Agriotes* spp. (wireworms, click beetle larvae), life cycle and economic relevance 261–2
- alkaloids, produced by filamentous fungi 25
- Allomyces macrogynus*, microtubule initiation 98, 99–100, 101
- antibiotics (fungal) 45–6
 - appressorium formation inhibitors 52, 53–6
 - autoinhibitors 52–3
 - classification by target stage 52
 - conidial germination inhibitors 52–3
 - echinocandins 51–2
 - fungal infection-related process inhibitors 52–6
 - lead compounds for fungicides 50–2
 - melanin biosynthesis inhibitors 55
 - pneumocandins 51–2
 - sordarins 50–1
- appressorium formation 25–7
 - fungal antibiotic inhibitors 52, 53–6
- Armillaria mellea*, bioluminescence-based biosensor 195–6
- Arthrobotrys oligospora*, nematode-trapping fungus 316–18
- Ashbya gossypii*, polarized growth 96
- aspergillosis 144–5
- Aspergillus* spp.
 - commercial use for enzyme production 145–7
 - ER stress responses 147–53
 - gene arrays 143–4
 - genome sequences 140–1
 - genome sizes and predicted gene numbers 141
 - polyketoid production 25–7
 - predicted genes and encoded products 142–3
 - protein secretion 140–1
 - protein secretion stress 147–53
 - proteins secreted 144–7
 - selective antibiotics produced 51–2
 - UPR in heterologous protein production 146–53
- Aspergillus* spp. secondary metabolism
 - biosynthetic gene pathways and products 84–7
 - clavines 86–7
 - fumitremorgins 85–6, 89
 - functional analysis of gene clusters 87–8
 - gliotoxin 85
 - pharmacological activity 89
 - secondary metabolic genes and clusters 83–4
 - tryprostatins 89
- Aspergillus flavus*, genomics 66 *see also* *Aspergillus* spp.
- Aspergillus fumigatus* 78–9, 82
 - aspergillosis 144–5
 - fumitremorgins 85–6, 89
 - genomics 141–4
 - gliotoxin 85
 - tryprostatins 89
 - see also* *Aspergillus* spp.
- Aspergillus nidulans*
 - genome 12–13
 - genomics 141–4
 - hyphal extension 95
 - microtubule initiation 98–102
 - microtubule plus-end dynamics 102–4

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)

Index

337

- polarized growth 96
- see also Aspergillus spp.*
- Aspergillus nidulans* secondary metabolism
 - dimethylallyltryptophan synthase (DMATS) 80, 82–3
- indole alkaloid biosynthetic pathways 80, 82–3
- information from genome sequencing 78–9
- major groups of metabolites 79–83
- non-ribosomal peptide synthetases (NRPS) 80–2
- PKS-NRPS hybrid genes 82
- polyketide synthases (PKS) 80, 81, 82
- terpene cyclases 80, 83
- see also Aspergillus spp.* secondary metabolism
- Aspergillus niger*
 - genome project 5
 - genomics 141–4
 - see also Aspergillus spp.*
- Aspergillus oryzae*
 - functional analysis of secondary metabolic clusters 87–8
 - genomics 141–4
 - see also Aspergillus spp.*
- Aspergillus terreus* 83
 - lovastatin production 25–7
- Athous spp.* (wireworms), life cycle and economic relevance 261–2
- auxostat technology *see* chemostat technology
- avermectin 45–6
- azoxystrobin 49
- Beauveria spp.*
 - biological control formulations 265–8
 - early research as biological control 263–4
 - fungal biocontrol agents 258–9
 - taxonomy 264–5
- Beauveria bassiana*
 - changing hosts 326, 327
 - leaf and stem endophytic behaviour 323–4
 - tenellin production 41–2, 43
- Beauveria brongniartii*, control of *M. melolontha* (European cockchafer) 268, 269
- benzene, degradation metabolic pathways 215–16
- bindweeds
 - perennial weeds of crops 275–6
 - potential biocontrols 276
 - problems with chemical control methods 276
- bioaccumulation of metals in fungi 246, 247
- bioaugmentation, role of white-rot fungi 206–7
- biocontrol agents (fungal) 263–5
 - Beauveria spp.* 263–5
 - early research on subterranean pests 263–4
 - formulations and application methods 265–8
 - Metarhizium spp.* 263–4, 265
 - risk assessment (*Stagonospora convolvuli* LA39 model) 282–7
 - soil pest control 258–9
 - taxonomy 264–5
- biofungicides *see* fungicides; *Trichoderma*-based biofungicides
- bioluminescence 190–1
 - in yeasts and filamentous fungi 189–90, 195–6
- bioluminescence-based biosensors
 - bacteria 191–2
 - comparative performance 196–8
 - contaminated soils applications 198–9, 200
 - environmental applications 198–9, 200
 - filamentous fungi 195–6
 - use in bioremediation 187–8
 - yeast (*S. cerevisiae*) 192–5
- bioprocess development, chemostat technology with filamentous fungi 7–8
- biorefinery
 - Aspergillus niger* genome project 5
 - contribution of fungal genomics 4–5
 - petroleum dependence reduction 3–4
 - process development with chemostat technology 7–8
 - role of fungal proteomics 6–7
 - role of fungi 3–4
 - systems biology approach 8–9
 - top 12 chemical building blocks 3–4
 - Trichoderma reesei* genome project 5–6
- bioremediation
 - bioaugmentation approach 206
 - bioluminescence-based biosensor applications 187–8, 198–9, 200
 - biostimulation approach 206
 - field trials of white-rot fungi 223–4
 - importance of mycorrhizal fungi 238
 - in situ* bioremediation 205–7
 - ligninolytic enzyme-redox mediator systems 207–13
 - mycoremediation approach 248–50
 - toxic metal contamination 237–8
 - xenobiotic contamination 205–7
 - see also* xenobiotic metabolism; xenobiotics; white-rot fungi
- biosensors
 - bioluminescence 189–90
 - microbial 188–90
 - roles in toxicity testing 188–90
 - yeasts and filamentous fungi 188–90

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)338 *Index*

- biosorption of metals to fungal cell walls 246
- black vine weevil *see* *Otiorhynchus sulcatus*
- Blumeria graminis* 53–4
- Botrytis cinerea*, fungicides used to control 45–6
- brownfield sites *see* bioremediation
- BTEX compounds, degradation metabolic pathways 215–16
- Calystegia sepium* (hedge bindweed)
 - perennial weed of crops 275–6
 - potential biocontrols 276
 - problems with chemical control methods 276
- Candida albicans*
 - action of sordarins 50–1
 - polarized growth 96
- cellulase production, *Trichoderma reesei*
 - genome project 5–6
- cerulenin 55
- chemostat technology, filamentous fungi
 - process development 7–8
- chlorophenols, degradation metabolic pathways 213–14, 215
- Cladosporium* sp. 53
- Claviceps purpurea* 82–3
 - cause of ergotism from contaminated rye 61–2
 - ergotamine biosynthesis 86–7
- clavines, in *Aspergillus* spp. 86–7
- Coccidioides immitis*, action of sordarins 50–1
- Cochliobolus heterostrophus* 62–3
 - T-toxin production 25–7
- Coleophoma empetri*, selective antibiotics produced 51–2
- Colletotrichum* spp., conidial germination inhibitors 52–3
- Colletotrichum lagenarium*
 - appressorium formation 25–7
 - polyketoid production 25–7
- Colletotrichum lindemuthianum*, mode of action of fungicides 52
- conidial germination inhibitors 52–3
- contaminated land *see* bioremediation
- continuous culture technology *see* chemostat technology
- Convolvulus arvensis* (field bindweed)
 - perennial weed of crops 275–6
 - potential biocontrols 276
 - problems with chemical control methods 276
- Coprinus heptemerus* 53
- Corymbites* spp. (wireworms), life cycle and economic relevance 261–2
- crop protection
 - development of new fungicides 45
 - resistant fungal pathogens 45
 - use of natural products 45–6
 - see also* fungicides; phytopathogenic fungi; *Trichoderma*-based biofungicides
- Cryptococcus neoformans*, action of sordarins 50–1
- cyclosporin 62–3
- Daktulosphaira vitifoliae* (grape phylloxera)
 - control by *Metarhizium anisopliae* 269
 - life-cycle and economic relevance 262
- Diabrotica virgifera virgifera* (western corn root worm)
 - control by *Metarhizium anisopliae* 269–70
 - life-cycle and economic relevance 263
- dimethylallyltryptophan synthase (DMATS), in *Aspergillus nidulans* 80, 82–3
- Drechmeria coniospora*, changing hosts 327
- dyes, degradation metabolic pathways 217
- dynein, role in hyphal extension 95
- echinocandins 51–2
- endophytes *see* *Epichloë* endophytes
- endoplasmic reticulum (ER)
 - protein folding 117–21
 - protein folding demand stress responses 122
 - RESS (repression under secretion stress) response 148–53
 - UPR (unfolded protein response) 122
 - UPR in *Aspergillus* spp. 147–53
- entomopathogenic fungi 310, 311, 312–13
 - canonical host barriers 314–15
 - comparisons with nematophagous fungi 324–6
 - evolution 328–9
 - examples of changing hosts 326–8
 - host physical barriers to overcome 313–16
 - insect host barriers 314–15
 - insect infection process 319–21
 - leaf and stem endophytes 323–4
 - multimodal behaviour and evolution 328–9
 - multimodal behaviour and host range 316, 317
 - multimodal behaviour and use as biocontrol 316, 317, 329–30
 - mycoparasitic behaviour 321
 - non-canonical host barriers 314, 315–16
- environmental toxicity *see* bioremediation
- Epichloë* endophytes
 - anti-herbivore effects 60–2
 - benefits for the host grass 60
 - benefits from mutualism 60
 - biological effects 60–2
 - engineering for desirable bioactivities 72

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)

Index

339

- ergot alkaloid synthesis 61–2
- exploitation of indole-diterpene
 - biosynthesis pathways 66, 70–2
- indole-diterpene biosynthesis 61, 62
- indole-diterpene biosynthesis model 65–7
- indole-diterpene chemical diversity 61, 65
- indole-diterpene gene clusters 66, 67–70
- loline alkaloid synthesis 61
- peramine synthesis 60–1
- selection for desirable chemotypes 72
- source of non-ribosomal peptides 63
- taxonomy 59–60
- Epichloë festucae*
 - molecular cloning of a peramine synthetase 61, 63–5
 - symbiotic fungal endophyte of grasses 59–60
 - synteny with *Fusarium graminearum* 63, 64
- Epichloë typhina* 59–60
- ER *see* endoplasmic reticulum
- ergot alkaloids, synthesis by *Epichloë*
 - endophytes 61–2
- ergotamine 61–2
- ergovaline 61–2
- ethylbenzene, degradation metabolic pathways 215–16
- European cockchafer *see* *Melolontha melolontha*
- explosive waste, degradation metabolic pathways 216
- fatty acids, appressorium formation
 - inhibitors 54–5
- fenpiclonil 45–6
- 'fescue toxicosis' in cattle 61–2
- Festuca* spp., symbiotic *Epichloë* endophytes 59–60
- filamentous actin (F-actin), role in hyphal extension 95
- flaviolin 53
- fludioxonil 45–6
- fumitremorgins, in *Aspergillus* spp. 85–6, 89
- fumonisin 25–7
- fungal antibiotics *see* antibiotics (fungal)
- fungal biocontrol agents *see* biocontrol agents (fungal)
- fungicides
 - development of new agrochemicals 45
 - emergence of resistant strains 45, 50
 - fungal metabolites as lead compounds 46–9, 50–2
 - lead finding process 56
 - natural antibiotics 45–6
 - scale of production 45
 - see also* *Trichoderma*-based biofungicides
- fusarin synthase 38–41
- Fusarium graminearum*
 - genome 12–13
 - synteny with *Epichloë festucae* 63, 64
- Fusarium moniliforme* 82
- fusarin C production 38–41
- Fusarium sporotrichioides* 83
- Fusarium venenatum*, fusarin C production 38–41
- garden chafer *see* *Phyllopertha horticola*
- gene clusters in fungal polyketoid
 - biosynthesis 31–3
 - investigation methodology 33–6
- genomics (fungal) 4–6
 - Aspergillus niger* genome project 5
 - Aspergillus* spp. genome sequences 140–1
 - Aspergillus* spp. genome sizes and predicted gene numbers 141
 - Aspergillus* spp. secondary metabolism 78–9
- contributions to industrial mycology 4–5
- fungal genome sequencing
 - projects 78–9
- identification of fungal secondary metabolites 78–9
- substrate range indications 4–5
- Trichoderma reesei* genome project 5–6
- Gibberella fujikuroi*, fumonisin
 - production 25–7
- Gliocladium roseum* 54–5
- gliotoxin, in *Aspergillus* spp. 85
- glisoprenins 54–5
- grape phylloxera *see* *Daktulosphaira vitifoliae*
- grasses (Pooidae), symbiotic fungal associations 59
- heptermone G 53
- heterologous protein production, *Aspergillus* spp. 146–53
- Histoplasma capsulatum*, action of
 - sordarins 50–1
- hyphal extension *see* polarized growth
- Hypocrea jecorina* 5–6
- indole alkaloid biosynthetic pathways, *Aspergillus nidulans* 80, 82–3
- indole-diterpene biosynthesis
 - exploitation of pathways 66, 70–2
 - Penicillium paxilli* model 65–7
- indole-diterpene gene clusters, in *Epichloë*
 - endophytes 66, 67–70
- indole-diterpenes
 - chemical diversity in *Epichloë*
 - endophytes 61, 65
 - mammalian tremorgens 62
 - synthesis by *Epichloë* endophytes 61, 62
- induced systemic resistance (ISR) 298–304

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)340 *Index*

- industrial mycology, contributions of fungal genomics 4–5
- insect fungal parasites *see* entomopathogenic fungi
- integrated pest management (IPM), for subterranean pests 258–9
- invertebrate fungal parasites *see* entomopathogenic fungi; nematophagous fungi
- kinesins, role in hyphal extension 95
- kresoxim methyl 49
- laccase 207–9, 210–11, 213
- Lecanicillium lecanii*, changing hosts 326
- lignin, breakdown by white-rot fungi 207–9
- lignin peroxidase 207–10, 211–13
- ligninolytic enzyme-redox mediator systems, in white-rot fungi 207–13
- Listronotus bonariensis* (Argentine stem weevil) 63–4
- loline alkaloids, synthesis by *Epichloë* endophytes 61
- lolitrem B, in *N. lolii*-perennial ryegrass associations 61, 65
- lolitrem 61, 62
- Lolium perenne*, symbiotic *Epichloë* endophytes 59–60
- lovastatin, biological properties 25–7
- lovastatin biosynthesis 29, 30
 - genes encoding PKS and post-PKS enzymes 31–3
- Magnaporthe grisea* (rice blast disease pathogen)
 - ace1 gene cluster 43
 - annual crop loss caused 10–11
 - appressorium formation 10–11, 16–18, 25–7
 - appressorium formation inhibitors 54–5
 - conidial germination inhibitors 53
 - evolution of pathogenicity 11–13
 - expansion of secreted proteins 13
 - functional analysis 14–18
 - functions of secreted proteins 15–16
 - G-protein-coupled receptors (GPCRs) 13–14
 - gene conservation 12
 - genome evolution 11–13
 - genome sequence 11
 - genome-wide transcriptional profiling 16–18
 - infection process 10–11
 - large-scale mutagenesis 15, 16
 - mode of action of fungicides 52
 - natural antibiotics used to control 45–6
 - novel signalling machinery 13–14
 - PAMPs (pathogen-associated molecular patterns) 13
 - polyketoid production 25–7
 - rapid development of pathogenic variants 10–12
 - rapid expansion of repetitive elements 11–12
 - synteny with other filamentous ascomycetes 12–13
- manganese-peroxidase 207–9, 210, 211–13
- May beetle *see* *Melolontha melolontha*
- melanin biosynthesis inhibitors 55
- Melolontha melolontha* (European cockchafer, May beetle)
 - control by *Beauveria brongniartii* 268, 269
 - life-cycle and economic relevance 259
- metal and mineral transformations 238–40
- immobilization by fungi 239–40, 246–7, 248, 249
- in the mycorrhizosphere 247–8
- interactions with mycorrhizal fungi 239–45
- solubilization by fungi 239–45
- metal-induced membrane damage 167–72
 - membrane-lipid peroxidation in Cd toxicity 168–72
- metal-induced protein damage 172–5
 - haploinsufficiency phenotypes 174–5
 - identification of protein targets of Cr toxicity 173, 174–5
 - protein (methionine) oxidation in Cr toxicity 172–5
- metal-induced ROS generation 163–5
 - oxidative damage to cell membranes 164–5
 - oxidative damage to DNA 165
 - oxidative damage to lipids 164
 - oxidative damage to proteins 164–5
- metal pollution
 - remediation 236
 - sources 236
- metal toxicity
 - conditions when heterogeneity is advantageous 178–9
 - evidence for ROS involvement 165–6
 - heterogeneous Cu resistance in *S. cerevisiae* 176–8
 - phenotypic heterogeneity in metal sensitivity 175–9
 - responses to environmental conditions 178–9
 - role of oxidative stress in *S. cerevisiae* 166–7
 - S. cerevisiae* model 162–3
 - suppression of heterogeneity in a stable habitat 178
- metals in the environment 161–2

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)

Index

341

- bioaccumulation in fungi 246, 247
- biosorption to fungal cell walls 246
- interactions with fungi 236
- interactions with microorganisms 162
- Metarhizium* spp.
 - biological control formulations 265–8
 - early research as biological control 263–4
 - taxonomy 264, 265
- Metarhizium anisopliae*
 - Dactulosphaira vitifoliae* (grape phylloxera) control 269
 - Diabrotica virgifera virgifera* (western corn root worm) control 269–70
 - fungal biocontrol agent 258–9
 - Otiorhynchus sulcatus* (black vine weevil) control 269
 - Phyllopertha horticola* (garden chafer) control 268
- 6-methyl salicylic acid 25–7
- 6-methyl salicylic acid synthase 29, 30
- microtubule-associated proteins 104, 105
- microtubule cytoskeleton, morphology, organization and dynamics 97–8
- microtubule-dependent motor proteins 105–8
 - dynein 105, 107–8
 - kinesins 105–7
 - role in hyphal extension 95
- microtubule filament lattice 104, 105
- microtubule organizing centre (MTOC) 95, 98–102
- microtubules
 - initiation 98–102
 - plus end dynamics 102–4
 - role in hyphal extension 95
 - role in polarized growth 103, 108–10
 - role of spindle pole bodies 98, 99–100, 101
- motor proteins *see* microtubule-dependent motor proteins
- mucidin 46
- Mycena citricolor*, bioluminescence-based biosensor 195–6
- mycoherbicide risk assessment (*Stagonospora convolvuli* LA39 model) 282–7
- mycorrhizal fungi
 - importance in bioremediation 238
 - solubilization of metal minerals 239–45
- myosins, role in hyphal extension 95
- nematode fungal parasites *see* nematophagous fungi
- nematophagous fungi 310, 311–12
 - canonical host barriers 314–15
 - comparisons with entomopathogenic fungi 324–6
 - evolution 328–9
 - examples of changing hosts 326–8
 - host physical barriers to overcome 313–16
 - multimodal behaviour and evolution 328–9
 - multimodal behaviour and host range 316, 317
 - multimodal behaviour and use as biocontrol 316, 317, 329–30
 - mycoparasitic behaviour 314, 321
 - nematode egg-destroying fungi 318–19
 - nematode host barriers 314–15
 - nematode infection process 316–19
 - nematode-trapping fungi 316–18
 - non-canonical host barriers 314, 315–16
 - root endophytes 322–3
- Neobulgaria pura* 54–5
- neobulgarones 54–5
- Neotyphodium* spp., evolution and taxonomy 60 *see also* *Epichloë* endophytes
- Neotyphodium lolii*
 - molecular cloning of a peramine synthetase 61, 63–5
 - symbiotic fungal endophyte of grasses 59, 60
- Neotyphodium lolii*–perennial ryegrass associations, production of lolitrem B 61, 65
- Neurospora crassa*
 - genome 12–13
 - microtubule dynamics 100–2
 - polarized growth 96
- non-ribosomal peptide synthetases (NRPS)
 - in *Aspergillus nidulans* 80–2
 - in filamentous fungi 62–3
- non-ribosomal peptides, biologically active therapeutic agents 62–3
- norsolorinic acid 25–7
- olive mill waste-water, degradation
 - metabolic pathways 216
- Omphalotus olearius*, bioluminescence 195
- organophosphate pesticides, degradation
 - metabolic pathways 215
- orsellinic acid 25–7
- orsellinic acid synthase 28–9
- Otiorhynchus sulcatus* (black vine weevil)
 - control by *Metarhizium anisopliae* 269
 - life-cycle and economic relevance 260–1
- Oudemansiella mucida* 46
- oudemansin 46, 49
- Paecilomyces* spp., changing hosts 326
- PAHs (polyaromatic hydrocarbons)
 - degradation metabolic pathways 217–20, 221

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)342 *Index*

- PAHs (polyaromatic hydrocarbons) (cont.)
 oxidation by fungal enzyme-redox mediator systems 208
Panellus stipticus, bioluminescence 195
 paxilline biosynthesis 65–7
 penicillin 62–3
Penicillium spp.
 polyketoid production of 25–7
 strobilurin-resistant species 46–9
Penicillium chrysogenum, source of penicillin 62–3
Penicillium paxilli, model of indole-diterpene biosynthesis 65–7
Penicillium roquefortii 83
 pentachlorophenol (PCP), degradation metabolic pathways 213–14, 215
 peramine 63
 synthesis by *Epichloë* endophytes 60–1
 peramine synthetase, molecular cloning 61, 63–5
 petroleum dependence reduction *see* biorefinery
Phanerochaete chrysosporium 206–7
 expression of LiP and MnP 211–13
 genome sequence 206–7
see also white-rot fungi
Phanerochaete sordida 208–9 *see also* white-rot fungi
 phenotypic heterogeneity
 causes and benefits 175–6
 in metal sensitivity 175–9
Phoma spp., squalenol production 25–7, 36–8
Phyllopertha horticola (garden chafer)
 control by *Metarhizium anisopliae* 268
 life-cycle and economic relevance 260
 phytopathogenic fungi
 economic losses and health risks 291–2
 fungal biopesticides 292–3
 problems with chemical controls 291–2
 PKS (polyketoid synthases), in *Aspergillus nidulans* 80, 81, 82
 PKS (polyketoid synthases) gene cloning
 fusarin synthase 38–41
 in fungal polyketoid biosynthesis 33–6
 squalenol synthase 36–8
 tenellin synthase 41–2, 43
see also polyketoid biosynthesis
 PKS-NRPS hybrid genes, in *Aspergillus nidulans* 82
 plant protection *see* crop protection;
 phytopathogenic fungi
 pneumocandins 51–2
Pochonia spp., nematode egg-destroying fungi 318–19, 325–6
Pochonia chlamydosporia, changing hosts 326
 polarized growth (hyphal extension) 95
 actin cytoskeleton 95
 cell end markers 95
 continuous tip elongation 96
 dynein 95, 105, 107–8
 kinesins 95, 105–7
 microtubules 95, 103, 108–10
 motor proteins 95, 105–8
 myosins 95
 polarisome 95
 vesicle supply centre (Spitzenkörper) 95, 96
see also microtubules
 polychlorinated dibenzo-p-dioxins, degradation metabolic pathways 214–15
 polychlorinated dibenzofurans, degradation metabolic pathways 214–15
 polyketoid biosynthesis (bacterial) 30, 31
 polyketoid biosynthesis (fungal)
 aflatoxin B1 30–1, 32
 chemical reactions 27–31
 fungal polyketide synthases (PKS) 27–8, 29
 gene cluster investigation methodology 33–6
 gene clusters 31–3
 genes encoding PKS and post-PKS enzymes 31–3
 highly reduced polyketoids 29, 30
 partially reduced polyketoids 29, 30
 phases of the process 27, 28
 PKS gene cloning 33–6
 post-PKS enzymes and reactions 27, 28, 30–1, 32
 programming of the PKS 28–30, 31
 unreduced polyketoids 28–9
see also PKS (polyketoid synthases)
 polyketoids
 aflatoxin B1 25–7
 fumonisin 25–7
 lovastatin 25–7
 6-methyl salicylic acid 25–7
 norsolorinic acid 25–7
 orsellinic acid 25–7
 produced by filamentous fungi 25
 range of structural complexity in fungi 25–7
 squalenol S1 25–7
 sterigmatocystin mycotoxin 25–7
 T-toxin 25–7
 tetrahydroxynaphthalene 25–7
 Pooidae grasses, symbiotic fungal associations 59
 protein folding in the endoplasmic reticulum 117–21 *see also* recombinant protein production

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)

Index

343

- protein secretion stress in *Aspergillus* spp. 147–53
- proteins secreted by *Aspergillus* spp. 144–7
- proteomics (fungal)
- benefits from chemostat research 8
 - methodologies 6–7
 - role in biorefinery 6–7
 - use of fungal genome information 6–7
- Pseudomonas putida* 206
- Pseudomonas pyrrocinia* 45–6
- Pyricularia oryzae* see *Magnaporthe grisea*
- pyromorphite (lead chlorophosphate), solubilization by fungi 244–5
- pyrrolnitrin 45–6
- quinoxifen 53–4
- recombinant protein production
- ER responses to folding demand stress 122
 - ER unfolded protein response (UPR) 122
 - process bottlenecks 117–18, 120
 - protein folding in the ER 117–21
 - see also UPR
- RESS (repression under secretion stress) response 148–53
- Resupinatus leightonii* 55
- rice blast disease (*Magnaporthe grisea* pathogen)
- annual crop loss 10–11
 - infection process 10–11
 - rapid development of pathogenic variants 10–12
 - see also *Magnaporthe grisea*
- risk assessment for fungal biocontrol
- agents 282–7
 - questions to be addressed 282–3
 - risk of metabolite entry into the food chain 285–6
 - risk of environmental contamination by metabolites 285–6
 - stability of metabolites in the environment 286–7
 - toxicity of metabolites 283–5
- ROS (reactive oxygen species)
- evidence for involvement in metal toxicity 165–6
 - metal-induced generation 163–5
 - oxidative damage to cell membranes 164–5
 - oxidative damage to DNA 165
 - oxidative damage to lipids 164
 - oxidative damage to proteins 164–5
- ‘ryegrass staggers’ in sheep 61, 62, 65
- Saccharomyces cerevisiae*
- bioluminescence-based biosensor 192–5
 - microtubule initiation 99
 - microtubule plus-end dynamics 102–4
 - polarized growth stage 96
- Saccharomyces cerevisiae* metal toxicity
- conditions when heterogeneity is advantageous 178–9
 - heterogeneous Cu resistance 176–8
 - membrane-lipid peroxidation in Cd toxicity 168–72
 - metal-induced membrane damage 167–72
 - metal-induced protein damage 172–5
 - metal-induced ROS generation 163–5
 - model 162–3
 - phenotypic heterogeneity in metal sensitivity 175–9
 - protein (methionine) oxidation in Cr toxicity 172–5
 - responses to environmental conditions 178–9
 - role of oxidative stress 166–7
 - suppression of heterogeneity in a stable habitat 178
- Schizosaccharomyces pombe*
- microtubule initiation 98, 99–100, 101
 - microtubule plus-end dynamics 102–4
 - polarized growth 96
- Scytalidium* spp. 55
- scytalol D 55
- secondary metabolites, range found in filamentous fungi 25 see also *specific fungi*
- Shewanella oneidensis* 206
- soil contamination see bioremediation
- soil pests see subterranean pests
- soja crop protection 49
- Sordaria arenaosa* 50–1
- sordarin 50–1
- spinosad 45–6
- squalestatin, biological properties S1 25–7
- squalestatin synthase 36–8
- Stagonospora convolvuli* (LA39 strain)
- assessment as bindweed mycoherbicide 276–8, 279
 - cercosporin toxin 278–81
 - efficacy in the field 278, 279
 - elsinochrome A toxin 278–81, 282–7
 - leptosphaerodione toxin 278–81
 - mass production, formulation and storage 277–8
 - mode of action against bindweed 276–7
 - toxin production in various isolates 278–81, 282
- Stagonospora convolvuli* LA39 biocontrol
- agent risk assessment 282–7
 - questions to be addressed 282–3
 - risk of environmental contamination by metabolites 285–6

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)344 *Index*

- Stagonospora convolvuli* (cont.)
 risk of metabolite entry into the food chain 285–6
 stability of elsinochrome A in the environment 286–7
 toxicity of elsinochrome A 283–5
 sterigmatocystin mycotoxin 25–7
 strobilurins (agricultural fungicides) 46–9
 discovery in Basidiomycetes 46
 mode of action 46–9
 structure 46–9
 synthesis 46–9
 synthetic strobilurin fungicides 49
Strobilurus tenacellus 46–9
 subterranean pest controls
 chemical pesticides 257–8
 early forms of control 257–8
 entomopathogenic fungi 263–5
 fungal biocontrol agents 258–9, 263–5
 integrated pest management (IPM) 258–9
 problems with controls 257–8
 subterranean pests and their economic relevance 259–63
Agriotes spp. (wireworms, click beetle larvae) 261–2
Athous spp. (wireworms) 261–2
Corymbites spp. (wireworms) 261–2
Daktulosphaira vitifoliae (grape phylloxera) 262
Diabrotica virgifera virgifera (western corn root worm) 263
 economic importance in Europe 257–8
Melolontha melolontha (European cockchafer, May beetle) 259
Otiorhynchus sulcatus (black vine weevil) 260–1
Phyllopertha horticola (garden chafer) 260
 T-toxin 25–7
 tenellin synthase 41–2, 43
 tenuazoic acid 53
 terpene cyclases, in *Aspergillus nidulans* 80, 83
 terpenoids, produced by filamentous fungi 25
 tetrahydroxynaphthalene 25–7
 tetrahydroxynaphthalene synthase 28–9
 toluene, degradation metabolic pathways 215–16
Tolyposcladium inflatum, source of cyclosporin 62–3
Tolyposcladium niveum 80–2
Trametes versicolor, expression of laccase 213
Trichoderma spp.
 beneficial fungal antagonists 294–7
 challenges in development of biopesticides 294
 diversity of habitats 293
 effects on plant growth improvement 304–6
 genetic and biological diversity 293
 genetically modified strains 294
 industrial and agricultural uses 294
 interactions with plants and pathogens 294–8
 interactions with the plant 298–302
 life-cycle 293
 molecular cross-talk with the plant 300, 302–4
 rhizosphere-competent strains 293–4
 versatility as biocontrol agents 293–4
Trichoderma-based biofungicides
 beneficial fungal antagonists 294–7
 effects on plant growth improvement 304–6
 induced systemic resistance (ISR) 298–304
 interactions with plants and pathogens 294–8
 interactions with the plant 298–304
 molecular cross-talk with the plant 300, 302–4
 range of commercial products 292–3
 range of uses 292–3
 stimulation of host plant defences 296–7, 298–302
Trichoderma atroviride 5–6
Trichoderma reesei
 discovery 5–6
 genome project 5–6
Trichoderma virens 5–6
 tryprostatins 89
 UPR (unfolded protein response) in filamentous fungi 118, 120, 124–5, 130–2
 engineering to increase protein productivity 118, 120, 132–3
 in *Aspergillus* spp. heterologous protein production 147–53
 in the ER 122
 mechanism of signal transduction 124–5, 130–1
 regulation of adaptive responses 131–2
 UPR in *S. cerevisiae* 122–5
 apoptosis 128
 decrease of influx of unfolded polypeptide chains 127–8
 ER associated degradation (ERAD) 126–7
 mechanism of signal transduction 122–5
 molecular chaperones and foldases 126
 nutrient sensor 128–30

Cambridge University Press

978-0-521-85935-6 - Exploitation of Fungi

Edited by G. D. Robson, P. van West and G. M. Gadd

Index

[More information](#)*Index*

345

- phospholipid biosynthesis 127
- regulation of adaptive responses to ER stress 126–8
- Ustilago* spp., actions of sordarins 50–1
- Ustilago maydis*
 - microtubule initiation 98, 99–100, 101
 - polarized growth 96
- Verticillium*, fungal pathogen evolution
 - 328–9 *see also* *Lecanicillium*; *Pochonia*
- western corn root worm *see* *Diabrotica virgifera virgifera*
- white-rot fungi
 - ability to break down lignin 207–9
 - degradation of xenobiotics 207–11
 - effects of soil environment 221–3
 - expression of ligninolytic enzyme systems 211–13
 - fate of breakdown products 220–1
 - field trials 223–4
 - interaction with native soil microbiota 223
 - laccase 207–9, 210–11, 213
 - lignin peroxidase 207–10, 211–13
 - ligninolytic enzyme–redox mediator systems 207–13
 - ligninolytic enzymes 206–7
 - manganese-peroxidase 207–9, 210, 211–13
 - role in bioaugmentation 206–7
 - see also* xenobiotic metabolism
- wireworms *see* *Agriotes* spp.; *Athous* spp.; *Corymbites* spp.
- xenobiotic metabolism
 - BTEX compounds (benzene, toluene, ethylbenzene, xylene) 215–16
 - chlorophenols 213–14, 215
 - dyes 217
 - explosive waste 216
 - fate of breakdown products 220–1
 - field trials of white-rot fungi 223–4
 - metabolic pathways associated 213–20, 221
 - olive mill waste-water 216
 - organophosphate pesticides 215
 - pentachlorophenol (PCP) 213–14, 215
 - polyaromatic hydrocarbons (PAHs) 217–20, 221
 - polychlorinated dibenzo-p-dioxins 214–15
 - polychlorinated dibenzofurans 214–15
 - see also* white-rot fungi
- xenobiotics
 - definition 205
 - degradation by white-rot fungi 207–11
 - in situ* bioremediation 205–7
 - oxidation by ligninolytic enzyme–redox mediator systems 207–11
- xylene, degradation metabolic pathways 215–16
- Zalerion arboricola*, selective antibiotics produced 51–2