

Index

- Abrikosov–Suhl resonance, 53
- acceptance ratio, 583, 618
- action, 483, 579, 646
- in ground-state path-integral Monte Carlo, 660
 - pair, 648
 - primitive approximation, 646
- adiabatic switching on, **200–202**, 710–712
- advanced correlation function, 95
- in frequency, 96
- aluminum
- electron lifetimes, 322
- analytic continuation, **96**, 108, 301, **666**
- Padé approximant, 301
- Anderson impurity model, **51–54**
- and DMFT, 436, 480
 - mean-field approximation, 79
 - numerical solution, 485
- anomalous diagrams, 215
- antiferromagnetic order, 43, 64, 78
- antiscreening, 326, 377
- in a carbon nanotube, 327
- argon
- optical absorption spectrum from the BSE, 373, 374
- atomic limit, 429–431
- not unique, 430
- atomic units
- Hartree, 2
- auxiliary bosons, 468
- auxiliary system, 423, 731–734
- DMFT, 436
 - examples, 432
 - interacting, 423, 732
 - Kohn–Sham, 67–72, 172, 431
- auxiliary-field quantum Monte Carlo, 485, 487, 662
- band offsets
- in GWA, 321
- band structure
- of silicon in QMC, 316
 - of silicon in the GWA, 316
- bandgap
- definition, 24
 - in G_0W_0 , 319
 - in QSGW, 319, 320
 - in the GWA, 319, 320
- bandgap problem, 24, 319
- bandwidth
- effect of self-consistency, 321
 - in QMC, 317
 - in the GWA, 317, 321
 - of silicon, 317
 - of sodium, 267
- basis
- for GW calculations, 304
 - for polarizability, 306
- Bayes’ theorem, 667
- Berry phase, 127, 128
- Bethe ansatz, 45, 52, 470
- Bethe lattice, *see also* semi-circular density of states, **446**, 455, 456
- bipartite, 449
 - definition, **446**
- Bethe–Goldstone approximation, 400
- Bethe–Salpeter equation (BSE), **230**, 371, **345–387**, 393
- argon, 373, 374
 - as time evolution, 382
 - as two-body equation, 357
 - beyond the TDA, 375
 - calculations in practice, 368
 - cancellations, 367
 - comparison to TDDFT, 382
 - diagram, 349
 - dynamical effects, 380
 - electron–electron channel, 405
 - electron–hole channel, 405
 - flowchart, 368
 - for the 4-point vertex, 349
 - for three channels, 404, 405
 - frequency structure, 354
 - GaAs, 373
 - graphene, 373

- GW-BSE approximation, 355, 357
 - history, 348
 - hole–hole channel, 405
 - mixing of transitions, 360
 - neglect of dynamical effects, 358
 - non-linear response, 383
 - overlap matrix, 360
 - particle–particle, 381, 382
 - problem of \mathbf{k} -points, 370
 - problems of limit $\mathbf{q} \rightarrow 0$, 369
 - scaling of calculations, 371
 - silicon, 372
 - spin structure, 353
 - spin structure in the GW-BSE approximation, 357
 - spin waves, 382
 - two-particle equation, 360
 - two-particle hamiltonian, 358
 - versus TDDFT, 399
 - without empty states, 371
- Betts cells, 459, 474
- bipartite lattice, 43, 45, 449
- Bloch's theorem
 - many-body, 125
- BN
 - DMC calculations cubic phase, 326, 637
 - GW corrections in the hexagonal phase, 326
 - total energy of the hexagonal phase, 339
- Born approximation, 103
- Born series, 195
- boron nitride, *see* BN
- Bose condensation, 599, 653, 654
- Bose statistics, 653
- boundary conditions, 124
 - fixed-node, 623
 - of functional differential equation, 227
 - open, 125
 - periodic, 125
 - twist averaged, 126, 601
 - twisted, 125
- branching, 614, 635
- Brillouin theorem, 198
- broken symmetry, 78–80
- bubble approximation, 252
- bubble expansion, 252
- cancellations
 - of dynamical effects in the BSE, 357, **380**
 - of self-consistency and vertex corrections, 398, 399
 - of vertex corrections in W and Σ , 396
- carbon nanotube
 - screening, 327
- cerium
 - DMFT calculations, 529–533
 - phase diagram, 530
 - photoemission, 27, 532
 - spectra, 532
- charge-transfer excitons, 374
- charge-transfer gap, 332
- chemical potential, 96
 - alignment in GW calculations, 293
- chocolate cake
 - and self-energy, 188
- cluster
 - defined as embedded cell, 458
 - optical spectrum, 378, 379
- cluster perturbation theory (CPT), **462**, 482
- coherent inelastic X-ray scattering (CIXS), 104
- coherent potential approximation (CPA), 434–435, 744–746
- COHSEX, **255**, 262, 279, 335
 - self-consistent, 294
- collective excitations, 29–32
- computational cost, *see* scaling
- conductivity, 101
- configuration interaction (CI), 366, 381
- conservation laws, 231–234
- conservation of
 - number, 232
 - total angular momentum, 232
 - total energy, 232
 - total momentum, 232
- conserving approximation, 187–189, 231
- constrained DFT, *see* DFT+DMFT method
 - isolated atoms and ions, 516
- constrained RPA, *see* GW+DMFT method
- constrained-path quantum Monte Carlo, 635
- continuity equation, 188, 417
- continuity principle, 7, 49, 54, 202, 228
- continuous-time quantum Monte Carlo, 487–498
 - auxiliary-field method, 490–491
 - dynamical interactions, 496–498
 - expansion in hybridization, 491–494
 - expansion in interaction, 487–491
- continuum excitons, 372
- contraction of operators, 206
- convergence
 - of GW calculations, 306, 307
 - of perturbation series, 228
- core polarizability, 289
- core polarization, 290
- core–valence separation, 289
- correlation
 - definition, 16
 - relation to unrestricted Hartree–Fock, 16, 78
- correlation energy, 8, **16**, 77
 - atomic, 138
 - DFT, 67
 - DFT-RPA, 75
 - of homogeneous electron gas, 271
 - of hydrogen molecule, 129–131
- correlation function, 86
 - advanced, 95
 - analytic structure, 97
 - current–current, 101
 - electron–electron, 119
 - hole–hole, 119

- particle–particle, 401
- retarded, 94
- spin autocorrelation
 - Fe, Ni, 538
 - NiO, 546
- time-ordered, 95
- two-particle, 116
- correlation function quantum Monte Carlo, 670
- Coulomb cut-off, 306
- Coulomb hole, 255
 - dynamical, 256
- coupled cluster, 198, 229
- coupled electron ion Monte Carlo, 662
- coupling constant integration, 114, 177, **186**, 218
- crossing symmetry, 416
 - GWA, 274
- crystal field splitting
 - definition, 509–510
 - different usages, 540
 - FeO, 549
 - MnO, 547–549
 - schematic diagram, 510
 - V₂O₃, 542–543
 - VO₂, 335
- crystal local field effects, 351
- Cu₂O
 - GW calculations, 290
 - optical absorption spectrum, 362, 363, 370
- cumulant, 276
- cumulant expansion, 410–415
 - application to HEG, 414
 - computational cost, 413
 - derivation, 411–413
 - diagrams, 412, 413
 - for graphite, 415
 - for HEG, 414
 - for silicon, 318, 415
 - link to DMFA, 412
 - spectral function, 413
- Curie–Weiss susceptibility, 20, 81
- current–current response function, 118
- cusplike condition, 648
- Δ -SCF, **63**, 260, 262, 328, 367, 516
- defects
 - optical spectrum, 379
- degeneracy
 - problem of perturbation theory, 418
 - problem of the GWA, 328
- density, 86, 112
 - diagram for, 205
- density functional theory, **65**, 72, **76**, 171–173, 431
 - before Hohenberg and Kohn, 65
 - current functional, 77
 - Hohenberg–Kohn theorem, 66, 171
 - Kohn–Sham method, *see* Kohn–Sham method
 - Levy–Lieb constrained search, 66
 - proof of invertibility, 729
- time-dependent, *see* time-dependent density functional theory
- density matrix, **88**, 112
 - and disorder, 114
 - one-body, 88, 112
 - single particle, 599
 - thermal, **645**
 - two-body, 120
- density matrix embedding, 442, 483
- density matrix renormalization group (DMRG), 443, 483, 499
- density of states, 111
- density–density response, 101
- density–density response function, 118
- detailed balance, **581**, 661
- determinantal Monte Carlo, 474, 662
- DFT+DMFT method, 515–522
 - constrained DFT, 516
 - double counting, 519
- DFT+Gutzwiller (DFT+G), 524
- DFT+U, 74, 260, 286, 522, 740
- DFT+U calculations
 - Ce, 531
 - NiO, 545
- DFT-RPA, 75
 - link to GWA, 264
 - total energy calculations, 302, **338–339**
 - total energy of H₂, 275
- diagrams
 - Bethe–Salpeter equation, 349
 - bubble, 207, 217, 238
 - closed loop, 207
 - connected, 205, 206
 - conversion into equations, 211
 - cumulant expansion, 412, 413
 - disconnected, 205, 206
 - G₀W₀, 413
 - GW₀, 413
 - Hedin’s equations, 249
 - in frequency space, 211, 215
 - in momentum space, 211, 215
 - ladder, 240
 - polarization, 210
 - RPA, 217, 252
 - scattering, 240, 402
 - skeleton, 216
 - T-matrix approximation (TMA), 241, 402
 - topologically distinct, 206
- dielectric function
 - inverse, 102, **247**, 346–347
 - Lindhard, 266
 - link to experiment, 103–105
 - macroscopic, 104
 - test-charge test-charge, 250
- diffusion Monte Carlo, 609, 611–621, 660
- disordered local moment (DLM) method, 334, **524**, 539, 746
- dispersion interaction, 18

- DMFT, *see* dynamical mean-field theory
- DMFT calculations
- actinides, 536
 - Ce, 529–533
 - CeIrIn₅, 533–535
 - Cm, 536
 - Fe, 538
 - MnO, 547–549
 - Ni, 538
 - NiO, 545–547
 - Pu, 536
 - V₂O₃, 542–543
 - VO₂, 336
 - Yb, 535
- DMFT+DFT, *see* DFT+DMFT method
- DMFT+GW, *see* GW+DMFT method
- DMFT+MBPT, *see* MBPT+DMFT methods
- double counting
- in DFT+DMFT, 519
 - in screened T-matrix approximation, 403, 560
- double excitations, 380
- dynamic correlation function, 93
- dynamic structure factor, **103**, 347, 350–351, 375–376, 666
- dynamical effects
- cancellations in the BSE, 358, **380**
 - GW calculations, 317
 - in the absorption spectrum of silicon, 381
 - in the BSE, 380
 - satellites, 380
- dynamical interactions, 496–498
- dynamical mean-field theory
- algorithm, **438–441**
 - cellular approx. (c-DMFA), 460–462
 - dual fermions, 572
 - dynamical cluster approximation (DCA), 463–466
 - causality, 466
 - dynamical vertex approximation, 572
 - equations, **435–444**
 - extended DMFT (EDMFT), 467–469, 567
 - Hubbard model calculations, 470–475
 - infinite dimensions, 445
 - antiferromagnetic solution, 449
 - metal–insulator transition, 448
 - paramagnetic solution, 447
 - overview, **425–429**
 - self-consistency condition, **428**, 438
 - self-consistent loop, 439
 - single-site approximation (DMFA), 427, **428**
 - single-site calculations, 445–453
 - two hybridized bands, 450
 - two interacting bands, 451
 - variational derivation, 441
- dynamical screening
- and embedding, 412
- Dyson amplitude, 108
- Dyson equation, 147, **148–150**, 212–213, 228–231
 - coupled equations, 555
 - figure, 164, 556, 558
 - overview, 163–165
- figure, 149
- for χ , 347
 - for screened interaction, 150
 - for T-matrix, 402
 - for the two-particle correlation function, 230
 - for the two-particle Green's function, 230
 - iteration, 297
 - kernel of, 231
 - solution of, 297
- effective energy technique, 307
- effective interaction, 235, 393
 - exchange–correlation, 235
 - classical, 249
 - in T-matrix approximation, 399
 - test charge–test electron, 396
- effective mass, 50, 159, 605
 - in the GWA, 321
- effective mass approximation
 - for Wannier–Mott excitons, 365
- effective two-particle interaction, 230
- elastic peak, 103
- electron addition, 107, 110
- electron energy loss spectroscopy (EELS), 29
- electron gas, *see* homogeneous electron gas
- electron removal, 110
 - approximate pictures, 391
 - schematic illustration, 22
- electron–boson coupling, 276, **410–413**
- electron–boson coupling model, **276**
 - spectral function, 397
- electron–electron correlation function, 119
- electron–hole correlation function
 - and experiment, 118
 - retarded, 117
 - time-ordered, 117
- electron–hole exchange, 359
- electron–hole interaction
 - analysis of effects, 361
 - direct, 359
 - effect on intensities, 362
 - effect on transition energies, 361
 - effects on plasmons, 375
 - exchange, 359
 - from the GWA, 355
 - mixing of transitions, 362
 - static approximation, 356
- electron–hole pairs, 116
- electron–hole propagators, 116
- electron–hole wavefunction, 365–366
 - and broken symmetry, 366
 - and spin, 366
 - and translational symmetry, 366
 - in LiF, 366
 - in MnO, 366
- empty states

- in GW calculations, 307
- energy loss spectrum
 - relation to optical spectrum, 353
- entropy, **114**, **115**, 172, 474, 475, 477, 515, 531
- equation of motion, 107, **223–226**
 - for an independent-particle Green's function, 225
 - for field operators, 224
- ergodic, **580**, 585, 603, 661
- Ewald image potential, 595
- exact constraints
 - use of, 415
- exact diagonalization, 481, 482
- exact exchange (EXX), **73**, 285, 732
- exchange splitting, 329
- excitations
 - double, 392
 - hole–hole, 408
 - neutral, 392
 - spin-flip, 392
- excited states
 - in quantum Monte Carlo, 669
 - in variational Monte Carlo, 603
- exciton, 29, 362
 - binding energy, 362
 - bright and dark, 362, 377, 378
 - charge transfer, 374
 - confined, 376
 - continuum, 373
 - dispersion, 376
 - in LiF, 376
 - Frenkel, 364–365
 - in LiF, 376
 - Wannier–Mott, 365, 373
- excitonic effects, 362
 - first order, 364
 - guidelines, 363
 - importance of, 363
 - scattering cross-section, 671
- expectation values, 85
- extended DMFT (EDMFT), 567
- extended Koopmans' theorem, 301
- extrinsic effects
 - in photoemission, 115
- f*-sum rule, 102
 - RPA, 259
- Fermi arc, 37
- Fermi liquid theory, 7, **49**, **51**, 605
- Fermi surface, 89
 - schematic illustration, 158
- fermion sign problem, 484, 489, 621, **622**
 - cases with no sign problem, 484, 487, 493
 - in auxiliary-field Monte Carlo, 634
 - in path-integral Monte Carlo, 654
- ferromagnetism, 20, 537
- Feynman diagrams, 202–206
- Feynman rules, 204, 209
- Feynman–Kac formula, 132, 135, 410, 630
- finite size effects, 601
- finite systems
 - GW calculations, 326–329
- fixed-node approximation, **623**, 641
 - in lattice models, 628
 - in path-integral Monte Carlo, 655
- fixed-phase approximation, 626
- FLEX, 403
 - and screening, 403
 - two-step, 403
- fluctuation amplitudes, 256
- fluctuation spectrum
 - of a correlation function, 95
- fluctuation–dissipation theorem, 101
- forward-walking of estimators, 629
- four-point vertex, 181, 403
 - Bethe–Salpeter equation, 349
- Frenkel excitons, **364–365**, 373
- frequency integrations
 - in GW calculations, 298–302
- Friedel sum rule, **55**, 735–737
- functionals
 - $\Psi[G, W]$, 179–181
 - conserving approximations, 187–189
 - definition, 724
 - derivative, 725
 - effective action, 182–185
 - free energy
 - proof of invertibility, 727
 - Klein *G* functional, 179
 - Luttinger–Ward, 441
 - Luttinger–Ward $\Phi[G]$, 174–179
 - of pieces of *G*, 417
 - of the interacting Green's function, 215
 - restricting domain of search, 441–442, 731–734
 - variational analysis, 733
 - self-energy, 441
 - spectral density, 433
 - table comparing density, Green's function, and Hartree–Fock functionals, 175
- G_0W_0 approximation, 285
 - diagrams, 413
 - problems of, 293
 - starting hamiltonians, 285
- GaAs
 - optical absorption spectrum from the BSE, 373
- Galitskii–Migdal total energy, **114**, 301
- gauge invariance, 233, 417
- Gell–Mann–Low theorem, 202
- grand partition function, 114
- grand potential, 85, **114**, 172, 178, 204, 441, 707
- grand-canonical ensemble, 85
- graphene
 - doped, 414
 - GW calculation, 321
 - optical absorption spectrum, 363, 373
- graphite

- cumulant calculation, 415
- electron energy loss, 351, 352
- Green's function
 - exponential form, 410
 - imaginary time, 213, 699–707
 - independent-particle, 107
 - non-interacting, 203, 705
 - one-body, 105–116
 - two-body, 115–119
- Green's function Monte Carlo, 611
- ground-state path-integral Monte Carlo, **659**, 670
- guiding function, 605, 616, 621, 670
- Gutzwiller approximation, 46, **741**
- Gutzwiller wavefunction, 46, **740**
- GW approximation, 237, 239, 253–259, **393–394**
 - and crossing symmetry, 416
 - and phase transitions, 342
 - calculation of quasi-particle properties, 287
 - charge density, 262
 - conservation laws, 259
 - definition, 253–259
 - G_0W_0 , 250, **285–286**
 - ground-state properties, 262–265
 - GW_0 , 259, **296**, 413
 - link to DFT-RPA, 264
 - occupation numbers, 263
 - problem of degenerate ground state, 328
 - self-energy, 282
 - size-consistency problem, 340
 - total energy, 263
- GW calculations
 - all-electron, 292
 - analytic continuation, 301
 - bandwidth, 317, 321
 - basis, 304
 - best W strategy, 297, 395
 - Ce, 532
 - confined systems, 326
 - convergence, 306, 307
 - core–valence exchange, 290
 - core–valence separation, 289
 - Cu_2O , 290
 - diamond, 313
 - dynamical effects, 317
 - empty states in, 307
 - evaluation of Φ -functional, 301
 - first-order quasi-particle equation, 288
 - flowchart, 284
 - frequency integration on the contour, 299
 - frequency integration using spectral functions, 299
 - frequency integrations, 298–302
 - fully self-consistent, 296
 - gaussian basis, 305
 - ground state, 337–341
 - GW_0 , 296
 - in a basis, 302
 - integration using Matsubara frequencies, 299
 - localized basis, 305
 - matrix elements, 302
 - models, 288
 - Ni, 538
 - NiO, 545
 - of band offsets, 321
 - of bandgaps, 319, 320
 - of effective mass, 321
 - of lifetimes, 322
 - of satellites, 324
 - of STM images, 325
 - PAW, 291
 - plane-wave basis, 304
 - problem $\mathbf{q} \rightarrow 0$, 304
 - problems of G_0W_0 , 293
 - scaling, 306, 308
 - self-consistency, 292–298
 - silicon, 312–319
 - space-time approach, 298, 300
 - total energy, 301, 339
 - transition metal oxides, 329–337
 - transition metals, 329–337
 - transport, 326
 - using a plasmon pole model, 298
 - valence only, 289
- GW_0 , 297
- GW+DMFT method, 561–568
 - constrained RPA, 568
 - diagrammatic expansion, 566
 - Luttinger–Ward functional, 566
- Hartree
 - atomic units, 2
 - potential, 62
- Hartree approximation, **61**, 228
- Hartree–Fock approximation, 16, **61–64**, 739
 - bandwidth, 267
 - derivation via functional derivatives, 225
 - equations, 62
 - Fermi surface singularity, 8, **64**, 81
 - restricted, **16**, 78, 129, 739
 - self-energy, 236
 - total energy functional, 174
 - unrestricted, **16**, 78, 79, 739
- Hartrees, D. R. and W. R.
 - calculations for atoms, 5
- He, 135
 - boson, 591, 654
 - droplets, 662
 - wavefunction, 143
- heat bath algorithm, 587
- heavy fermions, 32, 54, 533
- Hedin's equations, 246–251
 - diagrams, 249
- Hedin's pentagon, 250, 251, 393
- HEG, *see* homogeneous electron gas
- Heisenberg equation of motion, 224
- Heisenberg spin model, 45, **48**, 603, 651
- Hirsch–Fye algorithm, 485–487

- hole–hole correlation function, 119
- Holstein–Hubbard model, 496, 570
- homogeneous electron gas (HEG), 40–41
 - applications in path-integral Monte Carlo, 648, 659
 - applications in projector Monte Carlo, 636
 - correlation functions, 662
 - cumulant calculation, 414
 - GWA, 265–272
 - Jastrow factor, 136, 143
 - pair distribution, 91
 - wavefunction, 142
 - Wigner crystal, 139, 648, 652
- Hubbard approximations
 - I atomic limit, 743–744
 - III “alloy approximation”, 48, **434–435**, 744–747
 - resonant broadening correction, 746
- Hubbard atom
 - Green’s function, 743
 - self-energy, 743
- Hubbard clusters
 - TMA, 407
 - total energy in GWA, 272
- Hubbard dimer, **45**, 279, 535
 - broken symmetry, 79, 82
 - correlation functions, 58
 - DMFT solution, 444, 455
 - exact solution, 57, 58
 - exercise, 57, 58, 82, 455
 - GWA, 272–275
 - magnetic susceptibility, 57
 - TMA, 407
- Hubbard model, **44**
 - 1 dimension, 470
 - 2 dimensions, 471–474
 - doped, 472
 - spin correlations, 637
 - superconductivity, 473
 - 3 dimensions, 474–475
 - k-resolved spectra, 475
 - Néel temperature, 474
 - imaginary-time correlations, 662
 - in variational Monte Carlo, 603
 - semicircular DOS, 448
 - two hybridized bands, 450
 - two interacting bands, 451
- Hubbard–Stratonovich transformation, 633, 663
 - discrete, 664
- hybrid functionals, 74
 - link to the GWA, 261
- hybrid Monte Carlo, 617
- hybridization function, 151
- hydrogen
 - applications in path-integral Monte Carlo, 659
 - applications of diffusion Monte Carlo, 637
 - atom in diffusion Monte Carlo, 613
 - phase diagram, 662
- hydrogen molecule
 - GWA total energy, 275
- Heitler–London wavefunction, 130
 - ortho-para, 657
- image potential, 327
- image states, 327
 - in the GWA, 328
- impact ionization, 316
- importance sampling, 616, 641
- improved integration method, 307
- independent particle approximation
 - for one-body Green’s function, 107
 - for optical spectra, 350
 - response function, 361
- induced density, 101
- induced potential, 101, 394
 - exchange–correlation, 274, 394
- Inelastic X-ray scattering (IXS), 29, 103
 - resonant (RIXS), Yb, 535
- infinite dimensions, $d \rightarrow \infty$, 425, **445**, 446, 741
- insulators
 - charge transfer, 510, 543
 - Mott–Hubbard, 510
- interaction vertex, 204, 205
- interactions
 - effective screened, 512, 564, 570
 - typical values, 518
- interference effects
 - in photoemission, 115
 - in the BSE, 362, 372
- intrinsic effects
 - in photoemission, 115
- inverse dielectric function, *see* dielectric function
- inverse photoemission (IPES), 22
 - Ce, 532
 - Ge, 24
 - NiO, 543
- ionization energies
 - in the GWA, 327
- iron
 - lifetimes in the TMA, 407
- irreducible
 - polarizability, 150, 247
 - self-energy, 149, 176, 212
 - two-particle correlation function, 231, 348
 - vertex, 247
 - vertex, fully irreducible, 405
- iterated perturbation theory, 447, 498
- Janak’s theorem, 71
- kinetic energy, 89, 113
- Kohn–Sham method
 - auxiliary system, 67–72, 172, 431
 - equations, 67
 - Green’s function functionals, 173
 - hybrid functionals, 74
 - potential, 67
 - self-interaction correction (SIC), 74

- total energy functionals, 69–70, 172–173, 726–727
 bandgap problem, 72
 bands, 70
 exact exchange (EXX), 73
 generalized gradient approximations (GGAs), 69
 local density approximation (LDA), 68
 orbital-dependent functionals, 72
 spin density functionals, 75
 Kondo effect, **32–33**, 51, 528
 Co on Cu, 33
 Kondo insulator, 535
 Kondo temperature T_K , 32
 table, 33
 Koopmans' theorem, 17, 25
 as exact constraint, 417
 Kramers–Kronig relations, 100
 Kubo formulas, 101
- Lanczos method, 482
 Lang–Firsov transformation, 496
 lattice constants
 in DFT-RPA, 338
 lattice models
 fixed-node approximation, 628
 LCAO, 305
 Legendre transformation, 176, 725
 invertibility, 725
 Lehmann representation, 98
 LiF
 dynamic structure factor, 376, 376
 electron–hole wavefunction, 366
 exciton dispersion, 376
 optical absorption, 353
 lifetime, 25, 270
 due to excitation of phonons, 341
 in aluminum, 322
 in iron, 407
 in nickel, 407
 in noble metals, 323
 of quasi-particles, 316, 322
 linear extrapolation of estimators, 629, 642
 linear response, 99
 linearized augmented plane waves (LAPW), 292
 linked cluster theorem, 208
 Lippmann–Schwinger equation, 194
 lithium, 649
 LMTO, 292, 305
 local energy, **131**, 592
 local moments, 21, 421, 451, 503, **507**, 528
 Ce, 530
 CeIrIn₅, 534
 Fe, Ni, 539
 MnO, 549
 NiO, 546
 regime, 53, 79, 80
 V₂O₃, 542
 local potential
 defined, 61
 Kohn–Sham, 67
 locality approximation, 631
 localized basis
 for GW calculations, 305
 longitudinal–transverse splitting, 369
 loss function, 103, 347
 of silicon, 104
 low density limit, 400
 low-dimensional systems
 GW calculations, 326
 Luttinger theorem, 7, 44, 45, **54–55**, 158, 535,
 735–737
 and zeros of the Green's function, 737
 derivation, 735–737
 statement, **54**
 Luttinger–Ward Φ functional
 GWA, 264
 skeleton diagrams, 216
 Luttinger–Ward functional, 218
 calculations in the GWA, 340
- MacDonald's theorem, 604
 macroscopic dielectric function, 104
 magnetism
 in the GWA, 340
 overview, 20
 Stoner itinerant model, 82
 Zeeman field, 2
 Markov chain, *see* random walk
 Martin–Schwinger hierarchy, 226
 materials
 actinides, 18, 536
 Bi₂Sr₂CaCu₂O_{8+x}, 37
 BN, 637
 Ce, 27, 529–533
 CeIrIn₅, 533–535
 Fe, 20, 537–539
 FeO, 549
 gadolinium, 330, 341
 Ge, 23
 h-BN, 338
 lanthanides, 18, 529–536
 LiF, 30
 MnO, 340, 547–549
 Na, 27
 Ni, 20, 329–330, 537–539
 NiO, 31, 330–334, 376, 543–547
 silicon, 104, 342
 SrVO₃, 39
 transition metal oxides, 330–337, 540–549
 V₂O₃, 34–35, 542–543
 VO₂, 34, 334–337, 541
 Yb, 535
 Matsubara formalism, **699–709**
 difficulty to transform to real ω , 709
 grand potential Ω , 707
 Green's function at $T \neq 0$, 699–707
 non-interacting particles, 705

- poles on imaginary axis, 703
- Matsubara frequencies, **701**
- maximum entropy, 669
- MBPT+DMFT methods, 561–568
- mean-field theory, 59–65
 - broken symmetry, 78–80
 - comparison table, 432
 - Hartree, 61
 - Hartree–Fock, 61
 - Weiss, 64, 433
- Mermin functional, 172
- metal–insulator transition, **33–36**, 43, 46–48, 540
 - Brinkman–Rice–Gutzwiller approx., 47, 48, 747
 - FeO, 549
 - Hubbard approximation, 747
 - Hubbard III approximation, 47, 48
 - in DMFT, 448, 471
 - MnO, 547–549
 - V₂O₃, 34–35, 542–543
 - VO₂, 334–337, 541
- Metropolis Monte Carlo, 581–583, 646, 669
 - in diffusion Monte Carlo, 618
 - in ground-state path-integral Monte Carlo, 660
 - in variational Monte Carlo, 591
- mixed estimators in projector Monte Carlo, 629
- mixed valence, 535
- MnO
 - electron–hole wavefunction, 366
 - phase transition under pressure, 547–550
 - spin magnetic moment, 340
- modified Heisenberg representation, 483
- momentum distribution, 89, 113, 269, 654
 - computation in variational Monte Carlo, 598
 - electron gas, 89
 - GWA, 269
 - interacting, 113
 - QMC, 269
- Monte Carlo method, **579**
- Mott insulator, **33–36**, **43–44**, 55, 330, 449, 540, 543
 - doped, **35–36**, 472, 546
- Mott transition, **33–36**, *see also* metal–insulator transition, **43–44**, **46–48**, 445–450, 747
- MP2, *see* perturbation theory Møller–Plesset (MP)
- multiple solutions
 - of Dyson equations, 231
- multiplet effects, 367
- multiplets, 430, **506**, 512
 - Ce, 28, 29, 507
 - Kanamori parameters, 512
 - Racah parameters, 512
 - V, 506, 507
- Néel temperature, 64
- nearsightedness principle, 308, 427, 558
- nested cluster approximation, 467
- neutral excitation, 118
- neutron scattering, 104
- nickel, 407
- band structure in the GWA, 330
- DMFT, 537–539
- lifetimes in the TMA, 407
- satellite, 408
- TMA, 407
- NiO
 - analysis of excitation spectrum, 367, 544
 - DMFT calculations, 542–547
 - GW calculations, 330–337, 376
- noble metals
 - electron lifetimes, 323
- non-crossing approximation (NCA), 498
- non-linear response
 - from a Bethe–Salpeter-like equation, 383
- non-local potential
 - defined, 61
 - evaluation of matrix elements, 304
 - in projector Monte Carlo, 631
- normal order, 206
- occupation numbers, 88
 - GWA, 263
- one-body Green’s function, 105–116
- one-crossing approximation (OCA), 499, 534
- optical absorption, 347
- optical absorption spectrum, 104
- optical absorption spectrum, 29
 - BSE, 350
 - cluster, 378, 379
 - Cu₂O, 362, 370
 - defects in MgO, 379
 - graphene, 363
 - LiF, 353
 - metals, 380
 - organic crystals, 378
 - polymer, 377, 378
 - problem of **k**-points, 370
 - schematic illustration, 30
 - silicon as a function of temperature, 342
 - silicon with dynamical effects, 381
 - solids, 360
 - surface, 376, 377
- optical lattice, 475
- optimized effective potential (OEP), 73, 279, 732
- orbitals
 - localized, 513
- order parameter, 60, 75, 85, 92
- organic crystals
 - optical spectrum, 378
- pair correlation function, 91
 - electron gas, 91
 - normalized, 90
 - of a Wigner crystal, 93
- pair distribution, 90
- parquet approximation, 405
- parquet equations, 405
- parquet theory, 403

- partition function, 86
- partitioning states, 557
- path integrals, 483, 645
- path-integral Monte Carlo, 644–672
 - algorithm, 647
 - strengths and weaknesses, 671
- periodic table for transition elements
 - delocalized to localized, 504
- permutation operator, 652
- Perron–Frobenius theorem, 580
- perturbation theory
 - and degeneracy, 418
 - at $T = 0$, 199–213
 - at non-vanishing temperature, 213–215
 - Brillouin–Wigner, 195
 - in hybridization, 199
 - Møller–Plesset (MP), 197
 - problems of, 194–199
 - Rayleigh–Schrödinger, 196
- phonons
 - contribution to electron lifetimes, 341
 - coupling to, 111, 341, 415
 - in the GWA, 340
- photoemission, 21, 115
 - $(V_{1-x}Cr_x)_2O_3$, 36
 - angle resolved, 21
 - schematic illustration, 23
 - beyond the intrinsic spectral function, 319
 - $Bi_2Sr_2CaCu_2O_{8+x}$, 37
 - Ce, 27, 532
 - extrinsic effects, 115
 - Ge, 23
 - interference effects, 115
 - intrinsic effects, 115
 - MnO, 547–549
 - Na, 27
 - NiO, 543
 - of aluminum, 414
 - of lithium, 414
 - of silicon, 318, 415
 - of sodium, 414
 - satellites, 115
 - $SrVO_3$, 39
- pictures
 - Heisenberg, 684
 - interaction, 686
 - modified Heisenberg, 686, 700
 - Schrödinger, 684
- plasmaron, 269, 392, 414
- plasmon pole model, 286, 322
 - determination of parameters, 287
 - in GW calculations, 298
- plasmon satellites, 26–27
 - cumulant expansion, 410–415
 - GWA, 396
- plasmons, 26, 104, 351
 - effects of electron–hole interaction, 375
 - in the RPA, 351
- polarizability
 - irreducible, P , 248, 557
 - reducible, χ , 237, 249, 557
- polarization contribution
 - to the screened Coulomb interaction, 103
- polaron model, 276
- polymer
 - optical spectrum, 377, 378
- population control, 619, 642
- potential energy, 113
- pressure, 608
- projector auxiliary-field quantum Monte Carlo, 632–636
- projector quantum Monte Carlo, 609–641
 - strengths and weaknesses, 639
- Projector-Augmented Wave approach (PAW), 291
 - GW calculations, 291
- projectors, 610
- propagator, 106
- pseudogap, 37, 473
- pseudopotentials, 290
 - computation in variational Monte Carlo, 599
 - in projector Monte Carlo, 630
- Ψ -functional, 239
- QSGW, *see* quasi-particle self-consistent GW
- quantum impurity representation, 436
- quasi-particle self-consistent GW (QSGW), 271, 295, **316**
 - bandgaps, 319, 320
 - for NiO, 333
- quasi-particles, 23, **49–50**, 144, **155–163**
 - broadening, 157, 268
 - schematic illustration, 158
 - condition, 313
 - energies, 313
 - of silicon, 313
 - equations, 161–163
 - overview, 157–161
 - renormalization factor Z , 269, 294
 - shift, 314
 - strength, 89
 - weight Z , 159
- random walk
 - algorithm, **579**
 - branching, 614
 - efficiency, 586, 655
 - for excited states, 604
- random-phase approximation (RPA), 9, 203, **251–253**
 - and plasmons, 351
 - diagrams, 217
 - limits of, 352–353
 - performance of, 350–353
 - reduced Brillouin zone (RBZ), 459
 - reducible polarizability, 118
 - release-node algorithm, 627

- reptation quantum Monte Carlo, *see* ground-state path-integral Monte Carlo
- response-functions, 98–105
- restricted path integral identity, 655
- retarded correlation function, 94
 - in frequency, 96, 97
- Rydberg series, 373–374
- satellites, 26–29, 268
 - importance of self-consistency, 294
 - in nickel, 330, 408
 - in photoemission, 115
 - in photoemission of silicon, 318
 - in silicon, 324
 - in the GWA, 277, 324
 - in VO₂, 335
 - multiple, 396
- scaling of
 - DMFT solvers with temperature, 485–487
 - GW calculations, 306–308
 - MP2, 198
- scattering experiments, 103
- scattering theory, 194
- Schwinger–Dyson equations, 226
- scissor operator, 317, 325
- scissor shift, 317, 321
- screened Coulomb interaction W , 102, 237, 247
 - polarization contribution W^P , 103
- screened exchange approximation, 254, 261
- screened interaction approximation, 236–239
- second Born approximation, 402
- second quantization, 678–682
- self-consistency in GW, 258, 270, **292–298**, 396
 - cancellation with vertex corrections, 398, 399
 - effect on band width, 270, 321
 - effect on satellites, 270, 294
 - effects on VO₂, 335
 - guidelines, 297
 - in quasi-particle energies, 294
 - RPA and f -sum rule, 399
 - using COHSEX, 294
- self-correlation, 274
- self-energy, **146–147**, **212–213**, 229–230, 313–314
 - exchange–correlation part of, 229
 - and auxiliary systems, 423
 - DMFT, 426
 - functional of G^0 , 212
 - generalized, 147
 - GWA, 253–259
 - imaginary part, 316
 - improper, 212
 - interpretation of, 234
 - irreducible, 212
 - linear in frequency, 288
 - of silicon, 313
 - overview, 152–156
 - proper, 212
 - reducible, 212
 - spatial range, 257
 - self-energy corrections
 - in silicon, 315
 - self-interaction, 63, 285
 - self-interaction correction (SIC), 74, 261, 334, 523
 - self-screening, 273, 392, 394, 398, 417
 - semicircular density of states, **446**, 452
 - and Bethe lattice, 48, 446, 449
 - Sham–Schlüter relation, 732
 - Sherman–Morrison–Woodbury formula, 594, 665
 - shielded interaction approximation, 237
 - sidebands, 26–29
 - sign problem, *see* fermion sign problem
 - silicon
 - band structure, 317, 639
 - bandwidth, 317
 - cumulant, 318, 415
 - dynamic structure factor, 351
 - GW results, 312–319
 - loss function, 104, 375
 - optical absorption spectrum, 104, 372, 381
 - photoemission spectrum, 318
 - quasi-particle energies, 313–316
 - satellites, 318, 324
 - self-energy, 313
 - self-energy corrections, 315
 - spectral function, 313
 - simulation
 - definition, 578
 - singlet–triplet splitting, 357
 - size-consistency
 - of the GWA, 275, 340
 - size-extensive, 196
 - Møller–Plesset (MP) perturbation theory, 197
 - skeleton diagrams, **217**, 234
 - Slater integrals, 512
 - Slater transition rule, 517
 - Slater–Jastrow wavefunction, 410
 - smart Monte Carlo, 594, 635
 - sodium, 267
 - photoemission spectrum, 414
 - sodium clusters
 - in the GWA, 328
 - SOSEX, 403
 - space-time approach
 - for GW calculations, 298, 300
 - specific heat, 115
 - spectral density functional theory (SDFT), 433
 - spectral function, **96–98**, **109–111**, 155, 313, 314
 - cumulant expansion, 413
 - electron–hole, 111
 - interacting, 111
 - negative, 396, 415
 - of silicon, 313–315
 - of the one-body Green’s function, 109
 - of VO₂, 336, 337
 - positivity, 416
 - schematic illustration, 156

- sum rule, 109
- symmetric, 415
- spin liquid, **43**, 44, 55, 449, 652, 737
- spin transition
 - MnO, 547–549
 - two-band model, 451
- spin waves
 - Bethe–Salpeter equation, 382
- spin-flip, 392
- spin-orbitals, 61
- statistical error, **583**
 - binning analysis, 586
 - correlation time, 585
 - estimation, 583
 - standard deviation, 585
- STM images
 - in the GWA, 325
- structure factor, 91, 602
 - dynamic, *see* dynamic structure factor
- substrate
 - screening effect of, 328
- sudden approximation to photoemission, 115
- sum rules
 - in the GWA, 259
 - on spectral function, 96, 109
 - use of, 416
- supercell, **458**
- superfluid, 591, 599, 653
- surface
 - GW corrections, 321
 - optical spectrum, 376, 377
- T-matrix approximation (TMA), **239–241**, 274, 392, 399–402
 - calculations for nickel, 407
 - calculations of lifetimes, 407
 - diagrams, 402
 - electron–hole, 401
 - in practice, 406
 - particle–particle, 400, 407
 - screened, 403
 - spin-flip in, 402
 - time structure of, 400
- t-matrix in multiple scattering theory, 454, 745, 748–749
- Tamm–Dancoff approximation (TDA), 360
 - effects of, 375
- TDDFT, *see* time-dependent density functional theory
- temperature
 - in the GWA, 341–343
 - with path-integral Monte Carlo, 644
- test-charge test-charge dielectric function, 250
- test-charge test-charge interaction, 237
- test-charge test-electron interaction, 250
- thermodynamic limit, 196, 215, 695–697
- three-body scattering, 406
- three-peak spectrum, 29, **53**, 447, 530
- time
 - imaginary, 213
 - time step, 611, 646
 - time-dependent density functional theory, 76–77
 - and charge transfer excitations, 384
 - and the BSE, 383–385
 - and vertex corrections, 385
 - two-particle equation, 384
 - time-dependent density functional theory (TDDFT), **76–77**, 383
 - versus Bethe–Salpeter, 399
 - vertex corrections from TDLDA, 396
 - time-dependent Hartree approximation, 252
 - time-dependent Hartree–Fock, 355, 362
 - time-ordered correlation function, 95
 - in frequency, 97
 - topological order, 128
 - total energy
 - Galitskii–Migdal, 114, 302
 - in DFT-RPA, 302, 337–338
 - in GW, 263, 301
 - in terms of two-body density matrix, 120
 - of h-BN, 339
 - of homogeneous electron gas, 271
 - of hydrogen molecule, 275
 - transient estimate algorithm, 621, 641, 642
 - transition metal oxides, 330–337, 540–549
 - transition metals, 329–330, 537–539
 - transport
 - in the GWA, 326
 - trial energy, 611, 620
 - trial wavefunction, 616, 621, 659, 669
 - triplet excitations, 361
 - Trotter’s formula, 612, 615, 646
 - two-body density matrix, 120
 - two-particle correlation function, 116–118
 - as functional derivative, 231
 - irreducible, 231, 348
 - link to measurable quantities, 346
 - reducible, 349
 - two-particle correlations
 - electron–hole, 366
 - static, 90
 - two-particle Green’s function, 116–120
 - bound part, 349
 - two-particle hamiltonian
 - anti-resonant part, 359
 - coupling part, 359
 - pseudo-hermitian, 359
 - resonant part, 359
 - vacuum fluctuations, 206
 - vacuum state, 206
 - van der Waals dispersion interaction, 18, 136–138, 264
 - DFT-RPA, 265, 338
 - in MP2, 198
 - variance
 - in ground-state path-integral Monte Carlo, 661

- local energy, 133, 619
 - zero-variance property, 592, 605
- variational cluster approximation (VCA), 442, **466**
- variational energy, 133
- variational Monte Carlo, 591–607
 - algorithm, 593
 - in lattice models, 603
 - strengths and weaknesses, 606
- vertex, 247–250
 - four-point interaction, 181, 403
 - fully irreducible, 405
 - irreducible, 247
 - reducible, 247
- vertex corrections, 246, 247
 - by iteration, 393
 - cancellation with self-consistency corrections, 398–399
 - cancellations between W and Σ , 396
 - effects of, 394–399
 - effects on band width, 398, 399
 - effects on ionization potentials, 395
 - effects on lifetimes, 398
 - effects on spectral function, 396
 - first order, 393
 - from TDLDA, 396
 - in W , 394
 - in the self-energy, 395–397
 - non-locality, 395
 - of two arguments only, 396
- virial, 608
- Wannier functions, 513
 - SrVO_3 , 514
- Wannier–Mott excitons, 364, 373
- Ward identity, 417
- water, 649
- wavefunction
 - and localization, 127
- antisymmetry, 123
- atomic radial functions
 - Mn and Gd, 505
- backflow, **141**, 596, 603
- BCS, 140
- continuity, 124, 596
- cusplike condition, 124, 136
- electron–hole, 365–366
- geminal, 140
- Gutzwiller, 46, 603, **740**
- Heitler–London, 45, 58, **130**, 541
- insulators, 126
- Jastrow, 137, 596
- Jastrow correlation, 136
- local energy, 594
- many-body, 122–141
- measures of quality, 133, 597
- nodes, 625–626
- one-body, 597
- optimization, 596–598
- overlap, 134, 142, 629, 642
- pfaffian, 141, 596
- properties, 123–124
- quasi-force, 594
- real, 124
- Slater–Jastrow, **134**, 592, 604
- three-body, **141**, 596
- tiling property, 596, 625
- two electron, 129
- Weiss mean field, 64, 433
- Wick’s theorem, 206
- Wigner crystal, 87
 - density, 87
 - pair correlation function, 93
- Wigner transition, 40–43
- Wigner–Seitz method, 8
- winding number, 647, **653**