

Index

- abortions, previous, 4
 academic performance, 209–10, 255
 adverse outcomes, 210–14
 factors in, 226
 learning disability and, 218
 research needs, 216–18, 219
 testing, 211, 212
 acquisition parameters, for MRI, 44
 activated oxygen radicals, 21
 activation patterns, on fMRI, 120–1
 activities of daily living (ADLs), 143, 151
 ADC, *See* apparent diffusion coefficient (ADC)
 addictive behaviors, 33
 See also substance abuse
 adolescence
 altered neurodevelopment, 70
 behavioral outcome, 117
 brain plasticity, 117
 brain volume, 55
 caudate nucleus size, 57
 cerebellar volume, 60
 corpus callosal size, 56–7
 focal lesions, 58
 functional neuroimaging, 85–90, 86, 258
 MRI data, 69, 70–1
 normal brain development, 70–1
 ventricular enlargement, 55–6
 white matter abnormalities, 104
 adolescence to adulthood changes, 68, 70–3
 adolescents
 attention and hyperactivity problems, 63, 117
 behavioral outcomes, 245
 calculation difficulty, 62
 eating disorders, 119
 educational outcomes, 210
 educational support, 230, 233
 emotional outcomes, 118–19
 executive function, 201–3
 IQ and MRI, 61
 language development, 178–9
 learning difficulties, 219
 peer relationships, 116–17
 psychopathology, 120–1
 risk-taking, 119
 SES and outcomes, 232
 socialization difficulties, 119–20
 adulthood
 functional neuroimaging, 85–90
 long-term morbidity, 12
 MRI findings, 68
 adults
 alcohol use, 133
 bipolar disorder, 132
 drug dependence, 133
 eating disorders, 129–30
 non-psychotic disorders, 128–9
 postpartum psychoses, 132–3
 psychiatric disorders, 127–8
 etiology, 134–6
 family history, 128
 psychological distress, 128–9
 schizophrenia, 130–2
 SES and, 231
 suicide attempts, 129
 adverse experiences, hippocampal changes, 57
 age cohort, in study design, 91, 169, 173
 aggression, preterm and, 226–7
 air pollutants, 5
 air quality, in neonatal ICU, 238
 alcohol abuse, maternal, 4–5, 224
 alcohol use
 preterm adolescents, 119
 preterm adults, 133
 amniotic membranes, 3
 amphetamine sensitization, 135
 anatomical brain subdivisions, 54–5
 anisotropy, *See also* fractional anisotropy (FA)
 in diffusion imaging, 44–5, 48–9, 102
 anorexia nervosa
 adolescents, 119
 adults, 129–30
 risk factors, 254
 antecedent confounders, in studies, 165–6
 antenatal corticosteroids,
 See corticosteroids
 antenatal surfactant, *See* surfactant
 antibiotic resistance, in neonatal units, 20
 antibiotic treatment, preventive, 7–8
 antipsychotic medication, in pregnancy, 134
 anxiety
 in adolescence, 118
 in childhood, 115
 Apgar score, introduction of, 17
 APIP (Avon Premature Infant Project), 244, 247
 apparent diffusion coefficient (ADC), 44–5, 97
 abnormalities, 23, 48
 DEHSI and, 103–5
 white matter heterogeneity, 102–3
 arcuate fasciculus, 107–8
 arithmetical difficulties, 187
 arterial blood gas measurement, 18
 arteriovenous malformations (AVMs), 101
 assisted ventilation
 neurodevelopment and, 146–8
 treatment developments, 18
 ventilator management, 23
 atlases, white matter, 98
 attention allocation processing, 258
 attention deficit, 254
 abnormalities, 104
 academic performance, 218
 adolescents, 117–18
 children, 113–15
 executive functioning, 198
 MRI findings, 62, 63
 attention deficit disorder, white matter integrity, 57
 attention deficit hyperactivity disorder (ADHD), 254
 diagnosis, 113–14
 educational outcome, 216
 in adolescents, 117
 incidence, 113–15
 SES and, 226, 228–9
 attention deployment, 195
 attentional circuitry, 88–9

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- attentional control
 executive functioning, 195, 196
 impulse control, 195
 in adults, 202
 in school-age children, 199–200
 in young children, 197–8
 MRI findings, 62
 working memory deficits, 202
 attentional task, fMRI studies, 90
 auditory cortical activation, 80
 auditory language processing, 84
 auditory recognition memory, 188
 auditory tasks, corpus callosal damage, 85
 auditory tests, attention and attention control, 62
 Australia, mortality rates, 10
 autism, cerebral asymmetry, 55
 autism spectrum disorders (ASDs), 254
 cortical folding abnormalities, 47
 in adolescents, 120
 in children, 116
 automated image processing, 101–2
 AVMs (arteriovenous malformations), 101
 Avon Premature Infant Project (APIP), 244, 247
 axonal development
 changes over time, 102
 impaired, 24
 in adults, 72
 into adolescence, 104
 in white matter volume increase, 71
 myelination abnormalities, 23

 background variables, study design, 171, 216–17
 bacterial infections
 inflammation and, 5–6
 vaginitis, 3, 8
Bacterioides infection, 3
 basal ganglia lesions, 41
 Bayley Scales of Infant Development, 141–2, 151, 172
 BDNF (brain-derived neurotrophic factor), 135–6, 258
 Beery Developmental Test of Visual Motor Integration (VMI), 62
 Behavior Rating Inventory of Executive Function (BRIEF), 200
 behavioral interventions, 237
 behavioral outcomes, 113, 226–7, 254
 caudate nucleus volume, 63
 cognitive functioning and, 215–16
 environmental deprivation, 108
 in adolescence, 117
 in childhood, 113–15
 intensive follow-up, 245
 SES and, 228–9, 231
 Behavioral Theories, in neonatal development, 237–8
 Benton Judgment of Line Orientation test, 62
 bias, in research studies, 165
 bifrontal measurements, 46
 bilateral spastic cerebral palsy (BS-CP), 31, 35
 biological mechanisms, of preterm birth, 5
 biological risk factors, 171, 258–9
 educational outcomes, 214–15, 217, 219
 language function, 181–2
 neurodevelopmental outcomes, 164
 parental backgrounds, 169
 biparietal measurements, 46
 bipolar disorder, 132, 133–4
 birth weight
 CP risk, 30, 32–3
 school performance, 210–11
 subject populations, study design, 168
 blindness, oxygen therapy and, 17
 blocked-task paradigms, 78
 blood volume, PET measurement, 78
 blood-oxygen-level-dependent (BOLD) signal, 47, 76, 79–80
 bowel disorders, 20, 148
 BPD, *See* bronchopulmonary dysplasia (BPD)
 brain
 resting state networks, 79–80
 tissue characteristics, 44
 volume, 49, 64, 215
 quantitative MRI, 45–6, 54–5
 water content, 50
 brain-derived neurotrophic factor (BDNF), 135–6, 258
 brain development, 252–3
 early, 102–3
 gender differences, 149
 growth, 24, 257
 growth failure, 65
 maturation stages, 91
 brain hemorrhage, *See* intraventricular hemorrhage (IVH); periventricular hemorrhage (PVH)
 brain injury
 characteristic lesions, 103, 144
 differing long-term consequences, 92
 educational outcomes, 214–15
 executive functioning, 198–9
 language function, 180
 mechanisms, 20–1, 252
 psychiatric outcomes, 127–8
 qualitative MRI, 58–9
 structural abnormalities, 253–4
 brain organization, fMRI studies, 76
 brain plasticity, 253–4
 adolescent changes, 117
 adult changes, 69
 language function, 179–80
 need for research, 217–18
 neural system specialization, 181
 brain tumor risk, in CT scan, 42
 breast-feeding, 109, 240
 bronchopulmonary dysplasia (BPD)
 impaired cognitive function, 24
 neonatal morbidities, 146–8
 outcome in school-age children, 154, 156
 rates of, 11
 treatment developments, 18
 treatment side effects, 252
 Budin, Pierre, 17
 bulimia nervosa
 in adolescents, 119
 in adults, 129–30

 calculation difficulty, 62
Candida, postnatal infection, 20
 CANTAB tasks, 200, 201–2
 carbon dioxide, hypocarbia, 23
 cardiovascular disease, risk in adulthood, 1
 care procedures, in neonatal ICU, 239–40
 caregivers
 in long-term outcomes, 232–3
 in neonatal ICU, 238–9
 SES and outcomes, 229, 231
 support from, 256
 catechol-O-methyl transferase (COMT) genotype, 136, 258–9
 catheters
 indwelling, 20
 intravascular, 18
 caudate nucleus
 activation, 90
 behavior problems, 63
 cognitive ability, 61
 hyperactivity and, 204
 MRI data, 57, 61, 90
 cell damage, characteristic, 103
 cellular correlates of DTI, 98
 central executive working memory, 185, 187
 central nervous system development, 258
 cerebellar granule cells, 72

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- cerebellum
 - abnormalities, 59–60, 61, 72
 - development, 56, 73
 - volume
 - cognitive deficits, 60–1
 - quantitative MRI, 56
 - reduction over time, 71
 - ‘well-being’ reduced
 - visuo-perceptual ability, 62
- cerebral abnormalities
 - asymmetry, 55
 - undetected, 92
- cerebral blood flow
 - fluctuations in ICU, 21
 - regional (rCBF), 77–8
- cerebral (blood-to-brain) glucose transport (CTXglc), 79
- cerebral cortex
 - formation, 20–1
 - neurons of, 70
 - surface-based analyses, 46–7
- cerebral metabolic rate for glucose consumption (CMRglc), 79
- cerebral palsy (CP), 30
 - age at diagnosis, 31
 - clinical features, 31–2
 - corpus callosal size, 56
 - defined and classified, 30–1
 - dexamethasone and, 20
 - disability changes over time, 120
 - dyskinetic, 18
 - hemiparetic, 107
 - imaging, 34–5, 59–60, 253
 - in neonatal period, 40–1
 - neurodevelopmental outcome, 225
 - parenchymal lesions, 48
 - pathogenesis, 34–5
 - prevalence, 32–3, 34, 151–3
 - prognostic MRI, 48
 - PVL morphology-function in, 35
- cerebral perfusion, regional, 78
- cerebrospinal fluid (CSF)
 - fractional anisotropy, 97–8
 - isotropic diffusion, 97
 - MRI characteristics, 44
 - on ADC image sets, 44
- cervical cerclage, 7
- cervix
 - inflammation and, 5
 - mechanical stress, 7
- changes over time
 - educational outcomes, 213–14, 219
 - in adolescents, 120
 - IQ/DQ scores, 172
 - language function, 179
- childhood
 - caudate nucleus, 57
- cerebellar volume, 56
- congenital malformations, 64–5
- corpus callosum, 56–7
- corticosteroids, 63–4
- diffusion tensor MRI, 59
- focal lesions, 58
- functional neuroimaging, 80–4
- hippocampus, 57
- lateral cerebral ventricles, 55–6
- MRI findings, 59–65
 - qualitative, 58–9
 - quantitative, 54–5
- children, 143–56
 - attention deficit, 63, 113–15
 - cognitive outcomes, 60–1, 141–2, 143
 - developmental outcomes, 141–2
 - emotional outcomes, 115–16
 - executive functioning, 197–9
 - functional outcomes, 141, 143
 - hearing impairment, 152–3
 - hyperactivity problems, 113–15
 - indomethacin effects, 64
 - long-term outcomes, 173–4
 - morbidity, 1
 - motor deficits, 59–60, 61
 - pathways to psychopathology, 120–1
 - social difficulties, 116–17
 - vision impairment, 152
 - visuo-perceptual function, 61–2
- Chlamydia trachomatis* infection, 3
- chorioamnionitis, 5
- chorioamniotic membranes, 3, 5
- chronic lung disease (CLD), 149–50
 - corticosteroid prophylaxis, 63–4
 - post-discharge healthcare needs, 151
 - treatment history, 19, 20
- clinical practice, assessment of VPT/VLBW children, 205
- cognitive abilities
 - adolescent development, 117
 - decrease over time, 145
 - gender effects, 149
 - indomethacin and, 154
 - language development, 178
 - school-age children, 154, 178
 - SES and, 204–5, 225
 - visual impairment and, 62, 225
- cognitive conjunction, 79
- cognitive development
 - assessment, 141–2
 - early interventions, 230, 231–2, 242–3, 244
 - human milk and, 150–1
 - memory development and, 190–1
- cognitive flexibility
 - executive functioning, 195, 196, 197–8
- studies, 198–9
 - in adolescence, 201–2
- cognitive flexibility deficits
 - executive functioning, 200, 201, 202
 - working memory, 202
- cognitive impairment
 - ionizing radiation, 42
 - language development, 177–9
 - mild generalized, 209
 - MRI in childhood, 60–1
 - parenchymal lesions, 48
 - post-discharge environment, 151
 - school-age children, 153–6, 177–8
- cognitive interventions, 149–51, 237, 244–5, 256–7
- cognitive neuroscience
 - memory development, 185
 - memory types, 185–8
- cognitive outcomes, 141–2, 195, 254–5
 - behavior problems and, 215–16
 - BPD children, 146
 - cerebellum volume, 71
 - congenital anomalies, 153
 - corpus callosum growth, 71
 - factors in, 144, 156
 - biomedical and social factors, 226
 - environmental risk, 90
 - SES, 215, 231
 - gray matter changes, 69
 - in school-age children, 154
 - IQ in outcome measurements, 171–2
 - prognostic neuroimaging, 39
 - white matter changes, 69, 108
- cognitive readiness, in language skills, 177
- cognitive scores
 - early school age, 155
 - neonatal risk factors, 147
- cognitive strategies, 92
- cognitive subtraction, 78–9
- cognitive theories, in neonatal development, 237–8
- color naming skills, 179
- community support, 229, 245
- comorbidities, impact of, 151–3
- complex adaptive behavior, 108
- computed tomography (CT), 19, 42–3
 - guidelines, 43
 - outcomes and, 43
 - sensitivity of, 42–3
- conceptual reasoning, 201
- conceptualization in study design, 165–8
- confounding influences, in studies, 165–6

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- congenital anomalies
 maternal postpartum psychosis, 132, 133
 neurodevelopmental outcome, 257
 preterm birth and, 64–5
congenital hearing loss, 152–3
congenital heart disease, 64–5
conjunction analysis, 79
Contingency Naming Test (CNT), 200
 in adolescence, 201–2
 in childhood, 200
contingency-based developmental care, 240–1
continuous monitoring, 25
continuous positive airways pressure (CPAP), 18
control groups, in studies, 167–8
controlled articulatory rehearsal process, 186
conventional MRI, 44
conversational word use, 176
coordination problems, 61
coping strategies
 of children, 231
 parental, 241–2
corpus callosum (CC)
 abnormalities
 adolescence to young adulthood, 71–2
 in adolescence, 104–7
 in VPT adults, 72–3
 changes, 73, 107, 258
 COMT genotype, 136, 258–9
 damage, 85
 gene–environment interaction, 108–9
 plasticity of, 73, 107
 quantitative MRI, 56–7
 size
 cognitive deficits, 60
 in adults, 68
 language function, 61, 180, 204
 thinning
 language function, 85, 180
 minor motor problems, 61
 neural network plasticity, 49
 tractographic reconstruction, 99
 visuo-perceptual ability, 62
 VLBW and SGA compared, 107
 white matter connectivity, 107–8
cortex/cortices, T2 relaxation times, 58
cortical development
 abnormalities, 253
 neuronal injury and, 23–4
 order of maturation, 70
 structural reorganization, 68
 surface-based analyses of folding, 46
corticosteroids
 antenatal
 impact on mortality, 8
 IVH prevention, 22
 outcomes in children, 149, 154
 treatment history, 19
 neonatal, 63–4, 189
 postnatal, 149–50, 156
corticotropin-releasing hormone (CRH), 6–7
cortisol
 fetal inflammatory response, 5–6
 maternal stress response, 6–7, 251
costs of learning disabilities, 228
cranial radiation, white matter damage, 103
cranial ultrasound (CUS), 19, 40
 brain developmental changes, 120
 findings and outcomes, 40–1, 201, 253
 cerebral palsy and, 35
 PVL appearance, 22–3
 guidelines for, 41–2
 undetected lesions, 180, 92, 103
Creating Opportunities for Parent Empowerment (COPE), 241
criminal behavior, 226–7
CT, *See* computed tomography (CT)
cumulative risks, 90, 238
curiosity, importance of, 233
CUS, *See* cranial ultrasound (CUS)
cutaneous hemangioma, 42
cystic PVL
 CUS findings, 40, 41
 early pathology missed, 103
 sensitivity of CT scan, 42–3
cytokines, 5
Cytomegalovirus infection, 3–4
cytotoxic free radicals, 23
deferred imitation, 188
DEHSI, *See* diffuse excessive high-signal intensities (DEHSI)
dehydration, feeding practices and, 18
Delayed alternation (DA), 198
delayed match to sample tests, 89–90
dementia, perinatal brain injury, 73
deoxyhemoglobin, fMRI imaging, 47, 77
dependent behavior, 116
depression, 118
 in adolescence, 118
 in adulthood, 128–9
 in childhood, 115–16
deprivation of stimulation, 108
design issues, in research studies, 165–8
Detour Reaching Box (DRB) task, 198–9
development quotient (DQ), 227
developmental amnesia, 188–9
developmental assessment, 172
developmental coordination disorder (DCD), 121
developmental management, 246–7
developmental outcomes, 141–2
 delay hypothesis, 72
 educational, 213–14
 vision impairment, 152
developmental psychopathology, 259
dexamethasone, 19, 20, 149–50
diagnostic criteria, in long-term studies, 173
dietary deficiencies, 5
diffuse excessive high-signal intensities (DEHSI), 48, 103–5
diffusion tensor imaging (DTI), 59, 97–8, 105–6
 early brain development, 102–3
 information analysis, 98–101
 language function, 182
 MRI compared, 104
 region of interest, 99
 white matter tracks, 203
diffusion-weighted imaging (DWI), 44–5
diffusivity, 97, 102, 103–5
diplegia, 151–2
disability
 antenatal corticosteroids and
 surfactant use, 154
 changes over time, 120
 high-prevalence/low-severity
 dysfunctions, 164–5
 rates of, 145
dropout rates, in long-term studies, 169
drug dependence
 in adulthood, 133
 maternal, 135, 224
drug use, adolescents, 119
DTI, *See* diffusion tensor imaging (DTI)
dyskinetic cerebral palsy, 18
dyslexia, 213
dyspraxia, 61
ear protection, in MRI scanner, 47–8
early brain development, DTI and, 102–3
early childhood, *See* childhood; children
early interventions, 232–3
 and outcomes, 232, 251–2, 256–7
 disadvantaged children, 242
 international studies, 245–6
 programs, 230, 231–2, 242–5

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- studies needed, 218, 245–7, 259
 study design, 233
 early management preterm birth.
 See also intensive care units (ICUs)
 eating disorders
 in adolescents, 119
 in adults, 129–30
 economic costs, academic problems, 209, 255
 educational disability, *See* learning disability (LD)
 educational interventions, 230, 244–5
 studies needed, 247
 educational modifications, 210–11
 educational outcomes
 evidence of, 210–14, 217
 neuropsychological correlates, 215–16
 risk factors, 214–18, 231, 232
 educational support, 218
 vs. brain plasticity, 217–18
 emotional outcomes, 115–16, 226–7
 SES and, 228
 VLBW adolescents, 118–19
 encephalopathy of prematurity, 252
 endocrine stress response, 25
Enterococcus faecalis infection, 3
 environment
 cumulative risk, 90
 enrichment, 26, 72–3
 genetic influences and, 108–9
 in ICUs, 25, 238, 246
 parental behavior, 121
 post-discharge, 151
 environmental factors,
 See also socioeconomic status (SES)
 adolescent changes, 117
 adult psychiatric disorders, 135–6
 age at time of evaluation, 173
 brain development, 24–5
 central nervous system development, 258
 cognitive outcomes, 142, 145, 231
 executive function, 204–5
 in long-term studies, 169–70
 in neonatal development, 237–8
 incidence of PTB, 251
 language outcome, 180–1
 neurodevelopmental outcome, 164
 resiliency and risk after PTB, 224
 social outcomes, 231
 environmental stimuli
 deprivation of, 108
 in neonatal care, 237
 in neonatal development, 237–8
 environmental stress, 204
 environment-restriction hypothesis, 72
 epilepsy, CP and, 31, 32
 episodic buffer, working memory, 185
 episodic memory, 185, 187, 188–9
Escherichia coli infection, 3
 estimated date of delivery (EDD), 2
 ethamsylate, 22
 ethnicity
 folate metabolism, 5
 risk factor for PTB, 3
 SES and, 225, 228
 etiologies of preterm birth, 5
 European countries, CP rates in, 33, 34
 event-related potential (ERP) studies, 188
 event-related-tasks paradigms, 78
 executive control system, 196
 executive function
 central executive working memory, 187
 clinical practice, 205
 conceptual model, 196
 deficits, 203, 218
 development, 195, 203–4, 205
 gray and white matter changes, 69
 neural network adaptations, 49–50
 study methodology, 196–7
 executive functioning
 cognitive outcome, 254–5
 in adolescents, 118, 201–3
 in children, 197, 199–201,
 experience, white matter connectivity, 107–8
 explicit long-term memory, 187–8
 impairments, 189, 190
 MTL and, 189
 explicit memory, 185
 expressive language, 177, 178
 external stimuli, in NICU, 239
 externalization problems, 226–7, 254
 extremely low birth weight (ELBW)
 long-term morbidity, 141
 neonatal complications, 40, 141
 extremely preterm birth (EPT/EPTB), 1, 2
 childhood outcomes, 144–5
 CP rates, 33–4
 maternal infections and, 3
 neonatal morbidity, 11
 eye problems, *See* retinopathy of prematurity (ROP); visual impairment
 FA, *See* fractional anisotropy (FA)
 face–name association task, 258
 familiarity-based recognition, 187–8
 family background
 behavioral outcomes, 226
 cognitive outcomes, 227
 family functioning, remedial interventions, 233, 243
 family history
 psychiatric disorders in adults, 128, 131, 134
 risk factor for preterm birth, 3
 family income, 170
 family service plan, 245
 feedback processes
 environmental stimuli, 237–8
 executive functioning, 195
 feeding difficulties, early interventions, 243
 feeding practices
 and hypoglycemia, 18
 in neonatal units, 20
 femoral length, gestational age, 2
 fetal growth retardation, 1, 131–2
 fetal hypoxia, 132
 fetal programing hypothesis, 127
 fetal size measurement, 2
 fetal stress, 6
 fetal ultrasound, gestational age, 168
 fetus, inflammatory response in, 5–6
 fiber tracking, *See* tractography
 field of view
 in CT scan, 42
 in CUS, 40
 fluid-attenuated inversion recovery (FLAIR) imaging, 44, 58
 Flynn effect, IQ/DQ scores, 172
 fMRI, *See* functional magnetic resonance imaging (fMRI)
 focal lesions, 58
 folate intake, 5
 fractional anisotropy (FA), 49
 abnormalities, 104
 and DTI, 59, 97–8
 changes over time, 102
 IQ correlations, 108
 prognostic value, 104
 region of interest (ROI), 98
 tract anatomy, 98–101
 VBM template, 102
 VLBW and SGA compared, 07
 white matter heterogeneity, 98, 102–3
 free radicals, cytotoxic, 23
 friends, as protective factor, 229
 frontal lobes
 asymmetry, 55
 white matter connectivity, 108
 fronto-parietal network, 185–7

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- functional connectivity MRI (fcMRI), 47
- functional deficits
cerebellar damage, 72
in long-term studies, 170–1
- functional magnetic resonance imaging (fMRI), 47, 76
activation patterns, 120–1
basic principles, 76–7
limitations, 91
neural adaptations, 69
neural development, 79–80
neural plasticity, 257–8
neural specialization, 181, 182
- functional neuroimaging
methodology, 76–9, 90–2
studies, 79–80
adolescent and adult, 86
childhood, 83
neonatal, 81–2
- functional outcomes
children, 141, 143
early interventions, 149–51
infants, 144, 152–3
neonatal morbidities, 151
- functional skills, children, 153
- future research questions
academic performance and learning disability, 219
cerebellum and corpus callosum long-term, 73
memory development, 190–1
white matter development, 109
- future-oriented behavior, 195
- Gardnerella vaginalis* infection, 3
- gender
ADHD and, 115
age for gray matter peak volume, 70
aggression, 226–7
brain volume reduction, 55
depressive disorder, 128–9
developmental effects on brain function, 91–2
eating disorders, 119, 129–30
educational outcomes, 232
emotional vulnerability, 118–19
growth delay and social difficulties, 120
indomethacin exposure effects, 22, 64
internalizing behaviors, 128
language processing, 61, 71, 84
learning disability and SES, 228
neonatal morbidities, 148–9
- General Health Questionnaire (GHQ), 71
- genetic influences
emotional outcomes, 119
gestational age and, 258–9
in resilience, 108–9
preterm birth risk, 3, 5, 251
- germinal matrix (GM), and IVH, 21–2, 252
- germinal matrix hemorrhage (GMH), 103
- gestational age
“border of viability” 8–9
categorization, 2
CP risk, 30, 32, 33–4, 40–1
estimation, 1–2, 168
morbidity rates, 10, 11–12
subject populations, 168
- girls, eating disorders, 119
- glial cells, embryonic, 20
- Go/No-go task, 88–9, 90
- goal setting, executive functioning, 195, 196, 200–1
- goal-directed behavior, 195
- grade repetition, 210–11
- grammatical ability, 176, 177
- gray
white matter ratio, 68, 69, 92
- gray matter
fractional anisotropy, 97–8
isotropic diffusion, 97
normal development, 70–1
volume changes, 181
increase, 252
MRI data, 49, 54–5, 257
reduction, 23–4, 55, 72, 253
- growth, and nutrition, 150–1
- growth and development outcomes of Dutch children, 232
- growth delay, and social difficulties, 120
- head circumference
brain size and, 54
cognitive outcomes, 24, 215
gestational age, 2
- healthcare needs, post-discharge, 151
- health outcomes, intensive follow-up, 245
- hearing impairment, 18, 31, 152–3
- hemiparetic cerebral palsy, 107
- hemiplegia, 48, 151–2
- hemispheric lateralization, 79–80, 85
- hemodynamics of brain activity, 76–7
- hemorrhage, *See also* intraventricular hemorrhage (IVH); periventricular hemorrhage (PVH)
CT scan sensitivity, 42–3
CUS visualization, 40
- MRI value, 48
- high-prevalence/low-severity dysfunctions, 164–5, 218
- hippocampus
hypoxic-ischemic damage, 89
memory and, 187–9
quantitative MRI, 57
T2 relaxation times, 58
volume and cognitive ability, 61
- Holland, growth and development outcomes of Dutch children, 232
- home environment
educational outcomes, 214
executive function development, 204
neonatal risks, 224
SES and, 227
- home visitors, early interventions, 243, 244
- hospital services, post-discharge needs, 151
- hospital-based interventions, 238–42
- human breast milk, 150–1
- hyaline membrane disease, 18, 19
- hydrocephalus, 21
- hyperactivity
caudate nucleus and, 63, 204
in childhood, 113–15
of adolescents, 117–18
- hypocarbica, 23
- hypoglycemia, 18, 19–20, 24
- hypothalamic–pituitary–adrenal (HPA) axis, 251
- hypothermia, 25–6, 79
- hypoxia, BDNF and, 258
- hypoxia-ischemia
adult schizophrenia, 132, 135
in white matter injury, 23
PLIC abnormalities, 48
risk in VPTB, 72, 144
- iatrogenic brain injury, 17
- ICUs, *See* intensive care units (ICUs)
- IDEA (Individuals with Disability Education Act), 245, 247
- IHDP, *See* The Infant Health and Development Program (IHDP)
- illness severity scores, 171
- immune system, 5
- implicit learning, 189, 190
- implicit long-term memory, 189
- implicit memory, 185
- impulse control
executive functioning, 195
lacking, 199–200
- inattention, 186, 201
- incubators, early developments, 17, 18

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- individual variability
 - arteriovenous malformations (AVMs), 101
 - fMRI study confounder, 91, 92
 - neural plasticity, 92
 - physiological baseline, 90
 - white matter anatomy, 98, 104
- individualized developmental care, 240–1
- Individuals with Disability Education Act (IDEA), 245, 247
- indomethacin
 - gender effects, 149
 - IVH prevention, 22, 150
 - language delay, 80–4
 - outcome in childhood, 64, 154
- induced preterm birth, 5
- infancy
 - executive functioning, 197–8
 - memory development, 188
- Infant Development, Scales of, 141–2, 151, 172
- infant mortality, incidence, 8–10
- infants/toddlers, cognitive development, 141–2, 143
- infections
 - in neonatal units, 20
 - in white matter injury, 23
 - neonatal morbidities, 148
 - preterm birth risk, 3–4
- inflammation
 - genetic mechanism, 5
 - preterm risk, 5–6, 144
 - preventive strategy, 8
- information processing, 195, 196
- inhibitory control, 258
- insecure attachment, 119, 121
- intelligence quotient (IQ), *See* IQ
- intensive care (neonatal)
 - changes and long-term studies, 26
 - environment, 256
 - historical overview, 17
 - 1965–1985, 18–19
 - 1985–1995, 19–20
 - 1995 to present, 20
 - pre-1965, 17–18
 - neurodevelopmental outcome, 251–2
 - quality and survival rates, 9–10
 - variability, 168
- intensive care units (ICUs)
 - developmental management, 246
 - environment modification, 238
 - MRI difficulties, 47–8, 79
- interleukin-1 receptor antagonist, 5
- interleukin-6 (IL-6), promotor gene, 5
- interleukin-10 (IL-10), gene polymorphism, 5
- internal capsule, 107–8
- internal locus of control, 231
- internalizing difficulties, 115–16, 118
- inter-observer agreement, brain metrics, 46
- inter-observer variability, CUS findings, 40, 41–2
- interpregnancy interval, 4
- intrauterine environment, 144
- intrauterine growth restriction, 1, 47
- intrauterine infections, 3–4, 6
- intraventricular hemorrhage (IVH), 11
 - cranial ultrasound, 35, 41, 43
 - detection, 19, 39
 - DTI data, 103
 - executive functioning studies, 199
 - germinal matrix, 21–2, 252
 - neonatal morbidities, 145–6
 - pathogenesis, 21–2
 - prevention, 22, 150
 - rates of, 11
- introduction to treatment, 19
- ionizing radiation, 42
- IQ discrepancy, 212
 - changes over time, 213–14
 - learning difficulty assessment, 211
- IQ scores
 - and ADHD, 114–15
 - breast-feeding and, 109
 - cognitive outcome, 254
 - correlations with FA, 108
 - corticosteroid prophylaxis, 64
 - decline with age, 61
 - in childhood, 60–1
 - in CP children, 151
 - in school-age children, 154
 - language development, 178–9
 - limitations in, 171–2
- iron, free, 21
- isolette use, 47
- IVH, *See* intraventricular hemorrhage (IVH)
- jaundice, 18
- Judgment of Line Orientation Test, 62
- kangaroo mother care (KMC), 240
- kernicterus, jaundice and, 18
- language development
 - adolescents, 178–9
 - early interventions, 231–2
 - indomethacin and, 64
 - preschool years, 176–7
 - school years, 177–8
- study methodology, 179–80
- unilateral spastic cerebral palsy, 32
- language function, 176, 181–2
 - brain injury and, 180
 - cognitive development, 141–2
 - corpus callosum thinning, 85
 - environmental risk, 90, 180–1
 - gray and white matter changes, 69
 - indomethacin and, 154
 - neural network adaptations, 49–50
 - normal development, 176
 - studies, 179–80, 182
- language impairment
 - cognitive outcome, 255
 - delays, 80–4
 - hearing impairment and, 153
 - neural basis of deficits, 181
 - working memory problems, 187
- last menstrual period (LMP), 2
- lateral cerebral ventricles, 55–6
- lattice anisotropy index (LAI), 97–8
- learning disability (LD), 209–10
 - academic performance and, 218–19
 - cognitive outcome, 228, 255
 - in school-age children, 154
 - neurological abnormalities and, 257
 - rates and types, 211–13
 - research limitations and needs, 216–18
 - SES and, 228
 - vision impairment and, 152
 - visual paired-associated learning, 89
- learning readiness
 - deficits, 213
 - early interventions, 218, 230
 - importance of, 233
- left caudate nucleus, 204
- leiomyomata, uterine wall, 7
- letter trials, 85
- life course model, 238
- lighting, in neonatal ICU, 238
- listening vocabulary scores, 80, 84
- literacy, development, 176
- long-chain poly-unsaturated fatty acids LC-PUFA, 150–1
- long-term effects, PTB or FGR, 1
- long-term follow-up studies
 - conceptualization, 165–8
 - difficulties, 164
 - importance of, 174
 - language function, 182
 - methodology, 165, 171–4
 - need for, 109, 165, 217
 - subject populations, 168–71
- long-term memory, 185, 188–9

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- low achievement, 211, 212, 213
- low birth weight (LBW), 1
- adult psychological distress, 128–9
 - adult schizophrenia, 130–1
 - and ADHD, 114, 115
 - CP risk, 31, 32
- lung disease
- See also* bronchopulmonary dysplasia (BPD); respiratory distress syndrome (RDS)
 - chronic, 18
- magnetic resonance imaging (MRI)
- See also* diffusion tensor imaging (DTI); diffusion-weighted imaging (DWI); functional magnetic resonance imaging (fMRI)
 - biparietal diameter, 46
 - blood-oxygen-level-dependent contrast, 77
 - brain developmental changes, 120
 - brain growth and, 257
 - brain injury detection, 253–4
 - conventional, 44
 - guidelines, 50
 - in adolescence, 68, 120
 - in adulthood, 68–9
 - in childhood, 59–63
 - neurodevelopmental outcomes, 59–63, 257
- magnetization transfer ratio (MTR), 58–9
- major motor deficits, 59–60
- Malawi, 2, 4
- management of preterm birth, 63–4,
- See also* intensive care units (ICUs)
- mannose-binding protein 2 (MBL2), 5
- maps, white matter atlases, 98
- massage, 239
- maternal age, 4
- maternal education level, 228, 243
- maternal factors in outcomes
- cognitive function, 26, 181
 - cumulative risk, 238
 - educational outcomes, 214
 - neonatal risks, 224
 - post-discharge environment, 151
 - preterm birth risk, 4, 251
 - psychiatric outcomes in adulthood, 127–8
 - suicide in adults, 129
- maternal mental health
- bipolar disorder, 133–4
 - depression, 26
 - eating disorders, 130
 - schizophrenia, 133–4
- maternal parenting skills, 204
- maternal satisfaction, in neonatal ICU, 240
- mathematical ability
- academic skills and, 211, 212–13, 218
 - neuropsychological correlates, 215–16
 - shortcomings of tests, 217
- MDI (Mental Development Index), 151
- mean diffusivity (MD), 97, 98
- mechanical stress, uterus and cervix, 7
- medial temporal lobe (MTL), 188, 189
- medial temporal lobe-cortical circuit, 187–8
- mediator variables, 166
- medical advances, 169, 197
- medical care, SES and, 227–8
- medical complications
- age at evaluation, 173
 - neonatal, 11, 121
- medical interventions, 204
- medical management, 216
- medical risks, 90, 214–15
- membranes
- amniotic, 3
 - chorioamniotic, 3, 5
 - rupture, 5–6
- memory deficits
- cognitive outcome, 255
 - hippocampal damage, 89
 - visual paired-associated learning, 89
 - working memory, 89–90
- memory development, 190–1
- cognitive neuroscience, 185
 - in infancy, 188
 - phonological working memory, 177
- memory function, 49–50
- memory types, 185–8
- Mental Developmental Index (MDI), 141
- mental health risk factors, 120–1
- metabolic activity, functional neuroimaging, 79
- metabolic brain injury, 24
- microglia activation, 23
- middle childhood, *See* school-age children
- minor motor impairments, 61
- moderately preterm birth, 1, 2, 8
- moderator variables, 166
- monitoring neonates, with fMRI, 79
- morbidity
- in childhood, 1
 - increased survival and, 142, 144–5
 - long-term outcomes, 12, 251
 - neonatal, 1, 11
- mortality rates, 1, 8–10
- motivation, resilience and, 229
- motor deficits
- cerebral palsy, 32, 31
 - major, 59–60
 - minor, 61
 - parenchymal lesions, 48
- motor development, 90, 242
- motor learning task, 190
- motor response inhibition, 88–9
- motor skills, 141–2
- Movement Assessment Battery for Children (ABC) scores, 61
- MRI, *See* magnetic resonance imaging (MRI)
- MTL, *See* medial temporal lobe (MTL)
- multicenter networks, 168
- multimodal interventions, in ICU, 240–1
- multiple pregnancies/births, 4, 7, 144
- multi-search multi-location (MSML) task, 198
- Mycoplasma hominis* infection, 3
- myelination
- diffusion changes, 102
 - impairment, 103
 - MTR data, 58–9
 - white matter
 - diffusion imaging, 44–5, 98
 - volume increase, 71, 72
- naming skills, 179
- narcotics use, 4–5
- nasogastric tubes, 18
- necrotizing enterocolitis (NEC), 20, 148
- neighborhood conditions, 228, 229
- neonatal brain injury, undetected,
- See* undetected early lesions
- neonatal care, 25–6, 237
- neonatal corticosteroids,
- See* corticosteroids; neonatal
- neonatal development
- hospital-based interventions, 238–42
 - post-discharge interventions, 242–5
 - theoretical perspectives, 237–8
- neonatal functional neuroimaging studies, 79–80, 81–2
- neonatal intensive care, *See* intensive care (neonatal)
- neonatal interventions, and outcomes, 149–51
- neonatal medical complications
- academic performance, 217
 - long-term outcomes, 121
- neonatal medical risk factors, 214–15
- neonatal morbidity, 257
- combined, 153
 - post-discharge needs, 151

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- preterm infants, 1, 11
 rates, 10–12
 SES and, 225
 VLBW infants, 145–9
 neonatal mortality, 8–10
 neonatal risk factors, 147, 224–5
 neural activation, fMRI detection, 47,
 79–80, 91
 neural basis of language deficits, 181
 neural organization
 adolescent changes, 117
 developmental stages, 79–80
 fMRI data, 49–50, 69, 76
 neural plasticity
 early CC damage, 85
 fMRI data, 92, 257–8
 neural-maturational theory, 237
 neurobehavioral interventions, 243
 neurodevelopmental impairment
 infection and, 148
 neonatal morbidities, 153
 neurodevelopmental outcomes, ix,
 225
 adult psychiatric problems, 127
 assessment, 172–3
 attention deficit, 59–63
 brain growth and, 257
 breast milk feeding, 150–1
 early interventions, 251–2
 studies, 164–5
 neurodevelopmental processes, 186
 neurodevelopmental research, 259
 neuroendocrine stress response, 6–7,
 258
 neurogenesis, enhanced, 26
 neuroimaging, *See also* specific
 modality
 brain injury detection, 253–4
 CP pathogenesis, 34–5
 future trends, 26
 in research, 39
 neurological abnormalities
 and learning disability, 257
 dexamethasone and, 20
 neuronal activity, alterations in, 76
 neuronal injury, causal mechanisms,
 23–4
 neurons, of cortex, 20–1, 70
 neuropsychological correlates, 209,
 215–16
 neurosensory impairment, 48
 New Zealand, mortality rates, 10
 Newborn Individualized
 Developmental Care and
 Assessment Program
 (NIDCAP), 25, 48–9, 108,
 240–1, 256, 258
 noise, in neonatal ICU, 238
 non-cerebral malformations, brain
 MRI, 64–5
 non-psychotic adult disorders, 128–9
 non-violent suicide attempts, 129
 Norway, preterm birth risk, 4
 novelty preferences, 188
 nursing home visit, 243
 nutrition in the ICU, 20, 150–1, 246
 nutritional causes of brain injury, 24
 object/meaning association, 176
 obsessive compulsive disorder (OCD),
 109
 obstetric complications, 128, 129, 135
 occipital region brain tissue volumes,
 63
 occupational therapy, 243
 Oddball trials, 88–9
 old age
 corpus callosum size, 107
 perinatal brain injury, 73
 oligodendrocyte precursors, 23, 73,
 103
 oligodendrocytes
 injury to, 23
 numbers of, 102
 one-dimensional measurements, 45–6
 operant conditioning, 237–8
 operator bias, tractography, 99
 optimized VBM, 101
 organizational ability, 195, 196
 outcome measurement, 166, 171–4, 257
 in long-term studies, 171–3
 outcomes
 CT findings and, 43
 CUS images and, 40–1
 DTI findings and, 104
 MRI findings and, 48–50
 oxygen extraction, PET measurement,
 78
 oxygen flow, fMRI imaging, 76–7
 oxygen therapy, side effects, 17, 24, 252
 oxyhemoglobin, fMRI, 47, 77
 painful experiences, neonatal, 24–5
 parametric maps, 44, 45
 parenchyma
 on CUS, 40
 on MRI, 48
 parental backgrounds
 confounder in long-term studies,
 169–70
 educational outcomes, 214, 226,
 256
 resilience of child, 229
 social factors, 226, 229, 256
 parental education, 232
 parental mental disorder, 119
 parental stress and coping, 117, 241–2,
 246
 parent–child interactions
 cognitive enhancement, 218, 228,
 242–3, 256
 emotional outcomes, 119
 insecure attachment, 121
 SES and, 228
 parenting ability
 behavioral problems, 226–7
 developmental outcome, 121
 executive function, 204
 parenting interventions, 233
 parenting support, 226–7, 246
 parent-rated scores, 117–18, 217
 parietal lobe, 54–5
 parity, and maternal age, 4
Parvovirus B19 infection, 3
 passive listening language studies, 80–4
 patent ductus arteriosus (PA), 64
 paternal schizophrenia, 134
 patient access, in CT scan, 42
 PDI (Psychomotor Developmental
 Index), 141–2, 151–2
 Peabody Picture Vocabulary Test, 179
 pediatric follow-up care study, 244–5
 peer group difficulties, 116–17
 in adolescence, 120
 in childhood, 115
 social factors in, 226
 peer group socialization, 176
 perceptive selective attention
 allocation, 88–9
 perceptual information processing,
 187–8
 performance IQ scores, 60–1
 perinatal brain injury, 108–9, 259
 perinatal death, postpartum psychosis,
 132, 133
 perinatal interventions, outcomes,
 149–51
 periodontal disease, 3, 8
 peristriate cortex, 85
 periventricular hemorrhage (PVH),
 180
 periventricular hemorrhagic infarction
 (PHI), 103
 periventricular infarction (PVI), 41
 periventricular leukomalacia (PVL),
 35, 103, 145–6
 cognitive defects, 60
 CT scan sensitivity, 42–3
 early pathology missed, 103
 encephalopathy of prematurity,
 252
 long-term findings, 58, 59–60
 metabolic activity in primary
 sensorimotor cortex, 79

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- periventricular leukomalacia (PVL)
(*cont.*)
 morphology-function correlates,
 35, 36
 MRI images, 49
 myelination, 58–9
 rates of, 11, 22–3
 ultrasound detection, 35, 39
 visual impairment, 36, 62, 63
 white matter mapping, 59
- periventricular white matter, injury,
 22–3
- personality style, in adulthood, 72
- phenobarbitone, IVH prevention, 22
- phonetics, sound discrimination, 176
- phonological loop, 185, 186
- phonological processing
 language, 80–4
 neural basis, 181
 of adolescents, 178, 179
- phonological working memory
 deficits, 190, 255
 skills development, 177
- photic stimulation, 80
- physical outcomes, 116, 225
- physiological baseline, variability, 90
- physiological monitoring, 18
 developments, 18–19
 equipment problems with fMRI, 79
- Piagetian Theories, 238
- piano playing, 107–8
- placenta, impaired function, 6
- placental abruption, 6
- planning ability
 executive functioning, 195, 196,
 200–1
 in adolescence, 201–2
 school-age children, 200–1
- PLIC (posterior limb of the internal
 capsule), 48
- porencephalic cyst, 21
- Portage, 244
- positioning, in neonatal ICU, 239
- positron emission tomography (PET),
 77–8
- positron emitter, 78
- post-discharge environment, 151
 healthcare needs, 151
 interventions, 242–5, 246–7, 256–7
- posterior fossa, on CUS, 40
- posterior limb of the internal capsule
 (PLIC), 48
- posthemorrhagic ventricular dilatation
 (PHVD), 41
- postnatal infections, 20
- postnatal risks, SES and, 227–9
- postpartum psychoses, 132–3
- poverty, 227–9, 230–1
- practical issues, 47–8
- pragmatics, 176
- preeclampsia, 6
 heritability, 3
 preventive strategy, 8
- pregnancy characteristics, psychiatric
 outcomes in adults, 127
- pregnancy wheel, 2
- premature baby units, history, 17
- prematurity, *See* preterm birth
- prenatal care, lack of, 224, 225
- prenatal stress, 230–1
- preschool children
 cognitive assessment, 143, 173
 cognitive development, 227–8
 early development intervention, 230
 language development, 176–7
 memory development, 188–9
 SES, 227–8
- preterm birth, ix, 251
 behavioral outcomes, 117, 226–7
 childhood outcomes, 143–56
 classification, 168
 cognitive outcomes, 226
 defined, 1–2, 168
 early management, 63–5
 eating disorders, 130
 emotional outcomes, 115–16, 226–7
 incidence, 1, 251
 long-term studies, 170–1
 neurological sequelae, 30
 physical outcomes, 225
 postpartum psychosis, 132–3
 psychiatric outcomes, 130, 135–6
 psychological distress, 128
 rates of, 2–3
- preventive strategies, 7–8
- primary school, special education
 service needs, 209
- Probst bundle, 107
- procedural issues, in study design, 257
- procedural learning, 189
- process features of environment, 170
- prognostic information
 DTI, 101, 104
 MRI, 48–50, 253
 neuroimaging, 39–40
- pro-social behavior, 233
- prostaglandins, inflammation and, 5, 6
- protective factors, 229
- psychiatric morbidity, 254
 gene–environment interaction, 109
 in adolescence, 70
 in adulthood, 127–8, 134–6, 135
 future research, 73
 in childhood, 115–16
- psychological distress, in adults, 128–9
- Psychomotor Development Index
 (PDI), 141–2, 151–2
- psychopathology, pathways in, 120–1
- psychosocial development, 231–2
- public funding costs, of learning
 disability, 228
- quadriplegia, 151–2
- qualitative MRI, 58–9
- quantitative MRI, 54–7
- quantitative volumetric analysis, 45
- radioactive nuclide tracers, 77–8
- reactive oxygen species, 23
- reading ability
 academic performance and, 212–13,
 218
 neuropsychological correlates,
 215–16
 of adolescents, 178, 179
 of children, 178
 test shortcomings, 217
 white matter maturation, 98
- reading skills deficits, 80, 84, 255
- recall memory, in infancy, 188
- receptive vocabulary scores, 80, 84
- recollection-based recognition, 188
- region-of-interest (ROI)
 DTI data, 98, 99, 102–3
 MRI data, 68
 VBM compared, 101
- regional brain volumes, 54–5
- regional cerebral blood flow (rCBF)
 PET data, 78
 SPECT data, 77–8, 79
- rehospitalization, SES and, 225
- relative anisotropy (RA), 97–8
- religion, and outcomes, 229, 231
- remediation, language impairment, 177
- repeated preterm births, 3, 4
- representative samples, study design,
 168
- research needed, 1, 216–18, 245–6
- resilience, 229, 256
 SES and, 224, 231
 white matter and, 108–9
- respiratory distress
 executive function deficits, 202
 oxygen therapy risks, 17
 treatment developments, 18
- respiratory distress syndrome (RDS)
 morbidity long-term, 10, 12
 treatment impact, 8
 visual recognition memory, 188
- response inhibition, 202
- response selection, 88–9

- resting-state networks, 79–80
 results interpretation, study design, 257–9
 retinal damage, oxygen therapy, 17
 retinopathy of prematurity (ROP), 152
 neurodevelopmental outcome, 225, 257
 rates of, 11
 Rey-Osterrieth Complex Figure (RCF), 201
 planning ability, 201
 in adolescence, 201–2
 in childhood, 201
 risk aversion, 119, 133
 risk factors
 for preterm birth, 3–5, 252
 in neurodevelopmental outcome, 164
 SES and, 224
 Romanian orphans, 108
 ROP, *See* retinopathy of prematurity (ROP)

 sample selection, in studies, 197
 Scales of Infant Development, 141–2, 151, 172
 schizophrenia
 BDNF and, 258
 cerebral asymmetry, 55
 in adults, 70, 130–2
 maternal, 133–4
 risk factors, 254
 school
 as protective factor, 229
 behavioral disturbance in, 117–18
 quality in cognitive outcomes, 227
 school performance, teacher ratings, 210–11
 school-age children, 153–6
 cognitive assessments, 143, 173
 executive functioning, 199–201
 gender effects, 149
 language development, 177–8
 memory development, 188–9
 sedation, with fMRI, 79
 self-esteem, 229
 self-monitoring, lacking, 199–200
 self-regulation, 195
 semantic language processing, 80–4, 176, 181
 semantic memory, 185, 187
 sensorimotor function, 80
 sensorineural hearing loss, 18
 sensory development, 49–50, 80
 sensory stimuli, in neonatal units, 25
 sepsis, neonatal morbidities, 148
 serial reaction time task, 189–90

 serotonin levels, preterm children, 129
 SES, *See* socioeconomic status
 SGA (small for gestational age), VLBW compared, 107
 shifting tasks, 200
 short-term phonological memory store, 186
 sigmoid bundle, 107
 single modality interventions, in neonatal ICU, 239–40
 single photon emission computed tomography (SPECT), 77–8
 skill learning, 189
 skin-to-skin care (SSC), 240, 246
 sleeping, resting-state networks, 79–80
 sleep–wake cycling, in neonatal units, 25
 small for gestational age (SGA), VLBW compared, 107
 smoking
 maternal, 4, 224, 251
 preventive strategy, 8
 smoking and schizophrenia in pregnancy, 133, 134
 social abilities, in adolescence, 117
 social costs, academic problems, 255
 social factors
 executive function development, 204–5
 in academic performance, 219
 social skills
 functional assessment, 143
 hearing impairment, 153
 resilient children, 229
 social variables, multiple factors in, 171
 socialization difficulties
 in adolescence, 119–20
 in childhood, 116–17
 sociodemographic factors, cognitive development assessment, 142
 socioeconomic status (SES), 256
 cognitive outcomes, 181, 215, 227–9, 231
 disadvantaged, 8, 214–15, 227–9
 early interventions, 8, 230, 232–3
 educational outcomes, 214, 232
 environmental factors, 170, 229
 future studies, 259
 language function, 181–2
 long-term studies, 229–32
 preterm birth risk, 4
 protective factors, 229
 risk and resiliency, 224
 social outcomes, 231
 variability in cohort selection, 216–17
 socioemotional skills, 233

 sound discrimination, phonetics, 176
 sound exposure, in neonatal units, 25
 spastic cerebral palsy, 49, 31–2, 60
 spastic diplegia, 151–2
 MRI in childhood, 59, 60
 visuo-perceptual problems, 62
 spastic quadriplegia, 59
 spatial memory tasks, 89–90, 258
 spatial reversal (SR), 198
 spatial working memory
 deficits, 202, 255
 in adolescence, 201–2
 in young children, 198
 special education services
 Dutch long-term study, 232
 in primary school, 209, 210–11
 risk factors for, 227
 SES and, 226, 228
 specific language impairment (SLI), 178
 specific learning disabilities
 IQ discrepancy, 212
 low educational achievement, 213
 SPECT (single photon emission computer tomography), 77–8
 speech and language
 school-age children, 177–8
 SES and, 228
 speech articulation, 177–8
 speech development, 176
 speech impairment, 228, 255
 spontaneous preterm birth, 5
Staphylococcus infection, 20
 state modulation program, 243
 status features of environment, 170
 stimulation, deprivation of, 108
Streptococcus species infection, 3
 stress, parental, 121, 241–2
 stroke
 detection by DWI, 44
 early diffusion changes, 98
 white matter plasticity, 107
 structural abnormalities, on VBM, 101
 structural equation model, 215
 structural MRI, 54
 structural reorganization of brain, adult data, 69
 study design
 consistency needed, 216–18
 design issues, 165–8
 inconsistencies, 213–14
 result interpretation, 257–9
 subcortical leukomalacia, 41
 subcortical tissue loss, 56
 subject populations
 cohort selection, 216–18, 257
 loss in long-term studies, 169
 study design, 168–71

Cambridge University Press

978-0-521-87179-2 - Neurodevelopmental Outcomes of Preterm Birth: From Childhood to Adult Life

Edited by Chiara Nosarti, Robin M. Murray and Maureen Hack

Index

[More information](#)

Index

- subplate neurons, 23–4
- substance abuse
- preterm birth risk, 4–5, 251
 - preventive strategy, 8
 - suicide attempts, 129
- superior fasciculus, 107
- supplemental stimulation, 238–40
- support systems, SES and, 170, 231
- surface-based renderings, 47
- surfactant
- antenatal, 149
 - impact on mortality, 8
 - long-term outcomes, 154
 - treatment history, 19
- surgical intervention, for NEC
- children, 148
- survival rates, increased morbidity, 144–5
- Sweden
- infant mortality reduction, 8–10
 - mortality rates, 9, 10
 - preterm birth rates, 1, 2–3
 - preterm birth risk, 4
- synaptogenesis, fetal, 21
- T1/T2-weighted MRI, 44, 50, 58
- task performance differences, 91
- teacher/mentor, as protective factor, 229
- teacher-rated scores
- for adolescents, 117–18
 - for children, 210–11
 - shortcomings, 217
- teachers, training in child development, 233
- teenagers, *See* adolescents
- temporal lobe, quantitative MRI, 54–5
- term-equivalent MRI, 48–50
- thalamo-occipital measurement, 41
- The Abecedarian Project, 230
- The Infant Behavioral Assessment and Intervention Program (IBIAP), 243
- The Infant Health and Development Program (IHDP), 231–2, 244–5
- interventions, 256
 - studies needed, 246
- The Kauai Longitudinal Study, 230–1
- time of discharge MRI, 48–50
- time trends, *See* changes over time
- tissue correlates, of DTI, 98
- tobacco use, *See also* smoking
- touch stimuli, in neonatal ICU, 239
- Tower of Hanoi (TOH) test, 200–1
- Tower of London (TOL) test, 200–202
- tractography, 98–101, 104
- transportation, from ICU to MRI, 47
- transverse cerebellar measurements, 46
- treatment side effects, 252
- triplegia, 151–2
- tumor necrosis factor (TNF), 5
- twins, 4
- ultrasound (US), *See also* cranial ultrasound (CUS)
- fetal size, 2
 - gestational age, 168
- undetected early lesions
- DTI in prevention, 97
 - fMRI study confounder, 92
 - PVL, 103
 - ultrasound (US), 103, 180
- unilateral spastic cerebral palsy (US-CP), 31–2, 34
- United Kingdom (UK)
- neonatal intensive care study, 10
 - preterm birth, 2–3, 4
- United States (USA)
- early intervention national program, 245
 - ethnicity, 3, 225
 - preterm birth rates, 1, 2
 - SES effects, 4, 225
 - The Abecedarian Project, 230
 - The Infant Health and Development Program (IHDP), 231–2
 - The Kauai Longitudinal Study, 230–1
 - university education, of VPT/VLBW adults, 214
- Ureaplasma urealyticum* infection, 3
- uterus, mechanical stress, 7
- vascular mechanisms, preterm birth, 6
- vascular responses to neural activation, 79–80
- VBM, *See* voxel-based morphometry (VBM)
- ventilator use, *See* assisted ventilation
- ventricles, on CUS, 40, 41
- ventricular dilatation,
- posthemorrhagic, 21
- ventricular enlargement
- language function and, 180
 - prognostic MRI, 49
 - quantitative MRI, 55–6
 - VLBW adults, 68
- ventriculomegaly
- CT scan findings, 42, 43
 - neonatal morbidities, 145–6
 - VBM limitations, 101
- verbal fluency
- corpus callosum in, 61, 71, 204
 - deficits, 202, 204
 - development, 176
 - of adolescents, 178, 179, 202
 - SES and, 228
- verbal fluency processing
- functional neuroimaging, 85–8
 - letter trials, 85
- verbal IQ scores
- and fluency, 61
 - comprehension, 84
 - MRI in childhood, 60–1
- vermis, cerebellar, 56
- very low birth weight (VLBW)
- adult MRI data, 68
 - adult personality style, 72
 - CP risk, 30, 31, 32–3
 - long-term outcomes, 251
 - neonatal morbidities, 145–9
 - psychological distress in adulthood, 128–9
 - SGA compared, 107
- very preterm (VPT) infants, 1, 2
- adolescence to adulthood, 71
 - adult MRI data, 69
 - infections and, 3
 - long-term outcomes, 251
 - neurodevelopment, 70–3
 - rates, 2
 - reduced mortality rates, 8
- violent suicide attempts, 129
- viral infections, preterm birth and, 3–4
- visual expectancy task, 190
- visual function development, 80
- visual impairment, *See also* retinopathy of prematurity (ROP)
- changes over time, 120
 - CP and, 31, 32
 - MRI findings, 62–3
 - oxygen therapy, 17
 - PVL detection, 36
- Visual Motor Integration (VMI) test, 62
- visual paired-associated learning, 89
- visual recognition memory, 188
- visual system/visual function, PVL in, 35
- visual tests
- functional neuroimaging, 85
 - MRI findings, 62
 - visuo-perceptual function
 - MRI in childhood, 61–2
 - visual impairment and, 62
- visuospatial difficulties
- coordination and, 61
 - working memory deficits, 202
- visuospatial sketchpad, 185
- visuospatial working memory, 185–6, 190
- vitamin C lack, 5
- vitamin E, IVH prevention, 22

- VLBW, *See* very low birth weight (VLBW)
- volumetric brain studies, 204
- volumetric measurements, 45
- voxel-based morphometry (VBM)
- adult MRI data, 68, 69
 - brain developmental changes, 120
 - DTI analysis, 101, 107
 - on FA template, 102
 - VLBW and SGA compared, 107
- VPT, *See* very preterm (VPT) infants
- water
- brain tissue change over time, 102
 - brain tissue in the infant, 50
 - diffusion in DTI, 97–8
 - diffusion in DWI, 44
 - signal from H atoms in MRI, 43, 44
- Weschler Objective Numeric Dimensions test, 62
- white matter
- age-related changes in children, 181
 - anisotropic diffusion, 97
 - cell damage, 103
 - development, 71
 - cognitive function and, 108
 - DTI and, 102–7
 - microstructural abnormalities, 107
 - diffusion changes, 48–9, 98
 - environment enrichment and, 72–3
 - fractional anisotropy, 97–8
 - gender and volume, 55
 - heterogeneity, 98, 102–3, 104
 - myelinated, 44–5
 - plasticity, 107–8
 - regional cerebral blood flow, 79
 - resilience and, 108–9
 - spatial mapping of, 59
 - studies needed, 182
 - tract anatomy, 98–101
 - volume changes, 49, 54–5, 71–2, 257
 - vulnerability, 73
- white matter lesions, 65, 253
- cognitive delay, 36, 253
 - congenital heart disease, 64–5
 - CP and, 40, 59–60
 - CUS findings, 40
 - DEHSI, 48
 - executive functioning, 198–9, 203
 - in adolescence, 104–7
 - long-term findings, 58
 - mechanisms, 22–3
 - MRI
 - in childhood, 59–60
 - neonatal morbidities, 145–6
 - parenchymal lesions, 48
 - pathology detection, 103
 - pathology missed, 97, 103
 - periventricular damage, 21
 - prognostic, 48
 - qualitative, 58
 - Williams syndrome, 46–7
 - Woodcock–Johnson Tests of Achievement, 212
 - working memory, 185
 - changes over time, 202
 - deficits, 186, 190, 255
 - executive functioning, 195, 198–9
 - fMRI studies, 89–90
 - neural development, 185–7
 - neuroanatomical factor, 203–4
 - of adolescents, 179, 202
- X-ray beam, in CT, 42
- young adults
- educational outcomes, 214
 - executive function, 201–3
 - functional neuroimaging, 85–90
 - learning difficulties, 219
- young children,
- cognitive development, 141–2
 - neonatal care effects, 6