

Cambridge University Press

978-0-521-87375-8 - Clinical Magnetoencephalography and Magnetic Source Imaging

Andrew C. Papanicolaou

Index

[More information](#)

Index

- abnormal MEG waveforms
 - abnormal slow waves 75, 76
 - abnormal wave patterns 75–77
 - definition of MEG spikes and patterns 77–78, 79
 - epilepsy and MEG 78–81
 - epileptiform discharges 75–77
 - focal delta waves 75, 76
 - ictal discharges 81–82
 - polyspikes 75–78, 79
 - sharp waves 75–78, 79
 - spike-and-wave complexes 75–77
 - spikes 75–78, 79
- activation (function-specific brain activities) 4
- age-dependent changes in cognition, potential applications of MEG 174
- alpha waves 72
- Alzheimer's disease, potential applications of MEG 175–176
- arachnoidal cysts 88
- artifacts in normal spontaneous MEG
 - cardiac artifacts 63–65
 - dental implants 65–67
 - environmental noise 67
 - eye blinks 63, 64
 - eye movements 63
 - ferromagnetic dental implants 65–67
 - magnetized EEG electrodes 66, 67
 - muscle (EMG) activity 63, 64
 - respiratory artifacts 65
- Asperger's syndrome *see* autism spectrum disorder
- astrocytoma 88
- attention deficit hyperactivity disorder (ADHD), potential applications of MEG 181–182
- auditory evoked magnetic fields (AEFs)
 - aberrant AEFs 136–137
 - advantages over auditory evoked potentials (AEPs) 134
 - clinical applications 134
 - normative AEF data 136
 - referral questions that may be addressed 137
 - use in presurgical planning 137
- auditory evoked magnetic fields (AEFs)
 - recording 134
 - AEF sources 134–135
 - recording parameters 135
 - stimulation 135
- auditory evoked potentials (AEPs) 134
- autism spectrum disorder (ASD), potential applications of MEG 180–181
- axial gradiometers 19–21, 22
- axonal currents 7
 - contribution to magnetic flux 11–12
- baseline length, effects on magnetic flux recording 20
- beamformers
 - adaptive 44–45
 - advantages 45–46
 - characteristics 44
 - limitations 46
 - nonadaptive 44–45
 - relation to minimum-norm approaches 45–46
 - spatial filter functions 44
 - synthetic aperture magnetometry (SAM) 44–45
 - Tikhonov regularization 45
- benign epileptic transients of sleep (BETS) 70–71
- beta waves 72, 73
- bioelectromagnetic inversion 26
- biological magnetic fields, strength of 13, 15
- Biot–Savart Law 23

- boundary element models (BEMs) of the head 31–32
- brain activity
 - activation (function-specific activities) 4
 - baseline-activity profile 4
 - deviations from baseline activity 4
- brain mechanisms, identifying components 111–112
- brain mechanisms for language, detection of reorganization 154–156
- brain tumors
 - and lesional epilepsies 88
 - MEG spike localizations 88
 - potential applications of MEG 188
- Broca’s area, location using LRFs 145, 152–154
- cardiac artifacts in MEG 63–65
- cavernoma 88–89
- cavernoma with bleeding 88
- cerebral ischemia, potential applications of MEG 188–189
- cerebrovascular disease, MEG detection of functional reorganization 190–191
- clinical use of MEG/MSI
 - demagnetization of the subject 47–48
 - elimination of sources of noise 47–48
 - head positioning and fixing 47
 - integration of MEG/MSI and MRI images 48–50
 - length of the recording session 48
 - MEG analysis software 48, 49
 - monitoring the subject’s state of alertness 48
 - use in surgical neuronavigation systems 49–50
 - use of whole-head systems 47
 - see also* MEG recordings of spontaneous brain activity
- coherence analysis of MEG or EEG data 157–158
- coordinate system to define MEG measurement space 24–25
- corpus callostomy in children, role of MEG evaluation 98–100
- cortical malformations
 - and lesional epilepsies 89
 - MEG spike localizations 89
- delta waves 74
 - focal 75, 76
- dendritic currents 7
 - contribution to magnetic flux 11–12
- dental implants, artifacts in MEG 65–67
- depression, diagnostic potential of MEG 184
- dewar housing for magnetometers 13–16
- dipolar distributions 17–18, 19
 - extrema 18, 19
- dipole fitting *see* discrete source models
- direct transfer function (DTF) 158
- discrete source models
 - applications 35
 - characteristics 34
 - comparison with distributed source models 34–35
 - dipole fitting 34–35
 - global error minimisation 34
 - local estimator approach 34–35
 - multiple ECD (equivalent current dipole) model 34, 37–38, 39
 - reliability of estimated sources 38–39
 - single ECD (equivalent current dipole) model 34, 35–37, 38–39
 - spatial aliasing problem 39
 - spatial undersampling problem 39
 - spatiotemporal dipole modeling 37–38
- distributed source models
 - characteristics 40
 - comparison with discrete source models 34–35
 - dynamic statistical parametric mapping (dSPM) 43
 - EPIFOCUS algorithm 43
 - ill-posed (underdetermined) nature 40
 - lead-field normalization 41–42
 - LORETA method 42
 - magnetic field tomography (MFT) 43
 - minimum current estimates (MCE) 43
 - minimum-norm estimates (MNE) 43
 - minimum-norm models 40–43
 - nonlinear distributed source methods 43
 - regularization methods for minimum-norm models 41–43
 - sLORETA method 43
- drugs, use to elicit abnormal activity 61
- dynamic statistical parametric mapping (dSPM) 43
- dyslexia, potential applications of MEG 178–179
- ECD (equivalent current dipole), models *see* discrete source models

Cambridge University Press

978-0-521-87375-8 - Clinical Magnetoencephalography and Magnetic Source Imaging

Andrew C. Papanicolaou

Index

[More information](#)**202 Index**

- electroencephalogram (EEG) 3
 - measurement of volume currents 8–9
 - simultaneous MEG/EEG recordings 60
- EMFs *see* evoked magnetic fields
- environmental noise, artifacts in MEG 67
- EPIFOCUS algorithm 43
- epilepsy and MEG 78–81
 - ictal discharges 81–82
- epilepsy MEG investigations
 - extratemporal lobe epilepsies (ETLEs) 90–91
 - nonlesional temporal lobe epilepsies (TLEs) 90
 - see also* lesional epilepsy MEG investigations
- epilepsy surgery
 - description of the epileptogenic zone 83–84
 - epileptogenic zone localization with MEG 83–85
 - irritative networks 83–84
 - large network hypothesis 83–84
 - suitable patients 83
 - surgical outcome and MEG localizations 85
 - see also* pediatric epilepsy surgery
- epileptic activity provocation methods 85–86
- epileptic aphasia, potential applications of MEG 182
- epileptiform discharges 75–77
- epileptiform spike-and-wave discharge 4
- epileptogenic zone localization
 - alternatives to spike analysis 87
- epileptic activity provocation methods 85–86
- ictal MEG investigations 84
- interictal MEG compared with interictal EEG 84–85
- MEG localization 100
- planning for surgical procedures 83–84
- source localization techniques in adult epilepsy 86–87
- surgical outcome and MEG localizations 85
- temporal lobe epilepsy 87
- use of slow-wave activity 87
- equivalent current dipole (ECD) models *see* discrete source models
- event-related desynchronization (ERD) of MEG or EEG data 157
- event-related synchronization (ERS) of MEG or EEG data 157
- evoked magnetic field (EMF) data, future directions
 - coherence analysis 157–158
 - direct transfer function (DTF) 158
 - event-related desynchronization (ERD) 157
 - event-related synchronization (ERS) 157
 - spectral analysis of MEG data 157
 - temporal spectral evolution (TSE) 157
- evoked magnetic fields (EMFs)
 - clinical applications of MEG 111
 - generation by brain activity 111–112
 - identification of components of brain mechanisms 111–112
 - representation of brain activity 112
 - see also* auditory evoked magnetic fields (AEFs); language-related brain magnetic fields (LRFs); movement-related magnetic fields (MRFs); somatosensory evoked fields (SEFs); visual evoked magnetic fields (VEFs)
- evoked magnetic fields (EMFs) recording
 - artifact prevention 112
 - artifact removal 113–115
 - data storage 113
 - duration of sessions 112
 - general instructions for MEG activation studies 112
 - patient engagement in the activation task 112
 - removing “bad” channels 115–116
 - replications 113
 - signal averaging 117
 - signal filtering and electromagnetic-noise reduction 116–117
- evoked potentials (EPs) 3
- extratemporal lobe epilepsies (ETLEs), MEG investigations 90–91
- extrema of dipolar distributions 18, 19
- eye-blink artifacts in MEG 63, 64
- eye movement artifacts in MEG 63
- ferromagnetic dental implants, artifacts in MEG 65–67
- fetal MEG 178
- fidelity of MEG/MSI 4–6
- finite difference models (FDMs) of the head 32–33
- finite element models (FEMs) of the head 32–33
- first-order sensors 18–20
- focal cortical malformations 88
- forward problem in MSI 23–24, 27–28, 34
- frequency bands of normal spontaneous MEG 71

- functional magnetic resonance imaging (fMRI) 3–4
- functional reorganization following stroke 190–191
- future applications of clinical MEG
 - age-dependent changes in cognition 174
 - Alzheimer's disease 175–176
 - attention deficit hyperactivity disorder (ADHD) 181–182
 - autism spectrum disorder (ASD) 180–181
 - brain tumors 188
 - cerebral ischemia 188–189
 - cerebrovascular disease, functional reorganization 190–191
 - depression 184
 - dyslexia 178–179
 - epileptic aphasia 182
 - fetal MEG 178
 - functional reorganization following stroke 190–191
 - Lewy body dementia, differential diagnosis 176–177
 - mild cognitive impairment (MCI) 175–176
 - multiple sclerosis (MS) 187
 - neurodegenerative disorders 174–177
 - neurodevelopmental disorders 178–182
 - neurological disorders 186–189
 - normal aging process 174
 - obsessive-compulsive disorder (OCD) 185
 - Parkinson's disease 176–177, 187
 - post-traumatic stress disorder (PTSD) 185
 - psychiatric disorders 183–185
 - schizophrenia 183–184
 - stroke 188–189, 190–191
 - tinnitus 186–187
 - tremor syndromes 187
- ganglioglioma 88
- glioma 88
- global error minimization approach to source estimation 34
- gradiometers
 - magnetic flux recording 18–22
 - noise reduction 18–20, 21
- head models used in MEG
 - boundary element models (BEMs) 31–32
 - classes of head models 29, 30
 - finite difference models (FDMs) 32–33
 - finite element models (FEMs) 32–33
 - local spheres models 29–31
 - multisphere models 29–31
 - purpose 29
 - spherical models 29–31
- head positioning and fixing for MEG/MSI 47, 59
- hemimegalencephaly 88
- hemispheric dominance for language,
 - assessment using LRFs 145, 149–154
- hyperventilation, use to elicit abnormal activity 61
- hypnagogic hypersynchrony 71
- infinite mixture problem in MSI 26–27
- instrumentation problem in MSI 27
- inverse problem in MSI 23–24, 28
- invisible source problem in MSI 27
- K-complex waveforms 68
- lambda waves 69
- Landau-Kleffner syndrome 182
- language-related brain magnetic fields (LRFs)
 - aberrant LRFs 154–156
 - clinical applications 144
 - detection of reorganization of brain mechanisms 154–156
 - hemispheric dominance for language 145, 149–154
 - location of expressive language-specific cortex (Broca's area) 145, 152–154
 - location of receptive language-specific cortex (Wernicke's area) 145, 152–154
 - normative LRF data 149, 150
- language-related brain magnetic fields (LRFs)
 - recording 144–149
 - data processing 146–149
 - LRF sources 144–145
 - patient alertness and cooperation 146
 - recording parameters 146
 - stimulation 145
- lead-field normalization 41–42
- lesional epilepsy MEG investigations
 - brain tumors 88
 - cortical malformations 89
 - multiple lesions 88
 - pathologies associated with ictal-onset regions 88
 - presurgical localization and evaluation of lesions 88
 - vascular malformations 88–89

204 Index

- Lewy body dementia, differential diagnosis using MEG/MSI 176–177
- lobectomy candidates (paediatric), MEG evaluation 93–98, 125
- local estimator approach to source estimation 34–35
- local spheres models of the head 29–31
- LORETA (low resolution electromagnetic tomography) method 42
- magnetic field
 - components of 18–20
 - proportional dissipation 10–11
- magnetic field strength 10–11
 - biological magnetic fields 13, 15
- magnetic field tomography (MFT) 43
- magnetic flux
 - axonal current contribution 11–12
 - dendritic current contribution 11–12
 - direction relative to the current 9–10
 - right-hand rule 9–10
 - sources of 3–4, 8, 11–12
 - surface distribution 9–11
 - synaptic current contribution 11–12
- magnetic flux density 10–11
- magnetic flux recording
 - amplification of the induced currents 14–15
 - axial gradiometers 19–21, 22
 - components of the magnetic field 18–20
 - dewar housing for magnetometers 13–16
 - dipolar distributions 17–18, 19
 - effects of baseline length 20
 - extrema of dipolar distributions 18, 19
 - factors affecting recorded flux distribution 13, 14, 16–18, 19
 - first-order sensors 18–20
 - gradiometer sensors 18–22
 - magnetically shielded room 16
 - magnetometer construction and housing 13–16
 - magnetometer sensors 13, 14
 - noise reduction by gradiometers 18–20, 21
 - planar gradiometers 19–22
 - shape of magnetic flux lines at the head surface 13, 14
 - strength of biological magnetic fields 13, 15
 - superconducting magnetometer wires 13–14, 15
 - superconductive quantum interference devices (SQUIDS) 14–15
- magnetic permeability of tissues 11
- magnetic resonance imaging *see* MRI
- magnetic signals from neuronal signaling events 3–4
- magnetic source imaging *see* MSI
- magnetically shielded room for magnetic flux recording 16
- magnetized EEG electrodes, artifacts in MEG 66, 67
- magnetoencephalography *see* MEG
- magnetometers
 - construction and housing 13–16
 - magnetic flux recording 13, 14
- MEG (magnetoencephalography)
 - activation (function-specific activities) 4
 - baseline-activity profile 4
 - chronic deviations from baseline activity 4
 - definition 3
 - descriptions of the MEG/MSI process 26
 - fidelity of functional images 4–6
 - function-specific activities (activation) 4
 - nature of magnetic signals 3–4
 - neuronal signaling events 3–4
 - noninvasive nature 3–4
 - phasic deviations from baseline activity 4
 - reliability of results 4–5
 - source of magnetic signals 3–4
 - spatial resolution 5
 - spontaneous deviations from baseline activity 4
 - temporal resolution 5–6
 - uses 3
 - validity of results 5–6
- MEG analysis software 48, 49
- MEG dipole cluster, role in paediatric preoperative evaluation 93–98, 99
- MEG recordings of spontaneous brain activity
 - elicitation of abnormal activity via drugs 61
 - elicitation of abnormal activity via hyperventilation 61
 - head positioning 59
 - ISACM guidelines 59
 - recording during sleep 61
 - recording during the resting state 60–61
 - recording time 60
 - sampling rate 60
 - simultaneous MEG/EEG recordings 60
 - use of sedation 62
 - see also* clinical use of MEG/MSI
- mild cognitive impairment (MCI), potential applications of MEG 175–176

- minimum current estimates (MCE) 43
- minimum-norm estimates (MNE) 43
- minimum-norm models 40–43
 - as nonadaptive beamformers 45
- motor evoked fields (MEFs) *see*
 - movement-related magnetic fields (MRFs)
- movement-related magnetic fields (MRFs)
 - clinical applications 128
 - indications for MEG/MSI motor mapping 132–133
 - normative MRF data 130–132
 - referral questions that may be addressed 132–133
- movement-related magnetic fields (MRFs)
 - recording 128
 - elicitation 128–129
 - eye- and body-movement artifacts 130
 - recording parameters 129–130
 - triggering 129
- MRI (magnetic resonance imaging)
 - functional magnetic resonance imaging (fMRI) 3–4
 - integration with MEG/MSI 24–25, 48–50
 - nonlesional *see* pediatric epilepsy surgery (nonlesional MRI)
- MRI focal lesions, pediatric epilepsy surgery 92
- MSI (magnetic source imaging)
 - Biot–Savart Law 23
 - coordinate system defining the space of MEG measurements 24–25
 - coregistration of reference points 24–25
 - coregistration with structural MRI 24–25
 - definition 3
 - descriptions of the MEG/MSI process 26
 - estimation of attributes of the source 23
 - solution of the forward problem 23–24
 - solution of the inverse problem 23–24
 - use of fiducial points on the head surface 24–25
- MSI problems
 - forward problem 23–24, 27–28, 34
 - infinite mixture problem 26–27
 - inherent uncertainty in the process 26–27
 - instrumentation problem 27
 - inverse problem 23–24, 28
 - invisible source problem 27
- mu waves 69–70
- multiple ECD (equivalent current dipole)
 - model 34, 37–38, 39
- multiple sclerosis (MS), potential applications of MEG 187
- multisphere models of the head 29–31
- muscle (EMG) activity artifacts in MEG 63, 64
- neurocytoma 88
- neurodegenerative disorders, potential applications of MEG 174–177
- neurodevelopmental disorders, potential applications of MEG 178–182
- neurological disorders, potential applications of MEG 186–189
- neuronal signaling
 - abnormal deviations in activity 7–9
 - creation of electrical currents 7
 - primary (source) currents 7–9, 11–12
 - sources of magnetic flux 3–4, 8, 11–12
 - volume currents 8–9, 11
- noise reduction by gradiometers 18–20, 21
- nonadaptive beamformers, minimum-norm approaches as 45
- noninvasive nature of MEG/MSI 3–4
- nonlesional temporal lobe epilepsies (TLEs), MEG investigations 90
- nonlinear distributed source methods 43
- normal aging process, potential applications of MEG 174
- normal MEG waveforms 68–71
 - alpha waves 72
 - benign epileptic transients of sleep (BETS) 70–71
 - beta waves 72, 73
 - delta waves 74
 - frequency bands of normal spontaneous MEG 71
 - hypnagogic hypersynchrony 71
 - K-complex 68
 - lambda waves 69
 - mu waves 69–70
 - positive occipital sharp transients of sleep (POSTS) 69
 - sleep spindles 69, 70
 - slow waves 73, 74
 - theta waves 73, 74
 - vertex waves (V-waves) 68, 69
- obsessive-compulsive disorder (OCD), diagnostic potential of MEG 185
- oligodendro glioma 88

206 Index

- Parkinson's disease
 - differential diagnosis using MEG/MSI 176–177
 - use of MEG in tremor syndromes 187
- pediatric epilepsy surgery
 - candidates for evaluation 92
 - definition of nonlesional MRI 92–93
 - differences from adult epilepsy surgery 92
 - goals 92
 - outcomes with nonlesional MRI 92–93
 - with focal lesion on MRI 92
- pediatric epilepsy surgery (nonlesional MRI)
 - definition of nonlesional MRI 92–93
 - MEG evaluation of corpus callostomy candidates 98–100
 - MEG evaluation of lobectomy candidates 93–98, 125
 - MEG evaluation of topectomy candidates 93–98, 99
 - MEG localization of epileptogenic zones 100
- nonlesional refractory status epilepticus 100
- role of the MEG dipole cluster in
 - preoperative evaluation 93–98, 99
 - use of MEG in epilepsy surgery evaluation 100–101
- planar gradiometers 19–22
- polymicrogyria 88
- polyspikes (MEG waveforms) 75–78, 79
- positive occipital sharp transients of sleep (POSTS) 69
- positron-emission tomography (PET) 3–4
- post-traumatic stress disorder (PTSD),
 - diagnostic potential of MEG 185
- primary (source) currents 7–9, 11–12
- proportional dissipation of magnetic fields 10–11
- psychiatric disorders, diagnostic potential of MEG 183–185
- recording sessions for MEG/MSI, length of 48, 60
- reliability of MEG/MSI 4–5
- respiratory artifacts in MEG 65
- resting state, MEG recordings during 60–61
- SAM (synthetic aperture magnetometry) 44–45
- sampling rate for MEG recordings 60
- schizophrenia, diagnostic potential of MEG 183–184
- sedation
 - effects on beta waves 72
 - use during MEG recordings 62
- sharp waves (MEG waveforms) 75–78, 79
- single ECD (equivalent current dipole) model 34, 35–37, 38–39
- sleep, MEG recordings during 61
- sleep spindles 69, 70
- sLORETA method 43
- slow waves 73, 74
 - abnormal forms 75, 76
- somatosensory evoked fields (SEFs)
 - aberrant SEFs 123–124
 - advantages over somatosensory evoked potentials (SEPs) 118
 - clinical applications 118–119
 - indications for MEG/MSI somatosensory mapping 124–127
 - normative SEF data 120–123
 - referral questions that may be addressed 124–127
- somatosensory evoked fields (SEFs) recording 119–120
 - recording parameters 120
 - stimulation 119–120
- somatosensory evoked potentials (SEPs) 118
- source analysis 26
- source estimation 26
- source imaging 26
- source localization 26
- source models *see* discrete source models; distributed source models
- spatial aliasing problem, discrete source models 39
- spatial resolution of an image, definitions 5
- spatial undersampling problem, discrete source models 39
- spatiotemporal dipole modeling 37–38
- spectral analysis of MEG data 157
- spherical models of the head 29–31
- spike-and-wave complexes 75–77
- spikes (MEG waveforms) 75–78, 79
- stroke
 - MEG detection of functional reorganization 190–191
 - potential applications of MEG 188–189, 190–191
- structural MRI, coregistration with MSI/MEG-derived active source 24–25

Sturge–Weber syndrome 88	validity of MEG/MSI 5–6
superconducting magnetometer wires 13–14, 15	vascular malformations and lesional epilepsies 88–89
superconductive quantum interference devices (SQUIDS) 14–15	MEG spike localization 88–89
surgery <i>see</i> epilepsy surgery; pediatric epilepsy surgery	vertex waves (V-waves) 68, 69
surgical neuronavigation systems, use of MEG/MSI 49–50	visual evoked magnetic fields (VEFs) aberrant VEFs 141
synaptic currents 7	advantages over visual evoked potentials (VEPs) 138
contribution to magnetic flux 11–12	clinical applications 138
synthetic aperture magnetometry (SAM) 44–45	normative VEF data 141
	referral questions that may be addressed 141–143
temporal resolution of MEG 5–6	use in presurgical planning 141–143
temporal spectral evolution (TSE) analysis of MEG data 157	visual evoked magnetic fields (VEFs) recording 138–141
theta waves 73, 74	recording parameters 139–141
Tikhonov regularization 45	stimulation 139
tinnitus, potential applications of MEG 186–187	VEF sources 138
tissues, magnetic permeability of 11	visual evoked potentials (VEPs) 138
topectomy candidates (pediatric), MEG evaluation 93–98, 99	volume currents 8–9, 11
tremor syndromes, potential applications of MEG 187	
tuberous sclerosis 88	waveforms <i>see</i> abnormal MEG waveforms; normal MEG waveforms
	Wernicke’s area, location using LRFs 145, 152–154
uncertainty inherent in the MEG/MSI process 26–27	whole-head systems, clinical use of MEG/MSI 47
unilateral Ammon’s horn sclerosis 88	