

# Index

- Abnormal Involuntary Movement Scale (AIMS), 203
- N*-acetylcysteine, 163
- adolescent OCD, 100–2
- adolescent schizophrenia, 98–100
- clinical features of
    - obsessive–compulsive symptoms, 103–7
  - comorbidities, 107–8
  - differential diagnosis, 108–11
  - OCD-related comorbidities, 107–8
  - prevalence of
    - obsessive–compulsive symptoms, 102–3
  - tic disorders, 107–8
  - treatment, 111–13
- adolescents
- rate of onset of mental health disorders, 98
  - schizoid personality disorder, 109
  - schizotypal personality disorder, 109
- age of onset
- and severity of illness, 39
  - obsessive–compulsive disorder, 7
  - schizophrenia, 6
  - schizophrenia vs. OCD, 39–40
  - schizophrenic vs. obsessive–compulsive symptoms in schizo-obsessive patients, 40–3
- amisulpiride, 162, 197–9
- anterior cingulate cortex, 12
- antiglutamatergic agents, 204–5
- antipsychotic agents
- motor side effects, 203–4
  - OCD therapies, 16–17
  - schizophrenia therapies, 15–16
- antipsychotic-induced
- obsessive–compulsive symptoms, 213–14
  - diagnosis, 216–17
  - management, 217–21
  - mechanism, 214–16
- anxiety disorders, 123
- anxiety symptoms in schizophrenia, 4
- aripiprazole, 130, 162, 196–7, 213, 219
- Asperger's syndrome
- differential diagnosis, 159
- attention deficit hyperactivity disorder (ADHD), 89, 127
- atypical antipsychotic agents
- and obsessive–compulsive symptoms, 117
  - parkinsonian symptoms caused by, 117
  - See also* antipsychotic-induced obsessive–compulsive symptoms.
- autistic disorders
- differential diagnosis, 159
- autogenous obsessions, 152
- avoidant personality disorder
- differential diagnosis, 159–60
- bipolar disorder
- in children, 109
- bizarreness of delusions, 174–5
- Bleuler, Eugen, 2–4, 22, 29, 151
- body dysmorphic disorder (BDD), 123
- appearance preoccupations, 133
  - as a distinct disorder, 134
  - as related to OCD spectrum, 134–6
  - avoidance behavior, 133
  - case vignette, 133
  - comorbidities, 134
  - course of illness, 133
  - delusions, 133–4
  - development of DSM criteria, 132
  - distinction from schizophrenia, 134
  - DSM-IV-TR criteria, 132
  - in schizo-obsessive disorder, 139–40
  - in schizophrenia, 136–9
  - insight in patients, 133–4
  - neurocognitive impairments, 134
  - prevalence, 132–3
  - risk behaviors, 133
  - safety behaviors, 133
  - See also* dysmorphophobia.
- body dysmorphic disorder (BDD) in schizophrenia
- diagnosis and treatment, 141–4
- Bonn Scale for Assessment of Basic Symptoms (BSABS), 85
- borderline
- meanings of the term, 150–1
- borderline personality disorder, 150–1
- differential diagnosis, 159–60
- borderline schizophrenia, 150
- Brief Psychiatric Rating Scale, 195
- brief psychotic disorder, 6
- broadcasting of thoughts, 2, 5
- Broca's area, 11
- bromosis delusions, 132
- Brown Assessment of Beliefs Scale (BABS), 59–62, 104, 133, 171
- Brown Longitudinal Obsessive–Compulsive Disorder Study, 48
- buspirone, 161
- Calgary Depression Rating Scale, 204
- candidate genes
- obsessive–compulsive disorder (OCD), 8
  - schizo-obsessive disorder, 189–90
  - schizophrenia, 7
- cannabis, 8
- captopril, 115
- case reports
- adolescent
    - obsessive–compulsive symptoms with schizophrenia, 106–7
  - adolescent OCD preceding schizophrenia, 106

- case reports (cont.)  
 aggressive and sexual  
 obsessions, 78  
 amisulpiride therapy in  
 schizo-obsessive disorder,  
 198–9  
 aripiprazole therapy in  
 schizophrenia and OCD,  
 196–7  
 atypical obsessive–compulsive  
 neurosis, 176  
 BDD in a schizo-obsessive  
 patient, 140  
 BDD in a schizophrenia  
 patient, 140  
 bizarre content of obsessions  
 and compulsions, 174–5  
 body dysmorphic disorder  
 (BDD), 133  
 by Alfred Gordon, 23  
 by Erwin Stengel, 25  
 by Hoch and Polatin, 28  
 by Ismond Rosen, 27  
 by Sigmund Freud, 3  
 childhood OCD presenting  
 as schizophrenia,  
 110–11  
 cleaning compulsion in  
 schizophrenia, 70  
 clozapine therapy for  
 obsessions and  
 compulsions, 199–200  
 clozapine therapy in  
 treatment-resistant  
 schizoaffective disorder,  
 200  
 clozapine therapy in  
 treatment-resistant  
 schizophrenia, 199–200  
 clozapine-induced  
 obsessive–compulsive  
 symptoms, 217–19  
 compulsions related to  
 hallucinations and  
 delusions, 78  
 contamination obsession in  
 schizophrenia, 70  
 delusions caused by  
 interpretation of  
 obsessions, 23  
 delusions unrelated to  
 obsessions, 23  
 development of obsessive  
 delusions, 73–4  
 differential diagnosis  
 challenge, 158–9  
 dysmorphophobia, 133  
 dysmorphophobia and  
 schizophrenia, 137–8  
 early-onset schizophrenia  
 with obsessive–compulsive  
 symptoms, 103  
 electroconvulsive therapy  
 (ECT), 206–7  
 exposure and response  
 prevention (ERP) therapy  
 for OCD, 205–6  
 forbidden thoughts and  
 counting compulsion,  
 69–70  
 genetic risk deterioration  
 state, 92–3  
 high-risk prodromal  
 schizophrenia with  
 obsessive–compulsive  
 symptoms, 91–2  
 late-onset OCD in older  
 schizophrenia patient, 115  
 late-onset OCD related to  
 brain infarct, 115–16  
 late-onset OCD related to  
 brain lesion, 115  
 obsessional neurosis  
 (Freud), 3  
 obsessional neurosis to  
 paranoid schizophrenia, 25  
 obsessions abate when  
 delusions develop, 23  
 obsessions and formal  
 thought disorders, 78  
 obsessions and paranoid  
 schizophrenia, 44–6  
 obsessions and schizophrenic  
 disorder, 46  
 obsessions transformed into  
 delusions, 23  
 obsessive delusions, 75–6  
 obsessive hallucinations,  
 76–7  
 obsessive preoccupation  
 with a schizophrenic  
 delusion, 74  
 obsessive psychosis, 175–6  
 obsessive thoughts and  
 compulsive acts, 25  
 obsessive–compulsive  
 phenomena in  
 schizophrenia, 27  
 obsessive–compulsive  
 symptoms in elderly  
 schizophrenia patient,  
 114–15  
 obsessive–compulsive  
 symptoms with attenuated  
 positive symptoms, 91–2  
 OCD and brief intermittent  
 psychotic state, 93–4  
 OCD and schizophrenia,  
 69–70  
 OCD with comorbid  
 schizotypal personality  
 disorder, 163–4  
 OCD with paranoid  
 schizophrenia, 49  
 OCD with poor insight, 172–3  
 OCD with schizophrenia,  
 49–50  
 OCD with schizophrenic  
 disorder, 49  
 olanzapine monotherapy in a  
 schizophrenia patient,  
 195–6  
 olfactory reference syndrome  
 and schizophrenia, 137  
 paranoid schizophrenia  
 with aggressive  
 obsession, 46  
 primary dysmorphophobia,  
 138  
 pseudoneurotic  
 schizophrenia, 28  
 pseudo-hallucinations in  
 OCD, 177–8  
 psychotic deterioration of  
 obsessive symptoms,  
 176–7  
 ritualistic compulsive  
 behavior, 78  
 schizophrenia spectrum  
 disorders in an OCD  
 patient, 158–9  
 schizophrenia with a chronic  
 tic disorder, 131  
 schizophrenia with a chronic  
 tic disorder and OCD,  
 131–2  
 schizophrenia with comorbid  
 Tourette's syndrome and  
 OCD, 131  
 schizophrenia with good  
 insight into OCD, 62–3  
 schizophrenia with poor  
 insight into OCD, 63  
 schizophrenic disorder with  
 OCD, 46–7  
 schizotypal personality  
 disorder in OCD, 92–3,  
 155–6

- social skills training for  
 schizotypal OCD patient,  
 165
- SRI therapy for OCD in  
 schizophrenia, 202
- symmetry and repeating in  
 schizophrenia, 70–1
- symptom presentation in  
 late-onset OCD, 115–16
- therapeutic use of  
 glutamatergic drugs, 163
- treating clozapine-induced  
 obsessive-compulsive  
 symptoms, 218–19
- catatonia, 1
- Catatonia Rating Scale  
 (CRS), 203
- Center for the Assessment  
 and Prevention of  
 Prodromal States  
 (CAPPS) (UCLA), 87
- Chestnut Lodge Follow-up  
 Study, 30–1
- childhood-onset OCD, 8, 100–2
- childhood-onset schizophrenia,  
 98–100
- clinical features of  
 obsessive-compulsive  
 symptoms, 103–7
- comorbidities, 107–8
- differential diagnosis, 108–11
- OCD-related comorbidities,  
 107–8
- prevalence of  
 obsessive-compulsive  
 symptoms, 102–3
- tic disorders, 107–8
- treatment, 111–13
- children  
 bipolar disorder, 109
- delusions, 108–9
- delusions (non-psychotic),  
 108
- depression, 108–9
- differential diagnosis for  
 OCD, 109–11
- hallucinations, 108–9
- hallucinations  
 (non-psychotic), 108
- major depressive disorder,  
 108–9
- multidimensionally impaired  
 disorder (MDI), 109
- personality disorders, 109
- pervasive developmental  
 disorder (PDD), 109
- schizoid personality  
 disorder, 109
- schizotypal personality  
 disorder, 109
- children's Yale-Brown  
 obsessive-compulsive  
 Scale (CY-BOCS), 100
- Clinical Global Impression,  
 205
- clomipramine, 202
- clonazepam, 161
- clonidine, 130
- clozapine, 12, 130, 163–4,  
 199–202, 213–14
- cognitive-behavior therapy  
 (CBT), 144, 205–6
- cognitive dysfunction  
 schizophrenia compared to  
 OCD, 9–10
- Colorado Childhood-Onset  
 Schizophrenia Research  
 Program, 102
- Composite International  
 Diagnostic Interview  
 (CIDI), 90
- Comprehensive Assessment of  
 At-Risk Mental States  
 (CAARMS), 89
- compulsions  
 definition, 5
- features of, 5
- COMT gene  
 role in schizo-obsessive  
 disorder, 189–90
- course of illness  
 obsessive-compulsive  
 symptoms in  
 schizophrenia, 48–51
- schizo-obsessive disorder,  
 44–51
- temporal onset of symptoms,  
 44–7
- D-cycloserine, 16, 163
- delusional body dysmorphic  
 disorder (BDD),  
 133–4
- delusional disorder, 6
- delusions  
 bizarreness of, 174–5
- distinction from obsessions,  
 2–3, 57
- distinction from poor insight  
 obsessions, x
- dysmorphophobia, 132
- in children, 108–9
- non-psychotic, 108
- obsessive delusions, 73–6
- delusions of bromosis, 132
- delusions of control, 2
- delusions of parasitosis, 132
- delusions of reference, 134
- dementia praecox, 2
- dementias, 116
- demographic  
 obsessive-compulsive  
 disorder, 7
- schizophrenia, 6
- depression  
 in children, 108–9
- depressive symptoms in  
 schizophrenia, 4
- Diagnostic and Statistical  
 Manual. *See* DSM
- differential diagnosis  
 early-onset schizophrenia,  
 108–11
- OCD and tic disorders, 108
- OCD in children, 109–11
- schizotypal personality  
 disorder in OCD, 157–60
- tic disorders in  
 schizophrenia, 128–9
- Tourette's syndrome and  
 OCD, 108
- Dimensions of Delusional  
 Experience scale, 59
- disability  
 major causes worldwide, 8
- disconnection syndrome  
 schizophrenia as, 11
- Discriminate Response Task  
 (DRT), 185
- dopamine  
 dysfunction in OCD, 15
- dysfunction in  
 schizophrenia, 11
- dorsolateral prefrontal cortex  
 (DLPFC), 12
- hypofrontality in  
 schizophrenia, 11
- drug-drug interactions  
 in elderly patients, 117–18
- ways to avoid, 118
- drug-induced motor side  
 effects, 203–4
- drug-induced obsessive-  
 compulsive symptoms  
*See* antipsychotic-induced  
 obsessive-compulsive  
 symptoms.
- DSM-I, 30

- DSM-II, 30  
DSM-III, 30–1  
DSM-IV, 31  
    criteria for OCD, 4  
    criteria for schizophrenia, 4  
    criteria for schizotypal  
        personality disorder, 151  
    insight specifications, 171  
DSM-IV-TR  
    criteria for body  
        dysmorphic disorder  
        (BDD), 132  
    definition of schizophrenia, 2  
dysmorphomania, 139  
dysmorphophobia, 132  
    *See also* body dysmorphic  
        disorder (BDD).
- early-onset OCD, 100–2  
    similarities with early-onset  
        schizophrenia, 101  
early-onset of mental health  
    disorders, 98  
early-onset schizophrenia,  
    98–100  
    clinical features of  
        obsessive–compulsive  
        symptoms, 103–7  
    comorbidities, 107–8  
    differential diagnosis,  
        108–11  
    OCD-related comorbidities,  
        107–8  
    prevalence of  
        obsessive–compulsive  
        symptoms, 102–3  
    similarities with early-onset  
        OCD, 101  
    tic disorders, 107–8  
    treatment, 111–13  
eating disorders, 123  
ego-dystonic obsessions, 175  
ego-syntonic obsessions, 175  
elderly patients  
    drug–drug interactions,  
        117–18  
elderly schizophrenia patients  
    avoiding drug–drug  
        interactions, 118  
    diagnosis of  
        obsessive–compulsive  
        symptoms, 116–18  
    drug–drug interactions,  
        117–18  
    implications of late-onset  
        OCD, 116–18  
    obsessive–compulsive  
        symptoms, 113–16  
    treatment of  
        obsessive–compulsive  
        symptoms, 116–18  
electroconvulsive therapy  
    (ECT), 206–7  
environmental factors  
    schizophrenia and OCD  
        compared, 7–9  
escitalopram, 93, 115, 201  
Esquirol, J.E., 58
- Family History Research  
    Diagnostic Criteria  
    (FH-RDC), 157, 187  
family inheritance  
    schizo-obsessive disorder,  
        224  
    schizotypal personality  
        disorder in OCD,  
        156–7  
family studies  
    schizo-obsessive disorder,  
        187–90  
Fenton, W.S., 30–1  
Fixity of Beliefs Questionnaire  
    (FBQ), 171  
fluvoxamine, 106,  
    161, 201  
*folie avec conscience* (insanity  
    with insight), 2  
*folie de doute* (insanity of  
    doubt), 2  
formal thought disorders  
    and obsessions, 77–9  
    in schizophrenia, 4  
Freud, Sigmund, 3  
frontocortical–striatal–thalamic  
    loops disturbance in  
        schizophrenia, 11
- GABA (gamma-aminobutyric  
    acid)  
    dysfunction in  
        schizophrenia, 11–12  
genetic factors  
    obsessive–compulsive  
        disorder (OCD), 8  
    schizo-obsessive disorder,  
        187–90  
    schizophrenia, 7  
    schizophrenia and OCD  
        compared, 7–9  
genome-wide association  
    studies, 7–8
- glutamate  
    dysfunction in OCD, 15  
    dysfunction in  
        schizophrenia, 11  
glutamate transporter gene  
    SLC1A1, 8  
glycine, 16  
Gordon, Alfred, 22–4
- hallucinations  
    in children, 108–9  
    non-psychotic, 108  
    obsessive hallucinations,  
        76–7  
haloperidol, 16, 129,  
    162, 214  
Hamilton Rating Scale for  
    Depression (HAM-D), 114  
hebephrenia, 1  
heritability  
    schizophrenia and OCD  
        compared, 7–9  
Hillside Akathisia Scale (HAS),  
    203  
historical perspective  
    obsessive–compulsive  
        disorder (OCD), 2–3  
    relation between  
        schizophrenia and  
        OCD, 3  
    schizophrenia, 1–2  
history of schizo-obsessive  
    concept, 22  
    contemporary studies, 30–1  
    work of Alfred Gordon, 22–4  
    work of Emil Kraepelin, 22  
    work of Erwin Stengel, 24–6  
    work of Eugen Bleuler, 22  
    work of Hoch and Polatin,  
        27–30  
    work of Ismond Rosen,  
        26–7  
Hoch, Paul, 27–30  
hypochondriacal syndromes  
    (monosymptomatic  
        type), 132  
hypochondriasis, 123  
hypofrontality in  
    schizophrenia, 11
- ICD (International  
    Classification of Diseases)  
    (WHO), 30  
ICD-10  
    definition of schizophrenia, 2  
    schizotypal disorder, 151

- imaging findings
  - schizophrenia compared to OCD, 14
- immune system
  - possible role in OCD, 8
- indecisiveness in OCD, 5
- insight
  - in OCD patients, 5
  - lack in schizophrenia patients, 5
  - obsessive-compulsive symptoms in schizophrenia, 57–64
  - See also* OCD with poor insight.
- insight assessment, 171–2
- Interview for the Retrospective Assessment of the Onset of Schizophrenia (IRAOS), 85
- Janet, Pierre, 3, 25, 134–5
- Jaspers, Karl, 2, 58–9, 76
- ketamine, 11
- Kraepelin, Emil, 1–3, 22, 72, 132, 134–5, 151
- lamotrigine, 204–5
- latent inhibition, 152
- late-onset OCD
  - implications of, 116–18
- lateral orbital cortex, 12
- Legrand du Salle, Henri, 3
- Lewis, A., 24
- lithium, 161
- magical thinking, 5, 100, 151–2
- major depressive disorder (MDD), 89, 123
  - in children, 108–9
- manneristic catatonia, 204
- Mayer-Gross, W., 2, 24
- McGlashan, T.H., 30–1
- memantine, 16, 163, 204
- mental health disorders
  - early-onset, 98
- metformin, 115
- migration
  - risk factor for schizophrenia, 8
- Mini Mental State Examination (MMSE), 114, 117
- mismatch negativity (MMN), 185
  - molecular genetic studies
    - schizo-obsessive disorder, 189–90
- monosymptomatic
  - hypochondriacal syndromes, 132
- Montgomery-Asberg Depression Scale, 123
- Morselli, E., 134–5
- motor symptoms in schizophrenia, 4
- multidimensionally impaired disorder (MDI), 109
- National Comorbidity Survey Replication study (US), 36–7, 98
- negative symptoms in schizophrenia, 4
- neoplasms (brain), 116
- Netherlands Mental Health Survey and Incidence Study (NEMESIS), 90
- neurobiology of schizo-obsessive disorder, 181
  - brain functional alterations, 181–2
  - brain structural alterations, 181–2
  - genetic basis for schizo-obsessive disorder, 187–90
  - neurological soft signs, 183–5
  - neurophysiological alterations, 185–6
  - neuropsychological dysfunction, 182–3
- Neurological Evaluation Scale (NES), 184
- neurophysiological evaluation, 185–6
- mismatch negativity (MMN), 185
  - P300 event-related potentials (ERP), 185
  - P50 auditory-evoked response, 185
  - pre-pulse inhibition (PPI), 185
- neurotransmitters
  - alterations in OCD, 14–15
  - alterations in schizophrenia, 11–12
- obsessional neurosis, 3
- obsessions
  - and formal thought disorders, 77–9
  - definition, 2–3, 5
  - distinction from delusions, 2–3, 57
- features of, 5
- obsessive delusions, 73–6
- obsessive hallucinations, 76–7
- obsessive psychosis, 175–8
- obsessive-compulsive disorder (OCD)
  - age of onset, 7, 39–40
  - alternative names for, 2
  - brain structural and functional alterations, 12–15
  - burden of disease, 8
  - candidate genes, 8
  - clinical features, 7
  - cognitive dysfunction, 9–10
  - comorbid personality disorders, 150
  - comorbid schizotypal personality disorder, 150
  - comorbidities, 123
  - course of illness, 7
  - demographic, 7
  - differential diagnosis in children, 109–11
  - dopamine dysfunction, 15
  - DSM-IV criteria, 4
  - early-onset, 100–2
  - features shared with schizophrenia, 1
  - genetic factors, 8
  - glutamate dysfunction, 15
  - heritability, 8
  - historical perspective, 2–3
  - imaging findings, 14
  - indecisiveness, 5
  - insight in patients, 5
  - neurotransmitter alterations, 14–15
  - orbitofrontal-subcortical circuit, 12–14
  - origin of the term, 2
  - overlapping features with schizophrenia, 3, 6
  - pathological doubt, 5
  - pathological slowness, 5
  - possible autoimmune mechanisms, 8
  - prevalence, 7
  - prognosis, 7
  - serotonin dysfunction, 15
  - signs and symptoms, 4–5
  - treatment, 16–17
  - twin studies, 8
  - with psychotic features, 175–8
- See also* OCD.

- obsessive-compulsive neurosis, 2
- obsessive-compulsive
  - personality disorder (OCPD), 187, 189
- obsessive-compulsive phenomena
  - in prodromal schizophrenia, 85
- obsessive-compulsive symptoms
  - caused by atypical antipsychotics, 117
  - high-risk prodromal schizophrenia subgroups, 91–4
  - overlap with schizophrenic symptoms, 57
- obsessive-compulsive symptoms in schizophrenia
  - diagnosis in prodromal schizophrenia, 94–5
  - diagnostic challenge, 57
  - dimensions of symptoms, 66–71
  - effect on schizophrenia symptom severity, 71–2
  - high-risk prodromal subgroups, 91–4
  - insight, 57–64
  - management in prodromal schizophrenia, 94–5
  - obsessions and formal thought disorders, 77–9
  - obsessive delusions, 73–6
  - obsessive hallucinations, 76–7
  - prevalence, 33–9
  - psychotic-related
    - obsessive-compulsive symptoms, 72–3
  - recommendations for identification, 79–80
  - typical symptom characteristics, 64–6
- OCD with poor insight, 5, 171
  - assessment of insight in OCD, 171–2
  - bizarreness of content, 174–5
  - clinical correlates, 172–4
  - diagnostic
    - recommendations, 178–9
  - insight categories, 171
  - management
    - recommendations, 178–9
- OCD-related rituals
  - distinction from schizophrenia-related movement disorders, 57
- OCD-schizophrenia, x
- OCD-spectrum of disorders, 123
- olanzapine, 92, 115, 161–2, 194–6, 213–14
- olfactory reference syndrome, 132
- orbitofrontal-subcortical circuit in OCD, 12–14
- Overvalued Ideas Scale (OVIS), 172
- P50 auditory evoked response, 185
- Padua Inventory, 87
- panic disorder, 123
- paranoia (Kraepelin), 132
- paranoid personality disorder
  - differential diagnosis, 159–60
- parental age
  - and risk of schizophrenia, 7
- Parkinson’s disease, 117
- parkinsonian symptoms
  - caused by atypical antipsychotics, 117
- pathological doubt in OCD, 5
- pathological slowness in OCD, 5
- pathophysiology
  - schizophrenia and OCD compared, 10–15
- pathoplastically veiled view of schizophrenia, 24
- Pediatric Autoimmune Neuropsychiatric Disorders Associated with Streptococcal infections (PANDAS), 8, 101
- perinatal insults
  - risk factor for schizophrenia, 7
- perphenazine, 107, 115
- personality disorders
  - and OCD, 150
  - differential diagnosis, 159–60
  - in children, 109
- pervasive developmental disorder (PDD), 109
  - differential diagnosis, 159
- phencyclidine (PCP), 11
- pimozide, 130, 161
- Polatin, Phillip, 27–30
- poor insight obsessions
  - distinction from delusions, x
- Positive and Negative Syndrome Scale, 201
- positive symptoms of schizophrenia, 4
- pre-pulse inhibition, 14, 185
- prevalence
  - obsessive-compulsive disorder, 7
  - obsessive-compulsive symptoms in schizophrenia, 33–9
  - schizophrenia, 6
  - schizophrenia in OCD patients, 37–9
- Prevention through Risk Identification, Management and Education (PRIME) Research Clinic (Yale University, USA), 89
- prodromal phase of schizophrenia
  - at-risk mental state concept, 85–7
  - clinical significance of obsessive-compulsive symptoms, 87–91
  - clinical staging model, 84
  - definition of prodrome, 84
  - diagnosis of obsessive-compulsive symptoms, 94–5
  - early prodromal phase, 84
  - features of, 85
  - high-risk prodromal subgroups, 85–7
  - late prodromal phase, 84
  - management of obsessive-compulsive symptoms, 94–5
  - obsessive-compulsive symptoms and high-risk subgroups, 91–4
  - occurrence of obsessive-compulsive phenomena, 85
  - overt psychotic phase, 84
  - premorbid phase, 84
  - prevalence of obsessive-compulsive symptoms, 87–91
  - rating instruments, 84–5
  - reasons for study, 84
  - risk of conversion to psychosis, 85–7
  - vulnerability to develop schizophrenia, 85–7



prodrome definition, 84 prognosis obsessive–compulsive disorder, 7 schizophrenia, 7 progressive supranuclear palsy, 116 pseudoneurotic schizophrenia, 27–30 psychasthenia, 3, 25 psychiatric classification systems, 123 hierarchical approach, 2 psychotic disorders differential diagnosis, 157–9 psychotic disorders not otherwise specified, 6 psychotic-related obsessive–compulsive symptoms in schizophrenia, 72–3  quetiapine, 162, 213  reactive obsessions, 152 religious melancholy, 2 riluzole, 16, 163, 204 risperidone, 16, 94, 106, 130, 161–2, 194–7, 213–14 rituals related to OCD distinction from schizophrenia-related movement disorders, 57 Rosen, Ismond, 26–7  Scale for the Assessment of Negative Symptoms (SANS), 102 Scale of Prodromal Symptoms (SOPS), 89 Scale to Assess Unawareness of Mental Disorder (SUMD), 60 Schedule for the Assessment of Negative Symptoms (SANS), 62, 114 Schedule for the Assessment of Positive Symptoms (SAPS), 62, 114 Schedule to Assess Mental Disorders (SUMD), 104 schizoaffective disorder distinctive clinical features, 224 schizoid personality disorder differential diagnosis, 159–60 in children and adolescents, 109  schizo-obsessive disorder age of onset of schizophrenic vs. obsessive–compulsive symptoms, 40–3 as obsessive–compulsive subgroup of schizophrenia, 225–6 candidate gene studies, 189–90 comorbidities, 123–5 course of illness, 44–51 diagnosis of comorbid BDD, 141–4 family studies, 187–90 future research goals, 226–8 genetic basis, 187–90 molecular genetic studies, 189–90 neurobiological underpinnings, 225 obsessive–compulsive symptoms in schizophrenia, 223–4 provisional diagnostic criteria, 225–6 related disorders in first-degree relatives, 224 shared features of schizophrenia and OCD, 223 treatment, 225 variations in terminology, x with body dysmorphic disorder (BDD), 139–40 schizo-OCD, x schizophrenia age of onset, 6, 39–40 and Tourette’s syndrome, 125–7 anxiety symptoms, 4 brain structural alterations, 11 burden of disease, 8 candidate genes, 7 clinical features, 6 cognitive dysfunction, 9–10 comorbidities, 123–5 course of illness, 6 current definitions, 2 demographic, 6 depressive symptoms, 4 diagnosis of comorbid BDD, 141–4 dopamine dysregulation, 11 DSM-IV criteria, 4 DSM-IV-TR definition, 2 environmental risk factors, 8  features shared with obsessive–compulsive disorder (OCD), 1 first-rank symptoms (Schneider), 2 formal thought disorders, 4 GABA dysfunction, 11–12 genetic factors, 7 glutamate dysfunction, 11 heritability, 7 historical perspective, 1–2 ICD-10 definition, 2 imaging findings, 14 lack of insight in patients, 5 motor symptoms, 4 negative symptoms, 4 neurotransmitter alterations, 11–12 occurrence in OCD patients, 37–9 origin of the term, 2 overlap with OCD, 3 overlapping features with OCD, 6 positive symptoms, 4 prevalence, 6 prognosis, 7 second-rank symptoms (Schneider), 2 serotonin dysfunction, 12 signs and symptoms, 3–4 suicide risk, 4 symptoms overlap with OCD symptoms, 57 treatment, 15–16 twin studies, 7 un-understandability of, 2 schizophrenia and tic disorders, 125–7 case reports, 130–2 case vignettes, 130–2 clinical characteristics of tic disorders, 127–8 differential diagnosis of tic disorders, 128–9 treatment of tic disorders, 129–30 Schizophrenia Proneness Instrument-Adult Version (SPI-A), 85 schizophrenia-related movement disorders distinction from OCD-related rituals, 57 schizotaxia, 151 schizotypal disorder ICD-10 criteria, 151
---

- schizotypal OCD, x, 150  
 schizotypal personality disorder, 150–1  
   and subclinical  
     obsessive–compulsive symptoms, 151–2  
   comorbidity with OCD, 150  
   co-occurrence with other personality disorders, 159–60  
   diagnostic instruments, 151  
   DSM-IV criteria, 151  
   features shared with schizophrenia, 151  
   in children and adolescents, 109  
   subclinical, 151–2  
 schizotypal personality disorder in OCD  
   antiglutamatergic agents as potential treatment, 163–4  
   augmentation with low-dose antipsychotics, 161–2  
   clinical characteristics, 154–6  
   cognitive function, 157  
   diagnostic  
     recommendations, 165–6  
   differential diagnosis, 157–60  
   family inheritance, 156–7  
   neuroimaging, 157  
   non-pharmacological interventions, 164–5  
   prevalence, 152–4  
   treatment and responses, 160–1  
   treatment recommendations, 165–6  
   treatments, 160–5  
 Schizotypal Personality Questionnaire, 151  
 schizotypy, 151  
 Schneider, Kurt, 2, 4, 58, 72  
 scruples (disorder), 2  
 Seglas, Jules, 76  
 serotonin  
   dysfunction in OCD, 15  
   dysfunction in schizophrenia, 12  
 sertraline, 107  
 simple deterioration (Pick and Sommer), 1  
 Simpson–Angus Rating Scale (SAS), 203  
 single-nucleotide polymorphism (SNP) data, 7  
 social phobia, 123  
 SRIs  
   schizo-obsessive disorder treatment, 201–3  
 SSRIs, 130  
   motor side effects, 203–4  
   use in OCD, 16  
   use in schizophrenia, 15  
 Stengel, Erwin, 24–6  
 streptococcal infection and OCD, 8, 101  
 stroke, 116  
 Structured Clinical Interview for DSM-III-R Axis II Disorders, 187  
 Structured Clinical Interview for DSM-IV (SCID), 33, 36, 65, 187  
 Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I), 114, 157  
 Structured Clinical Interview for DSM-IV Axis II Disorders (SCID-II), 151, 157  
 Structured Interview for Prodromal Syndromes (SIPS), 87  
 subclinical schizotypal and obsessive–compulsive symptoms, 151–2  
 substance-use disorders, 123, 127  
 suicide risk  
   schizophrenia, 4  
 Taylor, Jeremy, 3  
 terminology  
   variations for schizo-obsessive disorder, x  
 thalamus, 12–14  
 thioridazine, 161  
 thiothixene, 161  
 thought insertion, 2, 5  
 thought-action fusion, 152  
 tic disorders, 123, 144  
   and schizophrenia, 125–7  
   distinction from OCD symptoms, 108  
   in early-onset schizophrenia, 107–8  
 tic disorders in schizophrenia  
   case vignettes, 130–2  
   clinical characteristics, 127–8  
   differential diagnosis, 128–9  
   treatment, 129–30  
 Tirat Carmel Mental Health Center (Israel), 41, 187  
 Tourette’s syndrome  
   and schizophrenia, 125–7  
   distinction from OCD symptoms, 108  
   *See also* tic disorders.  
 traumatic brain injury, 116  
 treatment  
   schizophrenia compared to OCD, 15–17  
 treatment of schizo-obsessive disorder  
   addition of serotonin reuptake inhibitors, 201–3  
   amisulpiride, 197–9  
   antiglutamatergic agents, 204–5  
   aripiprazole, 196–7  
   clozapine, 199–201  
   cognitive–behavior therapy (CBT), 205–6  
   drug-induced motor side effects, 203–4  
   electroconvulsive therapy (ECT), 206–7  
   monotherapy with antipsychotic agents, 194–201  
   non-pharmacological interventions, 205–6  
   olanzapine, 194–6  
   risperidone, 194–6  
   treatment challenges, 194  
   treatment recommendations, 207–9  
 twin studies  
   obsessive–compulsive disorder (OCD), 8  
   schizophrenia, 7  
 urban environment  
   risk factor for schizophrenia, 8  
 Westphal, C., 3, 29, 58, 72  
 withdrawal in schizophrenia, 2  
 Yale–Brown obsessive–compulsive Scale (Y-BOCS), 36, 46, 62, 65, 67, 100, 114, 171, 190  
 ziprasidone, 162  
 zopiclone, 115  
 Zucker Hillside Hospital Recognition and Prevention (RAP) Program (New York, USA), 89