

Index

Abbreviations

AAA – abdominal aortic aneurysm
 ANH – acute normovolemic hemodilution
 EGDT – early, aggressive goal-directed therapy
 GDT – goal-directed therapy
 HES – hydroxyethylstarch
 PVI – pleth variability index
 TUR – transurethral resection

Note: All index entries pertain to fluid therapy, unless otherwise noted. “Fluid therapy” is not listed in the index, and readers are advised to seek more specific topics.

- 4/2/1 rule, 66
- abdominal aortic aneurysm (AAA), hypovolemic shock, 168
- abdominal aortic surgery
 heart rate and blood pressures, 168
 HES and renal function, 145
 hypovolemic shock, 168
 “rules-of-thumb” for fluids, 25–6
- acetate, in Ringer’s solution, 3
- acute normovolemic hemodilution (ANH), 112, 184–5
- advantages, 112, 117
- anesthesia effects, 113, 114, 115
- definition/description, 112, 117
- efficacy, 116–17
 studies/literature reports, 117
 theoretical, 116–17
- limits, 113–16, 187
 cardiac conditions, 115–16
 hemostasis and, 116, 141
 physiological compensatory mechanisms, 112–13, 117
- tolerance to, 115, 185
 see also anemia, dilutional; hemodilution
- acute respiratory distress syndrome (ARDS), 74, 75, 123
- “fluids and catheter treatment trial”, 161
- adrenaline *see* epinephrine (adrenaline)
- adverse reactions of iv fluids, 137–47
 acid-base imbalance, 140–1
 excess tissue hydration and edema, 139–40
 hemostatic *see* hemostatic disturbances, fluid-associated
 hypersensitivity and anaphylaxis, 13, 123, 143–4
 hypervolemia and hypertension, 138–9
 see also fluid overload
 hypothermia, 140, 141
 local effects, 137–8, 145
 organ function impairment, 145
 summary, 145–6
 systemic, 138, 145
 temperature-/osmolality-associated, 138
 tissue deposition and pruritus, 144–5
- Aesculon system, 99
- albumin, 11, 123
 adverse effects, 12, 123
 clinical use, 12, 123
 in infants and neonates, 69
 iodated, 127–8, 130
 pharmacokinetics, 11–12
 plasma volume expansion, 123, 124
 “albumin debate”, 12
- ammonia, 150
- analgesia, multimodal, day surgery, 56
- anaphylactoid reactions, 143–4
- anaphylaxis, 143–4
 dextran associated, 13, 123, 143–4
- anemia
 in critically ill patients, 161
 preoperative, 20
- anemia, dilutional tolerance, 184–5, 189
 in anesthetized subjects, 186
 indirect signs for, 186
 limits, 185–6, 187, 189
 physiological mechanisms, 184–5
 therapeutic increase, 188
 see also hemodilution
- anesthesia
 compensatory mechanisms in hemorrhage and, 178
 depth, dilutional anemia tolerance, 188
 effects on ANH response, 113, 114, 115
 number of procedures, 103
 physiologic adjustments to, 113
 “rules of thumb” for perioperative fluids, 21–2
 in severe sepsis/septic shock, 158
 in trauma, hemorrhage resuscitation and, 182

- anesthesia (*cont.*)
 uncontrolled hemorrhage, management, 181
 volume kinetics affected by, 133
 anesthetic drugs, dilutional anemia tolerance, 188
 anthropometry, 128–9
 aortic reconstruction, hypertonic fluids, outcome, 78
 aortic surgery
 abdominal *see* abdominal aortic surgery
 hypertonic fluid benefits, 78
 “Arg-Arg” phenotype, 172
 arterial pressure, 103
 AAA surgery, 168
 hypovolemic shock, 166
 plasma volume expansion and, 124
 reduced during surgery, 133
 blood loss reduction, 178
 target in trauma resuscitation, 179
 in uncontrolled hemorrhage, 178, 179
see also mean arterial pressure (MAP)
 arterial pressure waveform analysis
 dynamic indices from, 85–6, 97, 98, 104
 limitations, 106
 PPV *see* pulse pressure variation (PPV)
 respiratory systolic variation test, 86
 SVV *see* stroke volume variation (SVV)
 systolic pressure variation, 86, 162
 goal-directed therapy, 97–8, 108
 limitations, 98
 arterial waveform pulse contour cardiac output *see* pulse contour analysis
 atrial natriuretic peptide (ANP), 46, 124, 169
 bariatric surgery, 61
 baroreceptors, 167
 basal fluid requirements, 18–19
 Bezold–Jarish reflex, 170
 bioimpedance (BIA), 128
 BioZ system, 99
 bleeding
 in hepatic surgery, 39
see also blood loss (surgical); hemorrhage
 blood
 preoperative autologous donation, 180, 184
 random-donor fresh whole, 180
 blood–brain barrier (BBB), 122
 disrupted, 122, 123
 normal functions, 122
 blood flow
 cerebral, in hypovolemic shock, 171–2, 173
 microcirculatory, autoregulation, early septic shock, 160
 microvascular, enhancement, 20, 24
 by colloids, 20, 24, 97
 by hypertonic solutions, 73
 redistribution, hemodilution effect, 113
 blood fluidity, 112–13
 blood groups, coagulation response to hemodilution, 116
 blood loss (surgical), 92
 children, 67
 fluid “efficiency” measurement, 130
 GDT using esophageal Doppler, 94–5
 management, 184
 reduction in hepatic surgery, 39
 “rules of thumb,” 21
see also hemorrhage
 blood pressure *see* arterial pressure
 blood products, fluids vs., 184–92
 blood transfusions
 antiviral testing, 180
 costs, 184
 deliberate hypotension in hemorrhage and, 178
 hemorrhagic shock and hypertonic fluid trials, 75
 indications, 20
 minimization strategies, 184
 blood loss management, 184
 dilutional anemia tolerance *see* anemia, dilutional
 intraoperative autologous, 184
 rapid, in hemorrhage, 182
 reduced by ANH, 112
 risk/benefits dilemma, 184
 in septic shock/severe sepsis, 161
 side effects, 125
 in uncontrolled hemorrhage, 180
 uncrossmatched type-O RBC, 180
 blood viscosity, 13, 115
 blood volume
 central *see* central blood volume (CBV)
 expansion *see* plasma volume expansion
 intrathoracic *see* central blood volume (CBV)
 reduced by hemorrhage, 177, 178
 body temperature, reduced *see* hypothermia
 body volume, 127–36
 calculation, using tracers, 127
 normal, 127
see also fluid volumes; volume kinetics
 bowel preparation, 23, 29, 52–3
 bradycardia, in stage II of hypovolemic shock, 168–9
 brain injury
 hypertonic fluid benefits, 73, 74
 clinical trials, 75–8
 Ringer’s solution in, 5
 bromide, extracellular fluid volume, 127
 calories, requirements by children, 66
 capillary fluid exchange
see transcapillary fluid exchange
 capillary leak, 12, 15, 19
 in sepsis, 157–8
 carbohydrates, preoperative, 58

Cambridge University Press

978-1-107-00098-8 - Clinical Fluid Therapy in the Perioperative Setting

Edited by Robert G. Hahn

Index

[More information](#)

- carbon monoxide, 128, 130
- cardiac index, 91
- improvement by hypertonic fluids, 78
- cardiac output, 177
- aortic blood flow velocity and, 39
- continuous, determination, 83–5
- dilutional anemia tolerance mechanism, 185
- hypovolemic shock, 170, 171–2
- increased, hemodilution effect, 112–13
- infants, 65
- inter-individual variation, 172
- maximization in surgery, 103
- measurement/calculation
- arterial waveform pulse contour, 84
- esophageal Doppler monitoring, 88
- fluid responsiveness in septic shock, 162
- transesophageal echocardiography, 87
- monitoring, preload dependence and, 105
- reduced, in hemodilution, anesthesia effects, 113, 114
- spinal anesthesia effect, 46–7
- surrogate parameters for monitoring, 104, 105
- see also* pulse pressure variation (PPV); stroke volume variation (SVV)
- thermodilution
- see* thermodilution
- cardiac resuscitation, hypovolemic shock and, 168, 169
- cardiac surgery
- acute normovolemic hemodilution, 115–16
- dilutional anemia tolerance, 186
- GDT using esophageal Doppler, 94
- cardiopulmonary bypass surgery, hypertonic fluids, 78
- cardiovascular response, absorption of irrigating fluids, 150–1
- carotid baroreceptors, 167
- catecholamines, 134
- see also specific catecholamines*
- cell membrane, 1
- central blood volume (CBV), 38, 166
- heart rate and arterial pressure response, 166
- in hypovolemic shock, 166, 170, 171
- Bezold–Jarish reflex and, 170
- pre-shock, 167
- stage II, 168–9
- stage III, 170
- surgical patients, 171–2, 173
- tilt table experiments, 171
- measurement, 129
- reduced, impact, 171
- central venous oxygen saturation (ScvO₂), 159, 162
- central venous pressure (CVP), 38
- in EGDT, 159
- fluid resuscitation in septic shock, 162
- hepatic surgery, blood loss reduction, 39
- in hypovolemic shock
- see* hypovolemic shock
- in liver transplantation, 39
- volume status assessment, 93, 104
- cerebral blood flow, in hypovolemic shock, 171–2, 173
- cerebral capillaries, 122
- cerebral damage, glucose solutions worsening, 6
- cerebral edema
- absorption of irrigating fluids and, 151
- reduction by hypertonic fluids, 74
- vasogenic, 122, 123
- cerebral oxygenation
- evaluation and maintaining, 172
- in hypovolemic shock, 171–2, 173
- Cesarean section, 45, 46, 54
- children, 65–72
- acute hyponatremia, 70
- blood loss (surgical), 67
- dehydration, 67
- fluid therapy for, 67
- fasting fluid deficit, 67
- fasting guidelines, 66
- fluid losses, 67, 68
- fluid volumes and renal function, 65
- intraoperative fluid management, 67–8
- clinical guidelines, 68
- glucose debate, 67–8
- isotonic hydrating solutions, 68
- quantity, 67
- metabolic requirements, 66
- oral fluid intake, 70
- perioperative hyperglycemia danger, 67–8
- postoperative fluid therapy, 70
- clinical guidelines, 70
- consensus on, 70
- controversies, 70
- postoperative hyponatremia, 70
- preoperative assessments, 66–7
- preoperative hypoglycemia, 67
- resuscitation, hydroxyethyl starch (HES), 69
- septic shock, 67, 69
- volume replacement
- crystalloids, 67
- hydroxyethyl starch (HES), 69
- see also* infants
- chloride ion levels, 140
- cholecystectomy, laparoscopic
- liberal fluids and outcome, 37
- perioperative fluids, 58
- postoperative nausea and vomiting, 24
- “rules-of-thumb” for fluids, 25
- circulation, function, 103
- citrate, 182
- coagulation
- cascade activation
- prevention in trauma, 20
- colloids affecting, 15

- coagulation (*cont.*)
 HES effect on in children, 69
 crystalloids effect, 141
 hemodilution effect, 116, 189
 management, after fluid replacement, 189
 procoagulatory drugs, 190
 temperature for,
 hypothermia avoidance, 140, 141
 in uncontrolled hemorrhage, 189
 coagulopathy
 colloid infusion-associated, 141
 dilutional, 189
 hypothermia-associated, 140, 141
 see also hemostatic disturbances, fluid-associated
 colloid fluids (colloids), 11–17
 adverse reactions, 11, 137, 138, 146
 hemostatic disturbances, 141–3
 local, 137
 organ function impairment, 145
 renal failure, 145
 tissue deposition, 144–5, 146
 in uncontrolled hemorrhage, 180
 albumin *see* albumin
 based on balanced electrolyte fluids, 141
 clinical use, 11
 crystalloids comparison *see under* crystalloid fluids (crystalloids)
 dextran *see* dextran
 elimination, 133
 fluid overload, 139
 in gastrointestinal surgery, 24
 gelatin *see* gelatin
 goal-directed therapy, 97
 intra-abdominal surgery, 40
 in hypovolemic shock, 172, 173
 indication, 11, 97
 in massive fluid requirements, 20
 for microvascular blood flow enhancement, 20, 24, 97
 miscellaneous effects, 15
 plasma volume expansion, 122–4, 138
 colloid osmotic pressure effect, 138
 reducing need for, 124–5
 preloading, in spinal anesthesia, 47, 48
 saline base, effect on chloride load, 141
 in severe sepsis/septic shock, 161
 starch *see* hydroxyethyl starch (HES); starch
 synthetic, plasma volume expansion, 123
 in uncontrolled hemorrhage, 180
 volume kinetics, 133
 volume replacement in infants, 69
 colloid osmotic pressure (COP), 11, 138
 hydroxyethyl starch (HES), 139
 maintaining, to avoid tissue edema, 20
 reduced, crystalloids and, 139, 140
 coload, in spinal anesthesia, 46
 crystalloid vs. colloid, 48
 support lacking for, 48
 colonic resection, 30, 36
 colorectal surgery, GDT using esophageal Doppler, 95
 compensatory mechanisms
 dilutional anemia tolerance, 184–5
 hemodilution, 112–13, 117, 184–5
 hemorrhage, and anesthesia, 178
 continuous cardiac output, 83–4
 coronary artery disease (CAD), 116
 coronary blood flow, 115, 116
 corrected flow time (FTc), 88, 94, 96
 creatinine, 14
 critically ill patients, 157–8
 anemia in, 161
 HES and renal failure risk, 145
 severe sepsis and septic shock, 157–8
 see also sepsis; septic shock
 crystalloid fluids (crystalloids), 1–10
 advantages, 4
 adverse reactions, 137, 138
 acid-base imbalance, 140
 edema and excess tissue hydration, 139–40
 hemostatic disturbances, 141, 142
 local, 137
 for children, 67, 69
 colloid fluids vs., 15–16
 gastrointestinal surgery, 23, 24
 GDT, 96–7, 98
 hemostasis affected by, 116
 hypovolemic shock, 172, 173
 response to ANH, 115
 severe sepsis/septic shock, 161
 spinal anesthesia, 46–7
 in uncontrolled hemorrhage, 180
 coload in spinal anesthesia, 48
 composition, comparisons, 2
 definition, 1
 distribution, 97, 123, 139
 elimination, 15
 fixed administration with GDT of colloids, 140
 fluid kinetics modeling, 47
 gastrointestinal surgery, 23
 glucose solutions, 5–8
 in hypovolemic shock, 172, 173
 indication, 97
 for infants, 69
 isotonic, in uncontrolled hemorrhage, 179–80
 mannitol, 8
 massive fluid resuscitation, adverse effects, 139–40
 normal saline *see* normal saline
 overhydration with, adverse effects, 29
 plasma volume expansion, 122–4, 133, 139, 171
 preload, in spinal anesthesia, 46, 47, 48, 54

- obstetric, 45–6, 47
 preoperative, geriatric surgery, 52–3
 in severe sepsis/septic shock, 161
 volume kinetics, 132–3
 volume needed, 139
 over estimated, 97
 weight gain in elderly, 52, 140
see also glucose solutions; Ringer's solution
 cystoscopy, 148
- D5W fluid, 5
 daily fluid requirements, 18–19
 day surgery, 56–64
 adverse events, 61
 benefits, 56
 discharge criteria, 61
 drugs impact on fluid balance, 61
 elderly patients, 56–7
 intermediate, clinical trials of fluid therapy, 58
 key messages, 62
 outcomes and safety, 56, 61–2
 perioperative fluids, 58–61
 postoperative fluids, 61
 preoperative fasting, 57
 preoperative nutrition
 correction of deficits, 57–8
 energy loading, 58
 dehydration, 52
 abdominal surgery, 29
 children, 67
 iatrogenic, gastrointestinal surgery, 29
 desmopressin, 189
 dextran (DEX), 12–13
 anaphylaxis risk, 13, 123, 143–4
 clinical use, 13, 123–4
 hemostatic disturbances due to, 141, 142, 143
 hypertonic solution with, in hemorrhagic shock, 74
 low-molecular weight dextran-1, 144
 metabolism, 144
 pharmacokinetics, 13, 123
 plasma volume expansion, 123–4
 rheological effect, 15
 safe use, 144
- solutions commercially available, 13
 Dextran 40, 138
 Dextran 70, 123, 138
 dextrose
 intraoperative, children, 67, 68
 postoperative fluids for children, 70
 diuresis
 osmotic *see* osmotic diuresis
 promoting/monitoring, 21
 diuretics, TUR syndrome treatment, 154
 DO₂
 critical, 185–6
 “luxury” (over-supply of oxygen), 185
 dobutamine, 163
 dopamine, 163
 Doppler-guided monitoring
 see esophageal Doppler monitoring (EDM)
- drugs
 impact on fluid balance, day surgery, 61
 pharmacokinetics, 131
- early, aggressive goal-directed therapy (EGDT), 158
 clinical trials, 158, 159–60
 importance of speed/timing, 159
 principles and benefits, 159
- echocardiography,
 transesophageal, 86–8
 costs, advantages and contraindications, 87
 multiplane probe, 87
 pulmonary artery catheter use vs., 87
- edema, 120
 adverse effects of infusion fluids, 139–40
 cerebral *see* cerebral edema
 crystalloids in plasma volume expansion, 123
 fluid replacement in sepsis, 160
 mechanisms, 121, 122
 perioperative fluid excess, 92
 pitting, 4
 prevention, colloid osmotic pressure, 20
 sequelae, 123
 elderly patients
- cardiac output in spinal anesthesia, 46–7
 colonic surgery, liberal vs. restricted fluids, 53
 critically ill patients, 158
 day surgery, 56–7
 restrictive vs. liberal fluids, 58
 definition, for trials, 52
 fluid retention, 52
 orthopedic surgery,
 restrictive vs. liberal fluids, 53
 perioperative cardiovascular events, 57
 spinal anesthesia in, 46–7, 48
 hypotension, 45, 46
 weight gain and fluid overload, 52, 140
see also geriatric surgery
- elective surgery
 fluid status assessment, 51
 fluid types to administer, 51
see also specific surgery types
- electrocautery, irrigating fluids, 148
- electrolytes
 glucose solutions with, 2, 5, 6, 7
 isotonic hydrating solutions for children, 68
 requirements by children, 66
- encephalopathy,
 hyperammonemic, 150
- endothelial cell edema, 73
- energy loading, preoperative nutrition, 58
- Enhanced Recovery after Surgery (ERAS)
 programs, 96
- ephedrine, 54
- epidural anesthesia *see* spinal anesthesia
- epinephrine (adrenaline), 163
 stage II of hypovolemic shock, 168
- esophageal Doppler monitoring (EDM), 51, 86, 87–8
 cardiac output calculation, 88
 contraindication, 40
 corrected flow time (FTc), 88, 94, 96
 in GDT *see* goal-directed therapy (GDT)

- esophageal Doppler monitoring (EDM) (*cont.*)
 intraoperative fluid management, 51
 intra-abdominal surgery, 37, 39
 limitations, 88
 principles/methods, 93
 protocols, 88
 volume status assessment, 86, 87–8
- ethanol, 1, 127
 irrigating fluid absorption measurement, 152, 153
- Evans blue, 128, 130
- evaporative loss, 23
- extracellular fluid (ECF)
 expansion
 mannitol, 8
 Ringer's solutions, 4
 hypo-osmotic irrigating fluids effect, 150
 volume, 127
 body weight relationship, 129
 children, 65
 measurement, 129
 normal adult, 127
 tracers, for measuring, 127
- extracellular fluid (ECF) space, 1
- extravascular space, 120
- factor XIII concentrate, 189
- “fast-track surgery,” 52, 53
- fasting, preoperative routines, 57
 fluid deficits in children, 67
 revised guidelines, 57, 62
 children, 66
- fatigue reduction, day surgery, 61
- fibrinogen, 189
- FiO₂ (inspiratory oxygen fraction), 188
- Flotrac/Vigileo, 38, 85, 98, 108
- fluid challenge
 Frank–Starling, 93, 104
 in severe sepsis/septic shock, 162
- fluid compartments, 127, 132
- fluid “efficiency,” 129–31
 measurement, 130
 hemoglobin dilution concept, 130
 hemoglobin levels, 130
 tracers, 130
- fluid exchange
 see microvascular fluid exchange
- fluid intake (oral)
 postoperative, in children, 70
 preoperatively, 57
- fluid kinetics *see* volume kinetics
- fluid losses
 children, 67, 68
 see also hypovolemia
- fluid overload
 adverse effects, 29
 avoidance strategy, 140
 colloids, 139
 crystalloids, 139–40
 perioperative complications, 92
 in elderly, 53
 postoperative, 51, 54
 pulmonary surgery, 54
 see also hypervolemia
- fluid retention, 51
- elderly, 52
- fluid status assessment, 51
- fluid volumes
 calculation, using tracers, 127
 kinetics *see* volume kinetics
- measurement
 anthropometry, 128–9
 bioimpedance, 128
 sodium method, 129
 tracers, 127–9
 sizes, normal, 127
- “fluids and catheter treatment trial,” 161
- Frank–Starling fluid challenge, 93, 104
- Frank–Starling relationship/
 curve, 85, 93, 97, 98
 absorption of irrigating fluids, 151
 cardiac function affecting, 104, 105
 description, 105
 in hemorrhage, 177
 preload dependence, 104, 105
 respiratory variations in arterial pulse pressure, 105
- fresh frozen plasma (FFP), 189, 190
- frusemide, 154
- FTc (corrected flow time), 88, 94, 96
- functional body fluid spaces, 132
- functional hemodynamic parameters, 103–6
 algorithm for monitoring in surgery, 109
 fluid responsiveness in septic shock, 162
 monitoring, clinical outcome and, 108
see also arterial pressure waveform analysis; pleth variability index (PVI); pulse pressure variation (PPV); stroke volume variation (SVV)
- gas exchange, hemodilution and anesthesia effects, 114
- gastrointestinal surgery
 accelerated care protocols, 29
 laparoscopic, 24–5
 liberal vs. restrictive fluids in elderly, 53
 open abdominal, fluid therapy, 23–4
 “rules-of-thumb” for perioperative fluids, 23–5
 laparoscopic surgery, 25
 open abdominal surgery, 24
- gelatin (GEL), 15, 124
 anaphylactoid/allergic reactions, 144
 hemostatic disturbances due to, 142, 143
 HES comparison, 143
 metabolism, 144
 plasma volume expansion, 124
 preloading, in spinal anesthesia, 47
 products and types, 144
 volume replacement in infants, 69
- general anesthesia, “rules of thumb” for fluids, 21
- geriatric surgery, 52–3
 perioperative fluid management, 53
 postoperative fluid management, 53

- preoperative fluid
 management, 52–3
 see also elderly patients
 Glasgow Coma Scale,
 hypertonic fluid
 administration, 75–8
 glucose
 isotonic hydrating solution
 effect (pediatric), 68
 renal threshold, 7
 glucose-free solutions, for
 children, 68
 “glucose loading,” 6
 glucose solutions, 5–8
 adverse reactions
 cerebral damage, 6
 local, 137
 volume overload and
 hypertension, 139
 clinical use, 6
 composition, 2
 contraindications, 6
 dosing and requirements, 7
 electrolytes with, 2, 5, 6, 7
 hyperosmolar, adverse
 effects, 139
 hypertonic, 6
 hyponatremia due to, 7
 insulin with, 6
 intraoperative fluids for
 children, 67–8
 “nitrogen-sparing effect,”
 6, 7
 osmolality, 137
 perioperative, in day surgery,
 61
 pharmacokinetics, 5–6
 rebound hypoglycemia, 7–8
 volume kinetics, 134
 glycine solution, 150
 elimination and absorption,
 150
 in TURP, 149
 glycosuria, 139
 “Gly-Gly” phenotype, 172
 goal-directed therapy (GDT),
 37, 51, 91–102
 in abdominal surgery
 see intra-abdominal
 surgery
 background to, 91
 benefits, 92, 99, 108
 colloids, with fixed
 crystalloid
 administration, 140
 definition/description, 91
 early aggressive *see* early,
 aggressive goal-directed
 therapy (EGDT)
 early, supranormal oxygen
 delivery, 91–2, 93, 99
 mechanisms, 92
 esophageal Doppler for, 93–6
 guidelines, 95
 improved outcome,
 studies, 96
 inotropes and vasodilators
 with, 96
 limitations, 96
 recommendations, 96
 hypovolemic shock, 172
 limitations, 92, 99
 modern, individualized
 volume optimization,
 92–9
 arterial pressure waveform
 analysis, 97–8, 108
 crystalloids vs. colloids,
 96–7, 98
 technologies (other), 98–9
 see also pulse pressure
 variation (PPV); stroke
 volume variation (SVV)
 monitors for volume status,
 93
 guidelines, fluid management,
 21–2
 esophageal Doppler for
 GDT, 95
 intraoperative fluids for
 children, 68
 postoperative fluids for
 children, 70
 see also “rules-of-thumb”
 Haemaccel™, 69
 Hartmann’s solution *see* lactated
 Ringer’s solution
 head injury, albumin-related
 mortality, 12
 head-up tilt (HUT), 166, 167,
 170
 heart rate
 AAA surgery, 168
 hypovolemic shock, 166, 167
 stage II, 168–9, 170
 stage III, 170
 hematocrit (Hct)
 critical, 185
 in dilutional anemia
 tolerance, 185
 hemodilution, 112–19, 172
 acute normovolemic
 see acute normovolemic
 hemodilution (ANH)
 anesthesia effects, 113, 114,
 115, 188
 compensatory mechanisms,
 112–13, 117, 184–5
 crystalloids, effect on
 coagulation, 141
 hemostasis affected by, 116,
 141, 189
 hypovolemic, 188
 limits, 113–16
 cardiac conditions,
 115–16, 186
 DO_{2crit}, 185–6
 hemostasis and, 116, 141,
 189
 normovolemic, 184, 185
 limits, 187
 tolerance to, 115, 185
 see also anemia, dilutional
 hemodynamic instability, 19
 hemodynamic monitoring, 108
 invasive *see* invasive
 hemodynamic
 monitoring
 non-invasive *see* non-
 invasive guidance, fluid
 therapy
 hemodynamic parameters
 fluid administration in intra-
 abdominal surgery, 38
 functional *see* functional
 hemodynamic
 parameters
 hemoglobin
 critical value, tissue
 oxygenation, 113, 125,
 185, 187, 189
 critically ill patients, 161
 postoperative, mortality,
 186
 dilution concept, 130
 fluid “efficiency”
 measurement by, 130
 measurements, volume
 kinetic analysis, 131,
 134
 normal, plasma volume
 expansion and, 125
 recommended levels, 20
 restrictive vs. liberal
 transfusion strategies, 20
 hemolysis-induced renal
 failure, 148

- hemorrhage
 cerebral blood flow and oxygenation, 171
 massive, procoagulatory drugs, 190
 pale skin and sympathetic activity, 169–70
 perioperative setting, 178
 uncontrolled, 177–83
 adjuvant therapies, 182
 blood transfusions, 180
 choice of resuscitation fluid, 179–81
 coagulation management, 189
 deliberate hypotensive *see under* hypotension
 goal of resuscitation, 181
 guidance on resuscitation, 181–2
 normothermia vs. hypothermia, 182
 oscillation of blood pressure, 179
 pathophysiology, 177–8
 plasma administration, 181
 primary resuscitative fluid, 181
 recommended management, 181–2
 in surgery, anesthesia and, 178, 179, 181
 in trauma, 179, 181, 182
 Type-O red blood cells, 180
hemorrhagic shock, 157, 177, 178
 hypertonic fluid effects, 73, 74–5, 180
 summary of trials, 76
 isotonic crystalloid fluids, 179–80
 late signs, 177
 see also hemorrhage, uncontrolled
hemostasis, 178
 ANH and, 116, 141
 in shock and injury, 180–1
 see also coagulation
hemostatic disturbances, fluid-associated, 141–3
 colloid-associated, 141–3
 crystalloid-associated, 141, 142
 mechanisms, 141
 see also coagulopathy
hepatic surgery, 39
 hypertonic fluids in, 78
 Hetastarch, 13, 73, 142
 high-flow, low-pressure state, 181
 hip surgery, “rules of thumb” for perioperative fluids, 22
HSD (hypertonic saline dextran), effect on ARDS, 74
hydroxyethyl starch (HES), 13, 14, 124
 130/0.4 preparations, 144
 safety aspects, 145
 anaphylactoid reaction rate low, 143
 anti-inflammatory effects, 20, 26
 bleeding association, 142
 clinical use, 14, 124, 141
 colloid osmotic pressure, 139
 first generation, hemostatic effects, 142
 gelatin comparison, 143
 in hemorrhage, 188
 hemostatic disturbances due to, 116, 141–3
 hyperoncotic, 14
 in hypertonic saline, 14
 metabolism, 144
 incomplete, tissue deposition, 144
 in neonates, 69
 pharmacokinetics, 14, 124
 in plasma-adapted solutions, 142, 145
 plasma volume expansion, 124
 preloading, in spinal anesthesia, 47
 preparations, characteristics, 141
 pruritus due to, 144–5
 renal failure risk after, 145
 third generation, hemostatic disturbances, 142–3
 volume replacement in infants/children, 69
 coagulation, effect on, 69
 quantity and types, 69
hyperchloremic metabolic acidosis, 46, 140
 normal saline causing, 1, 46, 150, 180
hypercoagulable state, dextran counteracting, 143
hyperdynamic state, 160, 162, 163
hyperglycemia, 139
 perioperative, in children, 67–8
hyperosmolar solutions, volume overload and hypertension, 139
hypersensitivity, reaction to iv fluids, 143–4
hypertension, adverse effect of iv fluids, 138–9
hypertonic fluids, 73–81
 in aortic surgery, 78
 benefits, 73
 in brain injury, 73, 74
 prehospital administration, outcomes, 75–8
 in cardiopulmonary bypass surgery, 78
 clinical trials, 74–8
 hemorrhagic shock, 74–5, 76
 traumatic brain injury, 75–8
 in hemorrhagic shock, 73, 74–5, 76
 in hepatic surgery, 78
 intraoperative studies, 78–9
 intracranial pressure reduction, 74, 75
 mechanism of action, 73–4
 in spinal surgery, 78
 types, 73
 in uncontrolled hemorrhage, 180
 see also hypertonic saline
hypertonic saline, 3, 73
 adverse reactions
 local, 137–8
 volume overload and hypertension, 139
 in aortic reconstruction, 78
 in aortic surgery, 78
 in hyponatremia, 7
 for increased intracranial pressure, 75
 osmolality, 137–8
 prehospital administration, outcome in brain injury, 75–8
 TUR syndrome treatment, 153

- volume kinetics, 132, 134
- hypertonic saline dextran (HSD), effect on ARDS, 74
- hypervolemia
 - absorption of irrigating fluids, 150–1
 - adverse effect of iv fluids, 138–9
 - see also* fluid overload
- hypo-osmolality, 151
- hypoalbuminemia, 12
- hypocalcemia, 182
- hypoglycemia
 - glucose solutions, 6
 - preoperative, in children, 67
 - rebound, 7–8
- hyponatremia, 150
 - absorption of irrigating fluids, 151
 - treatment, 153–4
- acute, management in children, 70
- brain damage, 153
- glucose solutions inducing, 7
- postoperative, in children, 70
- hypotension
 - anesthesia-induced, 21, 22
 - deliberate, in resuscitation, 178, 179, 181
 - barriers/problems, 179
 - drug-induced, 4
 - hemorrhage in trauma patients and, 179
 - maintenance, in hemorrhage, 178–9
 - in spinal anesthesia
 - see* spinal anesthesia
- hypothermia, 140, 141
 - benefits in resuscitation, 182
 - dilutional anemia tolerance, 188
- hypovolemia, 120
 - central, 168
 - consequences, 92
 - fluid status assessment and, 51
 - increased transcapillary escape rate, 122
 - plasma volume expanders
 - see* plasma volume expansion
 - prevention, perioperative fluids, 19
 - reversal, Ringer's solutions, 4
 - signs, 104
- Starling curve, LV stroke volume, 98
- tissue oxygenation and hemodilution effect, 114
- hypovolemic shock, 166–76
 - AAA surgery, 168
 - “cardiac arrest” in, 168
 - cardiac output, 170, 171–2
 - central blood volume
 - see* central blood volume (CBV)
 - central venous pressures, 170–2
 - normovolemia definition, 171
 - characteristics, 166
 - heart rate and arterial pressure, 166
 - hypertonic fluid benefits, 73
 - pre-shock, 167
 - stage II, 168–70
 - Bezold–Jarish reflex, 170
 - bradycardia, 168–9
 - pale skin and sympathetic activity, 169–70
 - to stop bleeding, 169
 - stage III, 168, 170
 - volume treatment, 166, 171–2
 - aim, 172
 - choice, 172–3, 174
 - titration for normovolemia, 172
- hypoxia, tissue, 185
- hysterectomy, hypertonic saline, 78
- immunosuppressive response, hypertonic fluids, 73–4
- indications and challenges, of fluid therapy, 18–21
 - basal fluid requirement provision, 18–19
 - microvascular blood flow enhancement, 19–20
 - oxygen transport adequacy, 20
 - promoting/monitoring of diuresis, 21
 - tissue edema prevention, 20
- indocyanine green (ICG), 128
- infants
 - fasting guidelines, 66
 - guidelines for intraoperative fluid therapy, 68
- volume replacement, 69
 - albumin, 69
 - crystalloids, 69
 - gelatins, 69
 - hydroxyethylstarch preparations, 69
 - lactated Ringer's solution, 69
 - see also* children; neonates; premature infants
- inflammation
 - early vs. late septic shock, 158
 - prevention in trauma, 20
- inflammatory cytokines, hypertonic fluid effects, 73
- infusion fluids
 - dilutional anemia tolerance, 188
 - as drugs, ix, 131
 - warming, importance, 140, 141
 - see also* colloid fluids (colloids); crystalloid fluids (crystalloids)
- inotropic agents, 82, 159, 163
- inspiratory oxygen fraction (FiO₂), 188
- insulin, glucose solutions with, 6
- interstitial fluid space, 3, 4, 11
- interstitial fluid volume, 1, 127
 - measurement, 128
 - normal adult, 127
- intra-abdominal surgery, 29–44
 - clinical trials, “fixed” fluid volumes, outcome, 30–7, 39
 - improved outcome with liberal fluids, 36–7
 - improved outcome with restricted fluids, 30–6
 - outcome differences lacking, 36
 - PVI monitoring, 108
 - summary of trials, 31
 - clinical trials, goal-directed fluid volume, outcome, 37–9, 108
 - benefits, 39
 - colloid boluses, 40
 - Doppler-guided fluid management, 37, 39
 - hepatic surgery, 39
 - optimization protocol, 38
 - restricted fluids vs., 38–9

- intra-abdominal surgery (*cont.*)
 - summary of trials, 33
 - duration, effect on fluid balance, 37
 - inadequacy of predefined strategy, 40
 - “rules of thumb” for fluids, 29
 - criticisms of, 29–30
 - summary of findings, 39–40
 - timing of fluid administration, 40
 - see also* gastrointestinal surgery
 - intracellular fluid (ICF)
 - hypo-osmotic irrigating fluids effect, 150
 - volume, 127
 - measurement, 128, 129
 - normal adult, 127
 - intracellular fluid (ICF) space, 1, 6
 - intracranial pressure
 - increase, causes, 122
 - reduction by hypertonic fluids, 74, 75
 - reduction by mannitol, 8, 75
 - intraoperative period
 - fluids for children
 - see* children
 - hypertonic fluids, 78–9
 - “rules of thumb” for
 - perioperative fluids, 21
 - intra-abdominal surgery, 29
 - laparoscopic cholecystectomy, 24
 - open gastrointestinal surgery, 24
 - orthopedic surgery, 22
 - vascular surgery, 26
 - intrathoracic blood volume (IBV) *see* central blood volume (CBV)
 - intravascular volume, 18, 51, 85
 - deficit, before surgery, 19
 - overload, colloid infusions, 139
 - replacement, in children, 67
 - invasive hemodynamic monitoring, 82–90, 92, 103
 - arterial waveform pulse contour cardiac output
 - see* pulse contour analysis
 - dynamic indices *see* arterial pressure waveform analysis
 - echocardiography, 86–8
 - esophageal Doppler
 - see* esophageal Doppler monitoring (EDM)
 - pulmonary artery catheterization
 - see* pulmonary artery catheterization (PAC)
 - transpulmonary
 - thermodilution cardiac output, 84
 - iodinated albumin, 127–8, 130
 - iohexol, 127
 - irrigating fluids, 148–56
 - absorption, 148
 - fluid comparisons, 151
 - incidence and presentation, 149
 - measurement, 151–3
 - mechanisms, 148–9
 - pathophysiology, 150–1
 - prevention, 153
 - treatment, 153–4
 - types of fluids used, 150
 - see also* transurethral resection (TUR) syndrome
 - complications, 148
 - composition, 148
 - extravasation, 149
 - massive, management, 154
 - hemolysis-induced renal failure, 148
 - types, 150
 - iso-osmotic fluid, 1
 - isotonic solution, 1
 - intraoperative, for children, 68
 - normal saline, 1–3
 - perioperative, 58
 - postoperative fluids for children, 70
- Jehovah's Witnesses, 114, 186
- knee surgery
 - perioperative restrictive vs. liberal fluids, 53
 - “rules of thumb” for perioperative fluids, 23
- labor, 8
- lactate
 - in resuscitation in hemorrhage, 182
 - in Ringer's solution, 3
 - serum levels, intra-abdominal surgery, 38–9
- lactated Ringer's solution, 3
 - composition, 2
 - perioperative, in day surgery, 58, 62
 - in uncontrolled hemorrhage, 179–80
 - volume kinetics, 132
 - volume loading, 5
 - volume replacement in infancy, 69
- laparoscopic cholecystectomy
 - see* cholecystectomy
- laparoscopic gastrointestinal surgery, 25
 - “rules of thumb” for fluids, 25
- left ventricular end-diastolic area, 87
 - assessment by
 - transesophageal echocardiography, 87
- left ventricular end-diastolic pressure (LVEDP), 98
- left ventricular end-diastolic volume (LVEDV), 82, 87
 - esophageal Doppler monitoring, 88
- left ventricular preload, 82–3, 85
- liberal fluid regimens
 - day surgery, 58
 - description, 52
 - in elderly, restrictive fluids vs. colonic surgery, 53
 - orthopedic surgery, 53
 - laparoscopic cholecystectomy outcome, 37
- LiDCO, 84–5, 98
- lithium dilution, calibrated pulse contour analysis by, 84–5
- liver transplantation, 39, 78
- lower body negative pressure (LBNP), 166, 170
- lower limb surgery, “rules-of-thumb” for fluids, 25
- lymphatic system, 120, 121–2
 - stimulation, 125

- malnutrition, preoperative management, 57–8
mannitol, 8, 75, 150
 elimination and absorption effects, 150
 hypertonic, 154
mass balance equation, 129
mean arterial pressure (MAP), 167
 in EGDT, 159
 noradrenaline use, 163
 reduced central blood volume, 167
 reduced, stage II of hypovolemic shock, 168, 173
 in uncontrolled hemorrhage, 179
 see also arterial pressure
mechanical ventilation, 104, 106
metabolic acidosis
 hyperchloremic
 see hyperchloremic metabolic acidosis
 hypochloremic, 1
 normal saline inducing, 1, 11, 46, 150, 180
microvascular blood flow
 see blood flow
microvascular fluid exchange, 120–6
 inside brain, 122
 lymphatic system, 121–2
 outside brain, 120–1, 122
 2-pore theory, 121, 124
 see also hypovolemia; plasma volume expansion
moderate-risk surgery, 103
 pulse oximeter waveforms, 106
morbidity, perioperative, 92
mortality, high-risk surgery, 91
muscular relaxation, dilutional anemia tolerance, 188
myocardial function, dilutional anemia tolerance, 188
neonates
 albumin use for volume expansion, 69
 HES use, 69
 see also infants
neutrophil action, 73
NICOM, 99
nitric oxide, 112
non-invasive guidance, fluid therapy, 103–11
 functional hemodynamic parameters, 104–6
 see also pulse pressure variation (PPV)
 pleth variability index
 see pleth variability index (PVI)
 plethysmographic waveform (Δ POP), 103, 106, 107
 preload dependence determination, 104
 pulse oximeter waveforms, 106–7
 static parameters, 104
 static vs. dynamic parameters, 104
 see also pulse oximeter waveforms
nonlinear least squares regression, 131
non-steroidal anti-inflammatory drugs (NSAIDs)
 day surgery, 61, 62
 impact on fluid balance, 61
noradrenaline (norepinephrine), 163
 stage II of hypovolemic shock, 168
normal saline, 1–3
 adverse effects, 3, 180
 composition, 2
 hyperchloremic metabolic acidosis due to, 1, 46, 150, 180
 for infants, 69
 irrigating solution, 150
 adverse effects, 151
 postoperative fluids for children, 70
normovolemia
 hemodilution
 see hemodilution
 maintenance, in anemia tolerance, 188
 plasma volume expanders
 for *see* plasma volume expansion
 “rules-of-thumb,” 19, 26
 in vascular surgery, 26
nutrition, preoperative day surgery and, 57–8
 energy loading, 58
obstetric spinal anesthesia
 see spinal anesthesia, obstetric
obstetric surgery, fluid management, 54
oedema *see* edema
oncotic pressure, plasma, 124
 reduced, 70, 123
oozing, 13
optimal fluid strategy, 18
 gastrointestinal surgery, 23–4
 see also under goal-directed therapy (GDT)
orthopedic surgery
 GDT using esophageal Doppler, 96
 restrictive vs. liberal fluids in elderly, 53
 “rules of thumb” for perioperative fluids, 22–3
osmolality, 1
osmolality-associated adverse effects, 137–8
osmotic diuresis, 7, 8
 glycine solution absorption causing, 150
osmotic pressure, colloid
 see colloid osmotic pressure (COP)
overhydration
 adverse effects, 29
 see also fluid overload
oxygen consumption
 hemodilution and anesthesia effects, 114
 increased, ANH, 113
 myocardial, 115, 116, 188
 VO_2 , 185, 186
 oxygen supply-dependency, 185
oxygen delivery, 103, 185
 critical DO_2 , 185–6
 factors affecting, 186
 myocardial, 115, 116
 optimization, 103
 over-supply (luxury DO_2), 185
 supranormal, early GDT, 91–2, 99
 targets, 91
oxygen dissociation curve, 113

Cambridge University Press

978-1-107-00098-8 - Clinical Fluid Therapy in the Perioperative Setting

Edited by Robert G. Hahn

Index

[More information](#)

- oxygen, inspiratory fraction (FiO₂), 188
- oxygen saturation
central venous (ScvO₂), 159, 162
measurement, pulse oximetry, 107
venous *see* venous oxygen saturation (SvO₂)
- oxygenation, tissue
adequacy, fluids for, 20
criticisms of hypothesis involving, 29
hemodilution effects, 112, 113–14
increased by liberal fluid use, 36
increased, hemodilution effect, 113–14
efficacy, 114–15
intra-abdominal surgery, 36, 40
- parasympathetic activation, hypovolemic shock, 168
- pediatrics, 65–72
see also children; infants
- pentastarch, 13, 144
- perfusion index (PI), 107
- perioperative fluids
day surgery, 58–61
geriatric surgery, 53
importance, 18
indications *see* indications and challenges
optimization, 18
principles, 19, 158
volume, 58
see also more specific topics
- pharmacokinetics
albumin, 11–12
crystalloids, 97
dextran, 13, 123
drugs, 131
glucose solutions, 5–6
hydroxyethyl starch (HES), 14, 124
infusion fluid kinetics and, 131–4
mannitol, 8
Ringer's solution, 3–4
starch, 14
- phenylephrine, 54
physiotherapy, 125
PiCCO, 85, 86, 98
pitting edema, 4
plasma
administration, 15, 184, 189
in trauma, 181
adverse reactions, 15
composition, 2
dilution, after infusion, 131
fresh frozen, 189, 190
in plasma volume expansion, 15
plasma colloid osmotic pressure *see* colloid osmotic pressure (COP)
Plasma-Lyte A, 2, 5
in uncontrolled hemorrhage, 179–80
plasma volume, 127
body weight relationship, 129
measurement, 127–8
normal adult, 127
plasma volume expansion
adverse effects, 124
colloid fluids, 11, 122–4, 138
albumin, 11–12, 123, 124
dextran, 13, 123–4
gelatin, 124
hydroxyethyl starch, 14, 124
reducing need for, 124–5
crystalloid fluids, 122–4, 133
efficiency of fluid, 129–31
ideal/"target," 134
normal arterial pressure and, 124
reduced renal clearance effect, 133
responders and non-responders, 134
Ringer's solution, 4
effectiveness, 133
pleth variability index (PVI), 103, 107–8
monitoring, clinical outcomes, 108
plethysmographic waveform *see* pulse oximeter waveforms
polycythemia, neonatal, 69
polyionique B66, 68
ΔPOP, 106, 107
2-pore theory, 121, 124
positive-pressure ventilation, 85, 97, 104
postoperative complications, "minor," 103
postoperative fluid management
in children/infants *see* children
day surgery, 61
geriatric surgery, 53
"rules of thumb," 21–2
laparoscopic surgery, 25
open gastrointestinal surgery, 24
orthopedic surgery, 22
vascular surgery, 26
postoperative nausea and vomiting (PONV), 169
day surgery, 62
laparoscopic gastrointestinal surgery, 24
prevention, 24–5
postoperative pain management, day surgery, 62
potassium, serum levels, absorption of irrigating fluids, 151
pre-eclampsia, 46, 48
preload dependence, 104, 105
assessment, 104
importance, 105
definition/description, 104
premature infants
albumin use for volume expansion, 69
volume replacement, 69
preoperative assessments, children, 66–7
preoperative fluids
geriatric surgery, 52–3
"rules of thumb," 21
laparoscopic surgery, 25–6
open gastrointestinal surgery, 24
orthopedic surgery, 22
vascular surgery, 26
pressure breathing, 166
protein exchange, microvascular, 120, 121
mechanisms, 121
convection, 121
passive, 121
2-pore theory, 121, 124
protein-sparing effect, 6, 7
pruritus, 144–5
hydroxyethyl starch (HES), 144–5
pulmonary artery catheterization (PAC), 82–4

- continuous cardiac output, 83–4
- early, GDT, 91
- thermodilution cardiac output, 83
- transesophageal echocardiography vs., 87
- usage, trends, 82
- pulmonary artery occlusion pressure (PAOP), 162
- pulmonary artery wedge pressure (PAWP), 82–3
- pulmonary capillary wedge pressure (PCWP), 104
- pulmonary edema, 29, 54, 123
 - fluid volume relationship, 54
 - rapid infusion of Ringer's solution, 4, 5
- pulmonary surgery, fluid management, 54
- pulse contour analysis, 84
 - calibrated, 84
 - by lithium dilution (LiDCO), 84–5, 98
 - by thermodilution (PiCCO), 85, 86, 98
 - continuous, 84–5
 - contraindications, 84
 - fluid responsiveness in septic shock, 162
 - non-calibrated, 84, 85
- pulse oximeter waveforms, 106–7
 - amplitude (Δ POP), 106, 107
 - components, 106
 - fluid responsiveness prediction, 107
 - principles, 106
 - respiratory variations, 106–7
- pulse oximetry, oxygen saturation measurement, 107
- pulse pressure, 38
- pulse pressure variation (PPV), 86, 97, 103
 - applications, conditions for, 106
 - fluid responsiveness in septic shock, 162
 - mechanical ventilation effect, 104
 - monitoring, clinical outcomes, 108
 - respiratory variations, pulse oximeter, 105, 106
 - tidal volume impact, 106
- red blood cells (RBCs)
 - hemostasis and, 116
 - labeling, mass calculation, 128
 - reduction, hemodilution, 112
 - transfusions, 180, 184
 - velocity, 113
- reflection coefficient, proteins, 120, 121
- regression equations, 129, 131
- rehydration, 19
- renal damage, absorption of irrigating fluids, 151
- renal failure
 - colloid-associated, 145
 - critically ill patients, 158
 - hemolysis-induced, 148
- renal function, children, 65
- renal insufficiency, Ringer's solutions cautions, 5
- renal threshold, glucose, 7
- respiratory systolic variation test, 86
- restrictive fluid regimens
 - description, 52
 - intra-abdominal surgery
 - effect on outcome, 30–6
 - GDT fluid volume effect on outcome vs., 38
 - perioperative, in elderly, 53, 58
- resuscitation
 - cardiac, hypovolemic shock and, 168, 169
 - children, hydroxyethyl starch (HES), 69
 - deliberate hypotension in *see* hypotension
 - hypertonic fluids in *see* hypertonic fluids
 - massive, crystalloid adverse effects, 139–40
 - in severe sepsis *see* sepsis
 - in uncontrolled hemorrhage *see* hemorrhage
- Resuscitation Outcomes Consortium, 75
- right ventricular filling, 83
- Ringer's solution, 3–5
 - acetated, 132
 - adverse effects, 4, 5, 29
 - in brain injury, 5
 - clinical use, 4
 - composition, 2
 - chloride, 141
 - lactate and acetate, 3
 - dosage/infusion volume, 4–5
 - for infants, 69
 - lactated *see* lactated Ringer's solution
 - pharmacokinetics, 3–4
 - Plasma-Lyte A *see* Plasma-Lyte A
 - plasma volume expander, effectiveness, 133
 - preloading, laparoscopic cholecystectomy, 24–5
 - in spinal anesthesia, 47
 - volume kinetics, 132–3
- “rules-of-thumb,” ix, 18–28
 - for adequate colloid osmotic pressure, 20
 - for adequate oxygen transport, 20
 - for basal fluid requirements, 19
 - general principles, 18–21
 - intra-abdominal surgery, 29
 - for normovolemia and hemodynamic stability, 19, 26
 - to prevent hypovolemia, 19
 - to prevent trauma-induced cascade systems, 20
- procedure-specific, 22
 - gastrointestinal surgery *see* gastrointestinal surgery
 - orthopedic surgery, 22–3
 - vascular surgery, 25
- to promote/monitor diuresis, 21
- for rehydration in fluid deficiency, 19
- standard (perioperative fluids), 21–2
- see also* guidelines, fluid management
- SAFE study, 161, 180
- saline
 - hypertonic *see* hypertonic saline
 - physiological (isotonic) *see* normal saline
- sepsis, 157
 - HES avoidance, 14
 - severe, 157–65
 - anesthesia in, 158
 - crystalloids vs. colloids, 161
 - definition, 157

- sepsis (*cont.*)
 early fluid resuscitation, 158–60
 fluid responsiveness, 162
 later fluid resuscitation, 160–1
 positive fluid balance and edema, 160
 summary/key messages, 163
 vasopressors and inotropic agents, 163
 septic shock, 157–65
 anesthesia in, 158
 children, 67
 definition, 157
 early vs. late, 158
 fluid management, 158, 160
 crystalloids vs. colloids, 161
 early, 158–60
 later, 160–1
 summary/key messages, 163
 fluid responsiveness, 162
 hydroxyethyl starch in children, 69
 vasopressors and inotropic agents, 163
 shock
 cold *see* hemorrhagic shock
 warm *see* septic shock
 smoking, 149
 sodium
 isotonic hydrating solution effect (pediatric), 68, 70
 for measuring irrigating fluid absorption, 151, 152, 153
 postoperative fluids for children, 70
 sodium method, 129
 sorbitol solutions, 150
 spinal anesthesia, 45–50
 cardiac output affected by, 46–7
 colloid fluid preloading, 47, 48
 coload, vs. preload, 46, 48
 crystalloid fluid preloading, 45–6, 47, 48, 54
 with gelatin or HES, 47
 crystalloids vs. colloids, 46–7, 54
 coload, 48
 in elderly, 45, 46–7, 48
 hypertonic fluids in, 78
 hypotension in, 45, 47, 54
 elderly, 45, 46
 etiological factors, 45
 fluid management summary, 48–9
 non-obstetric, hypotension in, 46
 obstetric, 45–6, 48
 hypotension, 45–6, 47, 54
 recommendations of fluids, 49
 “rules of thumb” for
 perioperative fluids, 21
 knee surgery, 23
 orthopedic surgery, 22
 timing of fluid administration, 48
 spinal surgery, intra-aortic hypertonic fluids, 78
 starch, 13–14
 adverse effects, 14
 clinical use, 14
 pharmacokinetics, 14
 preparations/formulations, 13–14
 see also hydroxyethyl starch (HES)
 Starling curve *see* Frank–Starling relationship/curve
 Starling fluid equation, 120, 121
 sterile water, 148, 150
 steroids, impact on fluid balance, 61
 Stewart–Hamilton equation, 83
 stroke, acute, 6
 stroke volume (SV), 38, 87
 increase in preload, 104
 increased, hemodilution effect, 113
 maximization, GDT for
 see goal-directed therapy (GDT)
 optimization, Starling curve, 93, 96
 pulse contour analysis, 84
 “recrutable”, in GDT, 93, 94
 Starling curve *see* Frank–Starling relationship/curve
 stroke volume variation (SVV), 104
 measurement, 86, 97
 monitoring, clinical outcomes, 108
 “Surviving Sepsis Campaign”, 159, 161, 162, 163
 sympathetic activation
 hypovolemic shock, 167, 168
 reduced, hypovolemic shock, 168, 169–70
 in severe hemorrhage, 169, 170
 systemic inflammatory response syndrome (SIRS), 157
 systemic vascular resistance (SVR), 162, 163
 systolic pressure, 38, 85, 86
 systolic pressure variation, 86
 fluid responsiveness in septic shock, 162
 T cells, hypertonic fluid effects, 73–4
 tachycardia, in hypovolemic shock, 170
 temperature-associated adverse effects, 137, 138
 tetrastarch, 142
 thermodilution, calibrated pulse contour analysis by, 85, 86
 thermodilution cardiac output, 83
 transpulmonary, 84
 thermodilution curve, 83, 84
 “third-day” transient circulatory overload, 140
 third space, 23, 29, 67, 97, 134
 uncontrolled hemorrhage, resuscitation, 179
 thyroid surgery, 58, 133, 134
 tidal volume, 106
 tilt table experiments, 166, 167, 171
 timing of fluid administration
 EGDT, 159
 intra-abdominal surgery, 40
 spinal anesthesia, 48
 tissue deposition, adverse reaction, 144–5
 tissue edema *see* edema
 tissue hydration, excessive, 139–40
 tissue oxygenation
 see oxygenation, tissue
 tonicity, 1
 total body water, 127, 129
 children, 65
 tracer methods
 fluid “efficiency” measurement, 130

- fluid volume measurement, 127–9
- limitations of use, 128
- transcapillary escape rate (TER), 120, 121
- increased, 122, 124
- transcapillary fluid exchange, 20, 120–1
- 2-pore theory, 121, 124
- transcapillary/transvenular hydrostatic pressure, 121
- transcervical resection of the endometrium (TCRE), 148, 149
- measurement of fluid absorption, 151
- transcytosis, 121
- transesophageal echocardiography *see* echocardiography
- transfusion-related acute lung injury (TRALI), 15, 190
- transfusion-related cardiac overload (TACO), 189, 190
- transpulmonary thermodilution cardiac output, 84
- transurethral resection of the prostate (TURP), 148, 149
- bipolar resection technique, 153
- glycine solution, absorbed fluid, 149
- irrigation fluid absorption prevention, 153
- transurethral resection (TUR) syndrome, 148
- death from, 151
- fluid absorption measurement, 151–3
- irrigating fluids causing, 150, 151
- pathophysiology, 150–1
- prevention, 153
- symptoms, 149
- treatment, 153–4
- transvascular fluid exchange, 120–1, 122
- see also* microvascular fluid exchange
- trauma
 - crystalloid vs. colloid fluids, 15, 16
 - hemorrhage, 179, 182
 - blood pressure target, 179
 - see also* hemorrhage
 - prevention of coagulation/cascade activation, 20
- tricuspid valve regurgitation, 83
- vagal activity, hypovolemic shock and, 169
- vascular surgery, “rules of thumb” for fluids, 25, 26
- vasoconstriction, in hemorrhage in trauma, 179, 182
- vasodilatation, 3, 133
- vasodilator peptides, 46, 47
- vasopressin, 169
- vasopressors
 - avoidance, plasma volume expansion and, 124
 - in septic shock, 163
 - in spinal anesthesia, 48–9
- Venofundin, 14
- venous oxygen saturation (SvO₂)
 - fluid responsiveness in septic shock, 162
 - hypovolemic shock, 171, 172, 174
 - inter-individual variation, 172
- vesicle transport, 121
- volume effect, infusion fluid, 129
- volume kinetics, 127–36
 - anesthesia/surgery effect, 133
 - clinical use, 134
 - colloid fluids, 133
 - crystalloid fluids, 47, 132–3
 - distribution, compartments, 131, 132
 - elimination, 131, 133
 - “functional” fluid spaces, 132
 - glucose solutions, 134
 - hemoglobin measurements for, 131, 134
 - hypertonic saline, 134
 - modeling, crystalloids distribution, 47
 - “nonfunctional” fluid spaces, 134
 - volunteer studies, 132–3
- volume status, assessment bedside, 82
- invasive *see* invasive hemodynamic monitoring
- volumetric fluid balance, irrigating fluid absorption, 151–2, 153
- Voluven, 14
- von Willebrand factor, 142, 143, 189
- von Willebrand-like syndrome, 142, 143
- warming, infusion fluids, 140, 141
- water isotopes, 127
- water requirements, children, 66
- weight, fluid volumes relationship, 129
- weight gain, 29, 30, 51
 - crystalloid fluids in elderly, 52, 140