

Contents

| | |
|---|------------------|
| <i>Preface</i> | <i>page</i> xvii |
| <i>Acknowledgements</i> | xix |
| 1 Introduction | 1 |
| 1.1 The role of MHD in fusion energy | 1 |
| 1.1.1 The plasma pressure in a fusion reactor | 1 |
| 1.1.2 The dimensionless pressure, β | 4 |
| 1.1.3 A variety of fusion concepts | 4 |
| 1.1.4 Structure of the textbook | 5 |
| 1.2 Units | 5 |
| References | 6 |
| Further reading | 6 |
| 2 The ideal MHD model | 7 |
| 2.1 Introduction | 7 |
| 2.2 Description of the model | 8 |
| 2.3 Derivation of the ideal MHD model | 12 |
| 2.3.1 Starting equations | 12 |
| 2.3.2 Two-fluid equations | 15 |
| 2.3.3 Low-frequency, long-wavelength, asymptotic expansions | 18 |
| 2.3.4 The single-fluid equations | 19 |
| 2.3.5 The ideal MHD limit | 22 |
| 2.4 Region of validity | 28 |
| 2.4.1 Overall criteria | 28 |
| 2.4.2 Conservation of mass | 30 |
| 2.4.3 Momentum equation | 30 |
| 2.4.4 Energy equation | 32 |
| 2.4.5 Ohm's law | 33 |
| 2.4.6 Summary of validity conditions | 34 |
| | vii |

| | | |
|------|--|----|
| viii | <i>Contents</i> | |
| | 2.5 Overall summary | 35 |
| | References | 35 |
| | Further reading | 35 |
| | Problems | 36 |
| 3 | General properties of ideal MHD | 39 |
| | 3.1 Introduction | 39 |
| | 3.2 Boundary conditions | 39 |
| | 3.2.1 Perfectly conducting wall | 40 |
| | 3.2.2 Insulating vacuum region | 40 |
| | 3.2.3 Plasma surrounded by external coils | 43 |
| | 3.3 Local conservation relations | 45 |
| | 3.3.1 Conservation of mass | 45 |
| | 3.3.2 Conservation of momentum | 45 |
| | 3.3.3 Conservation of energy | 46 |
| | 3.4 Global conservation laws | 47 |
| | 3.4.1 Perfectly conducting wall | 48 |
| | 3.4.2 Insulating vacuum region | 49 |
| | 3.4.3 Plasma surrounded by external coils | 51 |
| | 3.5 Conservation of flux: the “frozen-in field line” concept | 51 |
| | 3.6 Summary | 54 |
| | Further reading | 54 |
| | Problems | 55 |
| 4 | MHD equilibrium: general considerations | 58 |
| | 4.1 Introduction | 58 |
| | 4.2 Basic equilibrium equations | 58 |
| | 4.3 The virial theorem | 59 |
| | 4.4 The need for toroidicity | 61 |
| | 4.5 Flux surfaces | 62 |
| | 4.6 Surface quantities: basic plasma parameters and figures of merit | 66 |
| | 4.6.1 Fluxes and currents | 66 |
| | 4.6.2 Normalized plasma pressure, β | 68 |
| | 4.6.3 Kink safety factor, q_* | 70 |
| | 4.6.4 Rotational transform, ι , and the MHD safety factor, q | 70 |
| | 4.6.5 Summary | 72 |
| | 4.7 Equilibrium degrees of freedom | 72 |
| | 4.8 The basic problem of toroidal equilibrium | 73 |
| | 4.9 A single particle picture of toroidal equilibrium | 78 |
| | 4.10 Summary | 80 |

| | <i>Contents</i> | ix |
|---|-----------------|------------|
| References | | 81 |
| Further reading | | 82 |
| Problems | | 82 |
| 5 Equilibrium: one-dimensional configurations | | 85 |
| 5.1 Introduction | | 85 |
| 5.2 The θ -pinch | | 85 |
| 5.3 The Z-pinch | | 90 |
| 5.4 The general screw pinch | | 94 |
| 5.4.1 General properties | | 94 |
| 5.4.2 The parallel pinch | | 98 |
| 5.4.3 The perpendicular pinch | | 100 |
| 5.5 Inherently 1-D fusion configurations | | 102 |
| 5.5.1 The reversed field pinch | | 102 |
| 5.5.2 The low β ohmic tokamak | | 108 |
| 5.6 Summary | | 116 |
| References | | 117 |
| Further reading | | 117 |
| Problems | | 117 |
| 6 Equilibrium: two-dimensional configurations | | 123 |
| 6.1 Introduction | | 123 |
| 6.2 Derivation of the Grad–Shafranov equation | | 124 |
| 6.2.1 The $\nabla \cdot \mathbf{B} = 0$ equation | | 124 |
| 6.2.2 Ampere’s law | | 126 |
| 6.2.3 Momentum equation | | 126 |
| 6.3 Plasma parameters and figures of merit | | 128 |
| 6.3.1 Simple flux coordinates | | 128 |
| 6.3.2 The volume of a flux surface | | 130 |
| 6.3.3 The plasma beta | | 130 |
| 6.3.4 The kink safety factor | | 131 |
| 6.3.5 Rotational transform and the MHD safety factor | | 131 |
| 6.3.6 The MHD safety factor on axis | | 132 |
| 6.3.7 Alternate choices for $F(\psi)$ | | 134 |
| 6.4 Analytic solution in the limit $\varepsilon \ll 1$ and $\beta_p \sim 1$ | | 136 |
| 6.4.1 The coordinate transformation | | 137 |
| 6.4.2 The asymptotic expansion | | 138 |
| 6.4.3 The ε^0 equation: radial pressure balance | | 139 |
| 6.4.4 The ε^1 equation: toroidal force balance | | 140 |
| 6.4.5 Application to early reversed field pinches (RFP) | | 142 |

| | | |
|-------|--|-----|
| 6.4.6 | Application to early ohmic tokamaks and modern reversed field pinches | 144 |
| 6.4.7 | Summary | 151 |
| 6.5 | Analytic solution in the limit $\varepsilon \ll 1$ and $\beta_p \sim 1/\varepsilon$ (the high β tokamak) | 152 |
| 6.5.1 | The high β tokamak expansion | 153 |
| 6.5.2 | The circular high β tokamak | 154 |
| 6.5.3 | The flux conserving tokamak – avoiding the equilibrium β limit | 164 |
| 6.5.4 | The elliptic high β tokamak | 171 |
| 6.6 | Exact solutions to the Grad–Shafranov equation (standard and spherical tokamaks) | 176 |
| 6.6.1 | Mathematical formulation | 177 |
| 6.6.2 | Examples: TFTR and JET | 182 |
| 6.6.3 | Example: the spherical tokamak (ST) | 183 |
| 6.6.4 | The equilibrium β limit | 187 |
| 6.6.5 | Up–down asymmetric solutions | 189 |
| 6.6.6 | Example: the International Thermonuclear Experimental Reactor (ITER) | 192 |
| 6.6.7 | Example: the National Spherical Torus Experiment (NSTX) | 194 |
| 6.6.8 | Summary | 195 |
| 6.7 | The helical Grad–Shafranov equation (the straight stellarator) | 196 |
| 6.7.1 | Overview | 196 |
| 6.7.2 | The helical Grad–Shafranov equation | 197 |
| 6.7.3 | Low β analytic solution | 199 |
| 6.7.4 | The rotational transform | 207 |
| 6.7.5 | Summary | 209 |
| 6.8 | Overall summary | 210 |
| | References | 213 |
| | Further reading | 213 |
| | Problems | 215 |
| 7 | Equilibrium: three-dimensional configurations | 223 |
| 7.1 | Introduction | 223 |
| 7.2 | The high β stellarator expansion | 226 |
| 7.2.1 | Introduction | 226 |
| 7.2.2 | The basic equations | 227 |
| 7.2.3 | The high β stellarator expansion | 228 |
| 7.2.4 | Reduction of the equations | 230 |

| <i>Contents</i> | | xi |
|-----------------|--|-----|
| 7.3 | Relation of the high β stellarator expansion to other models | 236 |
| 7.3.1 | The high β tokamak | 237 |
| 7.3.2 | The straight stellarator | 237 |
| 7.4 | The Greene–Johnson limit | 240 |
| 7.4.1 | Comparison of expansions | 240 |
| 7.4.2 | The Greene–Johnson limit of the HBS model | 241 |
| 7.4.3 | The Greene–Johnson model | 243 |
| 7.4.4 | Summary | 244 |
| 7.5 | Vacuum flux surfaces | 244 |
| 7.5.1 | Single helicity – the limiting helical field amplitude | 244 |
| 7.5.2 | Multiple helicity stellarators | 249 |
| 7.6 | Effects of finite β | 251 |
| 7.6.1 | Low β single helicity solutions | 253 |
| 7.6.2 | Toroidal force balance in a current-free stellarator | 257 |
| 7.6.3 | How does a vertical field shift a stellarator with no net current? | 262 |
| 7.6.4 | The equilibrium β limit in a stellarator | 269 |
| 7.6.5 | The flux conserving stellarator | 273 |
| 7.6.6 | Multiple helicity, finite β stellarators | 275 |
| 7.7 | Neoclassical transport in stellarators | 277 |
| 7.7.1 | Review of transport in a tokamak | 278 |
| 7.7.2 | The problem with neoclassical transport in a stellarator | 282 |
| 7.7.3 | One solution – the omnigenous stellarator | 287 |
| 7.7.4 | The isodynamic stellarator | 292 |
| 7.7.5 | The symmetric stellarator | 297 |
| 7.7.6 | Boozer coordinates | 307 |
| 7.7.7 | Summary of neoclassical transport in a stellarator | 318 |
| 7.8 | Modern stellarators | 319 |
| 7.8.1 | The Large Helical Device (LHD) | 319 |
| 7.8.2 | The Wendelstein 7-X (W7-X) stellarator | 321 |
| 7.9 | Overall summary | 323 |
| | References | 324 |
| | Further reading | 325 |
| | Problems | 325 |
| 8 | MHD stability – general considerations | 327 |
| 8.1 | Introduction | 327 |
| 8.2 | Definition of MHD stability | 328 |
| 8.3 | Waves in an infinite homogeneous plasma | 330 |

| | | |
|--------|--|-----|
| 8.3.1 | The shear Alfvén wave | 332 |
| 8.3.2 | The fast magnetosonic wave | 332 |
| 8.3.3 | The slow magnetosonic wave | 333 |
| 8.3.4 | Summary | 334 |
| 8.4 | General linearized stability equations | 334 |
| 8.4.1 | Initial value formulation | 334 |
| 8.4.2 | Normal mode formulation | 336 |
| 8.5 | Properties of the force operator $\mathbf{F}(\xi)$ | 337 |
| 8.5.1 | Self-adjointness of $\mathbf{F}(\xi)$ | 337 |
| 8.5.2 | The “standard form” of δW | 338 |
| 8.5.3 | The “intuitive form” of δW | 340 |
| 8.5.4 | The “intuitive self-adjoint form” of δW | 342 |
| 8.5.5 | Real ω^2 | 344 |
| 8.5.6 | Orthogonality of the normal modes | 345 |
| 8.5.7 | Spectrum of $\mathbf{F}(\xi)$ | 345 |
| 8.6 | Variational formulation | 347 |
| 8.7 | The Energy Principle | 349 |
| 8.7.1 | Statement of the Energy Principle | 350 |
| 8.7.2 | Proof of the Energy Principle | 351 |
| 8.7.3 | Advantages of the Energy Principle | 354 |
| 8.8 | The Extended Energy Principle | 355 |
| 8.8.1 | Statement of the problem | 356 |
| 8.8.2 | The boundary conditions | 356 |
| 8.8.3 | The natural boundary condition | 358 |
| 8.8.4 | The surface energy | 359 |
| 8.8.5 | The vacuum energy | 360 |
| 8.8.6 | Summary of the Extended Energy Principle | 362 |
| 8.9 | Incompressibility | 363 |
| 8.9.1 | The general minimizing condition | 363 |
| 8.9.2 | Ergodic systems | 364 |
| 8.9.3 | Closed line systems | 365 |
| 8.9.4 | Summary and discussion | 366 |
| 8.10 | Vacuum versus force-free plasma | 367 |
| 8.10.1 | The nature of the problem | 367 |
| 8.10.2 | Vacuum vs. force-free plasma: the same results | 367 |
| 8.10.3 | Vacuum vs. force-free plasma: different results | 368 |
| 8.10.4 | The real situation: a resistive region | 368 |
| 8.11 | Classification of MHD instabilities | 369 |
| 8.11.1 | Internal/fixed boundary modes | 370 |
| 8.11.2 | External/free boundary modes | 370 |

| <i>Contents</i> | | xiii |
|-----------------|--|------|
| 8.11.3 | Pressure-driven modes | 370 |
| 8.11.4 | Current-driven modes | 373 |
| 8.12 | Summary | 375 |
| | References | 376 |
| | Further reading | 377 |
| | Problems | 377 |
| 9 | Alternate MHD models | 381 |
| 9.1 | Introduction | 381 |
| 9.2 | Transition from collision dominated to collisionless regimes | 382 |
| 9.3 | General formulation | 384 |
| 9.4 | Ideal MHD closure | 385 |
| 9.5 | Kinetic MHD | 385 |
| 9.5.1 | Basic assumptions | 385 |
| 9.5.2 | Derivation of the kinetic MHD model | 386 |
| 9.6 | The Chew, Goldberger, Low (CGL) double adiabatic model | 391 |
| 9.6.1 | Formulation of the problem | 392 |
| 9.6.2 | Derivation of the double adiabatic model | 393 |
| 9.6.3 | The modified double adiabatic model | 395 |
| 9.7 | Summary | 396 |
| 9.7.1 | Ideal MHD | 396 |
| 9.7.2 | Kinetic MHD | 396 |
| 9.7.3 | The double adiabatic model | 397 |
| | References | 398 |
| | Further reading | 398 |
| | Problems | 398 |
| 10 | MHD stability comparison theorems | 400 |
| 10.1 | Introduction | 400 |
| 10.2 | Ideal MHD equilibrium and stability | 401 |
| 10.3 | Double adiabatic MHD equilibrium and stability | 402 |
| 10.3.1 | Relation between the perturbed quantities and ξ | 403 |
| 10.3.2 | The double adiabatic MHD Energy Principle | 404 |
| 10.3.3 | Summary of CGL stability | 406 |
| 10.4 | Kinetic MHD equilibrium and stability | 406 |
| 10.4.1 | Equilibrium | 407 |
| 10.4.2 | Stability of a closed line cylindrical system | 408 |
| 10.4.3 | Stability of an ergodic cylindrical system | 411 |
| 10.4.4 | Stability of a general toroidal configuration | 414 |
| 10.5 | Stability comparison theorems | 422 |
| 10.5.1 | Closed line cylindrical geometry | 422 |

| | | |
|-----|--|-----|
| xiv | <i>Contents</i> | |
| | 10.5.2 Ergodic cylindrical geometry | 423 |
| | 10.5.3 Closed line toroidal geometry | 424 |
| | 10.5.4 Ergodic toroidal geometry | 425 |
| | 10.6 Summary | 426 |
| | References | 427 |
| | Further reading | 427 |
| | Problems | 427 |
| 11 | Stability: one-dimensional configurations | 428 |
| | 11.1 Introduction | 428 |
| | 11.2 The basic stability equations | 430 |
| | 11.2.1 The Energy Principle | 430 |
| | 11.2.2 The normal mode eigenvalue equations | 430 |
| | 11.2.3 Incompressible MHD | 431 |
| | 11.3 Stability of the θ -pinch | 431 |
| | 11.3.1 Application of the Energy Principle to the θ -pinch | 432 |
| | 11.3.2 Minimizing δW_F for a θ -pinch | 433 |
| | 11.3.3 Continuum damping in a “slab” θ -pinch | 434 |
| | 11.4 Stability of the Z-pinch | 443 |
| | 11.4.1 Energy Principle analysis of $m \neq 0$ modes | 444 |
| | 11.4.2 Energy Principle analysis of the $m = 0$ mode | 447 |
| | 11.4.3 Double adiabatic Energy Principle analysis of the $m = 0$ mode | 450 |
| | 11.4.4 The hard-core Z-pinch | 452 |
| | 11.5 General stability properties of the screw pinch | 458 |
| | 11.5.1 Evaluation of δW for a general screw pinch | 459 |
| | 11.5.2 Suydam’s criterion | 465 |
| | 11.5.3 Newcomb’s procedure | 470 |
| | 11.5.4 The normal mode eigenvalue equation | 477 |
| | 11.5.5 The oscillation theorem | 482 |
| | 11.5.6 The resistive wall mode | 486 |
| | 11.6 The “straight” tokamak | 494 |
| | 11.6.1 Reduction of δW for the straight tokamak | 495 |
| | 11.6.2 Sawtooth oscillations – the internal $m = 1$ mode | 497 |
| | 11.6.3 Current-driven disruptions – external low m modes | 500 |
| | 11.6.4 Density-driven disruptions – external low m modes | 512 |
| | 11.6.5 Resistive wall modes – external low m modes | 518 |
| | 11.6.6 Edge localized modes (ELMs) – external high m modes | 519 |
| | 11.6.7 Summary of the straight tokamak | 527 |

| <i>Contents</i> | | xv |
|-----------------|--|-----|
| 11.7 | The reversed field pinch (RFP) | 529 |
| 11.7.1 | Physical parameters describing an RFP | 530 |
| 11.7.2 | Instability of an RFP without a B_z reversal | 532 |
| 11.7.3 | The $m = 0$ instability | 534 |
| 11.7.4 | Suydam's criterion | 535 |
| 11.7.5 | Internal modes | 537 |
| 11.7.6 | Taylor's theory | 538 |
| 11.7.7 | Ideal external modes | 552 |
| 11.7.8 | Overview of the RFP | 557 |
| 11.8 | Summary | 558 |
| | References | 562 |
| | Further reading | 562 |
| | Problems | 563 |
| 12 | Stability: multi-dimensional configurations | 570 |
| 12.1 | Introduction | 570 |
| 12.2 | Ballooning and interchange instabilities | 572 |
| 12.2.1 | Introduction | 572 |
| 12.2.2 | Shear, periodicity, and localization | 574 |
| 12.2.3 | General reduction of δW for ballooning modes | 578 |
| 12.3 | The ballooning mode equations for tokamaks | 581 |
| 12.4 | The ballooning mode equation for stellarators | 585 |
| 12.5 | Stability of tokamaks – the Mercier criterion | 588 |
| 12.5.1 | Introduction | 588 |
| 12.5.2 | Cylindrical limit: the Suydam criterion | 589 |
| 12.5.3 | Toroidal geometry: the Mercier criterion | 591 |
| 12.5.4 | Analytic limits of the Mercier criterion | 596 |
| 12.5.5 | Summary | 598 |
| 12.6 | Stability of tokamaks – ballooning modes | 598 |
| 12.6.1 | Introduction | 598 |
| 12.6.2 | The s - α model for ballooning modes | 599 |
| 12.6.3 | β limits due to ballooning modes | 605 |
| 12.6.4 | Summary | 608 |
| 12.7 | Stability of tokamaks – low n internal modes | 608 |
| 12.7.1 | Introduction | 608 |
| 12.7.2 | Low n internal modes with finite shear | 609 |
| 12.7.3 | Low n internal modes with small shear | 610 |
| 12.7.4 | Summary | 611 |
| 12.8 | Stability of tokamaks – low n external ballooning-kink modes | 611 |
| 12.8.1 | Introduction | 611 |

| | | |
|---------|--|-----|
| 12.8.2 | Simplification of δW_F by the high β tokamak expansion | 613 |
| 12.8.3 | High β stability of the surface current model | 615 |
| 12.8.4 | Numerical studies of ballooning-kink instabilities | 627 |
| 12.9 | Stability of tokamaks – advanced tokamak (AT) operation | 630 |
| 12.9.1 | Introduction | 630 |
| 12.9.2 | Bootstrap current profile – the β_N limit | 632 |
| 12.9.3 | Wall stabilization in an advanced tokamak | 635 |
| 12.9.4 | “Infernal” modes | 636 |
| 12.10 | Stability of tokamaks – $n = 0$ axisymmetric modes | 637 |
| 12.10.1 | Introduction | 637 |
| 12.10.2 | Vertical instabilities in a circular plasma | 637 |
| 12.10.3 | Horizontal instabilities in a circular plasma | 640 |
| 12.10.4 | Vertical instabilities in an elongated plasma | 641 |
| 12.11 | Overview of the tokamak | 647 |
| 12.12 | Stellarator stability | 649 |
| 12.12.1 | High n modes in a stellarator | 650 |
| 12.12.2 | The parallel current constraint | 651 |
| 12.12.3 | The relation between average curvature and magnetic well | 652 |
| 12.12.4 | Average curvature of a straight helix | 656 |
| 12.12.5 | Shear stabilization of a straight helix | 658 |
| 12.12.6 | Stabilization of a toroidal stellarator | 659 |
| 12.12.7 | Numerical results | 665 |
| 12.13 | Summary | 666 |
| 12.14 | The final word | 669 |
| | References | 670 |
| | Further reading | 671 |
| | Problems | 672 |
| | <i>Appendix A Heuristic derivation of the kinetic equation</i> | 678 |
| | <i>Appendix B The Braginskii transport coefficients</i> | 688 |
| | <i>Appendix C Time derivatives in moving plasmas</i> | 691 |
| | <i>Appendix D The curvature vector</i> | 695 |
| | <i>Appendix E Overlap limit of the high β and Greene–Johnson stellarator models</i> | 697 |
| | <i>Appendix F General form for $q(\psi)$</i> | 706 |
| | <i>Appendix G Natural boundary conditions</i> | 707 |
| | <i>Appendix H Upper and lower bounds on δQ_{KIN}</i> | 711 |
| | <i>Index</i> | 718 |