

Index

- abdominal injuries
 epidemiology, 563–5
 imaging strategies, 599–600
 lower rib fractures, 186–7, 190, 596
 lumbar spine injuries, 532, 535
 mortality, 565
 visceral, 565–89
- abdominal radiography, 599
 duodenal hematoma, 571
 pancreatic injuries, 578
 small bowel perforation, 568–9
- absorbed dose, radiation, 669–70
- abusive head trauma (AHT), 343, 345–53
 accidental vs., 351–3, 403, 423–4, 478–9
 biomechanics, 347–8
 cervicospinal junction and spine, 494–553
 clinical diagnosis, 346–7
 cost to society, 353
 editor's note, 343
 epidemiology, 346
 extra-axial hemorrhage/
 nonhemic collections, 394–441
 facial injuries, 375–82
 imaging patterns, 363–70
 imaging protocol, 491
 imaging strategies, 386–8, 487–92
 legal challenges to expert testimony, 648–51
 media scrutiny of legal proceedings, 650
 parenchymal brain injury, 453–81
 retinal hemorrhage, 350–1
 scalp, subscalp, and cranium, 357–88
 shaking as mechanism of injury, 349–50
 skull fractures. *See* skull fractures
 soft tissue injuries, 363–4
 strength of evidence, 315–16
 terminology and history, 345–6
- accidental trauma, 268–75,
See also falls
 head, 351–3, 403, 423–4, 478–9
 imaging findings, 272–5
 lower extremity, 272
 perineal region, 599
 ribs, 274–5
 skull fractures, 372–4, 382–3
 sternum, 275
 subdural hemorrhage, 421, 423–4
 upper extremity, 273–4
- acetabular fractures, 101, 117
- acoustic impedance, 680
- acromial fractures
 imaging features, 199–203
 mechanisms of injury, 198
 normal variants vs., 299
 projectional effect mimicking, 300, 303
 seizure disorders causing, 198, 219, 221
- acromion process, 197–8
 developmental variant, 300, 303
 ossification variants, 298–301
- adrenal injuries/hemorrhage, 574, 581, 584–6, 598
- adversarial allegiance, culture of, 320
- age of injury. *See* dating
- age, patient
 callus formation and, 212
 dating fractures and, 210–11
 inflicted skeletal trauma and, 27, 209
- AHT. *See* abusive head trauma
- alanine aminotransferase (ALT), serum, 581
- alar ligaments, 505–6
- American Academy of Pediatrics (AAP)
 global skeletal imaging recommendations, 332
 skull radiography guidelines, 386
- terminology of abusive head trauma, 346
- American College of Radiology (ACR), 684
 global skeletal imaging recommendations, 332
- imaging abusive head trauma, 487
- skull radiography guidelines, 386
- standards for skeletal surveys, 325, 336
- American Health Information Management Association (AHIMA), 641
- amylase, serum, 578
- anal dilation, reflex, 621
- anal injuries, 576–7
- aneurysms, traumatic
 intracranial, 474–5
- angioneurotic edema, 364
- anogenital injuries
 accidental, 599
 sexual abuse, 620–1
- anoxic brain injury, 467, 469–70
- anterior longitudinal ligament, 508, 532
- anticoagulated patients,
 intracranial hemorrhage, 431
- anti-scatter grid, 681–2
- anxiety, giving evidence in court, 662
- aorta
 pseudoaneurysm, 597–8
 transection, 544, 546, 597
- apical ligament, 505–6, 508
- apnea
 factitious, 618
 trauma-induced, 471–2, 523–5
- apophyseal bone, development, 16
- apoptosis, brain cells, 456
- apparent diffusion coefficient (ADC), 696
- apparent diffusion coefficient (ADC) maps, 704
 anoxic brain injury, 470
 cerebral contusions, 463
 cerebral edema, 459
- delayed necrosis, 455
 hyperemia, 471
 ischemic brain injury, 473
 normal, acute/subacute phase, 457
 parenchymal brain lacerations, 467–8
 subdural effusion, 419
- arachnoid cysts, 431
- arachnoid mater, 395–6
- arterial spin labeling (ASL), 490–1, 705
 cardiac arrest, 457
 hyperemia, 469
 ischemic brain injury, 473
 near hanging, 470
- arteriovenous fistulas (AVFs), cerebral, 436
- arteriovenous malformations (AVMs), cerebral, 432, 436
- arthrography
 distal humeral physeal injuries, 139
 proximal femoral injuries, 55, 59, 284
- arthrogryposis
 birth-related fractures, 279, 286
 orthopedic manipulations, 223, 225
- ascites
 abdominal injuries, 578, 584
 chylous, 579, 585–6
- aspartate aminotransferase (AST), serum, 581
- atelectasis, 595
- atlanto-axial ligament, 506
- atlanto-occipital ligaments, 505–6, 508
- atlanto-occipital subluxation/
 dislocation, 513
- atlas (C1), 504–5
- attachments, parent–child, 634
- attenuation, radiation, 669
- attorneys. *See* lawyers
- autophagy (autophagocytosis), 456

- axis (C2), 505
congenital spondylolysis, 547–8, 550
spondylosis fracture, 509, 513–15
traumatic spondylolysis. *See* hangman's fracture
- axonal stretch injury, traumatic.
See diffuse axonal injury
- baby bouncers, 273
- baby box, patient immobilization, 713
- baby walker-related falls, 478
- babygram, 325, 709
inadequacy, 325, 336, 712
- bacteremia, induced, 618–19
- banding artifact, MRI, 698, 703
- barium upper gastrointestinal (uGI) series. *See* upper gastrointestinal (uGI) series
- battered child syndrome, 1, 3, 23
- batteries, ingested, 615
- beak, metaphyseal, 294
- Becquerel, Antoine Henri, 668
- benign enlargement of subarachnoid spaces in infancy (BESSI), 418, 420–3
- b-factor, diffusion imaging, 703
- bias
clinical research, 313–14, 319
expert medical testimony, 663
- bile duct injuries, 582
- biomechanics
abusive head trauma, 347–8
accidental head trauma, 351–3
craniocervical junction/cervical spine injuries, 508–12
retinal hemorrhages, 350–1
shaking of infants, 349–50
skull fractures, 360–3
subdural hemorrhage, 407–8
- birth injuries. *See* obstetric trauma
- bladder
hypertrophy, 588
needle implantation, 588
rupture, 587–8
- Bloch equations, 690
- Block, Robert W., 650
- Bloembergen, Nicolaas, 695
- blood vessels, bones, 13
- blood volume, MRI, 704–5
- blue sclerae, 255
- bone, 9–13
anatomy, 9–13
cancellous, 10, 12
cortical, 10–12
formation and growth, 15–20
lamellar, 10–11
modeling, 20
remodeling, 20, 213–14
sites vulnerable to trauma, 21 types, 10
vascular and nerve supply, 13
woven, 10–11
x-ray imaging, 672
- bone age, delayed, 610
- bone cells, 13–14
- bone demineralization, 240.
See also osteopenia
- fracture risk, 248
- osteogenesis imperfecta, 254–7, 260, 262–3
- premature infants, 244
- bone densitometry, 248
- osteogenesis imperfecta, 263
- bone dysplasias, inherited, 217–18
- bone marrow, MRI, 698
- bone mineral density (BMD) fracture risk and, 248
“temporary brittle bone disease”, 318
- bone scintigraphy, 330.
See also positron emission tomography
- availability, 329
- child in a cast, 331
- clavicular fractures, 193–4
- cost, 330
- distal femoral epiphyseal separation, 71, 81
- distal humeral physeal injuries, 139, 145
- distal radial fractures, 146, 153
- femoral shaft fractures, 63, 65, 70
- global skeletal assessment, 325
- hand fractures, 151, 159–60
- humeral fractures, 133, 135–6, 148
- interpreter dependency, 330
- mandibular fractures, 381, 385
- need for additional imaging, 330
- radiation dose, 329, 680
- radiographic skeletal surveys vs., 328–30
- radius and ulna shaft fractures, 143, 145, 153
- recommendations, 332
- renal abnormalities, 586–7
- rib fractures, 180, 182–4, 186, 189
- scapular fractures, 200
- sedation requirements, 329
- sensitivity, 329
- skull fractures, 362, 388
- soft tissue injuries, 54, 57
- specificity, 329
- spinal trauma, 527–8, 531, 534–5, 551
- SPNBF, 30, 32
- technical factor dependency, 330
- tibial shaft fractures, 97
- bowed legs, physiologic, 295–7
- bowel injuries, 567–77
- Bradford Hill criteria, 311
- brain
catch-up growth, 364–5, 611
- contusions. *See* cerebral contusions
- liquefaction, 456–8, 468
- MRI techniques, 697–8, 701, 704–5
- parenchymal injuries, 453–81
accidental vs. abusive, 478–9
differential diagnoses, 478–81
direct mechanical, 461–6
follow-up imaging, 457–8, 478
normal CT and MRI, 478
outcome, 481
pathophysiologic principles, 454–61
patterns indicating abuse, 477
role of repetitive injury, 476
secondary, 467–76
parenchymal lacerations, 464–6, 468
- brainstem injuries, 517, 522–9.
See also craniocervical junction (CCJ) injuries
- mechanisms, 525–7
- pathology, 522–5
- retroclival epidural hematomas and, 530
- breast calcification, 619–20, 622
- bridging veins, cerebral cortical, 395–6, 409–10
- thrombosis, 410–11
- torn, 404, 409–11
- bright cerebellum sign (reversal sign), 467, 469
- bright fat effect, fast spin echo (FSE), 701
- brittle bone disease. *See* osteogenesis imperfecta
- brittle bone disorders, other, 257
- Bruck syndrome, 257
- bruising. *See also* soft tissue injuries
- abusive head trauma, 346–7
- facial, 375–8
- immune-thrombocytopenic purpura, 434–5
- sexual abuse, 622
- bucket handle fractures, 35, 42
- distal femur, 67, 69–70, 73–4
- distal humerus, 141
- distal radius and ulna, 146, 155–7
- distal tibia, 89, 92, 99, 101, 103, 105–6
- fibula, 94, 109
- histologic studies, 37, 40
- proximal femur, 59, 61
- proximal humerus, 124, 130
- proximal tibia, 75–9, 85–6, 89
- rib fractures resembling, 179, 182
- buckle fractures, cortical, 31–2
- distal humerus, 147
- distal radius, 145, 153
- femoral shaft, 64, 67
- fibula, 95–6
- first metatarsal, 97, 110
- hand phalanges, 152, 159
- proximal humeral metaphysis, 134, 147
- proximal radius, 141, 150
- proximal tibial metaphysis, 82, 91
- ribs, 170, 176
- scapula, 203
- supracondylar distal humerus, 137, 139
- bunk beds, falls from, 479
- burden of proof, court proceedings, 646
- burns, 620–1
- C1 (atlas), 504–5
- C1–2 capsular ligaments, 506–7
- C2. *See* axis
- cadaveric studies. *See* postmortem research
- Caffey, John, 2, 23, 34–5, 345–6
- Caffey–Kempe syndrome, 346
- Caffey disease, 218, 230–1
- calcification. *See also* heterotopic bone formation
- fat necrosis and, 619–20, 622
- necklace distribution, 619
- paraspinal, 542, 544, 594
- precervical, 594
- calcifying cartilage zone of growth plate. *See* mineralizing zone of growth plate
- calcium
homeostasis, 240–3
metabolism, disorders of, 240–9
- calibration, radiographic equipment, 709–12
- callus
clavicular fractures, 193, 280
- dating fractures, 209, 214–15
- femoral shaft fractures, 64, 66, 68
- formation, 212
- humeral shaft fractures, 135–6
- obstetric trauma, 280–1
- osteogenesis imperfecta, 257, 262
- rib fractures, 168, 173, 176–7, 182
- scapular fractures, 201, 203
- calvaria, 357
- canaliculi, bone, 13
- cancellous bone, 10, 12
- capillary telangiectasia, intracranial, 436
- capsular ligament, C1–2, 506–7
- caput succedaneum, 363–4

Index

- cardiac arrest, brain injuries, 457, 468–9
 cardiac enzymes/troponins, 597
 cardiac injuries, 597, 617–18
 cardiopulmonary resuscitation (CPR), injuries due to, 189, 598
 cardiovascular trauma, 597
 care and protection cases, 662
 caregivers. *See* parents
 cartilage
 endochondral ossification, 15
 tidemark, 20
 cartilage anlage, 15
 cartilage hair hypoplasia, 218, 221
 casts, skeletal imaging of children in, 330–1
 causation, 311
 Bradford Hill criteria, 311
 evidence-based radiology, 312
 caustic ingestions, forced, 615–16
 cavernous malformations
 intracranial, 436
 multiple, 436
 celiac disease, 613
 cell death, mechanisms, 454–8
 central nervous system (CNS)
 infections, 436, 476, 480
 metabolic disorders involving, 434, 437–8, 480
 cephalohematoma (subperiosteal hematoma), 363–4, 372
 birth-related, 374–5, 384
 cerebellar tonsillar herniation, 518
 cerebral contusional clefts, 464
 cerebral contusional tears, 464
 cerebral contusions, 454, 461–2
 cerebral edema, 458–60
 abusive head trauma, 471
 acute necrosis, 454
 cytotoxic, 458–9
 hyperemic, 469–71
 hypernatremic dehydration, 437–8
 interstitial, 459–60
 intramyelinic, 458–9
 malignant unilateral, mixed
 attenuation SDH, 425
 severe diffuse, accidental head injury, 479
 vasogenic, 458–9
 cerebral hemorrhage,
 parenchymal.
 See intracerebral hemorrhage
 cerebral palsy, 218, 222–4
 cerebral perfusion, 469–71
 cerebral sinovenous thrombosis (CSVT), 431, 435–6
 cerebral swelling. *See* cerebral edema
 cerebral venous anomalies,
 developmental, 436
 cerebral venous infarcts, 474–5
 cerebrospinal fluid (CSF)
 MRI, 697
 volume depletion, subdural
 hemorrhage, 427, 431–2
 cervical spine injuries, 501–31
 extra-axial collections, 513,
 515–22
 fractures, dislocations, and disk
 herniations, 509, 511,
 513–15
 hyperflexion fracture-
 dislocations, 512–13,
 515
 imaging strategies, 551–3
 importance of timely
 diagnosis, 501–4
 ligamentous strains and
 disruptions, 508, 512–13
 mechanisms of injury,
 508–12
 pseudosubluxation, 515
 relevant anatomy and
 function, 504–8
 retroclival collections, 518–20,
 529–31
 skeletal and ligamentous,
 508–15
 spectrum, 508
 spinal cord, 522–9
 clinical issues, 527–8
 imaging, 517, 528–9
 mechanisms, 525–7
 pathology, 522–5
 cervical spondylolysis,
 congenital, 547–8
 cesarean section, 279
 clavicular fractures, 279
 femoral fractures, 283–6
 humeral fractures, 281–3
 Chance-type fractures, 532,
 541–5
 chemical shift selective
 suppression (CHESS),
 700, 702
 chest radiography, 326, 596
 pharyngeal perforation, 592–3
 chest trauma. *See* thoracic
 trauma
 child abuse
 defined, 1
 incidence and demographics,
 1–2
 child abuse pediatricians (CAPs),
 633, 660
 child abuse pediatrics, 1, 633
 Child Abuse Prevention and
 Treatment Act
 (CAPTA), 1
 child protection
 aiding decision-making,
 324–5
 courtroom testimony, 662
 reporting statutes, 641–3
 state criminal statutes, 646
 Child Protection Teams (CPTs),
 633, 636, 638
 consultation with, 642
 Child Protective Services (CPS)
 contacts with, 660
 disclosure of information to,
 640–1
 reporting suspected abuse,
 642–3
 children. *See also* minors
 characteristics of abused,
 633–4
 consent by, 639–40
 immobilization, skeletal
 surveys, 329, 713–14
 interaction with at-risk, 636
 records retention, 641
 risk factors for abuse, 634
 chiropractic manipulations, 189,
 193, 275
 choking, dysphagic, 441, 471
 chondrocytes
 bone formation, 15–16
 growth plate, 17–19
 chondro-osseous junction (COJ),
 classic metaphyseal
 lesion, 38, 40–2
 chronic fractures, dating, 211–12
 chronic subdural hemorrhage
 (cSDH), 416–17, 425–31
 dating, 398–400, 417
 differential diagnosis, 430–1
 imaging features, 414–15, 417,
 423, 426–9
 internal membranes, 414–15,
 423, 427, 429
 neomembrane formation, 417,
 425, 427
 origins, 417, 422, 425–6
 re-bleeding, 428–30
 innocent or trivial causes,
 425–6, 429–30
 re-injury/inflicted trauma,
 426, 430
 chronically ill children, 608–9
 chylothorax, 596
 chyloous ascites, 579, 585–6
 civil legal proceedings, 645–6
 burden of proof, 646
 courtroom testimony, 662
 pre-trial depositions, 651–2
 classic metaphyseal lesion
 (CML), 34–47
 age of injury, 39–42, 45–6
 anatomic site, 39
 as manifestation of rickets,
 247
 distal femur, 65–71, 73–4, 76,
 78
 distal humerus, 141, 149
 distal radius and ulna, 146–8,
 155–6
 distal tibia, 88–93, 99–100,
 103–5
 distribution, 34, 36, 72
 epiphyseal separations vs.,
 47–8
 femoral shaft fractures
 accompanying, 65, 71
 fibula, 94–6, 109
 fracture extent, 39, 43–4
 histologic studies, 35–7
 mechanisms of injury, 43–4
 normal metaphyseal step-off
 contour vs., 293
 obstetric trauma, 281–6
 prevalence, 34–5
 proximal femur, 59, 61
 proximal humerus, 124–30
 proximal tibia, 73–9, 83, 85–6,
 88
 radiographic projection, 39, 43
 radiographic views, 325–6
 radiologic correlates of
 healing, 42, 44–5
 radiologic–pathologic
 correlates, 36–40
 clavicle, 190
 congenital pseudoarthrosis, 306
 inflicted fractures, 189–95
 imaging features, 192–5
 mechanisms of injury, 190–3
 normal variants, 306–7
 obstetric trauma, 195, 213–15,
 279–80
 clay shoveler's fracture, 536
 clinical history. *See* history,
 clinical
 club foot deformity, serial casting
 for, 223, 225
 CML. *See* classic metaphyseal
 lesion
 coagulopathies, intracranial
 hemorrhage, 430, 432–4
 codfish vertebrae, osteogenesis
 imperfecta, 262
 cognitive biases, 313–14
 Cohen, Jonathon, 208
 coherent scatter, 669
 cohort studies, 310
 coiled-spring appearance, small
 bowel hematomas, 571–2,
 574–5
 coins, ingested, 613–14
 COL1A1/COL1A2 gene
 mutations, 258–60
 collagen fibers
 bone, 10
 tendoligamentous, 13
 collagen type I, 258–9, 261
 defects in osteogenesis
 imperfecta, 259–60
 collimation, x-ray beam, 668,
 670, 712
 colonic injuries, 576–7
 comminuted fractures
 femoral shaft, 63, 66
 skull, 364, 370
 Commonwealth v. Martin, 648–9
 commotio cordis, 597

- Compton scatter, 669
 computed radiography (CR),
 675–6, 684
 radiation dose management,
 682
 skeletal surveys, 709, 712
 computed tomography (CT),
 677–9
 abusive head trauma, 487–9
 adrenal injuries, 585
 angiography (CTA),
 neurovascular injury, 489
 benign enlargement of
 subarachnoid spaces in
 infancy, 418, 423
 chest trauma, 596
 classic metaphyseal lesion,
 37–40, 44
 clinical role, 330
 craniocervical and cervical
 spine injuries, 551–2
 bony and ligamentous,
 510–11, 513, 515
 retroclival hematoma, 519,
 531
 spinal subdural hemorrhage,
 522
 epidural hemorrhage, 401–3
 femoral injuries, 60–1, 66
 future developments, 684
 helical, 679
 humeral injuries, 124, 127, 131
 hypoperfusion complex, 570,
 582, 584, 598
 intracranial extra-axial
 hemorrhage, 397–8
 kidney injuries, 586–7
 liver injuries, 581–2
 mandibular fractures, 381
 multi-detector (MDCT),
 678–9
 pancreatic injuries, 578–9
 parenchymal brain injuries
 contusions, 462–3
 diffuse axonal injury, 464
 follow-up, 457–8
 lacerations, 464, 466
 necrotic cell death, 454–6
 normal findings, 478
 postmortem imaging, 335, 337
 pseudo-subarachnoid
 hemorrhage, 406
 quantitative (QCT), 248
 radiation dose, 671, 682–4
 Effective Dose (D_{eff}), 683–4
 metrics, 683–4
 reduction strategy, 489, 679,
 682–3
 rib fractures, 174, 180, 182,
 184, 188, 190, 274
 rib variants, normal, 302–4
 scalp injuries, 362–3, 369–70
 skull fractures, 364, 387
 small bowel hematoma, 572–5
 small bowel perforation, 569–70
 specimen high-resolution, 337
 splenic injuries, 583–4
 sternal injuries, 196–7
 subarachnoid hemorrhage,
 404–6
 subdural hemorrhage, 397,
 408, 411, 488
 chronic, 398–9, 414–15, 417,
 424, 427–8
 dating, 398
 mixed attenuation, 398, 422,
 424–5
 subdural nonhemic
 collections, 413–16,
 419
 thoracolumbar spine trauma,
 532, 540
 tibial CML, 92, 103
 visceral trauma, 564, 599
 whole body, 326
 computed tomography (CT)
 number, 678
 computed tomography dose
 index (CTDI), 683
 concussion, 476
 confessions, by perpetrators, 349
 confidentiality, medical records,
 640–1
 congenital insensitivity to pain,
 218, 221–3
 consent, informed, 638–40
 contrast discrimination
 CT, 678
 x-ray image, 672–4
 contrast studies, oral, 600
 contrecoup contusion, 461
 contusion, brain. *See* cerebral
 contusions
 contusional clefts, brain, 464
 contusional tears, brain, 464
 copper deficiency, 218, 233–4
 coracoid fractures, 198, 202–3
 Cormack, Allan, 677
 corner fractures, 35, 39, 42
 distal femur, 67, 69–70, 73, 75
 distal radius and ulna, 146–7,
 155
 distal tibia, 92, 100–1, 103–4
 fibula, 94
 first metatarsal, 111
 histopathologic studies, 37–8
 proximal femur, 59, 61
 proximal humerus, 124–5, 127,
 130
 proximal tibia, 75–9, 85–6,
 89
 spondylometaphyseal
 dysplasia (SMD) with,
 217, 219
 corporal punishment, shock due
 to, 620
 cortical bone, 10–12
 normal variants, 293, 298–300
 cortical buckle fractures.
 See buckle fractures
 cortical laminar necrosis, 460–1
 cortical thickening
 after intestinal transplantation,
 231
 Caffey disease, 230–1
 radial and ulnar shafts, 145,
 152
 costochondral junction (CCJ)
 bulbous configuration, 301–2,
 304
 fractures, 172, 176–9, 182–3,
 186–7
 costovertebral articulations,
 170
 rib fractures near, 168, 170, 187
 coughing, paroxysmal, 441, 471
 coup contusion, 461
 court order, for records, 646–7
 courtroom testimony, 654–6,
 662–3. *See also* expert
 medical testimony
 coxa vara deformity, 55,
 57–9
 cranial injuries, 358–60,
 See also scalp injuries;
 skull fractures
 imaging patterns, 364–5
 repeat, 369, 375
 cranial meninges, 395–6
 cranial sonography, 487
 cranial injuries, 375, 384, 388
 parenchymal brain injuries,
 456
 parenchymal brain lacerations,
 464–5
 subarachnoid hemorrhage, 406
 cranial sutures, 357, 359
 biomechanics of fractures,
 360–1
 fissures intersecting, 370–1
 normal variants, 372, 382
 splitting/widening
 catch-up brain growth,
 364–5, 611
 increased intracranial
 pressure, 364, 374
 vitamin A intoxication,
 228–9
 traumatic diastasis, 365, 373
 craniocervical junction (CCJ)
 injuries, 501–31,
 See also neck injuries
 extra-axial collections, 515–22
 fractures, dislocations, and disk
 herniations, 513–15
 imaging strategies, 551–3
 importance of timely
 diagnosis, 501–4
 ligamentous strains and
 disruptions, 508, 512–13
 mechanisms of injury, 508–12
 medullary and spinal cord
 injuries, 522–9
 relevant anatomy and
 function, 504–8
 retroclival collections, 518–20,
 529–31
 skeletal and ligamentous,
 508–15
 spectrum, 508
 craniofacial ratio, 358
 cranium. *See* skull
 criminal legal proceedings,
 645–6
 burden of proof, 646
 courtroom testimony, 662
 witness depositions, 652
 cross-examination, expert
 witnesses, 655–6
 cruciate ligament of atlas, 505–8
 crushing injuries, head trauma,
 475, 479
 CT. *See* computed tomography
 CT number, 678
 cultural issues, 635
 Curie, Pierre and Marie, 668
 cut back zone, 20
 dates, courtroom appearances,
 661–2
 dating
 fractures, 208–15,
 See also fracture dating
 intracerebral hemorrhage, 399,
 466
 subdural hemorrhage,
 397–400, 417, 422, 426–7
 Daubert hearings, 648–9
 Davis, Christopher, 648
 defendants, 646. *See also*
 perpetrators
 aiding assessment of intent,
 647
 deferoxamine, 234
 definition of child abuse, 1
 deformities, residual. *See* growth
 disturbances/residual
 deformities
 dehydration, hypernatremic,
 437–8
 demarcation problem, 319–20
 demineralization. *See* bone
 demineralization
 demographics, child abuse, 1–2
 dentinogenesis imperfecta (DI),
 254, 256–7
 depositions, expert witness,
 651–2
 detective quantum efficiency
 (DQE), 681–2
 developmental dysplasia of hip
 (DDH), 55, 59
 diagnosis of child abuse
 additional imaging after, 661
 challenges to, 3–4
 evidence-based approach,
 310–13
 hierarchy of efficacy, 312
 medical, vs. medico-legal, 324
 radiology report, 659

Index

- diaphysis, 10
 growth and development, 20
 diets, fad, 611–12
 diffuse axonal injury (DAI),
 462–4
 accidental head trauma, 479
 diffuse correlation spectroscopy,
 492
 diffusion imaging, MRI,
 703–4
 diffusion tensor imaging (DTI)
 basic principles, 696–7, 704
 diffuse axonal injury, 464
 diffusion-weighted imaging (DWI),
 490. *See also* apparent
 diffusion coefficient
 (ADC) maps
 anoxic brain injury, 470
 basic principles, 696–7, 703–4
 cerebral contusions, 461
 delayed necrosis, 455
 diffuse axonal injury, 465
 near hanging, 470
 normal, acute/subacute phase,
 457
 subdural effusion, 419, 476
 DiGeorge syndrome, acromial
 variant, 300, 303
 digital radiographic technologies,
 675–7
 flat-panel systems. *See* digital
 radiography
 future developments, 684
 radiation dose management,
 682
 selection of best detector,
 676–7
 skeletal surveys, 709
 stimutable phosphor
 technique. *See* computed
 radiography
 digital radiography (DR), 675–6
 direct flat-panel detectors, 676
 indirect flat-panel detectors,
 675–6
 radiation dose management,
 682
 skeletal surveys, 709, 712
 system selection, 676–7
 digital subtraction angiography
 (DSA), 677
 digital tomosynthesis (DT), 684
 1,25-dihydroxyvitamin
 D (1,25D₂), 240–2
 disabled children, 608–10, 634
 discitis, inflammatory, 549
 discovery, pre-trial, 646–7, 651
 disprovability (falsifiability) test,
 319–20
 distal femur, 65–71
 birth-related injuries, 283–6
 classic metaphyseal lesion,
 65–71, 73–4, 76, 78
 irregularities in toddlers,
 295–8
 metaphyseal spur, 293
 metaphyseal step-off
 configuration, 294
 nonspecific metaphyseal
 fractures, 65, 72
 normal anatomy, 65–7, 73
 physeal injuries and epiphyseal
 separations, 70–1, 80–1
 distal forearm, 145–8, 153
 buckle fractures, 145, 153
 classic metaphyseal lesion,
 146–8, 155–6
 normal anatomy, 146, 154
 physeal injuries, 148, 157
 distal humerus, 136–41
 accidental trauma, 273
 classic metaphyseal lesion, 141,
 148–9
 obstetric trauma, 281–3
 lateral condylar fractures, 140,
 147, 273
 medial condylar fracture, 140,
 146
 physeal injuries and epiphyseal
 separations, 138–41,
 144–6
 birth-related, 281–4
 residual deformities, 140,
 146
 supracondylar fractures,
 137–9, 273–4
 distal tibia, 88–93
 classic metaphyseal lesion,
 88–93, 99–100, 103–5
 normal anatomy, 89, 98
 physeal injuries and epiphyseal
 separations, 93, 107
 simple metaphyseal fractures,
 88, 98
 district attorneys, contacts with,
 660
 Dixon techniques, MRI, 701, 703
 documentation. *See* records
 dogs, attacks by, 613
 domestic violence. *See* intimate
 partner violence
 Doppler sonography, infantile
 hemangioma, 233
 dose length product (DLP),
 683
 double inversion recovery (DIR)
 imaging, 701
 Down syndrome, 609–10
 drowning, 620
 drug-induced bone changes,
 232–4
 dual-energy x-ray
 absorptiometry (DXA),
 248
 duodenal injuries
 intramural hematomas, 570–3,
 583, 598, 610
 pancreatic injuries and,
 579
 perforation, 567–70, 576
 strictures complicating, 575–6
 dura mater, 395–6
 endosteal or periosteal layer,
 395
 meningeal layer, 395–6
 dural border cell layer (DBCL),
 396, 409
 hemorrhage within, 411–12
 dural venous plexus, 409
 dynamic contrast-enhanced
 (DCE) MRI, 704–5
 dysautonomia, familial, 221, 223
 dysphagic choking, 441, 471
 dysplasias, inherited bone,
 217–18
 ear injuries, 377
 echo planar imaging (EPI),
 703–5
 echo time (TE), 693–4
 gradient-echo imaging, 701
 spin-echo imaging, 699
 Edmunds, Audrey, 649–50
 Effective Dose, radiation, 670–1
 CT scanning (D_{eff}), 683–4
 effective echo time (ETE), 699
 elbow dislocation, 138
 electronic medical records
 (EMR), 651
 emails, as evidence, 651
 emancipated minors, 639
 emergencies, consent during, 639
 emphysema, subcutaneous
 mediastinum, 591–2
 neck, 590–4
 encephalitis, viral, 436, 476, 480
 endochondral ossification, 15–20
 epidemiology, child abuse, 1–2
 epidural hemorrhage (EDH),
 401–3
 anatomic aspects, 395, 401
 imaging considerations, 402–3
 retroclival, 518, 530–1
 spinal, 513, 515–17, 532
 epiphyseal separations, 47–8,
See also physeal injuries
 distal femur, 70–1, 80–1
 distal humerus, 138–41, 146
 distal radius, 157
 distal tibia, 93
 imaging strategies, 330
 obstetric trauma, 281–6
 proximal femur, 55–61
 proximal humerus, 128–34
 proximal tibia, 80–5, 90, 92
 epiphysis, 10
 blood vessels, 13
 development, 16
 equivalent absorbed dose,
 radiation, 670
 esophago-aortic fistula, 613
 esophagus
 burns from hot liquids, 615–16
 caustic burns, 616
 foreign bodies, 613–14
 perforations, 590–4, 613, 622
 evidence
 expert medical. *See* expert
 medical testimony
 Federal Rules of, 647
 hierarchy of, 311
 legal proceedings, 647
 levels of, diagnostic studies,
 313
 testimonial, 647
 evidence-based medicine (EBM),
 309–11
 common pitfalls, 313–15
 criticism of, 315–16
 evidence-based radiology (EBR),
 311–20
 common pitfalls, 313–15
 “temporary brittle bone
 disease” and, 316–19
 expert medical testimony, 645,
 647–51, 662–3
 avoiding bias, 663
 care and protection cases, 662
 challenges to reliability, 648–50
 collateral: motions for new
 trial, 649–50
Daubert, 648–9
 habeas corpus, 650
 charts/diagrams/other aids,
 653, 663
 courtroom procedure, 654–6
 courtroom strategy, 663
 criminal cases, 662
 effects of increased scrutiny,
 650–1
 media attention, 650
 opinion, 648, 653–5
 reliability and *Daubert*, 648
 respecting the jury, 663
 expert witnesses, 647–57
 “battle” between, 648, 663
 consulting (opinion), 648,
 653–5
 cross-examination, 655–6
 direct examination, 654–5
 financial compensation, 653–4
 helping the trier of fact, 647
 hostile, 652
 pre-trial depositions, 651–2
 pre-trial preparation, 652–4,
 661–2
 qualification, 647–8
 record keeping, 651
 re-direct examination, 656
 scheduling trial dates, 661–2
 sequestration, 653, 656
 exposure time, x-ray, 668
 extra-axial hemorrhage and
 nonhemic collections
 intracranial, 394–441
 spinal, 515–22
 extracorporeal membrane
 oxygenation (ECMO),
 SDH due to, 430
 extradural hemorrhage.
See epidural hemorrhage

- eye injuries. *See also* retinal hemorrhages
 ruptured globe, 384
- facet joint disruption and dislocation, 539–44
- facial fractures, 378–82
- facial injuries, 375–82
- fact witnesses, 647–8
- factitious illness, inflicted.
See Munchausen syndrome by proxy
- factor VIII deficiency, 433
- factor IX deficiency, 433
- factor X deficiency, 434
- failure to thrive (FTT)
 nonorganic, 610–11
 organic, 609–10, 613
- falls
 head injury
 biomechanics, 351–3
 clinical data, 351
 parenchymal brain injuries, 478–9
 skull fractures, 372–4, 382–3
 skeletal injuries, 269–72
- false falx sign, 412–13
- falsifiability (disprovability) test, 319–20
- family. *See also* parents
 characteristics, 634–5
 interacting with, 635–6
- FAST (focused assessment with sonography for trauma), 599
- fast spin echo (FSE), 699
 bright fat effect, 701
 single-shot (SSFSE), 698, 700
- fat
 embolism, 475–6
 MRI, 688–9, 695, 701–3
 x-ray imaging, 672
- fat necrosis
 and calcification, 619–20, 622
 medullary, 579
- fat saturation pulses, MRI, 700
- fatal child abuse. *See also* postmortem research
 differential diagnosis, 338
 epidemiology, 1–2
 postmortem skeletal imaging.
See postmortem skeletal imaging
- Federal Rules of Evidence, 647
- feet
 fractures, 96–7, 110–12
 osteolytic lesions complicating pancreatitis, 579–80
 radiographic skeletal surveys, 327
- femur, 53–71
 accidental trauma, 270–1, 273
- birth-related trauma, 55, 58–9, 71, 283–6
- distal. *See* distal femur
- neck and trochanteric region
 fractures, 61–3
- physiologic SPNBF, 291, 293
- predictors of abuse, 56–7
- proximal. *See* proximal femur
- shaft fractures, 63–5, 67, 70
 accidental, 273
 bilateral, 64, 67
 birth-related, 283–4, 286
 CML with, 65, 71
 through pre-existing
 physiologic SPNBF, 64, 69, 291
- soft tissue injuries, 54, 57
- ferritin, chronic subdural hematoma, 400, 429
- fetus
 abuse, 620
 brain MRI, 698
 calcium homeostasis, 241–3
 movements, “temporary brittle bone disease” and, 318
 rib fractures, 166, 264
 spontaneous fractures, 279
- fibroblast growth factor-23, 14
- fibula, 94–6
 accidental trauma, 270, 272
 classic metaphyseal lesion, 94–6, 109
 metaphyseal fractures, 94, 108
 metaphyseal step-off configuration, 294
 shaft fractures, 87, 92, 94–5, 107
- finger fractures, 148–52, 159–60
- finite element (FE) modeling
 abusive head trauma, 348
 retinal hemorrhages, 350–1
- first rib
 fractures, 178, 184–6, 188
 ossification defects/
 pseudarthrosis, 300, 303
- fissures, cranial, 370–1, 378–9
- FLAIR (fluid-attenuated inversion recovery)
 imaging, 490, 701
- flip angles (α), 693
- gradient-echo imaging, 701
- spin-echo imaging, 699
- fluoroscopy, 677
- follow-up skeletal surveys, 327
- fontanels, 358
- foot. *See* feet
- football sign, 566, 568–9
- foramen magnum, 504
- forearm, 141–8
 accidental trauma, 272
 bilateral fractures, 143, 152
 distal. *See* distal forearm
- midshaft fractures, 141–5, 151–2
- proximal fractures, 141, 149–50
- foreign bodies
 genitourinary, 618–19, 621
 orally inserted, 590, 613–16
 rectal, 577, 621–2
 tracheobronchial tree, 596, 615
 vaginal, 621–2
- Fourier transformation, 692
- fractional anisotropy (FA),
 cerebral edema, 458
- fracture dating, 208–15
 age of child and, 209–11
 chronic and remote fractures, 211–12
- classic metaphyseal lesion, 39–42, 45–6
- clinical history, 209–10
- forensic significance, 208–9
- guidelines, 215
- pathophysiology, 209–12
- radiographic phases, 213–14
- repetitive injury and, 210–11
- skull fractures, 369–70, 377
- spinal trauma, 551
- fracture healing
 factors influencing, 209–12
 histologic phases, 212–13
 induction phase, 212
- radiographic phases, 213–14
- skull fractures, 369–70, 377
- fractures, 23–48. *See also* skeletal trauma; *specific types and locations of fractures*
 accidental, 268–75
 bone mineral density and, 248
 congenital rickets, 245
 dating. *See* fracture dating
 hospitalized children, 189, 192, 316–17
 intrauterine, 279
 line definition, dating role, 214–15
 lower extremity, 53–117
 metabolic bone disease of prematurity, 244
 obstetric trauma, 279–87
 osteogenesis imperfecta, 254–5, 261–2
 radiology report, 659
 sexually abused children, 623
 “temporary brittle bone disease”, 316–18
 thorax, 164–203
 upper extremity, 121–52
 vitamin D deficiency rickets, 246–8
 without bruising/external signs, 316, 318
- frontal radiographic views,
 skeletal surveys, 326
- functional magnetic resonance imaging (fMRI)
 basic principles, 704–5
 diffuse axonal injury, 464
- gadolinium (Gd), 704–5
- gamma camera, 668, 679
- gamma rays, 668–9
 interactions with matter, 669
 properties, 669
 radiation dose terminology, 669–71
 sources, 668–9
- gastric artery pseudoaneurysm, 597–8
- gastric distention, neglected children, 611
- gastric hematoma, 566–7
- gastric rupture, 565–6, 611
- gastric ulcer, 566–7
- gastrointestinal tract trauma, 565–77
- genitourinary tract
 foreign bodies, 618–19, 621
 injuries, 586–9, 621–2
- genu varum, physiologic, 295–7
- glenohumeral dislocation, 132
- glenoid fractures, inferior, 198–9, 201, 203
- global skeletal assessment, 324–30
 modalities, 326–30
 recommendations, 332
 sexual abuse, 622–3
 special situations, 330–2
 unexplained femoral shaft fractures, 65
- glutaric aciduria type I (GAI), 434, 438, 476, 480
- gradient echoes, 693–4
- gradient-echo imaging, 695–6, 701–3
- Gray (Gy), 670
- gray matter, MRI, 697–8
- greenstick fractures, 32
- groove of Ranvier, 16–17
- growth (and development)
 nonorganic failure to thrive, 610–11
 skeletal, 15–20
- growth disturbances/residual deformities, 212–13
- distal femoral physeal injuries, 71
- distal humeral physeal injuries, 140, 146
- meningococemia, 224, 228
- osteomyelitis, 224, 227
- proximal femoral physeal injuries, 58, 61
- proximal humeral physeal injuries, 126, 131, 134
- proximal tibial physeal injuries, 83, 92
- growth hormone (GH), reduced secretion, 610–11
- growth plates (physes), 10, 15–20
 closure, 20
 growth factors and hormones regulating, 17
 injuries. *See* physeal injuries

Index

- growth plates (physes) (cont.)
 ribs, 170, 177
 vertebrae, 531
 vulnerability to trauma, 21
 zones, 17–18
 growth recovery (arrest) lines,
 147–8, 211
 guardians, legal. *See* parents
- habeas corpus challenges, 650
 hair pulling, 364, 619
 hairline fractures, femoral shaft,
 63
 half-value layer (HVL), 669
 hands
 fractures, 148–52, 158–60
 radiographic skeletal surveys,
 327, 712
 hanging, near, 467–8, 470
 hangman's fracture, 511, 515
 differential diagnosis, 547–8
 mechanism of injury, 511–12
 HASTE sequences, MRI, 700
 Haversian systems, 11–12
 head trauma
 abusive. *See* abusive head
 trauma
 accidental, 351–3, 403, 423–4,
 478–9
 healing
 fracture. *See* fracture healing
 physal injuries, 213
 Health Insurance Portability and
 Accountability Act
 (HIPAA), 640–1
 hearing loss, osteogenesis
 imperfecta, 256
 hemangiomas, infantile, 232–3
 hematologic disorders,
 intracranial hemorrhage,
 434–6
 hematologic malignancies, 599,
See also leukemia
 hematuria, 586
 hemoglobin, oxidation state, 399
 hemoperitoneum, liver injuries,
 582
 hemophagocytic
 lymphohistiocytosis
 (HLH), 438–9, 599
 hemophilia, 433
 hemorrhage
 classic metaphyseal lesion, 39
 fracture site, 212
 intracranial. *See* intracranial
 hemorrhage
 subperiosteal, 27, 29–32, 212
 hemorrhagic disease of newborn,
 430, 432–3
 hemosiderin
 chronic subdural hematoma,
 400, 429
 intracranial extra-axial
 hemorrhage, 399
 hemothorax, 595
- hereditary hemorrhagic
 telangiectasia, 436
 hereditary sensory and
 autonomic neuropathies
 (HSAN), 219–21
 herpes encephalitis, 436, 476, 480
 heterotopic bone formation.
See also calcification
 pelvis and thigh, 101, 117
 precervical, 594
 spinal trauma, 542, 544
 high-chair accidents, 273
 hip. *See also* proximal femur
 developmental dysplasia
 (DDH), 55, 59
 dislocation, pelvic fracture
 with, 101, 116
 soft tissue injuries, 54, 57
 hired guns, 653, 656
 histologic analysis, postmortem
 specimens, 337
 historical aspects
 abusive head trauma, 345–6
 child abuse, 2–3
 history, clinical
 abusive head trauma, 347
 fracture dating, 209–10
 hole-in-the-rib appearance, 178,
 182
 homicide cases, 647, 649–50
 Hounsfield Unit (HU), 678
 Hounsfield, Godfrey, 677
 Howship lacunae, 14
 humerus, 121–41
 accidental trauma, 273–4
 birth-related injuries, 123–4,
 131, 138, 280–4
 distal. *See* distal humerus
 physiologic SPNBF, 291–2
 proximal. *See* proximal
 humerus
 shaft fractures, 135–6
 accidental, 273–4
 birth-related, 280–1
 Hunter syndrome, 234–5, 549,
 551
 hydrocarbon poisoning, 616
 hydrocephalus, benign idiopathic
 external. *See* benign
 enlargement of
 subarachnoid spaces in
 infancy
 hyperemia, cerebral, 469–71
 hyperextension toddler's
 fracture, 272
 hyperflexion cervical spine
 injuries, 515
 hypernatremic dehydration, 437–8
 hyperostosis, infantile cortical
 (Caffey disease), 218,
 230–1
 hypertrophic osteoarthropathy,
 218, 231
 hypertrophic zone of growth
 plate, 18
- classic metaphyseal lesion,
 40–2
 hypervitaminosis A, 218, 228–32
 hypoperfusion complex, CT, 570,
 582, 584, 598
 hypopharyngeal perforations,
 590–4, 614
 hypophosphatasia, infantile/
 childhood, 257
 hypopituitarism, transient,
 610–11
 hypoxic ischemic
 encephalopathy (HIE),
 471–4
 as cause of SDH/retinal
 hemorrhages, 440–1
 bilateral, 471–2
 medullary/cervical cord
 injuries and, 523–5
 unilateral, 472–4, 477
- ifosfamide, 232
 ileum, intramural hematoma,
 570
 iliac fractures, 101, 115
 images, radiologic.
See radiologic images
 imaging, principles and
 technologies, 667–84
 immobilization, patient, skeletal
 surveys, 329, 713–14
 immune thrombocytopenic
 purpura (ITP),
 idiopathic, 434–5
 immunodeficiency, thymic
 involution and, 596, 613
 impacted fractures
 long bones, 33, 36
 tibial shaft, 86, 95
 impalement injuries, 616–17
 impeachment, 652, 656
 Inca bone, 371, 380
 incidence, child abuse, 1–2
 infants (<1 year). *See also*
 neonates; premature
 infants
 biomechanics of falls, 352
 cranium, 357–61
 epidemiology of skeletal
 injuries, 25, 27
 fracture healing, 210–11
 physiologic SPNBF, 290–1
 radiographic skeletal surveys,
 713
 infections
 CNS, 436, 476, 480
 skeletal features, 223–6
 inferior vena cava (IVC)
 thrombosis, 582
 inflammation, fracture site, 212
 information
 protected health (PHI),
 disclosure, 640–1
 withholding, from suspected
 perpetrators, 641
- informed consent, 638–40
 informed permission, 639
 intensive care, skeletal imaging of
 children in, 331
 intent, 1, 647
 interleukin-1 receptor antagonist
 deficiency, 224
 internal jugular vein laceration,
 594
 interparietal bone (Inca bone),
 371, 380
 interspinous ligament, 508, 532
 intestinal injuries, 567–77
 intestinal transplantation, 231
 intimate partner violence (IPV),
 634–5
 intracerebral hemorrhage, 466
 dating, 399, 466
 intracranial aneurysms,
 traumatic, 474–5
 intracranial hemorrhage. *See also*
specific types
 coagulopathies causing, 430–4
 extra-axial, 396–412
 estimating age, 397–400
 hematologic disorders, 434–6
 intracranial hypotension
 MRI features, 428, 432
 subdural hemorrhage, 427,
 431–2
 intracranial imaging
 protocol, 491
 strategies, 487–92
 intracranial injury (ICI)
 epidemiology, 346
 extra-axial hemorrhage and
 nonhemic collections,
 394–441
 minor accidental falls, 478
 parenchymal brain injury,
 453–81
 shaking as a mechanism,
 349–50
 skull fracture association,
 358–60, 367
 spinal injury association, 501
 intracranial pressure, raised, split
 cranial sutures, 364, 374
 intradural hematoma (IDH),
 440–1
 intramembranous ossification,
 15, 20
 intraparietal suture, 372, 382
 intraventricular hemorrhage
 (IVH), 416, 418–19
 intussusception, small bowel,
 576–7
 inversion time (TI), 700
 ionizing radiation, 669–70
 iron oxidation products
 chronic subdural hematoma,
 400, 429
 intracranial hemorrhage, 399
 ischial avulsion fracture, 66, 101,
 116

- isolated sulfite oxidase deficiency (ISOD), 477
- jejunum
 intramural hematomas, 570–5
 perforations, 567–9
 strictures, 575
- judges, 654, 662
- jury, 649, 653, 663
- Keller, K. A. and Barnes, P. D., “Rickets vs. abuse” paper, 247–8
- Kempe, C., 3, 23
- kidney injuries, 581, 586–7
 ectopic kidney, 586, 588
 sexual abuse, 621, 623
- kinky-hair syndrome.
See Menkes disease
- knuckle beating, repeated, 152, 160
- k-space, MRI, 692
- k-value, CT, 683
- kVp (peak kilovoltage), 668, 672
 skeletal surveys, 713
- kyphosis, congenital, 548, 550
- labeling, radiographic images, 714
- lactate dehydrogenase (LDH), serum, 580
- lamellae, interstitial, 12
- lamellar bone, 10–11
- Langerhans cell histiocytosis (LCH), 438
- Larmor frequency, 690–1
- lateral radiographic views, skeletal surveys, 326
- Lauterbur, Paul, 691
- law enforcement agencies.
See police/law enforcement agencies
- lawyers (attorneys), 646
 issuing subpoenas, 661
 pre-trial conferences, 652–4, 661–2
 pre-trial depositions, 651–2
- lead shielding, 714
- left gastric artery
 pseudoaneurysm, 597–8
- legal issues, 638–43
 consent, 638–40
 medical records, 640–1
 radiologist in court, 645–57
 reporting abuse, 641–3
- legal proceedings, 645–57
 burden of proof, 646
 civil vs. criminal cases, 645–6
 evidence, 647
 giving evidence. *See* expert medical testimony
 media interest, 650, 654
 motion hearings, 647
 pre-trial discovery, 646–7, 651
 trial dates, 661–2
- leptomeningeal cysts, 365, 388
- leptomeningeal tumors, 435, 438
- leptomeninges, 395
- leukemia
 intracranial hemorrhage, 434–5
 skeletal features, 218, 232
 visceral changes, 599
- leukemic bands/lines, 78, 87, 232
- ligamentum flavum, 506, 532
- limbus vertebrae, 535
- line scan diffusion imaging (LSDI), 696
- lipids. *See* fat
- liquefaction, brain, 456–8, 468
- liver function tests (LFTs), 580–1
- liver injuries, 580–3
 imaging, 581–3
 lower rib fractures, 187, 190, 581–2
 penetrating, 580
- long bones
 cortical irregularity and indistinctness of shafts, 298
 epidemiology of injuries, 25–7, 29–30
 fractures in osteogenesis imperfecta, 256–7, 260–1
 impacted fractures, 33, 36
 injuries, 30–48
 oblique fractures. *See* oblique fractures of long bones
 obstetric trauma, 280–6
 physiologic SPNBF, 291
 shaft fractures, 31–4
 sites vulnerable to injury, 21
 spiral fractures. *See* spiral fractures of long bones
 transverse fractures.
See transverse fractures of long bones
- Looser zones, 243
- low-birthweight infants, 609
- lower extremity
 accidental trauma, 272
 fractures, 53–117
 soft tissue injuries, 54, 57
- lower urinary tract injuries, 587–8
- lumbar puncture complications
 intracranial subdural hemorrhage, 432
 spinal epidural hemorrhage, 516
 spinal subdural hematoma, 517
 vertebral body compression fracture, 549
- lumbar spine injuries, 532, 535,
See also thoracolumbar spine injuries
- lung injuries, 595–6
- lye ingestion, forced, 616
- lytic osseous lesions.
See osteolytic lesions
- M (magnetic moment vector), 690
- mA (tube current), 668
 skeletal surveys, 713
- macrocephaly of infancy, benign.
See benign enlargement of subarachnoid spaces in infancy
- magnetic field gradient coils, 689, 691–3
- magnetic field gradients, 691
- magnetic field strength (B₀), 689–90
- magnetic resonance angiography (MRA)
 head and neck, 489, 491
 principles, 697, 702
- magnetic resonance imaging (MRI), 489–91, *See also* specific pulse sequences and techniques
 abusive head trauma 489
 adrenal injuries, 585
 benign enlargement of subarachnoid spaces in infancy, 418, 422
 biologic principles, 696–8
 clavicular fractures, 192–3
 craniocervical/cervical spine injuries, 552–3
 bony and ligamentous, 508, 510–11, 513, 515
 cord injuries, 517, 522, 528–9
 epidural hematoma, 513, 517
 retroclival hematoma, 518–20, 531
 subdural hemorrhage, 514, 516, 520–2
 epidural hemorrhage, 403
 femoral injuries, 56, 58, 60, 62, 66, 71
 humeral injuries, 128, 133, 137, 139, 142, 144
 infantile hemangioma, 233
 intracranial extra-axial hemorrhage, 397–400
 intracranial hypotension, 428, 432
 obstetric trauma, 284
 osteomyelitis and septic arthritis, 224–6
 parenchymal brain injuries
 anoxic injury, 470
 cerebral edema, 458–60
 contusions, 461–2
 cortical laminar necrosis, 460–1
 diffuse axonal injury, 464–5
 follow-up, 457–8, 478
 hematoma, 466
 lacerations, 464–8
 necrosis, 454–7
 normal findings, 478
 pelvic fractures, 115
- physical principles, 689–96
 pulse sequence approaches/
 parameter settings, 698–705
 retinal hemorrhages, 468
 sacral fractures, 545, 549
 scanner components, 689–91
 skeletal injuries, 330
 soft tissue lower extremity injuries, 54, 57
 spinal imaging strategies, 491, 551–3
 subarachnoid hemorrhage, 403, 405–6
 subdural hemorrhage, 407, 411, 426
 birth-related, 418, 420
 chronic, 400, 415, 417, 423, 428–9
 dating, 399–400
 subdural nonhemic collections, 412–13, 415–16, 419
 thoracolumbar spine injuries, 529, 531, 534–5, 540–1
 tibial CML, 93, 106
 toddler’s fracture, 270
 whole body. *See* whole body magnetic resonance imaging
- magnetic resonance spectroscopy, ¹H.
See proton magnetic resonance spectroscopy
- magnetic resonance venogram (MRV), brain, 491
- male genital injuries, 588–9, 622
- malnutrition, 611–13
- mandibular fractures, 378–81, 385
 birth-related, 287
- mandibular subperiosteal new bone formation, 381, 386
- Mansfield, Sir Peter, 691, 703
- Martin, Raymond, 648
- masking, electronic, 712
- mature minor doctrine, 639
- media interest, legal proceedings, 650, 654
- mediastinum
 abscess, 591–3
 avulsed tooth in, 593
 emphysema, 591–2
 pseudocyst, 592
- medical records. *See* records (medical)
- medulla oblongata injuries, 522–9
 mechanisms, 525–7
 pathology, 522–5
- medullary canal, 10
- medullary cavity, formation, 16
- MELAS (mitochondrial myopathy encephalopathy lactic acidosis and stroke-like episodes), 480

Index

- Mendosal suture, 359, 372
 meningeal artery injuries, 401–2
 meninges, cranial, 395–6
 meningitis, bacterial
 hemorrhagic complications, 436
 subdural effusion, 415, 419, 436
 subdural empyema, 419
 meningococemia, infantile, 224, 228
 Menkes disease
 intracranial features, 437, 440, 477, 480
 skeletal features, 218, 234–5
 mental health sequelae, long-lasting, 634
 mesenteric hematomas, 575–6, 583
 meta-analysis, 310
 metabolic bone disease, 240–9
 of prematurity, 243–4
 metabolic disorders
 birth-related fractures, 286
 CNS involvement, 434, 437–8, 480
 miscellaneous, 235
 metacarpal cloaking, 151, 158
 metacarpal fractures, 148–51, 158
 metaphyseal dysplasias (MD), 217–18
 McKusick type, 218, 221
 Schmid type, 218, 220
 metaphyseal fractures
 (nonspecific)
 distal femur, 65, 72
 distal radius, 146, 154
 distal tibia, 88, 98
 fibula, 94, 108
 iatrogenic, 223, 225
 osteogenesis imperfecta, 256–7, 261
 proximal humerus, 132–4
 proximal tibia, 85–6, 93–4
 “temporary brittle bone disease”, 316
 metaphyseal lesion, classic.
 See classic metaphyseal lesion
 metaphyses, 10, 20
 bands, proximal tibial CML, 78, 87
 beak contour, 294
 blood vessels, 13
 distal femoral, 65–7, 73
 distal tibial, 89, 98
 distal ulnar cupping, 294–5
 irregularities
 differential diagnosis, 218
 distal femur and proximal tibia in toddlers, 295–8
 rickets, 241, 243
 normal radiologic variants, 292–8
 postmortem specimen resection, 337
 proximal humeral, 126–7
 proximal tibial, 73–5, 83
 spur, 294–6
 step-off configuration, 292–5
 subphyseal radiolucency in normal infants, 298–9
 vulnerability to trauma, 21
 metatarsal fractures, 96–7, 110–12, 151
 methemoglobin
 extra-axial/subdural hemorrhage, 399–400, 428
 intracerebral hematoma, 399, 466
 methotrexate, 232
 middle meningeal artery injury, 401–2
 migraine, hemiplegic, 481
 mineralizing zone of growth plate, 18–19, *See also* zone of provisional calcification
 minors. *See also* children
 emancipated, 639
 mature minor doctrine, 639
 medical care without parents’ consent, 640
 records retention, 641
 mitochondrial myopathy
 encephalopathy lactic acidosis and stroke-like episodes (MELAS), 480
 modulation transfer function (MTF), 681
 motion hearings, 647
 motion, patient
 CT scanning, 679
 nuclear imaging, 680
 prevention, skeletal surveys, 329, 713–14
 x-ray imaging, 674
 motions for a new trial, 649–50
 multiple hereditary exostoses (MHE), 218–19, 222
 Munchausen syndrome by proxy (MSBP), 613, 617–19
 muscle injuries, severe
 widespread, 586–7, 620
 myelin vacuolization (intramyelinic cerebral edema), 458–9
 myelodysplasia, 218, 221–3
 birth-related fractures, 279
 myoglobinuria, 586–8
 myositis ossificans.
 See heterotopic bone formation
 naphtha poisoning, 616
 nasal injuries
 fractures, 381–2
 soft tissue, 377–8
 nasal septal hematomas, 381–2
 National Incidence Study of child abuse and neglect, fourth (NIS-4), 2
 near infrared spectroscopy, frequency domain, 492
 neck injuries, 590–4, *See also* craniocervical junction (CCJ) injuries
 abusive head trauma, 347
 pharyngeal and esophageal perforations, 590–4
 spine. *See* cervical spine injuries
 necklace calcification, 619
 necroptosis, 456
 necrosis (neuronal), 454–7
 acute, 454–6
 cortical laminar, 460–1
 delayed, 455–7
 needles, inserted, 580, 588, 597, 616–18
 neglect, 608, 610–13
 nonorganic failure to thrive, 610–11
 reporting suspected, 642
 neonates
 calcium homeostasis, 243
 hemorrhagic disease, 430, 432–3
 obstetric injuries. *See* obstetric trauma
 skeletal features, 211
 neoplasia
 differential diagnosis, 232
 inflicted lower extremity trauma vs., 55, 57
 inflicted upper extremity trauma vs., 122, 126
 intracranial hemorrhage vs., 435, 438
 nerve supply, bones, 13
 neuroblastoma, 598
 neurocentral synchondrosis
 vertebral injuries, 536–40
 posterior ligamentous disruption with, 544, 546
 neurofibromatosis type 1 (NF1), 235
 neurologic outcomes, 353, 481
 neuromuscular disorders, birth-related fractures, 279, 286
 neurons
 cell death mechanisms, 454–8
 developmental pruning, 454
 nightstick injuries, 142
 nonaccidental injury, 1, 268
 nuchal ligament, 508
 nuclear imaging, 679–80,
 See also positron emission tomography
 nature and properties of
 gamma rays, 668–9
 radiation dose, 329, 680
 skeletal. *See* bone scintigraphy
 small bowel hematoma, 574
 technology and dose-related issues, 680
 nutrient arteries, 13
 nutrient canals, 291–2, 298–300
 nutritional deficiencies, 611–12
 nutritional rickets, 245–6, 611–12
 obesity, extreme childhood, 623
 objects, falling, 272–3
 oblique fractures of long bones, 32–3, 35
 accidental, 270, 272–4
 femur, 61, 63–4
 humerus, 135
 obstetric trauma, 281
 tibia, 72, 87, 95–6
 oblique radiographic views, 325–6
 obstetric trauma, 279–87
 cervical spine epidural hematoma, 517
 clavicular fractures, 195, 213–15, 279–80
 femoral fractures, 55, 58–9, 71, 283–6
 fracture healing, 210
 humeral fractures, 123–4, 131, 138, 280–4
 long bone injuries, 280–6
 rib fractures, 286–7
 risk factors, 279, 286
 scalp and cranium, 287, 374–5, 384
 spinal injuries, 287, 549
 subdural hemorrhage, 418, 420
 occipital bone, 504–5
 biomechanics, 360
 developmental variants, 371
 fractures, 365–7
 O’Connor, John F., 208
 odontoid process, 505
 olecranon fractures, 141, 144, 149, 151
 oral trauma, 590, 622
 orally inserted foreign bodies, 590, 613–16
 orbital injuries, 382–6
 orthopedic deformities.
 See growth disturbances/residual deformities
 orthopedic manipulations, for clubfoot deformity, 223, 225
 Osler–Weber–Rendu disease, 436
 ossification
 enchondral, 15–20
 intramembranous, 15, 20
 ossification centers
 primary, 15–16
 secondary, 16–17
 secondary-like, 16
 osteoblast-lineage cells, 13–14
 osteoblasts, 13

- osteoclasts, 14
 resorption of mineralized cartilage, 18
- osteocytes, 13–14
- osteocytic osteolysis, 14
- osteogenesis imperfecta (OI), 218, 254–64
 abuse of children with, 263, 609–10
 biochemical and molecular aspects, 258–60, 263
 chondro-osseous pathology, 258
 classification, 254–5
 clinical features, 255–7
 genetics, 257
 imaging features, 260–3
 intracranial hemorrhage, 439
 overview of diagnosis, 263–4
 transient. *See* “temporary brittle bone disease”
 type I, 254, 256, 260, 262
 type III, 255, 261
 type IV, 255, 257, 259–60
 type V, 255
- osteogenic sarcoma, inflicted injury simulating, 55, 57
- osteolytic lesions
 after traumatic pancreatitis, 579–80
 comminuted fractures resembling, 63, 66
- osteomalacia, 240
 maternal, congenital rickets, 244–5
- osteomyelitis, 218, 223–6
 abused infant, 224, 228
 inflicted injury simulating, 141, 149
 meningococemia with, 224, 228
- osteons (Haversian systems), 11–12
- osteopenia, 240. *See also* bone demineralization
 causes, 218
 chemotherapeutic drug-induced, 232
 neglected children, 612
 of prematurity, 244
 paralytic neurologic disorders, 221–2
- osteopetrosis, 218–19, 221
- osteoporosis, 240, 248
- osteoprogenitor cells, 13
- osteoprotegerin (OPG), 14
- pain, congenital insensitivity to, 218, 221–3
- pancreatic injuries, 577–80
- pancreatic pseudocysts, 577–9
- pancreatitis, 577–9
 chronic, 579
 osseous changes, 579–80
- parallel imaging technique, MRI, 692, 701
- paralytic neurologic disorders, 221–3
- parathyroid hormone (PTH), 241–3
- parenchymal brain injuries. *See under* brain
- parent–child attachments, 634
- parent–infant traumatic stress syndrome (PITS), 346
- parents (or caregivers or guardians). *See also* perpetrators
 as victims of abuse, 635
 characteristics of abusive, 634–5
 consent issues, 639–40
 interactions with, 635–6, 660
 withholding information from, 641
- parietal fractures, 365–7, 369–70
 biomechanics, 360–1
 wrap around (biparietal), 364, 366, 373, 383
- patients. *See* children
- peak kilovoltage (kVp), 668, 672
- skeletal surveys, 713
- pelvic mass, after sexual assault, 621, 623
- pelvis
 fractures, 97–101, 114–16
 normal variants, 307–8
 soft tissue injuries, 101, 117
- perfusion imaging, MRI, 704–5
- perichondrium, 12–13, 15, 30
- perineal injuries, accidental, 599
- periosteal blood vessels, 13
- periosteum, 10, 12–13, 30
- skull, 357
- periportal tracking, liver injuries, 581–2
- perpetrators. *See also* defendants;
 parents
 aiding assessment of intent, 647
 confessions by, 349
 withholding information from, 641
- pertussis (whooping cough), 441, 471
- PET. *See* positron emission tomography
- phalangeal (finger) fractures, 148–52, 159–60
- pharyngeal perforations, 590–4, 613, 615, 622
- phosphorus metabolism, disorders of, 240–9
- photoelectric effect, 669
- physeal injuries, 47–8. *See also* epiphyseal separations;
 Salter–Harris (SH) injuries
- distal femur, 70–1
- distal humerus, 138–40, 144–5
- distal radius, 148, 157
- distal tibia, 93, 107
- healing, 213
- obstetric trauma, 281–6
- proximal femur, 55–61
- proximal humerus, 129–34
- proximal tibia, 80–5, 90
- ribs, 170–1
 Salter–Harris classification, 47
- physes. *See* growth plates
- physics survey, annual, 709–12
- pia mater, 395–6
- placental parathyroid-related protein (PTHrP), 241–2
- plaintiff, 646
- plastic bowing injuries, 32–3
 fibula, 87, 92, 94
 radius and ulna, 143, 152
- pleural effusions, 591, 595
 chyloous, 596
- pneumatosis
 colonic, 576
 gastric, 566–7
 intestinalis, 576–7
- pneumomediastinum, 591, 595
- pneumoperitoneum, 568–9
- pneumothorax, without rib fractures, 595
- poisonings, 613–16, 618
- police/law enforcement agencies
 contacts with, 660
 disclosure of information to, 640–1
 reporting suspected abuse, 643
- popping rib, 189, 191
- portal vein
 gas, 576–7, 583
 thrombosis, 582
- positioning, patient, for skeletal surveys, 713–14
- positron emission tomography (PET), 679–80,
See also bone scintigraphy
- clavicular fractures, 194
- cranial injuries, 362
- distal femoral epiphyseal separation, 81
- femoral CMLs, 60–1, 70, 79
- femoral fractures, 63, 65
- fibular shaft fractures, 107
- humeral injuries, 128, 132–3
- pelvic fractures, 115–16
- radiation dose, 329, 680
- radiographic skeletal surveys vs., 328–9
- rib fractures, 180, 183–4, 186
- role in child abuse, 680
- sensitivity, 329
- spinal trauma, 534, 551
- SPNBF, 30, 32
- sternal injuries, 197
- posterior longitudinal ligament, 508, 532
- postmortem research
 abusive head trauma, 347–8
 craniocervical junction/
 cervical spine injuries, 524–5
 fracture distribution, 25, 29
 liver injuries, 581
 proximal femoral epiphyseal separations, 55, 59
 skull fractures, 361, 365
- postmortem skeletal imaging, 335–8. *See also* specimens, postmortem
 data analysis, 338
 differential diagnosis of fatal abuse, 338
 histologic analysis of resected material, 337
 skeletal surveys, 331, 335–7, 714
- Power ratio, 513
- precession, 690–1
- pregnancy. *See also* fetus
 abuse during, 620
- premature infants
 metabolic bone disease, 243–4
 risk of abuse, 609
 “temporary brittle bone disease”, 316
- prenatal rib fractures, 166, 264
- programmed cell death (PCD), developmental, 454
- proliferating zone, growth plate, 18–19
- prosecutors, 646
- prostaglandin E1, 232
- protected health information (PHI), disclosure, 640–1
- proton density (PD) spin-echo imaging, 699–701
- proton magnetic resonance spectroscopy (¹H-MRS), 490–1
 concussion, 476
 diffuse axonal injury, 464
- proximal femur, 57–61,
See also hip
 birth-related injuries, 283–6
 classic metaphyseal lesion, 59, 61
 physeal injuries and epiphyseal separations, 55–61
- proximal forearm fractures, 141, 149–50

Index

- proximal humerus, 126–34
 classic metaphyseal lesion, 124–30
 birth-related, 281–3
 overlap with epiphyseal separation, 134
 fracture-dislocation, 130, 134
 metaphyseal beak, 294
 metaphyseal nonspecific fractures, 132–4
 normal anatomy, 123, 126–7
 physeal injuries and epiphyseal separations, 128–34
 birth-related, 281–3
 deformities complicating, 131, 134
 overlap with CML, 134
 spurious SPNBF, 291, 294–5
 step-off contour, 293, 295
- proximal tibia, 72–86
 classic metaphyseal lesion, 73–9, 83, 85–6, 88
 cortical irregularity, 293, 298
 irregularities in toddlers, 295–8
 metaphyseal beak, 294
 metaphyseal step-off configuration, 294
 nonspecific metaphyseal injuries, 85–6, 93–4
 normal anatomy, 73–5, 83
 physeal injuries and epiphyseal separations, 80–5, 90, 92
- pseudoscience, 319–20
 pseudo-subarachnoid hemorrhage, 406–7
- psychosocial considerations, 633–7
- psychosocial dwarfism, 610–11
- psychosocial sequelae, long-lasting, 634
- pulmonary consolidation, hydrocarbon poisoning, 616
- pulmonary edema, 596
- pulmonary trauma, 595–6
- Purcell, Edward, 695
- qualification, expert witnesses, 647–8
- quality assurance (QA), 708–15
- quality control (QC), 708
- quality of published studies, 315
- quantitative computed tomography (QCT), 248
- quantum mottle (quantum noise), 673
- questions
 cross-examination, 656
 direct examination, 654–5
 evidence-based medicine, 310–11
- radiation dose, 669–71
 absorbed dose, 669–70
- clinical aspects of managing, 682–4
- CT. *See under* computed tomography
- Effective Dose, 670–1
- equivalent absorbed dose, 670
- global skeletal assessment, 329
- radionuclide imaging, 329, 680
- units of expression, 670
- x-ray image contrast and, 673
- radiation, ionizing, 669–70
- radicular vein, lacerated, 517
- radio-frequency (RF) pulses, MRI, 690
- fat saturation, 700
- inversion, 700–1
- refocusing, 695
- transmitting and receiving hardware, 689, 692–3
- radiographic image. *See* x-ray image
- radiography, 675–7
 craniocervical and cervical injuries, 512–13, 515
 digital technologies. *See* digital radiographic technologies
- film-screen systems, 675
- fracture dating, 213–14
- maintaining and calibrating equipment, 709–12
- radiation dose management, 682
- rib fractures, 173, 180–2, 326
- skeletal surveys. *See* skeletal surveys, radiographic specimen, 337
- radiologic images
 formation, 667–84
- informal advice to colleagues, 660
- labeling, 714
- pre-trial review, 653
- quality, 681–4
- quality assurance, 708–15
- radiologists
 as expert witnesses. *See* expert witnesses
- contacts with abuse investigators, 660
- expert testimony. *See* expert medical testimony
- general legal issues, 638–43
- interactions with parents, 635–6, 660
- precautionary measures, 660
- reporting suspected abuse, 642, 660
- response to child abuse, 658–63
- subpoenas, 646–7, 661
- radiology report, 640, 651, 659–60, *See also* records
- radiology team/staff
 best practice guidelines, 636–7
- manual immobilization of patients, 714
- responsibility for quality assurance, 714–15
- specific challenges for, 635–6
- training, 636
- radionuclide bone scans. *See* bone scintigraphy
- radionuclide imaging. *See* nuclear imaging
- radionuclides, 668
- radiopharmaceuticals, 668–9
- radius, 141–8
 distal fractures, 145–8, 153
 CML, 146–8, 155–6
 metaphyseal nonspecific, 146, 154
 normal anatomy, 146, 154
 physeal injuries, 148, 157
 midshaft fractures, 141–5, 151–2
 physiologic SPNBF, 291
 proximal fractures, 141, 150
 traumatic dislocation of head, 141, 151
- randomized controlled trials (RCTs), 310, 315
- RANK, 14
- RANK ligand (RANKL), 14
- Rapunzel syndrome, 566, 613
- reasonable suspicion standard, 641–2
- records (medical), 640–1
 confidentiality, 640–1
 expert opinion testimony, 653
- guidelines for creating, 640, 651, 659–60
- pre-trial discovery, 646–7
- pre-trial review, 652, 662
- retention, 641
- rectum
 foreign bodies, 577, 621–2
 injuries, 576–7, 620–1
- red blood cells (RBCs), subdural hematoma, 400
- relaxation times, 694–6
- religious issues, 635
- remodeling, bone, 20, 213–14
- remote fractures, dating, 211–12
- renal artery occlusion, acute, 597
- renal failure, 586–8, 620
- repetition time (TR), 692
- gradient-echo imaging, 701
- spin-echo imaging, 699
- repetitive injuries, 211
 fracture dating and, 210–11
- reporting child abuse, 641–3, 660
- liability aspects, 642
- practical aspects, 642
- process, 642–3
- reasonable suspicion standard, 641–2
- reserve zone, growth plate, 17
- retinal hemorrhages (RH), 350–1, 466, 468
- biomechanics, 350–1
- choking/coughing as explanation, 441, 471
- hypoxic brain damage as cause, 440
- legal aspects, 648–9
- 13-*cis*-retinoic acid, 232
- retinoschisis, traumatic, 350
- retroclival hematomas/other collections, 518–20, 529–31
- retroperitoneal free air/oral contrast, 569–70
- retropharyngeal abscess, 590
- reversal sign (bright cerebellum), 467, 469
- rhabdomyolysis, renal failure, 586–7, 620
- rib cage, 164–89
- rib fractures, 164–89
 abdominal visceral injuries, 186–7, 190, 596
 accidental, 274–5
 advanced and multimodality imaging, 180, 182–4, 186
 anterior and lateral, 170, 172, 175–7, 182
 birth-related, 286–7
 cardiopulmonary resuscitation (CPR) and, 189
 costochondral junction (CCJ), 172, 176–9, 182–3, 186–7
 distribution, 166, 168–70
 epidemiology, 25–7, 29–30, 164–5
 first rib, 174, 178, 184–6, 188
 historical aspects, 164–5
 hospitalized children, 189, 192, 317
 imaging features, 180–4
 intrathoracic injuries, 319, 595
 intrauterine, 166, 264
 liver injuries and, 187, 190, 581–2
 midposterior, 170, 176
 multiple hereditary exostoses vs., 219, 222
 near costovertebral articulations, 168, 170, 187
 osteogenesis imperfecta, 261–2, 264
 patterns, pathology, and mechanisms, 167–78
 posterior, 166, 170–6, 181–2, 187
 pseudo, 305
 premature infants, 244
 radiography, 173, 180–2, 326
 rib head, 167, 170–1, 173–4, 181
 rib neck, 168, 171–6, 181–2
 solitary, 188, 191
 special focus topics, 184–9
 “temporary brittle bone disease”, 316, 318–19

- ribs
 anatomic divisions, 168
 double inferior cortex, 302–4
 fusion anomalies, 300–1, 303–4
 lateral notch, 305–6
 normal anatomy, 165–7
 normal variants, 300–6
 ossification defects/
 pseudarthrosis, 300, 303
 popping, 189, 191
 postmortem specimen
 resection, 337
 pseudo-SPNBF, 295, 305
 thin, osteogenesis imperfecta, 261
- rickets, 240, 243–8
 child abuse vs., 247–8
 congenital, 244–5
 nutritional, 245–6, 611–12
 premature infants, 244
 radiographic features, 241, 243
 vitamin D deficiency, 243, 245–6
- Rigler sign, 568
 Riley–Day syndrome, 221, 223
 ring of Lacroix, 16–17
 risk factors for abuse, child, 634
 Röntgen, Wilhelm Conrad, 668
- sacral fractures, 544–5, 547–8
 Salter–Harris (SH) injuries, 47–8,
See also physeal injuries
 classification, 47
 type I, proximal humerus, 129
 type II, 48
 distal radius, 148, 157
 distal tibia, 93, 107
 proximal humerus, 128, 131
 proximal radius, 141, 150
 proximal tibia, 82, 91–2
 type III, distal humerus, 140
- scalp
 anatomy, 358
 nontraumatic swelling, 364
- scalp injuries
 dating skull fractures, 370
 differential diagnosis, 372
 hair pulling, 364, 619
 imaging, 362–5
 obstetric, 374–5, 384
- SCALP mnemonic, 358
 scapular fractures, 197–203
 anatomy and mechanisms of
 injury, 197–8
 imaging features, 199–203
- scatter, radiation, 669, 672
 scintigraphy. *See* nuclear imaging
- sclerosis, bony
 diffuse, involving long bones, 33
 first metatarsal fractures, 97, 111
 proximal humeral CML, 127, 130
 proximal tibial CML, 78–9, 88
 rib fractures, 181
 scapular fractures, 199, 203
 scrotal trauma, 588–9, 622
 scurvy, 218, 226–9
 SDH. *See* subdural hemorrhage
- second impact syndrome (SIS), 476, 479
 sedation, for global skeletal
 assessment, 329
- seizures
 causing acromial fractures, 198, 219, 221
 causing proximal femoral
 physeal injuries, 223
- sepsis, induced, 618–19
 septic arthritis, 223–5, 227
 sequestration orders, witness, 653, 656
- sexual abuse, 620–3
 esophageal perforation, 593
 pelvic injuries, 101, 117
 rectal and anal injuries, 576–7, 620–2
 reporting, 643
- shaken baby syndrome (SBS), 346
 shaking (of infants), 3
 biomechanics, 349–50
 challenges to expert testimony, 648–51
 craniocervical junction/
 cervical spine injuries, 512, 523–5, 527
 evidence-based approach, 315
 historical descriptions, 345–6
 intracranial injuries, 349–50
 media scrutiny of legal
 proceedings, 650
 retinal hemorrhages, 350–1
 subdural hemorrhage, 407
 terminology, 346
- Sharpey's fibers, 13
 sharpness, x-ray image, 673
 shear stiffness, pediatric brain, 348
- shear-strain axonal injury.
See diffuse axonal injury
- shielding, patient, 714
 shock, due to corporal
 punishment, 620
 shopping carts, falls from, 474, 478–9
- short tau inversion recovery
 (STIR) imaging, 701–2
- shoulder
 dislocation, 132
 fracture-dislocation, 130, 134
 shoulder dystocia, 279–80, 286
 shoulder girdle injuries, 198, 202
 sickle cell crisis, 231–2
 Sieverts (Sv), 670
 Silverman, Fred N., 3, 23, 358
- single photon emission
 computed tomography
 (SPECT), 679
- single-shot fast spin echo
 (SSFSE), 698, 700
- sinus of Breschet injury, 402
 sinusitis, subdural empyema
 complicating, 416
- Size-Specific Dose Estimates
 (SSDE), 684
- skeletal assessment, global.
See global skeletal
 assessment
- skeletal imaging, 324–32
 child in a cast, 330–1
 critically ill child in intensive
 care, 331
 global assessment, 324–30
 postmortem, 331, 335–8
 specific injuries, 330
- skeletal scintigraphy. *See* bone
 scintigraphy
- skeletal surveys, radiographic,
 326–7, 708–12
 after hours, 331–2
 availability, 329
 child in a cast, 330
 cost, 330
 critically ill child in intensive
 care, 331
 definition and background,
 709
 digital acquisition, 709
 equipment maintenance and
 calibration, 709–12
 examination, 712–14
 follow-up, 327
 fracture distribution, 29–30
 image identification, 714
 indications, 25, 708–9
 infants with skull fractures,
 387–8
 interpreter dependency, 330
 patient immobilization and
 positioning, 329, 713–14
 postmortem, 331, 335–7
 indications, 335–6
 initial radiologist review, 337
 technique, 336–7, 714
 principles/imaging protocol,
 326–7, 710, 712
 quality assurance, 708–15
 radiation dose, 329
 radiation protection concerns,
 682
 recommendations, 332
 responsibility for quality
 assurance, 714–15
 scintigraphy vs., 328–30
 sedation requirements, 329
 sensitivity, 329
 specificity, 329
 spine, 327, 549–50
 standards, 325, 336
 technical factors, 330, 713
- skeletal trauma, 23–48
 dating fractures, 208–15
 differential diagnosis
 accidental trauma, 268–75
 diseases, dysplasias, and
 syndromes, 217–35
 metabolic bone disease,
 240–9
 normal variants, 290–308
 obstetric trauma, 279–87
 osteogenesis imperfecta,
 254–64
 epidemiology and distribution,
 24–30
 factors predicting abuse, 24, 28–9
 lower extremity, 53–117
 thorax, 164–203
 upper extremity, 121–52
 vulnerable sites, 21
- skeleton, 9–21
 growth and development, 15–20
 normal variants, 290–308
 structure and function, 9–14
- skull
 anatomy and development,
 357–9
 fissures, 370–1, 378–9
 normal variants, 370–2
 osteogenesis imperfecta, 259,
 262
 vascular grooves, 371, 380
- skull fractures, 358–75
 accidental, 372–4, 382–3
 basilar, 364, 369, 372
 bilateral, 365–6, 373
 biomechanics, 360–3
 biparietal, 364, 366, 383
 birth-related, 287, 374–5, 384
 comminuted, 364, 370
 complex, 361–4, 367, 370
 depressed, 364–5, 367, 369–70
 differential diagnosis, 373–4,
 383
 diastatic, 365
 differential diagnosis, 370–5
 elevated, 364
 epidemiology, 25–7, 29–30,
 358–60
 growing, 365, 367, 388
 healing, 369–70, 377
 imaging patterns, 364–5
 imaging strategies, 386–8
 indicating need for skeletal
 survey, 387–8
 linear (simple), 361, 364, 369
 patterns in abuse, 365–9
 ping-pong, 369, 373, 383
 repeat, 369, 375
 soft tissue injuries, 362–5
- skull radiographs, 386
 basilar skull fractures, 364
 follow-up, 388
 mandibular fractures, 381, 385
 skeletal surveys, 327
 subgaleal hematoma, 362–3

Index

- small intestinal injuries, 567–76
 association with abuse, 564, 568
 intramural hematomas, 570–5
 miscellaneous, 575–6
 perforations, 567–70
 strictures, 574–6
- Smith, Shirley Ree, 650
- social isolation, 635
- social services, reporting
 suspected abuse, 642–3
- societal costs, abusive head
 trauma, 353
- Society for Pediatric Radiology
 (SPR), 684
 global skeletal imaging
 recommendations, 332
 Image Gently campaign, 682,
 684
- soft tissue injuries
 cranial, 363–4, 370
 facial, 375–8
 lower extremity, 54, 57
 pelvis, 101, 117
 radiographic fracture dating,
 213–14
 sexual abuse, 622
 shock due to, 620
 upper extremity, 122, 126
 whole-body MRI, 600
- sonography, 680
 abdominal visceral trauma,
 599
 clinical role, 330, 680
 cranial. *See* cranial sonography
 distal femoral CML, 70
 distal femoral epiphyseal
 separation, 71, 80–1
 distal humeral physeal injuries,
 139, 142, 145
 liver injuries, 582
 obstetric trauma, 281–5
 osteomyelitis and septic
 arthritis, 224–5
 pancreatic injuries, 578–9
 proximal humeral epiphyseal
 separation, 128, 133
 rib ossification defects, 300
 small bowel hematoma, 572–5
 spinal subdural hemorrhage,
 522
 SPNBF, 29, 32
 spatial resolution, x-ray image,
 673–4
- SPBC. *See* subperiosteal bone
 collar
- special needs, children with,
 608–10, 634
- specimens, postmortem
 high-resolution CT, 337
 histologic analysis, 337
 radiography, 337
 resection, 337
- SPECT (single photon emission
 computed tomography),
 679
- sphenoparietal venous sinus
 injury, 402
- spin echoes, 693–4
- spinal cord injuries
 cervical, 522–9
 clinical issues, 527–8
 imaging, 517, 528–9
 mechanisms, 525–7
 pathology, 522–5
 without radiographic
 abnormality
 (SCIWORA), 527–8
- spinal dysraphism (myelodysplasia),
 218, 221–3
 birth-related fractures, 279
- spinal trauma, 343, 494–553
 birth-related, 287, 549
 cervical spine. *See* cervical
 spine injuries
 craniocervical junction.
See craniocervical
 junction (CCJ) injuries
 dating, 551
 differential diagnosis, 547–9
 editor's note, 343
 epidemiology and patterns,
 495–502
 historical descriptions, 495
 imaging strategies, 549–53
 outcome, 495
 paraspinal calcification, 542,
 544
 sacral fractures, 544–5
 thoracolumbar.
See thoracolumbar spine
 injuries
 vertebral body fractures.
See vertebral body
 fractures
- spine. *See also* vertebrae;
 vertebral body fractures
 MRI, 491
 postmortem specimen
 resection, 337
 prevalence of fractures,
 495–502, 504
 relevant anatomy, 504–8,
 531–2
 three-column concept, 533
- spine radiographs
 cervical spine injuries, 508, 551
 facet joint disruption and
 dislocation, 540
 hypopharyngeal perforation, 591
 osteogenesis imperfecta, 256,
 259, 262
 sacral fractures, 544
 skeletal surveys, 327, 549–50,
 552
 thoracolumbar spine injuries,
 527–8, 531, 534
- spin-echo imaging, 695–6,
 699–701
- spinous process fractures, 533,
 536
- spiral fractures of long bones, 33
 accidental, 271, 273
 femur, 63–4
 humerus, 135
 tibia, 72, 87
- Spivack, Betty, 649
- splenic injuries, 583–4
- SPNBF. *See* subperiosteal new
 bone formation
- spoiling, gradient-echo imaging,
 702, 706
- spondylometaphyseal dysplasia
 (SMD), 218
 Japanese form, 218
 Sutcliffe/corner fracture type,
 217, 219–20
- spongiosa
 primary, 18–19
- spur, metaphyseal, 294–6
- stairways, falls down, 271,
 479
- starvation, 612–13
- State v. Edmunds*, 649–50
- steady state free precession
 (SSFP) imaging, 702–3
- step-off configuration, normal
 metaphyseal cortex,
 292–5
- step-off deformity, cortical,
 proximal tibia, 79, 90
- sternal fractures, 195–7
 accidental, 275
- sternoclavicular dislocation,
 280
- STIR imaging, 701–2. *See also*
 whole-body magnetic
 resonance imaging
- stomach
 injuries, 565–7
 neglected children, 611
- strangulation
 brain injuries, 467–8, 470
 necklace calcification, 619
- stress, family, 634–5
- stroke, adult, imaging features,
 454, 460
- study designs, 310
- study quality, 315
- subarachnoid hemorrhage
 (SAH), 403–7
 imaging, 405–6
 neoplasms mimicking, 434,
 438
 nontraumatic causes, 404,
 406
 pseudo- and other mimics,
 406–7
 skull fracture association,
 359
- subarachnoid space, 396
 benign enlargement, in infancy
 (BESSI), 418, 420–3
 enlarged, differential diagnosis,
 421
 tumor cell deposits, 438
- subdural effusion, 415, 417–19
 chronic SDH vs., 430–1
 hemorrhagic complications, 436
- subdural empyema, 416, 419
 chronic SDH vs., 431
- subdural hematoxygroma, 412,
 414–15
 evolving into chronic SDH,
 417, 426
 mixed attenuation,
 nonenhanced CT, 424
- subdural hemic and nonhemic
 collections, 397, 407–19
 challenges in interpreting,
 420–31
 imaging strategies, 488
 spinal, 491, 547
- subdural hemorrhage (SDH),
 407–12
 accidental vs. abusive, 421,
 423–4
 acute interhemispheric,
 412–13
 as indicator of abuse, 407
 benign enlargement of
 subarachnoid spaces in
 infancy and, 419–23
 biomechanics, 407–8
 birth-related, 418, 420
 challenges in interpreting,
 420–31
 choking/coughing as
 explanation, 441, 471
 chronic. *See* chronic subdural
 hemorrhage
 clinical differential diagnosis, 347
 compartmentalization, 408
 dating, 397–400, 417, 422, 426–7
 historical descriptions, 345–6
 hypoxic ischemic
 encephalopathy (HIE)
 and, 440–1
 imaging strategies, 488
 ipsilateral hypoxic-ischemic
 injury, 472–4, 477
 legal aspects, 648–9
 mixed attenuation,
 nonenhanced CT, 398,
 422, 424–5
 neuropathologic aspects,
 407–12
 nontraumatic causes and
 mimics, 412, 427,
 430–9
 retroclival, 519–20, 530–1
 skull fracture association, 359
 spinal, 515, 517–22, 547, 553
 anatomy and mechanisms,
 517
 differential diagnosis, 549
 imaging, 514, 516–22
 spontaneous, in infants
 (SSDHI), 439
 unsubstantiated theories of
 origin, 440–1

- subdural hygroma, 413–16
 chronic SDH vs., 417, 430
 evolving into chronic SDH, 417, 426, 428
- subdural space (or compartment), 396, 408–9
- subgaleal hematoma, 362–4, 372, 619
- subgaleal space, 357
- subperiosteal bone collar (SPBC), 16–17
 classic metaphyseal lesion, 37–9
 distal radius and ulna, 146, 154
 distal tibia, 89
 normal variants, 292–3
 proximal humerus, 123, 127
 proximal tibia, 75, 83
 rickets, 243
 vulnerability to trauma, 21
- subperiosteal hematoma, cranial (cephalohematoma), 363–4, 372
 birth-related, 374–5, 384
- subperiosteal hemorrhage, 27, 29–32, 212
- subperiosteal new bone formation (SPNBF), 30–2
 alternative causes, 32, 218, 228–32
 Caffey disease, 231
 classic metaphyseal lesion, 69
 clavicle, 194–5, 280
 dating fractures, 209, 214–15
 distal femoral epiphyseal separation, 71
 distal humeral physeal injuries, 139, 144–5
 distal radius and ulna CMLs, 148, 155
 distal tibial CML, 92, 105
 fibular fractures, 94
 healing fractures, 210, 212
 healing physeal injuries, 213
 humeral shaft fractures, 135–6
 mandibular, 381, 387
 obstetric trauma, 280–1
 physiologic, 32, 290–3
 fractures through pre-existing, 64, 69, 291
 prostaglandin E1-induced, 232
 proximal humeral epiphyseal separation, 128–30, 134
 proximal tibial epiphyseal separation, 82
 radiographic features, 28, 32
 radionuclide bone scans, 30, 32
 radius and ulna, 144, 148, 151
 ribs, 174, 176–7, 182
 sonographic features, 29, 32
 spurious/pseudo
 humerus, 291, 294–5
 ribs, 295, 305
 supracondylar distal humerus, 137, 140
- tibial shaft fractures, 88
 timing of appearance, 210
 vitamin A intoxication, 228
- subphyseal metaphyseal radiolucency, normal infants, 298–9
- subpial hemorrhage, 396, 417, 420
- subpoenas, 646–7, 661
- subscalp
 anatomy, 358
 injuries, 363–4
- sudden infant death syndrome (SIDS), 336, 338
- sudden unexpected infant death (SUID), 336, 338
- suffocation, 467, 469
- sulfite oxidase deficiency, isolated (ISOD), 477
- superior pubic ramus (SPR) fractures, 101, 114
 lucencies in osteopetrosis, 219, 221
 normal ossification variants, 307–8
- supraspinous ligament, 508
- susceptibility-weighted imaging (SWI), 489–90
- swirl sign, epidural hemorrhage, 401, 403
- synchondroses, cranial, 358–9
 normal variants, 372
 traumatic diastasis, 365
- synchrotrons, 681
- Syndrome de Silverman-Ambroise Tardieu, 3
- syphilis, congenital, 218, 226, 228
- systematic reviews, 310
- T1, 695
- T1-weighted gradient-echo imaging, 701–3
- T1-weighted spin-echo imaging, 697, 699–701
- T2, 694–5
- T2*, 694–5
- T2*-weighted gradient-echo imaging, 701–3
- T2-weighted spin-echo imaging, 697, 699–701
- Tardieu, Ambroise, 3, 358
- technetium 99m (^{99m}Tc) methylene diphosphonate (MDP)/medronate scans. *See also* bone scintigraphy
 radiation dose, 329
 radiographic skeletal surveys vs., 328
 rib fractures, 182–3, 189
 toddler's fracture, 270
- tectorial membrane (of atlanto-axial joint), 507–8
 injuries, 529–31
- telephone contacts, abuse
 investigators, 660
- televisions, falling, 272–3, 475, 479
- “temporary brittle bone disease” (“TBBD”), 316–20
 Miller, 317–19
 Paterson, 247, 316–17
- tetanus, neonatal, acromial fractures, 198
- thalassemia, 234
- thigh
 skeletal trauma. *See* femur
 soft tissue injuries, 54, 57
- thoracic compression, manual, 168, 171–4, 176, 178
- thoracic spine fractures, 527–8, 534–5. *See also* thoracolumbar spine injuries
- thoracic trauma. *See also* rib fractures
 bony, 164–203
 cardiac injuries, 597
 esophageal perforation, 593
 internal injuries, 319, 594–7
- thoracolumbar spine injuries, 531–44
 chronic changes, 535
 fracture-dislocations, 536–44, 546
 imaging features, 527–8, 534–5
 imaging strategies, 549–53
 isolated posterior ligamentous injuries, 544, 547
 mechanisms of injury, 532
 patterns of injury, 533–4
 posterior elements, 535–44
 relevant anatomy, 521, 531–2
 spinous process fractures, 533, 536
 vertebral body fractures. *See* vertebral body fractures
- three column spine concept, 533
- three-dimensional (3D)
 constructive interference in steady state MRI, 491
- three-dimensional (3D) CT models
 rib fractures, 182, 184, 188
 scalp injuries, 362–3
 skull fractures, 364–5, 367, 371–2
- three-dimensional Fourier transform (3D-FT) MRI, 692
 gradient-echo imaging, 701
 spin-echo imaging, 700
- thymic involution, 596, 613
- tibia, 71–93
 accidental trauma, 270, 272
 distal. *See* distal tibia
 physiologic SPNBF, 291–3
 proximal. *See* proximal tibia
 shaft fractures, 86–8, 95–6
- tidemark cartilage, 20
- time-of-flight (TOF) magnetic resonance angiography (MRA), 697, 702
- tin ear syndrome, 377
- tissue weighting factors (w_T), 671
- toddlers (1–3 years)
 biomechanics of falls, 352
 distal femoral and proximal tibial metaphyseal irregularities, 295–8
 epidemiology of skeletal injuries, 27
- toddler's fracture, 87, 97, 270, 272
 hyperextension, 272
- toe fractures, 97, 112
- tomosynthesis, digital (DT), 684
- tooth, in mediastinum, 593
- torture, child, 623
- Toxocara* infection, 480
- trabeculae, bone, 12
 pediatric skull, 360–1
- trabecular transition zone, 19, 21
- tracheal rupture, 594
- training, radiology staff, 636
- transaminases, serum, 580–1
- transverse fractures of long bones, 33–4, 36
 femur, 63, 65
 humerus, 135–6
 obstetric trauma, 281
 radius and ulna, 143, 150–1
- transverse ligament of atlas, 505–6, 508
- traumatic brain injury (TBI). *See* abusive head trauma; intracranial injury
- trauma-X, 1
- trials. *See* legal proceedings
- trichobezoar, 566, 613
- trichopolydystrophy. *See* Menkes disease
- troponins, cardiac, 597
- tube current (mA), 668
 skeletal surveys, 713
- Tuerkheimer, Deborah, 650
- tumors. *See* neoplasia
- turbo spin echo (TSE). *See* fast spin echo
- two-dimensional Fourier transform (2D-FT) MRI, 691–2
 spin-echo imaging, 700
- tympanic membrane, perforated, 376
- ulcerative colitis, neglected, 613
- ulna, 141–8
 distal
 CML, 146–8, 155
 fractures, 145–8, 154
 normal anatomy, 146, 154
 midshaft fractures, 141–5, 151–2

Index

- ulna (cont.)
 normal distal cupping, 294–6
 physiologic SPNBF, 291
 proximal fractures, 141, 149
 ultrasonography. *See* sonography
 upper extremity
 accidental trauma, 273–4
 fractures, 121–52
 soft tissue injuries, 122, 126
 upper gastrointestinal (uGI)
 series, 599–600
 duodenal perforation, 570
 gastric hematoma, 566–7
 gastric ulcer, 567
 small bowel hematoma, 571–4
 urinary tract injuries, 586–9
 urinoma, 587
 Uscinksi, Ronald, 648–9
- vaginal delivery, difficult/
 instrumented, 279–81,
 283, 285, 287
 vaginal injuries, 621–2
 vascular grooves, cranial, 371,
 380
 vascular injuries, 597
 liver, 582
 neck, 594
 vascular malformations,
 intracranial, 432, 436
 venous infarcts, cerebral, 474–5
 ventricular septal defect (VSD),
 traumatic, 597
 ventricular shunting, SDH
 complicating, 427
- vertebrae
 facet disruption and
 dislocation, 539–44
 fracture-dislocations, 536–44,
 546
 limbus, 535
 neurocentral synchondrosis
 injuries, 536–40, 544, 546
 relevant anatomy, 531
 spinous process fractures, 533,
 536
 vertebral artery hemorrhage, 594
- vertebral body
 normal, 521–2, 531
 wandering, 538–9
- vertebral body compression
 fractures, 533–4
 imaging features, 527–8, 532,
 534–5
 osteogenesis imperfecta, 262
 prevalence in abuse, 501
 radiologic–histopathologic
 correlates, 523, 534
- vertebral body fractures
 cervical, 510, 515
 chronic effects, 533, 535
 imaging features, 534–5
 pattern 1 (compression
 deformity). *See* vertebral
 body compression
 fractures
 pattern 2 (anterosuperior end
 plate), 525, 534
 pattern 3 (combined lesions),
 526, 534
 prevalence in abuse, 495–501
 radiologic–histopathologic
 correlates, 523, 525–6, 534
- vertebral notching
 abusive, 533, 535
 normal, 549, 551
- vertically orientated fractures,
 tibial shaft, 87, 97
- viral encephalitis, 436, 476, 480
- visceral trauma, 563–600
 abdominal, 565–89
 differential diagnosis, 598–9
 epidemiology, 563–5
 imaging strategies, 599–600
 lower rib fractures with, 186–7,
 190, 596
 mechanisms of injury, 565
 mortality, 565
 neck and thorax, 590–7
 occult, 565
- visual aids, to expert testimony,
 653, 663
- vitamin A intoxication, 218,
 228–32
- vitamin C deficiency (scurvy),
 218, 226–9
- vitamin D, 240–2
 deficiency
 abuse with, 249
 maternal, congenital rickets,
 244–5
 prenatal, lack of effect,
 242–3
 rickets, 243, 245–6
 “temporary brittle bone
 disease”, 317
 neonates, 243
 prenatal function, 242–3
 supplements during
 pregnancy, 243
- vitamin D receptor (VDR),
 prenatal function, 242
- vitamin K deficient bleeding
 (VKDB), 430, 432–3
- von Willebrand disease, 433
- wandering vertebral body, 538–9
- water molecules, in MRI, 688–9,
 697
 diffusion in tissue, 696
 distinction from fat, 701–3
 rotational and translational
 motions, 694–5
- whiplash injuries, 407, 527,
See also shaking
- whiplash shaken infant
 syndrome, 346
- white matter injuries
 bilateral symmetric, anoxia,
 467
 parenchymal lacerations,
 464–6
 shear-strain. *See* diffuse axonal
 injury
- white matter, MRI, 697–8
- whole-body magnetic resonance
 imaging (WB-MRI), 326,
 701
 femoral fractures, 62–3
 rib fractures, 184, 188
 visceral injuries, 581, 600
- whooping cough (pertussis), 441,
 471
- Wimberger sign, 226
- witnesses
 expert. *See* expert witnesses
 fact, 647–8
 sequestration, 653, 656
- Wormian bones, 371–2
 normal young children, 371,
 381
 osteogenesis imperfecta,
 261–2
- woven bone, 10–11
 classic metaphyseal lesion, 42
 healing fractures, 212–13
- wrap-around artifact, MRI, 700
- wrap-around skull fracture, 364
- x-ray image, 672–4
 contrast, 672–4
 physiologic motion and
 voluntary movement,
 674
 quality, concepts, and metrics,
 681
 spatial resolution, 673–4
- x-ray tube, 668
- x-rays, 668–79
 detection systems, 674–9
 discovery, 668
 interactions with matter, 669
 plain film imaging.
See radiography
 polychromatic vs.
 monochromatic, 681
 properties, 669
 radiation dose terminology,
 669–71
 sources, 668
- zone of provisional calcification
 (ZPC)
 histologic (hZPC), 18–19
 rachitic changes, 243
 radiologic (rZPC), 19
 CML, 36–9
 zones, growth plate, 17–18