

# Index

- 100 Hue test, 252
- ABC model of anosognosia, 166
- abnormal involuntary movement, 43  
 classification, 43  
 permanent movement disorders, 46  
   dystonia, 46–8  
   tremor, 48–50  
   vascular generalized chorea, 52  
   vascular parkinsonism, 50–2  
 transient movement disorders, 43  
   asterixis, 45–6  
   episodic paroxysmal dyskinesia, 46  
   hemiballism–hemichorea, 43–5  
   *see also specific conditions*
- abulia, 335  
 abulia major, *see* akinetic mutism  
 abulia minor, 337
- acalculia, 23, 101  
 assessment, 110  
 dissociating numbers and language, 108–9  
 Gerstmann syndrome, 109–10  
 mental calculation deficits, 106–8  
   approximate vs. exact calculation, 107–8  
   conceptual processing, 108  
   elementary calculation, 107  
   executive control and sequencing, 107  
   input/output processes, 106  
   spatial layout of arithmetic problems, 106–7  
 number-transcoding deficits, 105  
 quantity-processing deficits, 105–6  
   core semantic deficits, 105–6  
   neglect and the number line, 106, 145  
   numerosity perception, 106  
   rehabilitation, 110  
   spatial acalculia, 104
- achromatopsia, 39, 234, 250
- activation regulation function  
 assessment, 267
- activities of daily living  
 apraxia impact, 60
- assessment, 417–18  
 fatigue impact, 381  
 standardized observation, 59  
 training, 60
- acute confusional state (ACS), 315, 340  
 characteristics, 340–1  
 differential diagnosis, 342  
 management, 346–7  
 pathophysiology, 341–2  
*see also* delirium
- aggressive behavior, 358–9
- agnosia, 35, 247  
 apperceptive, 214, 247, 248  
   form agnosia, 248  
   integrative agnosia, 248  
 associative, 247, 248–50  
   from disconnection, 249  
   from loss of semantic knowledge, 249–50  
 auditory, 114, 118–19, 120  
 object, 248–50  
 visual, 247  
   assessment, 251–3  
   causes, 250  
   color, 250  
   management, 253  
   prognosis, 253  
   strokes and, 250–1
- agraphia, 86, 89, 90  
 assessment, 93–4  
 causative strokes, 90–1, 93  
   MCA lesions, 92–3  
   clinical presentation, 89  
   management, 95  
   prognosis, 94–5  
   with alexia, 87
- akinetic mutism, 23, 25–6, 38, 317, 334, 335  
 associated strokes, 336–7  
 causes, 336  
 differential diagnosis, 335–6  
 fronto-diencephalic form, 334  
 management, 337  
 thalamic form, 334
- Albert Test, 144
- alexia, 23, 35, 86, 90
- assessment, 93–4  
 causative strokes, 90–1, 93  
   MCA lesions, 92–3  
   PCA lesions, 91–2  
 clinical presentation, 89  
 management, 95  
 prognosis, 94–5  
 with agraphia, 87  
 with anomia, 88–9  
 without agraphia, 87  
*see also* dyslexia
- Alice in Wonderland syndrome, 181
- alien hand phenomenon, 33, 37
- allocentric neglect, 87
- alteration of level of consciousness, 312–13  
 acute conditions, 315–17  
 evaluation, 312–13  
 frequency, 320  
 mechanisms, 318–20  
 prognostic significance, 320  
 stroke syndrome associations, 320–30  
   anterior cerebral artery lesions, 330  
   MCA lesions, 330  
   midbrain lesions, 321–3, 327–9  
   pontine lesions, 327–9  
   rostral basilar artery occlusion, 320–1  
   thalamic lesions, 321–3, 324–7  
   subacute/chronic conditions, 317
- alternating motion rate, 79
- Alzheimer disease, 3–4, 397  
 treatment, 427
- Alzheimer Disease Assessment Scale-Cognition (ADAS-Cog), 417
- amnesia, 24, 38  
 assessment, 287  
 management, 288–9  
 prognosis and recovery, 288  
 thalamic lesions and, 36  
*see also* episodic memory; memory disturbances

- amusia, 114, 119–20
- $\beta$ -amyloid (A $\beta$ ), 4, 397
- amyloid angiopathy, 2, 12
- amyloid precursor protein, 4, 397
- anarthric tetraparesis, 336
- anomia, with alexia, 88–9
- anomic aphasia, 35, 67
- anosodiaphoria, 38
- anosognosia, 35, 38, 127, 158, 172–3
- anatomical correlates, 142
- as prognostic marker, 162–3
- assessment, 163–4
- causative strokes, 161–2
- clinical features, 158–9
- degree of severity, 159
- for blindness, 209
- incidence after stroke, 159–61
- pathogenesis, 164–6
- ABC model, 166
- concomitant disorders, 164
- disconnection mechanism, 165
- discovery theory, 165
- feedforward theory, 165–6
- motor awareness theory, 166
- psychological motivation, 165
- two-factor theory, 166
- unilateral neglect relationship, 164–5
- recovery, 162
- therapy, 166–7
- Anosognosia for Hemiplegia Questionnaire, 163
- anterior cerebral artery, lesions
- alteration of level of consciousness, 330
- delirium and, 344–5
- dysexecutive syndrome, 263
- hemineglect association, 138
- left, 33–5
- right, 37
- anterior choroidal artery, lesions
- left, 35
- right, 37
- anterior communicating artery
- aneurysm, 264–5, 282–3, 337
- antidepressant therapy, 367–9
- antihypertensive therapy, 424–5, 426–7
- antipsychotic therapy, 353
- antithrombotics, recurrent stroke
- prevention, 424
- Anton syndrome, 39, 209
- anxiety, fatigue association, 377
- apallic syndrome, *see* persistent vegetative state
- apathy, 38, 256–7, 358
- CADASIL and, 394
- Apathy Evaluation Scale, 417
- aphasia, 23, 65, 284–5
- aphasic syndromes, 65–6, 68–9
- crossed aphasia, 69
- mixed transcortical aphasia, 68
- subcortical aphasia, 69
- assessment, 71–2
- imaging techniques, 70–1
- oral comprehension, 65
- oral production, 65
- causative stroke, 69–70, 284–5
- MCA infarct association, 35, 69
- characteristics, 69
- fluent, 35, 67–8
- anomic aphasia, 67
- conduction aphasia, 68
- transcortical sensory aphasia, 68
- Wernicke aphasia, 67–8
- mimicking conditions, 71
- non-fluent, 66–7, 336
- Broca aphasia, 67
- global aphasia, 66
- transcortical motor aphasia, 67
- optic, 35, 247, 249
- prognosis, 72
- treatment, 72–3
- pharmacotherapy, 73
- speech and language therapy, 72–3
- transcranial magnetic stimulation, 73
- aphasic mutism, 23
- apperceptive agnosia, 214, 247, 248
- form agnosia, 248
- integrative agnosia, 248
- apperceptive prosopagnosia, 236
- Apple test, 145
- apraxia, 23, 285
- assessment, 57
- gestural impairment, 58–9
- object/tool use impairment, 59
- classification, 56–7
- constructional, *see* visuoconstructive disorders
- functional implications, 59–60
- limb apraxia, 55
- neural basis, 61
- of speech, 35
- prognosis and recovery, 60
- treatment, 60–1
- apraxia screen of TULIA, 58–9
- arousal, 313
- ascending reticular activating system (ARAS), 313–15
- asomatognosia, 170–1
- assessment, 183–5
- documentation, 184
- hemiasomatognosia, 171–3
- phantom limbs, 173–6
- supernumerary, 174–6
- rehabilitation, 185
- Assessment of Intelligence in Dysarthric Speakers, 78
- Association Internationale Polar la Recherche et “Enseignement en Neurosciences” (AIREN), *see* NINDS-AIREN
- associative agnosia, 247, 248–50
- from disconnection, 249
- from loss of semantic knowledge, 249–50
- associative prosopagnosia, 236
- asterixis, 45–6
- asymmetry, *see* functional asymmetry
- ataxic dysarthria, 80–1
- athymormia, 334
- associated strokes, 337
- see also* akinetic mutism
- attention, 297, 302
- dorsal attention network, 302
- ventral attention network, 302
- attention disorders, 258
- attentional process assessment, 267
- audiometry, *see* auditory perception disorders; deafness
- auditory agnosia, 114, 118–19, 120
- auditory neglect, 120
- auditory perception disorders, 114, 116
- anatomical correlations, 115–16, 120
- audiometric assessment, 121–2
- dichotic listening test, 122
- evoked auditory potentials, 122
- oto-acoustic emissions, 122
- sequences of auditory signals, 122
- standard pure tone audiometry, 121
- word discrimination test, 121
- see also* deafness
- auditory hallucinations, 121
- brainstem lesions and, 120–1
- causative strokes, 116
- delayed disorders, 115
- diagnostic circumstances, 114–15

## Index

- auditory hallucinations (*cont.*)  
   neuropsychological assessment, 122–3  
   auditory lexical decision tasks, 123  
   linguistic and speech perception tests, 122–3  
   non-speech sounds, 123  
   phonemic identification and discrimination, 123  
   receptive musical ability, 123  
   semantic access and processing, 123  
   prognosis, 123  
   transient disorders, 115  
   treatments, 123–4
- autoscopic phenomena, 182–3
- Babinski–Nageotte syndrome, 318
- Baking Tray task, 145
- Bálint syndrome, 24, 39, 218–19, 302–3  
   assessment, 214, 225–7  
   associated deficits, 221–2  
   causes, 224  
   complete, 219  
   differential diagnosis, 224  
   gaze apraxia, 220  
   in stroke, 224–5  
   lesion locations, 222–4  
   minor forms of, 222  
   numerosity perception deficit, 106  
   optic ataxia, 219–20  
   prognosis, 227  
   rehabilitation strategies, 227  
   simultanagnosia, 220–1
- ballism, 44  
   *see also* hemiballism–hemichorea
- Barthel Index, 417
- basal forebrain lesions, memory impairment, 282–3
- basic activities in daily living, *see* activities of daily living
- basilar artery occlusion, 320–1, 329
- Beck Depression Inventory (BDI), 364
- Behavioral Dyscontrol Scale, 267
- Behavioral Inattention Test, 143
- Behavioural Assessment of the Dysexecutive Syndrome, 267
- bell cancellation test, 144
- Benton Face Recognition Test, 236, 241
- bilateral anterior opercular syndrome, 39
- bilateral deafness, 23
- bilateral lesions, 38–9
- Binswanger disease, 392–3
- bipolar disorder, 354  
   structural brain abnormalities, 355  
   *see also* depression; mania
- Birmingham Object Recognition Battery (BORB), 240, 252
- blindness  
   differential diagnosis, 214  
   psychogenic, 213  
   *see also* cortical blindness; *specific conditions*
- blindsight, 210
- Block-Tapping Task, 205
- bodily self, 170–1  
   image, splitting of  
   *see also* asomatognosia
- body disorders, 180  
   *see also specific disorders*
- Boston Diagnostic Aphasia Examination, 65, 71, 416
- Boston Naming Test, 415
- brain atrophy, 3  
   fatigue and, 378–9
- brainstem evoked auditory potentials, 122
- brainstem lesions  
   hearing disorders and, 120–1
- Brief Visual Memory Test, Revised, 415
- Brixton Test, 258
- Broca aphasia, 67
- Broca's area  
   working memory and, 301  
   written language disorders and, 93
- CADASIL (cerebral autosomal dominant arteriopathy with subcortical infarcts and leukoencephalopathy), 2, 12, 393–6, 400–1  
   clinical features, 4  
   white matter abnormalities, 394–6
- calcarine artery occlusion, 212
- calculation, 102  
   *see also* acalculia; number processing
- California Verbal Learning Test (CVLT), 415
- caloric vestibular stimulation, 147, 166, 185
- Cambridge Face Memory Test, 241
- Canadian Stroke Newton (CSN), *see* NINDS-CSN
- cancellation tasks, 144–5
- Capgras syndrome, 37, 352, 353
- capsular genu lesion, vascular dementia and, 392
- CARASIL (cerebral autosomal recessive arteriopathy with subcortical infarcts and leukoencephalopathy), 396
- catastrophic reaction, 36
- Catherine Bergego Scale, 164
- caudate nucleus lesions  
   delirium and, 345  
   vascular dementia and, 392
- CELEB test, 241
- Center for Epidemiologic Study, Depression Scale (CES-D), 416
- central executive, 298  
   deficits, 303–5
- central rostrocaudal deterioration syndrome, 318–20
- centromedian nucleus lesions, 48
- cerebellar tremor, 48–9
- cerebral amyloid angiopathy, *see* amyloid angiopathy
- cerebral hemorrhage, 265  
   vascular dementia and, 397–8
- cerebral microbleeds, 3
- cerebral venous thrombosis, 265
- Chicago Multiscale Depression Inventory, 416
- cholesterol-lowering therapy, 425
- cholinergic deficiency, delirium and, 342
- cholinesterase inhibitors  
   Alzheimer disease treatment, 427  
   CADASIL treatment, 12  
   delirium treatment, 347  
   vascular dementia treatment, 402–3, 427–8
- chorea, 44  
   vascular generalized chorea, 52  
   *see also* hemiballism–hemichorea
- clock drawing test, 198, 199
- Cloninger Temperament and Character Inventory, 359
- Cognitive and Graded Activity Training trial, 383

- cognitive behavioral therapy  
 depression, 369  
 fatigue treatment, 383
- cognitive impairment, 22  
 acute focal cognitive deficits, 22–6  
 central visual disturbances, 24  
 dysexecutive syndrome, 25–6  
 language disorders, 22–3  
 memory disturbances, 24–5  
 right hemisphere syndrome, 24  
 cerebrovascular disease and, 2–4  
 diagnosis, 28  
 diffuse cognitive disturbances, 26–8  
 mixed pathology, 3–4  
 post-stroke impairment detection, 29  
*see also* vascular cognitive impairment
- cognitive strategy training, 60
- Collier's sign, 321
- color agnosia, 250
- color perception, 214  
 assessment, 252
- color recognition disturbances, 250  
 achromatopsia, 250  
 color agnosia, 250
- coma, 313, 317, 335  
 somnolent, 321  
 vigilant, 321  
*see also* alteration of level of consciousness; Glasgow Coma Scale
- Composite International Diagnostic Interview (CIDI), 364
- conceptual apraxia, 57  
*see also* apraxia
- conduction aphasia, 35, 68
- confabulations, 25, 282, 283
- configural face processing, 236
- configuration, 236
- confusion, *see* acute confusional state; delirium
- Confusion Assessment Method, 342
- Confusional State Evaluation, 343
- consciousness, 313  
 clouding of, 315  
*see also* alteration of level of consciousness
- constructional apraxia, *see* visuoconstructive disorders
- continuous positive airway pressure, 383
- convergence disorders, 321
- corpus callosum  
 lesions, 33–5, 104  
 number-processing role, 103, 104
- Corsi Block Tapping Test, 307
- cortical blindness, 24, 39, 209  
 assessment, 213–15  
 electroencephalography, 215  
 visual evoked potentials, 215  
 causes, 210  
 clinical symptoms, 209  
 management, 215  
 prognosis, 211–12  
 recovery, 212  
 residual vision, 209–10  
 stroke-related, 211  
 causative strokes, 212–13
- cortical deafness, 23, 114, 116–17, 120
- Cotard syndrome, 181, 182, 352
- C-reactive protein, 379
- crossed aphasia, 69
- deafness  
 bilateral, 23  
 cortical, 23, 114, 116–17, 120  
 pure word, 114, 117–18, 120  
 subtypes, 118  
*see also* auditory perception disorders
- Déjerine syndrome, 318
- delayed-response tests, 259
- delirium, 26, 315, 340  
 assessment, 342–3  
 causative strokes, 344–6  
 anterior cerebral artery infarcts, 344–5  
 head of the caudate nucleus, 345  
 inferior genu of the interior capsule, 345  
 MCA infarcts, 38, 345  
 PCA infarcts, 344  
 subarachnoid hemorrhage, 345–6  
 thalamic stroke, 344  
 characteristics, 340–1  
 diagnostic criteria, 340, 341  
 differential diagnosis, 340–1  
 incidence in stroke, 344  
 management, 346–7  
 natural history in stroke, 344  
 prognostic significance, 346  
 risk factors in stroke patients, 346  
 subtypes, 340  
*see also* acute confusional state
- Delirium Assessment Scale, 343
- Delirium Index, 343
- Delirium Rating Scale, 343
- Delirium Symptom Interview, 343
- delusion, 38, 351  
 assessment, 353  
 causative strokes, 352  
 frequency in stroke patients, 352  
 misidentification delusion, 352–3  
 treatment, 353, 354
- dementia  
 CADASIL and, 394  
 cerebrovascular disease and, 2  
 pre-stroke dementia detection, 29  
*see also* cognitive impairment; vascular dementia
- demonic possession, 181
- depersonalization, 180–1
- depression, 36  
 assessment, 416–17  
 fatigue association, 377  
 frequency in stroke patients, 366  
 future directions, 370  
 impact after stroke, 367  
 management  
 electroconvulsive therapy, 370  
 pharmacotherapy, 367–9  
 psychological therapies, 369–70  
 personality change relationship, 359  
 routine screening in stroke patients, 365–6  
 symptoms, 364–5  
 natural history, 366  
 vascular depression, 366–7
- diabetes mellitus, 4  
 fatigue and, 377  
 management, 425–6  
 recurrent stroke prevention, 423
- Diagnostic and Statistical Manual of Mental Disorders* (DSM), 10, 340, 351, 363, 388–9
- dichotic listening test, 122
- digit span test, 307
- Digit Symbol Substitution Test, 414
- directional hypokinesia, 130
- disconnection syndromes, 33–5  
 anosognosia, 165
- discovery theory of anosognosia, 165
- disengagement deficit, 146
- disinhibition, 256
- disintegration syndrome, 35
- disorientation, 26

## Index

- divided attention deficit, 259  
 dopamine 1 receptor stimulation, 300  
 Doppelgänger, 183  
 dorsal attention network, 302  
 double homonymous hemianopia, 211  
 double monocular elevator palsy, 321  
 double simultaneous stimulation, extinction on, 144  
 drawing tasks, 145  
 visuoconstructive disorder, 193–4  
 DSM, *see Diagnostic and Statistical Manual of Mental Disorders*  
 dysarthria, 22, 35, 76, 77  
   associated stroke types, 82  
   ataxic dysarthria, 80–1  
   classification, 77–8  
   evaluation, 78  
   flaccid dysarthria, 81  
   spastic dysarthria, 79  
   treatments, 82  
   unilateral upper motor neuron dysarthria, 79–80  
 dysexecutive syndrome, 25–6, 38, 255  
   assessment, 266–8  
   bedside assessment, 266  
   behavioral disturbances, 266–7  
   executive cognitive functions, 267  
   optional assessments, 267–8  
   behavioral dysexecutive syndrome, 255–7, 266–7  
   characteristics, 260  
   cognitive dysexecutive syndrome, 257, 258, 267  
   divided attention and task coordination, 259  
   episodic memory impairment, 259–60  
   impact on daily life, 261  
   information generation deficit, 258–9  
   management, 268–9  
     cognitive rehabilitation, 268–9  
     compensatory approach, 269  
     pharmacological treatment, 268  
   response initiation and suppression, 257–8  
   response slowing, 260  
   rule use and problem-solving deficits, 258  
   stroke and, 260–5  
     determinants, 261  
     lacunar infarct, 264  
     prevalence, 260–1  
     subarachnoid hemorrhage, 264–5  
     territorial infarct, 262–4  
   sustained attention deficit, 259  
   working memory, 259  
   *see also* executive function  
 dysgraphia  
   surface, 88  
 dysgraphia, surface, 87  
 dyskinesia, episodic paroxysmal, 46  
 dyslexia  
   associated lesions, 91  
   deep, 88  
   neglect, 87  
   phonological, 88  
   surface, 87, 88  
   *see also* alexia  
 dyslipidemia management, 427  
   *see also* hypercholesterolemia  
 dystonia, 46–8  
   midbrain lesions, 48  
   myoclonic, 47  
   striato-pallidal lesions, 46–7  
   thalamic lesions, 47–8  
 electroencephalography, cortical blindness, 215  
 Ekblom syndrome, 181–2  
 electroconvulsive therapy, depression, 370  
 emotional disturbances, 256  
 emotional incontinence, 38, 358  
 environmental agnosia, 37  
 environmental dependency syndrome, 256, 257  
 episodic buffer, 298  
 episodic memory, 277–8  
   assessment, 205–6, 287  
   impairment, 24–5  
   dysexecutive syndrome, 259–60  
   management, 288–9  
   prognosis and recovery, 288  
   vascular dementia, 398–9  
   strokes affecting executive systems, 286–7  
   strokes affecting memory-processing systems, 284–6  
   language dominant hemispheric lesions, 284–5  
   left hemispheric lesions, 285  
   right hemispheric lesions, 285–6  
   strokes causing amnesia, 278–84  
   basal forebrain lesions, 282–3  
   fornix lesions, 280  
   medial temporal damage, 278–80  
   retrosplenial cortex lesions, 284  
   thalamic lesions, 280–2  
 episodic paroxysmal dyskinesia, 46  
 error analysis, 251  
 evoked auditory potentials, 122  
   of brainstem, 122  
   of late latency (cortical), 122  
   of middle latency (semi-early), 122  
 executive function, 255  
   CADASIL and, 394  
   vascular dementia effects, 398  
   *see also* dysexecutive syndrome  
 exercise  
   fatigue management, 383  
   fatigue relationship, 378  
 explicit memory, 277  
 extinction  
   on double simultaneous stimulation, 144  
   perceptual, 129–30  
   anatomical correlates, 141–2  
 eye movement disorders, 214, 220  
   assessment, 226  
   *see also* gaze apraxia  
 Eysenck Personality Inventory, 359  
 face inversion effect, 237  
 face recognition, 231  
*see also* prosopagnosia  
 familiarity, sense of, 352  
 fatigue, 375  
   after stroke, 375, 376, 379  
   diagnosis, 380, 381  
   epidemiology, 376  
   severity assessment, 380–1  
   after subarachnoid hemorrhage, 382  
   correlated factors, 382  
   epidemiology, 382  
   associated factors, 376–9  
   anxiety, 377  
   depression, 377  
   neurological impairment, 377  
   neurotransmitter/neuroendocrine changes, 379  
   physical fitness, 378  
   pre-stroke fatigue, 377  
   sleep disturbance, 378  
   sociodemographic factors, 377  
   somatic comorbidities, 377  
   stroke subtype, 378  
   systemic inflammation, 379  
   white matter changes/brain atrophy, 378–9  
   central fatigue, 376  
   classification, 379–80



- activity-dependent fatigability, 380  
 primary fatigue, 379  
 secondary fatigue, 379–80  
 impact on patients, 382  
 activities of daily living, 381  
 institutionalization, 381  
 patients' perception, 381  
 quality of life, 381  
 return to work, 381  
 peripheral fatigue, 376  
 post-stroke case definition, 380  
   community patients, 380  
   hospital patients, 380  
 treatment, 382–3
- feeling of a presence, 183
- fitness, fatigue relationship, 378
- Five Minute Protocol, 414
- flaccid dysarthria, 81
- fluent aphasia, 35, 36, 67–8  
*see also* aphasia
- Foix–Chavany–Marie syndrome, 39
- foreign accent syndrome, 35
- form agnosia, 248
- fornix lesions, memory impairment, 280
- Fregoli syndrome, 352
- Frontal Behavioral Inventory, 267, 401
- frontal lesions  
   memory impairment, 286–7  
*see also* dysexecutive syndrome
- Frontal Systems Behavior Scale, 417
- frontoparietal control system, 303
- Frontotemporal Behavioral scale, 401
- Full Outline of UnResponsiveness (FOUR) score, 313, 317
- functional asymmetry, 32–3  
   development, 33  
   left hemisphere syndromes, 33–6  
   right hemisphere syndromes, 34, 36–8
- functional independence measure, 417
- fusiform face area, 233, 301
- Gage, Phineas, 357
- galvanic vestibular stimulation, 147
- gaze apraxia, 220
- gaze paresis, 321
- generalized anxiety disorder, 38
- Gerstmann syndrome, 23, 109–10
- gestures, 55–6  
   apraxia impact on gestural communication, 59–60  
   gestural impairment assessment, 58–9  
   gestural training, 60
- Glasgow Coma Scale, 313, 317
- global aphasia, 66
- global hyperactivity, 256
- global hypoactivity, 256
- glucocorticoid hormones, 341–2
- Goal Management Therapy, 269
- GREFEX study, 260, 261
- Grober and Buschke Test, 394
- Groupe d'Etude sur la Rééducation et l'Evaluation de la Négligence, 143
- Hachinski Ischemic Score, 389, 390
- hallucinations, 26, 38  
   auditory, 121  
   autoscopic, 182  
   in cortical blindness, 209  
   kinesthetic, 174
- Halstead–Reitan Neuropsychological Battery, 412
- Hamilton Depression Rating Scale, 364
- health-related quality of life, 367
- hearing disorders, *see* auditory perception disorders; deafness
- heautoscopy, 182
- hemianacousia, 117, 122
- hemianopia, 128–9, 211–12  
   double homonymous, 211  
*see also* cortical blindness
- hemiasomatognosia, 171–3
- hemiballism–hemichorea, 43–5  
   associated lesions, 44  
   causes, 44  
   mechanisms, 44–5
- hemichorea, *see* hemiballism–hemichorea
- hemiconcern, 178, 179  
   acute, 37
- hemi-depersonalization, 172
- hemineglect, 24, 36–7, 87, 126  
   anatomical correlates, 136–7  
   cerebrovascular topographic associations, 137–8  
   chronic spatial neglect, 142
- distinct neglect components, 140–2
- egocentric vs. allocentric neglect, 140–1
- near vs. far space neglect, 141
- role of distributed brain networks, 138–40
- anosognosia relationship, 164–5
- assessment, 143–6
- causative strokes, 136
- characteristics, 134
- clinical manifestations, 126–8  
   dissociations between neglect subtypes, 133–4  
   frames of reference, 131  
   imaginal representation, 133  
   implicit processing, 134  
   motor deficits, 130–1  
   perceptual deficits, 128–30
- functional neurophysiological anomalies, 143
- occurrence in stroke, 135–6
- prognosis, 147
- sectors of space, 132–3  
   extrapersonal space, 132  
   peri-personal space, 132  
   personal space, 132
- therapeutic approaches, 147–8  
*see also* neglect
- hemiplegia, unawareness of, *see* anosognosia
- hereditary cerebral hemorrhages with amyloidosis of Dutch type, 397
- hereditary cystatin C amyloid angiopathy, 397
- herniation  
   central (transtentorial) downward, 318–19, 320  
   central (transtentorial) upward, 320  
   cingulate, 320  
   lateral (uncal), 318
- herpes simplex encephalitis, 71
- holistic face processing, 236  
   assessment, 236–7
- Holmes tremor, 48, 49
- Hopkins Verbal Learning Test (HVLT), 415
- horizontal segment of the intraparietal sulcus, 103
- Horner syndrome, 324–5
- hyperactivity, global, 256
- hypercholesterolemia, 423  
   management, 425

## Index

- hyperhomocysteinemia, 423–4  
 management, 426
- hyperlipidemia, 4
- hypersomnia, 317, 321
- hypertension, 4  
 fatigue association, 377  
 management, 424–5, 426–7  
 recurrent stroke prevention, 423
- hypoactivity, global, 256
- ICD-10, *see International Statistical Classification of Diseases and Related Health Problems*, 10th revision
- ideational apraxia, 57  
*see also* apraxia
- ideomotor apraxia, 57  
*see also* apraxia
- illusion des sosies, 352
- illusionary movements, 174
- imaging studies, 4–5  
 aphasia, 70–1  
 vascular cognitive impairment diagnosis, 16
- imitation, 55
- implicit memory, 277
- implicit processing, in spatial neglect, 134
- infarcts  
 silent, 2  
 subcortical, 2
- inferior occipital gyrus, face recognition role, 233
- inflammation, fatigue and, 379
- Informant Questionnaire on Cognitive Decline in the Elderly, 29
- information generation deficit, 258–9
- instrumental activities of daily living, *see* activities of daily living
- Instrumental Activities of Daily Living Scale, 29
- insulin resistance, 4
- integrative agnosia, 248
- interleukin, 379
- International Classification of Diseases and Related Health Problems*, 10th revision (ICD-10), 10, 341, 363–5, 389, 399
- interpeduncular arteries, 322
- Inventory of Behavioral Dysexecutive Syndrome, 267
- Iowa Gambling Test, 414
- jargon aphasia, 35
- Judgment of Line Orientation, 416
- kinesthetic hallucinations, 174
- Kocher–Cushing signs, 320
- Koro syndrome, 182
- lacunar infarct, dysexecutive syndrome, 264
- Landmark Task, 145
- language  
 dissociating numbers and language, 108–9  
 linguistic perception tests, 122–3  
 tests, 415–16
- language disorders, 22–3  
 written language impairment, 23  
*see also specific disorders*
- language therapy *see* speech and language therapy
- large vessel disease, 11
- lateralization, *see* functional asymmetry
- lateralized disorders of bodily consciousness, 172
- Lausanne Emotion in Acute Stroke Study, 260
- left hemisphere syndromes, 33–6  
 cognitive syndromes, 33–6  
 neuropsychiatric syndromes, 36
- letter cancellation test, 144
- leukoaraiosis, 2–3, 11
- limb kinetic apraxia, 57  
*see also* apraxia
- limb praxis, 55–6
- limb shaking phenomenon, *see* episodic paroxysmal dyskinesia
- line bisection, 145
- lipid-lowering therapy, 425
- locked-in syndrome, 317–18, 327, 335  
 associated lesions, 337  
 management, 337
- long-term memory, 297
- loss of psychic autoactivation, 263
- lycanthropy, 181
- macrosomatognosia, 172
- magnetic misreaching, 219
- major hemispheric syndromes, 399
- malignant hypertensive encephalopathy, 213
- mammillothalamic tract, 282
- mania, 38, 354  
 assessment, 356  
 associated lesions, 356  
 management, 356  
 post-stroke mania, 355  
 associated lesions, 355–6  
 primary versus secondary mania, 354  
 treatment, 356  
*see also* bipolar disorder
- matching tests, 252
- maze tests, 206
- MCA, *see* middle cerebral artery
- medial frontal lobe lesions, 33
- medulla oblongata lesions, 318
- memantine, 403  
 Alzheimer disease treatment, 427  
 vascular dementia treatment, 427–8
- Memorial Delirium Assessment Scale, 343
- memory, 277  
 assessment, 287, 415  
 explicit, 277  
 implicit, 277  
 semantic, 277  
*see also* episodic memory; memory disturbances; working memory
- memory disturbances, 24–5, 277–8, 284  
 CADASIL, 394  
 episodic memory, 24–5  
 assessment, 205–6, 287  
 dysexecutive syndrome, 259–60  
 strokes affecting executive systems, 286–7  
 strokes affecting memory-processing systems, 284–6  
 strokes causing amnesia, 278–84  
 management, 288–9  
 PCA infarcts and, 36  
 prognosis and recovery, 288  
 short-term memory, 24  
 spatial memory assessment, 205  
 working memory, 24  
 central executive deficits, 303–5  
 dysexecutive syndrome, 259  
 phonological deficits, 300–1

- stroke deficits, 303  
 visual working memory, 301–3
- mesiofrontal stroke, 26
- Messerli Test, 258
- metabolic syndrome, 4
- metacognitive process assessment, 267–8
- microsomatognosia, 172
- midbrain lesions  
 alteration of level of consciousness, 321–3, 327–9  
 dystonia and, 48
- middle cerebral artery (MCA), lesions  
 agraphia association, 92–3  
 alexia association, 93  
 alteration of level of consciousness, 330  
 delirium and, 345  
 dysexecutive syndrome, 262–3  
 hemineglect association, 137–8  
 left, 35  
   agraphia association, 93  
   alexia association, 92–3  
   neglect dyslexia association, 91  
 right, 37, 38
- middle fusiform gyrus, face recognition role, 233
- migraine, 71
- Mini Mental State Examination (MMSE), 28, 401, 413
- minimally conscious state, 317, 318
- mirror asomatognosia, 185
- misidentification delusion, 352–3
- misoplegia, 178, 179
- mixed transcortical aphasia, 68
- molecular biomarkers, 16
- Montgomery–Åsberg Depression Rating Scale, 364
- Montreal Cognitive Assessment (MoCA), 28, 413–14
- Montréal–Toulouse Battery, 65, 71, 241
- mood disorder with a general medical condition, 354
- motivational interviewing, 369
- motor awareness theory of anosognosia, 166
- motor neglect, 130–1, 173  
 assessment, 145
- movement disorders, *see* abnormal involuntary movement
- multi-infarct dementia, 1, 26, 392  
 history of terminology, 9
- multiple sclerosis, 71
- musical perception, 119–20
- mutism, 23  
 aphasic, 23  
*see also* akinetic mutism
- myoclonic dystonia, 47
- myoclonus, negative, *see* asterixis
- n-back test, 303, 307
- National Institute of Neurological Disorders and Stroke (NINDS), *see* NINDS-AIREN; NINDS-CSN
- negative autoscapy, 182
- negative myoclonus, *see* asterixis
- neglect  
 auditory, 120  
 dissociations between subtypes, 133–4  
 dyslexia, 87  
   associated lesions, 91  
 functional neurophysiological anomalies, 143  
 motor, 130–1, 145, 173  
 object-centered, 302  
 personal, 146, 173  
 representational, 133, 142, 146  
*see also* hemineglect
- NEO five-factor inventory, 359
- neuroimaging, *see* imaging studies  
 vascular cognitive impairment diagnosis, 16
- Neuropsychiatric Inventory, 267, 417
- neuroticism, 359
- National Institutes of Health stroke scale, 28
- NINDS-AIREN criteria for vascular dementia, 388–9, 410
- NINDS-CSN  
 cognitive test list, 413  
 harmonization conference, 411  
 initial validation, 418–20  
 neuropsychological protocols, 413, 414, 416
- non-conscious hemiasomatognosia, 172
- number processing, 101–2, 104  
 anatomical implementation, 102–3  
 calculation, 102  
 deficits, 106–8  
 hemispheric specialization, 103–4
- input/output, 102  
 quantity-processing deficits, 105–6  
 transcoding, 102  
 triple code, 101  
*see also* acalculia
- Nun Study, 2, 3
- object agnosia, 248–50
- Object Location Test, 206
- object recognition  
 non-visual, 251  
 prosopagnosia relationship, 234–6  
*see also* face recognition; visual agnosia
- object use, 56  
 impairment assessment, 59
- object-centered neglect, 302
- obsessive-compulsive disorder, 38
- obtundation, 315
- occipitotemporal cortex, 103  
 role in prosopagnosia, 232–3
- oculoleptomeningeal amyloidosis, 397
- optic aphasia, 35, 247, 249
- optic ataxia, 219–20, 222  
 assessment, 226
- Organic Brain Syndrome Scale, 343
- orthography, impaired access to, 87  
 from vision, 87
- orthography-to-phonology  
 correspondence, impaired access to, 87–8  
 combined deficits, 88
- Ota's task, 144
- oto-acoustic emissions, 122
- out-of-body experience, 183
- PACE trial, 383
- palipsychism, 36
- pantomime, 55
- paramedian artery, 263–4, 281–2  
 infarct effects on consciousness, 321–3  
 territory, 321–2
- Patient Competency Rating Scale, 164
- pattern span test, 307
- PCA, *see* posterior cerebral artery
- peduncular hallucinosis, 121
- perceptual extinction, 129–30  
 anatomical correlates, 141–2



## Index

- perseverations, 256, 257, 258  
 persistent vegetative state, 317, 335  
   associated lesions, 337  
 personal neglect, 132, 142, 173  
 anosognosia relationship, 164–5  
   assessment, 146  
 personality, 356  
 personality change, 356–7  
   apathy, 358  
   assessment, 5, 359  
   brain damage association, 357  
   emotional incontinence, 358  
   management, 359, 360  
   post-stroke depression risk  
     relationship, 359  
   receptive aprosodia, 358  
   stroke association, 357–8  
   violent and aggressive behavior, 358–9  
 phantom limbs, 173–6  
   supernumerary, 174–6  
 phonemic disarticulation, 35  
 phonemic identification and discrimination, 123  
 phonological dyslexia, 88  
 phonological loop, 298  
 phonological representations, impaired  
   access to, 88–9  
 polyopic heautoscopy, 182  
 pontine lesions, alteration of level of  
   consciousness, 327–9  
 post-stroke psychosis, *see* delusion;  
   psychosis  
 posterior cerebral artery (PCA), lesions  
   alexia association, 91–2  
   cortical blindness and, 212–13  
   delirium and, 344  
   left, 35–6  
   memory impairment, 278–80  
   neglect dyslexia association, 91  
   right, 37  
   visual agnosia and, 250–1  
 posterior parietal cortex  
   Bálint syndrome lesions, 223–4  
 posterior reversible encephalopathy, 26–8  
 post-stroke dementia, 11, 387, 423  
   treatment  
     cognitive decline prevention, 424–6  
     dementia progression prevention, 426–7  
     recurrent stroke prevention, 423–4  
     symptoms, 428  
     vascular risk factors, 424–7  
     *see also* dementia  
 post-stroke depression  
   determinants, 366  
 post-stroke depression, 36  
   diagnosis, 363–5  
   frequency, 366  
   future directions, 370  
   impact, 367  
   management, 367–70  
     electroconvulsive therapy, 370  
     pharmacotherapy, 367–9  
     psychological therapies, 369–70  
   natural history, 366  
   routine screening, 365–6  
   *see also* depression  
 prefrontal cortex, working memory  
   and, 300, 303–5  
 premotor neglect, 130  
 presenile dementia with cerebral  
   hemorrhage, 397  
 Present State Examination, 364  
 pre-stroke dementia, 29  
 prism adaptation, 148, 185  
 problem solving deficits, 258  
 problem-solving therapy, 369  
 PROFESS trial, 424–5  
 PROGRESS trial, 402  
 prosopagnosia, 231  
   frequency, 232  
   lesion localization, 232–3  
   laterality, 232  
   nature of, 236–40  
   neuropsychological assessment, 240–1  
   specificity, 233–6  
   subtypes, 236  
 pseudobulbar syndrome, 358  
 pseudo-extinction, 146  
 pseudo-hemianopia, 128  
 pseudo-sixth phenomenon, 321  
 psychogenic blindness, 213  
 psychosis, 38  
   diagnostic criteria, 351  
 pupillary reaction to light, 213  
 pure tone audiometry, 121  
 pure word deafness, 114, 117–18, 120  
   subtypes, 118  
 quality of life  
   fatigue impact, 381  
   health-related quality of life, 367  
 reading, 86  
   assessment, 93–4  
   *see also* alexia; dyslexia  
 receptive aprosodia, 358  
 Recognition Memory Test, 241  
 reduplicative paramnesia, 25, 37  
 representational neglect, 133  
   anatomical correlates, 142  
   assessment, 146  
 response initiation and suppression  
   deficits, 257–8  
   vascular dementia, 398  
 retrosplenial cortex lesions, memory  
   impairment, 284  
 reverse digit span test, 307  
 Riddoch's phenomenon, 209–10  
 right hemisphere syndromes, 34, 36–8  
   cognitive syndromes, 36–7  
   neuropsychiatric syndromes, 38  
 rule use deficits, 258  
 screening tests, 28  
 semantic memory, 277  
 semantic representations, impaired  
   access to, 88  
   combined deficits, 88  
 senile dementia, 9  
 sequences of auditory signals, 122  
 serial 7's test, 307  
 short-term memory, 297  
   impairment, 24  
   spatial memory assessment, 205  
 short-term recall test, 308  
 silent infarct, 2  
 simultanagnosia, 214, 220–1  
   assessment, 227  
 Six Elements test, 258  
 sleep disturbance, 378  
 small vessel disease, 11  
 social behavior impairment, 256  
 somatoparaphrenia, 176–80, 353  
   anatomical findings, 179  
   varieties of, 177  
 spasm of fixation, 220  
 spastic dysarthria, 79

- spatial acalculia, 104
- spatial attention disorders  
 Bálint syndrome, 220–1
- spatial neglect, *see* hemineglect; neglect
- speech  
 apraxia of, 35  
 disturbances, 33  
 production, 76–7, 81  
 speech perception tests, 122–3  
*see also specific conditions*
- speech and language therapy  
 aphasia, 72–3  
 auditory perception disorders, 123–4  
 dysarthria, 82
- spelling, 86, 89
- splitting of the body image, 181
- star cancellation test, 144
- statins, 425
- stato-kinetic dissociation, 210, 214
- stereotypic behavior, 256
- stimulus-centered neglect, 87
- stimulus–rewards association  
 assessment, 267
- stress hormones, 341–2
- striato-pallidal lesions, dystonia and, 46–7
- stroke  
 recurrent stroke prevention, 423–4  
 antithrombotic drugs, 424  
 diabetes mellitus, 423  
 hypercholesterolemia, 423  
 hyperhomocysteinemia, 423–4  
 hypertension, 423  
*see also subarachnoid hemorrhage; specific lesions*
- Stroop Test, 267, 307, 414, 418
- Structured Clinical Interview for DSM-IV, 364
- stupor, 317
- subarachnoid hemorrhage  
 delirium and, 345–6  
 dysexecutive syndrome and, 264–5  
 fatigue after, 382  
 correlated factors, 382  
 epidemiology, 382
- subcortical aphasia, 69
- subcortical ischemic vascular dementia, 2, 11–12, 393, 412
- subcortical syndrome, 11
- superior longitudinal fascicle  
 hemineglect association, 140
- supernumerary phantom limbs, 174–6
- supplementary motor area lesions, 33
- surface dysgraphia, 87, 88
- surface dyslexia, 87, 88
- sustained attention deficit, 258, 259
- Syst-Eur trial, 402
- task coordination deficit, 259
- test for upper limb apraxia, 58
- thalamic lesions  
 akinetic mutism, 26  
 alteration of level of consciousness  
 anteromedian, central and posterolateral thalamic infarcts, 325–7  
 combined polar and paramedian infarct, 323  
 paramedian infarcts, 321–3  
 rostral mesencephalic–hypothalamic–thalamic infarct, 324–5  
 delirium and, 344  
 dysexecutive syndrome, 263–4  
 dystonia and, 47–8  
 left, 36  
 memory impairment, 280–2  
 tremor and, 49–50  
 vascular dementia and, 391–2
- theory of mind, 257
- token tests, 415
- tool use, 56  
 impairment assessment, 59
- top of the basilar syndrome, 211, 320–1
- topographical disorientation, 202–3  
 assessment, 205–6  
 clinical examination, 205  
 neuropsychological assessment, 205–6  
 classification, 203  
 management, 206  
 prognosis, 206  
 prosopagnosia association, 234  
 stroke relationships, 203–5
- Tower of London, 258
- Trail Making Test, 414  
 Part B, 267, 308
- transcortical aphasia, 35  
 motor aphasia, 67  
 sensory aphasia, 68
- transcranial direct current stimulation, 308
- transcranial magnetic stimulation, aphasia treatment, 73
- transitivity, 178
- tremor, 46, 48–9  
 cerebellar tremor induced by a pure thalamic lesion, 49–50
- trimodal extinction, 129
- tuberothalamic artery, 263
- two-factor theory of anosognosia, 166
- unconsciousness, *see* alteration of level of consciousness
- unilateral upper motor neuron dysarthria, 79–80
- vascular cognitive disorder, 10, 410
- vascular cognitive impairment, 1, 5, 9, 410  
 assessment  
 behavioral disturbances, 416–17  
 daily function, 417–18  
 etiologic heterogeneity and, 411–12  
 executive domain tests, 414  
 language and visuospatial tests, 415–16  
 memory domain tests, 415  
 neuropsychological assessment, 412–17  
 screening examinations, 413–14  
 classification, 5, 410  
 clinical presentation, 4, 12–18  
 case vignettes, 13–16  
 diagnosis, 5  
 diagnostic criteria, 5, 10, 410–11  
 molecular biomarkers, 16  
 neuroimaging role, 16  
 epidemiology, 11  
 evaluation, 13  
 history of terminology, 9–10  
 imaging studies, 4–5  
 mild impairment, 5  
 neuropathological subtypes, 11–13, 411–12  
 large vessel disease, 11  
 non-infarct pathology, 12  
 small vessel disease, 11  
 subcortical ischemic vascular dementia, 11–12  
 treatment, 17–18  
*see also cognitive impairment*
- vascular cognitive impairment, no dementia (VCI-ND), 9, 410
- vascular dementia, 1, 9, 387  
 assessment, 399–401  
 cerebrovascular disease, 399–401

Index

- vascular dementia (*cont.*)
  - markers, 401
  - neuropsychological assessment, 401
  - Binswanger disease, 392–3
  - CADASIL, 393–6, 400–1
  - categories of, 390–1
  - causes, 390
    - rare causes, 396–7
  - diagnostic criteria, 387–90
  - hemodynamic mechanisms, 398
  - multi-infarct dementia, 392
  - multiple brain hemorrhages and amyloid angiopathies, 397–8
  - neuropsychology of, 398–9
  - prevalence, 1
  - single stroke dementia, 391–2
    - cortical stroke, 391
    - subcortical stroke, 391–2
  - subcortical ischemic, 2
  - treatment, 402–3, 427–8
    - blood pressure reduction, 402
    - pharmacotherapy, 402–3
  - see also* dementia
- vascular depression, 366–7, 416
- vascular generalized chorea, 52
- vascular parkinsonism, 50–2
  - diagnostic criteria, 50
  - pathophysiology, 52
- ventral attention network, 302
- ventral intermediate nucleus lesions, 47–8
  - tremor and, 49–50
- vestibular caloric stimulation, *see* caloric vestibular stimulation
- violent behavior, 358–9
- visual acuity measurement, 214
- visual agnosia, 247
  - assessment, 251–3
    - color perception and categorization, 252
    - copying and drawing from memory, 251–2
    - error analysis, 251
    - matching tests, 252
  - non-visual object recognition, 251
  - causes, 250
    - strokes, 250–1
  - management, 253
  - prognosis, 253
  - standardized testing, 252–3
- visual attention deficit assessment, 226–7
- visual cueing paradigm, 146
- visual disturbances, 24, 39
  - see also* cortical blindness; *specific conditions*
- visual evoked potentials and cortical blindness, 215
- visual feedback, 185
- visual field assessment, 214
- visual grasp reflex, 220
- Visual Object and Space Perception Battery, 240, 252
- visuoconstructive disorders, 193–6
  - assessment, 199
  - causative strokes, 199
  - in stroke, 198–9
    - frequency, 198
  - prognosis, 199–200
- visuospatial sketchpad, 298
  - dorsal, 302–3
  - ventral, 301–2
- Wallenberg syndrome, 81, 318
- Wechsler Adult Intelligence Scale, 240
- Wechsler Memory Scale, 240, 241
- Wernicke aphasia, 35, 67–8
- Wernicke area, working memory and, 300–1
- Western Aphasia Battery, 65, 416
- Wisconsin Card Sorting Test, 258, 267, 414, 418
- word discrimination test, 121
- word span test, 307
- working memory, 297–8
  - capacity, 305–7
  - cognitive models, 298–9
  - impairment, 24
    - central executive deficits, 303–5
    - cognitive rehabilitation, 308
    - dysexecutive syndrome, 259
    - phonological deficits, 300–1
    - stroke deficits, 303
  - visual working memory, 301–3
  - neuroscientific studies, 299–300
  - tests of, 306, 307–8
- writing, 86
  - assessment, 93–4
  - see also* aphasia
- xenomelia, 178