

Index

- Amazon, 21, 24, 378–379
 overbank, crevasses processes, 25
 physical and biogeochemical processes, 92
 plume dispersals, 55
 sediment accumulation rates, 91
 subaqueous delta clinoform system, 91–96
 tidal-regime changes, sediment dynamics model, 92–94
- Arctic Ocean, riverine organic and inorganic carbon fluxes, processing, fate
 acids to aldehydes ratios, 534–535
 aragonite undersaturation, 542–543
 Arctic riverine DIC during transport, sources and fluxes, 538–543
 Arctic riverine organic matter exported to coastal systems
 sources, composition, fluxes, 533–538
 Arctic riverine tPOC, tDOC fluxes, 533–534
 Arctic rivers, hydrology and fluxes vs. global rivers, 530–532
 CaCO₃ concentration, 542
 CDOM absorption, DOC concentration, total lignin concentrations, 535–537
 dissolved inorganic C system, six largest Arctic rivers, seasonal variation, 539
 flow weighted DOC concentrations for the five largest Arctic rivers, 543–544
 freshwater discharge patterns, six biggest Arctic rivers, 530–533
 HCO₃ concentration, 539–542
 optical measurements vs. catchment area-normalized daily discharge, six largest Arctic rivers, 535–537
 overview, 530, 546
 pCO₂ values, 539–541
 riverine DOM fate, degradation, transport, constraining flux models, 544–546
 riverine OM utilizing abundance ¹⁴C and ¹³C, sources and cycling, 543–544
 six biggest Arctic rivers, catchments, 530, 533, 539
 tDOC degradation, 545–546
 tDOM, tPOM biomarker compositions, 534–535
 tDOM optical properties, 535–538
- black carbon (BC)
 atmospheric BC, 204–205
 BC, 214–218
 defined, 200
 degradation, 222–224
 dissolved, 206–207
 fate of, 220–224
 global cycling of, 224
 global pyrogenic PAH discharge vs., 218–220
 isolation, dissolved phase, 210–213
 isolation, quantification, source discrimination, 207–214
 in large river systems, 214–220
 overview, 200–202
 quantification, chemical and thermal methods, 209–210
 quantification, visual methods, 209
 soil, 205–206
 source discrimination, 213–214
 sources, 200–208
 storage BC in ocean, 220–222
- Changjiang (Yangtze) and Huanghe (Yellow) Rivers, 21
 accumulation/erosion, Changjiang delta sensitive areas, 131–133
 anthropomorphic changes since 1050's, 125–129
 Changjian river basin, sediment load, 125–126
 Chinese Dynasties, capital migration, 122–125
 climate change, human activities roles, 127–132
 Huanghe delta topology, sea level rise, 133
 Huanghe river basin, sediment flux since 1950's, 127–129
 human impacts, land-use change, 122–125
 hydrologic settings, 121–122
 Indian monsoon rainfall, annual sediment flux sine 1970, 128–130
 mega-deltas future, 131–134
 monthly SST anomalies, South Oscillation Index (SOI), 130–132
 regional settings, 118–122
 river basins, surface lithology, 122
 topography, information, 118
 water discharge, warm/cold phase linkage, 127–128

- Changjiang (Yangtze River) Watersheds across East China Sea, carbon biogeochemistry
- biochemical budgets, feed-backs open to ocean and atmosphere, 257–259
 - budget calculation of DOC, case studies, 257–258
 - carbon cycle, influence factors, major pathways, 252–257
 - carbon fate inferred from pigments, 256–257
 - CH₄ and CO₂ fluxes, PN Section, 262
 - CH₄ (nmol/L), salinity, temperature (°C) along PN Section, vertical distributions, 244–245
 - CH₄/CO₂ ratio, PN Section, 262
 - Chl-*a* to POC ratio, 264
 - chlorophyll *a*, pheophorbide profile, 260
 - CO₂ and HCO₃⁻, long term mean monthly discharge, 238
 - Δ*p*CO₂ data, 241
 - DIC and CH₄, 252–253
 - DIC in the ECS distribution, 241–243
 - dissolved CH₄, preliminary budgets, 259
 - dissolved inorganic carbon, budget calculations, 258
 - dissolved inorganic carbon (DIC), CH₄, 238–245
 - dissolved organic carbon (DOC), particulate organic carbon (POC) flux, 238–252
 - δ¹⁵N variation trend, 260–262
 - DOC, spatial distribution, 248
 - DOC and POC, 253–256
 - DOC and POC, annual flux data, 245–246
 - DOC and POC distribution in East China Sea, PN Section, 248
 - DOC distributions, Ra-isotopes vs. observations, 254
 - DOC to POC ratio, surface waters, 264
 - DOC vs. salinity, Changjiang Estuary, 246–248
 - ECS, carbon biogeochemistry, 262–266
 - future work, 266–267
 - historical records evidence, 259–262
 - land-source influx, open boundary process characters, 238–252
 - monitoring data, history, 238
 - monthly CH₄ concentrations at Xuliujing, water discharges at Datong Hydrographic Station, 243–244
 - nitrogen flux, bloom events, 262
 - organic carbon degradation rates (K), comparisons, 255
 - organic carbon degradation rates (K) comparisons, 244–245
 - organic carbon (OC), main stream and estuary of Changjiang and adjacent coastal waters, 247–248
 - overview, 237–238, 266
 - p*CO₂, long term annual mean, 238–239
 - phytoplankton, estimated (%) contributions to POC, 253–254
 - pigments, 250–252, 256–257
 - POC derived from phytoplankton, estimated contributions (%), 253–254
 - preserved diagnostic pigments, 260–262
 - surface CH₄, Changjiang Estuary and East China Sea, 243–244
 - surface CH₄ and salinity, Changjiang Estuary, adjacent coastal areas, 244–245
 - surface DIC, TALK, pH and *p*CO₂ compilation, 240–241
 - surface *p*CO₂ and salinity, Changjiang estuary, 240–241
 - tides, climate, seasons and, 237–238
 - coastal waters influenced by river plumes, carbon dioxide dynamics
 - Amazon river plume, CO₂, *p*CO₂, 165–166
 - Changjiang plume, 160–161
 - DIC and TALK vs. salinity plots, northern Gulf of Mexico, 162–164
 - global CO₂ degassing fluxes, inner estuaries to atmosphere, 155–158, 168
 - global river-plume CO₂ flux, tentative estimate, 166–169
 - Mississippi-Atchafalya river plume, CO₂, *p*CO₂, 161–162
 - p*CO₂, pH changes during plume mixing, high carbonate river example, 160–161, 165
 - p*CO₂ distribution pattern in river plumes, control patterns, 158–166
 - p*CO₂ vs. salinity, northern Gulf of Mexico, 162
 - river bicarbonate (HCO₃⁻) concentrations, latitudinal distribution, 158–159
 - TALK, DIC, fugacity of CO₂ (*f*CO₂) vs. salinity distributions, 165
 - coastal zone
 - human civilizations and, xiv
 - LOICZ definition, 3
 - net change in total organic compounds (OC), 4
 - particulate organic carbon (POC) storage, transit time, 3
 - Congo River, estuary and plume geochemistry
 - anthropogenic influences, 554
 - carbon fluxes, organic matter geochemistry, 566–571
 - channel morphology of Congo Canyon, associated channel-levee system, 575–576
 - chemical, physical denudation rates, 563–565
 - chlorophyll *a*, POC, and DOC in estuary and plume plotted against salinity, 573–574
 - Congo Canyon, associated channel-levee system, 575–578
 - Congo Canyon and Channel, bathymetry map, 576–578
 - Congo Fan, 575–576
 - Congo River mainstem, organic matter geochemistry, 571
 - Congo watershed, map, 555–556
 - discharge overview, 556
 - dissolved solutes, soil CO₂ chemical neutralization, 558–560
 - dissolved trace elements, organic footprint, 560–561
 - DOC, SUVA₂₅₄ and A₈ box plots, Epulu River, 569–571
 - Epulu River, organic matter geochemistry, 569
 - erosion rates, chemical weathering, fluxes, 563
 - estuarine and plume process, 571–575
 - hydrology, source to sea, 554–556
 - inorganic and sediment geochemistry, 556–566

- inorganic poor waters of system, 556–558
- monthly average DOC concentration, mean monthly water discharge relationship, 567
- monthly discharge data, Kinshasa-Brazzaville, 556
- normalized abundance of elements, 562–563
- organic carbon (OC) export to Atlantic Ocean, 567
- organic carbon (OC) flux via major tributaries, 566–567
- organic matter deposition, 576–578
- organic matter geochemistry, Congo tributaries, 567–571
- organic matter in surface plume, 572–574
- overview, 554
- particle load, inorganic chemistry, 562–563
- particle-rich estuarine bottom waters, 574–575
- silica *in-situ* biogeochemical cycling, 560
- Sr isotopes and Sr/Na ratio relationship, mixing diagram, 558–559
- surface photo-reactor, 574
- temporal variability, inorganic geochemistry, 561
- turbidity distribution, Congo estuary and across shelf axial to Congo Canyon, 571, 574
- Ubgani River, organic matter geochemistry, 567–571
- weathering regimes, 565–566
- European Environmental Agency, estuary definition, 32
- Ganges-Brahmaputra, 21
- Bay of Bengal, fluvial sediment reworking and deposition, organic carbon (OC), 356–359
- Bengal Fan, 358–359, 365, 367
- Brahmaputra floodplain carbon sources, transformation and storage, 364
- bulk organic carbon $\delta^{13}\text{C}$ values, 361–362, 363
- bulk organic carbon (OC) radiocarbon content, Lower Meghna sediments, Bengal Fan core top sediments, 365–366
- carbon export, Ganges-Brahmaputra to Bengal Fan, 364–365
- combination of influences, 96–100
- Ganges floodplain carbon sources, transformation and storage, 363
- global warming and, 368
- Himalayan range carbon sources, transformation and storage, 360–363
- hydrography, Bangladesh, 354
- major depocenters for sediment and organic carbon, 356 map, 353, 359
- organic carbon (OC), Ganges and Brahmaputra rivers and Bengal Fan, 365
- organic carbon (OC) content, Bengal Fan sediments, 365–367
- organic carbon (OC) fate, Bengal Fan, 365–367
- overbank, crevasses processes, 25
- overview, 353–355, 367–368
- sediment dispersal system basemap, 96
- shelf sedimentary system, 96
- suspended, bed sediments vertical depth profile, 360–361, 364
- Swatch of No Ground (SoNG), 98, 100, 358–359
- Godavari River, 379–380
- Huanghe (Yellow) rivers, derived materials flux and fate, climate change, human activities impacts
- Bohai and Yellow seas, Huang river sediments fate, 146–149
- coastline changes, 1976–2006, 144–145, 146, 151
- deltaic progradation since middle Holocene, 143–144, 146
- distal mud depocenter, longshore transport to Yellow Seas, 148–149
- historical course shifts, 142, 146–149
- Holocene sediment and accumulation, 144
- Huanghe river watershed, 138
- mean annual sediment load, annual deltaic land increase relationship, 146, 151
- no-flow dried-up days, yearly distribution 1972–2012, 141
- proximal deltaic depocenter, rapid sediment accumulation, 146–148
- recent decrease of water and sediment discharge, causes, 145–146
- recent-year water discharges, decreasing sediment loads impacts, 146
- river basin landcover distribution, 138–140
- river-derived sediment to coast and sea, conceptual model, 149–151
- riverine POC ^{14}C -content (Fmod), suspended load relation, 148–149
- sediment fluxes, decadal scale, 144–145
- sediment fluxes, millennial scale, 142–144
- water discharge, sediment load, 140–141, 145–146
- western Bohai to South Yellow Sea, suspended matter, plume, 146–148
- International Geosphere Biosphere Programme (IGBP), xiii
- Land Ocean Interaction in the Coastal Zone (LOICZ), xiii, 3
- large delta-front estuaries (LDE). *See also* Atchafalaya River and Mississippi River delta-front estuaries
- biogeochemical processes, xiv
- total organic carbon (OC), xiii
- large deltaic systems, shelf and slope sedimentation
- Amazon
- physical and biogeochemical processes, 92
- sediment accumulation rates, 91
- subaqueous delta cliniform system, 91–96
- tidal-regime changes, sediment dynamics model, 92–94
- continental-margin sedimentation factors, 87
- fluvial, marine controls, 104–106
- Ganges-Brahmaputra
- combination of influences, 96–100

- large deltaic systems, shelf and slope sedimentation (*cont.*)
 sediment dispersal system basemap, 96
 shelf sedimentary system, 96
 Swatch of No Ground (SoNG), 98, 100
 geological framework, human influence, 106–107
 human populations and, 86
 large river-sediment dispersal systems, data and
 classifications, 88–89
 Mississippi-Atchafalya River System (MARS)
 drainage basin, 100–102
 hurricane deposit thicknesses, Katrina and Rita,
 103–104
 proximal accumulation-dominated system, 100–104
 Multidisciplinary Amazon SEDiment Study (AmasSeds),
 91
 overview, 87–91
 larger river plumes, freshwater and sediment dispersal
 Amazon, 55
 buoyancy effect, 69–70
 carbon, 78
 estuary, 59–60
 evaporation, 78–79
 far-field plume, 63–64
 flocculation, schematic plot, 70
 flocculation, hindered settling, 70–75, 79
 flocculation, hindered settling, 70–75, 79
 mid-field plume, bulge, 62–63
 moderate to large systems, recent studies, 55–57
 near-, mid-, and far-field dispersal, 76–78
 near-field plume, 61
 river plume anatomy, 57–65
 river plume classification, 65–68
 riverine input, carry capacity, 76
 salinity space, 57–59
 sediment dynamics, 68–75
 small systems, recent studies, 57
 surface-trapped plume, features, 57–58
 surface-trapped river plume, dynamical regions, mixing
 processes, 55
 Teignmouth, 55
 33 g kg⁻¹ surface salinity contour, monthly locations,
 58–59
 transparent exopolymer particles (TEP), 72–73, 74,
 75
 25 largest rivers, 55
 very far field plume, 64–65
 lower Mississippi River. *See* Mississippi-Atchafalya River
 System, lower Mississippi River
 Mekong River, nutrient chemistry and matter cycling
 influence in Vietnam coast zone
 biochemistry, productivity, 299–312
 coastal upwelling systems, 296
 coastal waters hydrography, 297–299
 compared to other tropical estuaries, 313–314
 $\delta^{13}\text{C}$ values, particulate organic matter (POM) from filter
 samples, 308–309
 $\delta^{15}\text{N}$, amino acids from increasing plankton size fractions,
 310
 $\delta^{15}\text{N}$ patterns, particulate organic matter (POM) from filter
 samples, 310
 drainage area map, 299
 food web, zooplankton nutrition, 308–312
 long-term seasonal mean of Mekong runoff, 299
 Mekong plume, 299
 Mekong River Commission (MRC), 314
 Mekong River estuary, catchment future, 314–315
 nitrogen fixation, 303–305, 312, 315, 377, 460
 nitrogen fixing, 303–308
 overview, 296–297, 315
 pelagic primary productivity rates, 304–306
 phytoplankton species composition, 299–303
 primary production, nitrogen fixation, 303–308
 research, monitoring, 296–297
 salinity, nutrient distribution, 304–305
 salinity zonation mesohaline, nutrient ratios N:P, Si:N, N2
 fixation rates, 306–308
 South China Sea, map and topography, 297–298, 299,
 313
 southeast Asian monsoon region, climatology, 297
 Mississippi River Delta (MRD), climate change effects on
 ecology
 anthropomorphic impacts, Louisiana coastal zone, 427
 anthropomorphic impacts, Mississippi River Basin (MRB),
 423–427
 climate change effects, 421
 biochemical recycling, Mississippi Delta, 428–430
 delta vegetation, 430–439
 MRB, 428
 common plant species, historic changes, 432
 ecology of Mississippi River Delta, 421–423
 emergent vegetation, projected change under future climate
 scenarios, 435–437
 future scenarios for 2012 Master Plan models, 435–437
 habitats, 422
 human impacts, climate change synergistic interaction,
 435–439
 Hurricanes and, 434
 marsh types, 430
 mature baldcypress diameter increase, 434
 MRB, agricultural activity and, 424–425
 MRB, stream flow modification and, 426–427
 salinity, flooding stress, 434–435
 swamp forest trees production as function of hydrology,
 433
 2001 MRB land cover classifications, bottom water
 hypoxia in NGOC, 423

- Mississippi River (MR) and Atchafalaya River (AR)
 delta-front estuaries (LDE), sedimentary carbon transport dynamics. *See also* Mississippi River Delta, climate change effects on ecology
 algal organic carbon across shelf/slope, transport and sources, 488
 algal pigment biomarkers fucoxanthin (Fuco), diadinoxanthin (Diadino), chlorophyll-*a* (Chl *a*) in surface sediments, 481–482
 bulk organic carbon burial rates, 486
 controls on phytoplankton sources, carbon diagenesis across depositional gradient, 482
 diagenesis, OC sources and transport rates effects, 485–487
 hurricanes effect on sediment, OC transport, 480–482
 lambda 8 (mg lignin/100 mg OC), (Ad/Al)_v (vanillic acid/vanillin ratio) in surface sediments, 489
 major hurricanes (category 3 at landfall), Louisiana or Mississippi coastlines, 1851–2003, 480–481
 1996 hypoxia region relative to plume regions, maps, 484–485
 organic matter decay sources, nutrients and transport processes linkages, 482–492
 overview, 473, 492–493
 Pb-210 based accumulation rates, surface sediments compositional characteristics, 490
 sediment and carbon dynamics across shelf/slope, linkages, 476–479
 sediment input, deposition, 473–475
 sediment remobilization mechanisms, timing, 475–476
 sedimentary processes, upper Mississippi Canyon regions, 479–480, 481–482
 stations sampled, Mississippi Canyon regions, 477–479
 terrestrial OC sources across shelf/slope, transport and sources, 488–492
 zooplankton grazing, 485
- Mississippi-Atchafalaya River System (MARS), lower Mississippi River (LMR). *See also* Mississippi River Delta
 agriculture intensification and, 412
 alga organic matter, 403
 annual DOC, POC fluxes, 397, 401–402
 biogeochemistry future, global warming, land-use change, extreme climates, 412–413
 dissolved silicon loading, 409
 flow, concentrations of SPM, DOC, nitrate, SRP and dissolved silica, 399
 Gulf coastal hypoxia and, 410
 levees, 398–399
 Mississippi and Atchafalaya ramped pyrolysis data, 404–405
 nutrient export, Mississippi River to Gulf of Mexico, 408–412
 organic carbon (OC) sources, transport, transformation, 401–407
 overview, 413–414
 ramped pyrolysis thermograph for LMR POC, 403–405
 seasonal variations, 409–410
in situ recycling, nutrient dynamics, 410
 trace metal geochemistry, complexation with organic matter, 407–408
 water, sediment, and water quality monitoring stations, 397
 water, sediment dynamics, 398–401
- Mississippi-Atchafalaya River System (MARS), nutrient and carbon dynamics
 bacterial-mediated transformations, 460
 biogeochemical processes along river-ocean mixing gradient, diagram, 458–459
 carbon system properties, air-sea flux of CO₂, 462–463
 climate-, human-related drivers of long term change, 463
 DIN and DON, time-series, 452
 DIN flux, SeaWiFS, MODIS Aqua chlorophyll *a* products, 458
 discharge vs. dissolved inorganic nitrogen flux, 452
 dissolved inorganic compounds to orthophosphate or total phosphorus, molar ratios, 452–455, 459
 human impact, 448–449
 mean discharge, 2003–2011, 449–450
 nutrient enhancement of primary and secondary productivity, outflow region, 458–461
 nutrients and carbon changes, retrospective analysis, 451–458
 organic and inorganic C, N, P delivered to Gulf of Mexico, average concentrations and annual fluxes, 454–455
 outflow regions, MODIS greyscale image, 449
 overview, 448–451
 oxygen and respiration patterns, subsurface waters, 461–462
 POC flux, 455–457
 Total P and orthophosphate, time-series, 452–453
- Nile delta in Anthropocene, coast change drivers and impacts on land-material transfer
 agricultural farming, 594
 airborne nitrogen deposition, 600–601
 anthropogenic drivers, 586, 588, 590–596
 aquaculture, 594–595
 artificial fertilizers, 594–595
 Aswan High Dam (AHD) impact, 587, 589–590, 592–593, 600
 brackish water lakes, 586
 classic Nile bloom, pre-Dam conditions, 596–598
 classic vs. modern phytoplankton Nile bloom, 596–600
 coastal pollution, water quality degradation, 590
 dams and barrages locations, construction dates, 588–589
 delta population density, urbanization, 586, 595–596
 future directions, research, 600–601
 geography, 585
 historical change, 587

- Nile delta in Anthropocene, coast change drivers and impacts
 on land-material transfer (*cont.*)
 Long Aswan Dam (LAD) impact, 590–592
 major distributaries, 584–585
 marine biological productivity, Nile phytoplankton, 593
 modern Nile bloom, post-Dam conditions, 598–600
 Nile delta satellite image, 584–587
 northwestern Egyptian Mediterranean coast development,
 600
 origin of delta, 584–587
 overview, 584–587
 pelagic to demersal (P/D) ratio, 599–600
 sea level rise (SLR), land subsidence, 587–588
 sediments, fertile soil, 584–587
 threats, major environment problems, 587–590
 tidal range, 585–586
 water discharge and suspended sediment load, long-term
 variations, 590–593, 596
 waterways, irrigation construction, 593–594
- Pearl River (Zhujiang) estuary, phytoplankton blooms and
 nutritional limitation
 chl *a*, monthly average, 282–283
 chl *a*, multivariate ENSO index (MEI) correlation, 288
 climate change, eutrophication, 286–289
 conceptual model, 285
 ecosystem buffering process, 276
 estuarine, coastal waters processes, 277–278
 geographic, oceanographic settings, 276–278
 main estuary map, 274
 nutrient, nutrient limitation variability dynamics, 285–289
 overview, 274–276, 289
 phytoplankton biomass, spatial and seasonal variability,
 281–285
 rainfall, 277–278
 river outflow controlled variability, 281–283
 salinity changes, NO₃, SiO₄, PO₄, 286–289
 surface salinity, temperature change, 280–282
 tidal cycle regulated variability, 283–284
 wind direction, speed, 277–278
 wind event driven variability, 284–285
- Pearl River (Zhujiang) plume, physical dynamics,
 biogeochemistry, 341
 average surface velocity vectors, seasonally averaged, 340
 biogeochemistry, 341–344
 2001 vs. 2008, 343–344
 August 2001 case, 343
 May 2001 case, 341–344
 mixing model, different water masses, 344–347
 MODIS color index (CI), May 2001, 342–343
 net surface velocity vectors, simulated result, 341
 North China Sea Shelf
 downwelling circulation, 339
 overview, 336–338
 shelf circulation, 338–339
 surface velocity vectors, 338–339, 340
 upwelling circulation in summer (wet season), 338–339
 nutrients (DIN, DIP, and Si(OH)₄) and DIC dynamics, 347
 overview, 321–322, 349–350
- Pearl River
 basics, 322–325
 characteristics, hydrochemistry, 322–327
 drainage basin maps, 322, 325–327, 331, 332, 336
 precipitation, river discharge, 325
 river discharge, 325, 341–342
- Pearl River estuary
 biogeochemistry controls, conceptual paradigm, 332
 circulation, 327–332
 gravitational circulation, 327–328
 hydrology, biochemistry, 332–336
 intra-tidal circulation, 329–332
 mixing behavior, conceptual summary, 336
 overview, 325–327
 salinity distributions, 332–335, 336
 sub-tidal circulation, 328–329
 surface elevations variation, barotropic vectors, 329,
 339–340
 velocity vectors, salinity, 331
 water density as function of water depth, 327–328
 wind speed vectors, surface chlorophyll *a* (Chl *a*)
 concentrations, 328
 physical dynamics, biogeochemistry coupling, 344–349
 plume effect on shelf circulation, 340–341
 plume over shelf, 339–340
 surface temperature distribution and salinity, northern
 South China Sea shelf, August 2008, 343
 temperature vs. salinity in northern South China Sea
 (NSCS), 345–347
- plumes. *See* coastal waters influenced by river plumes, carbon
 dioxide dynamics; large river plumes, freshwater and
 sediment dispersal; Pearl River plume
- Po River delta plain, channel bed aggradation, 25
- Rhine delta, anthropogenic changes in sediment and nutrition
 retention
 climate change and, 642
 contemporary sediment deposition, lower Rhine delta,
 636–637
 deforestation, 631
 deposition after embankment, 1350–1850 AD, 634–636
 economic development, population, 631–632
 floodplain layout scenarios, 642–643
 future nutrient retention, 643–644
 future retention changes, 642–644
 future role of Rhine Delta, 644–646
 future suspended sediment trapping, 643
 Holocene trapping amounts, 634
 Holocene variations in sediment trapping, 632–636
 International Commission for the Protection of the Rhine
 (IKSR), 631–632

- land use change, 630–631
- landscaping measures, 646
- nitrogen, phosphorus concentrations in Rhine at Lobith, 638–639, 641
- nitrogen, phosphorus emissions in Germany, 639
- nitrogen, phosphorus emissions in Rhine basin, 639
- nutrient fluxes, decadal trends and variability, 637–639
- nutrient retention, lower Rhine distributaries, 639–642
- overbank fines deposited, 635–636
- overview, 629, 644–646
- post-normalization deposition rates, 1850 AD-present, 636
- present-day suspended sediment trapping, 637
- Rhine basin, 629
- Rhine basin, historic changes, 629–632
- Rhine-Meuse delta, Waal formation, 632–635
- river engineering, 632
- suspended sediment and nutrients retention, 637
- river-coastal systems, biogeochemistry
- Arctic rivers, 10–11
 - canyons, 8
 - climate change and, 10–11
 - CO₂ fluxes, coastal waters, 8
 - coagulation, flocculation, 7
 - coastal water CO₂ flux, riverine loadings links, 8
 - diagenesis, diagenetic studies, 7–8
 - dissolved inorganic nitrogen (DIN), 8
 - dissolved organic carbon (DOC), 5–6
 - future research directions, 10–11
 - interactive domains, 4–5
 - largest 25 rivers, 3
 - lower, middle latitude coast zone, 11
 - Margin zone, 4
 - missing organic carbon (OC) conundrum, 5–6
 - NSF MARGINS Source-to-Sink program, 8
 - organic matter (OM) degradation “priming,” 6
 - riverine inputs of particulate (POC), dissolved organic matter (OM), 5–6
 - River-Ocean Mixing Zone, 4–5
 - terrestrial total OC (TDOC), 5
- riverine organic carbon (OC), watershed process impacts
- climate impacts, 182, 190
 - $\delta^{13}\text{C}$ (top) and $\delta^{14}\text{C}$ (bottom) values of coarse (63–2,000 μm) and fine (0.1–63 μm) POC, 185–186
 - exported riverine organic signature prediction, 183–189
 - fossil (rock-derived) OC content of riverine POC vs. the ¹⁴C-content of the non-fossil component, 188–189
 - in-channel and lowland sources, processes, 181–182
 - overview, 174–175
 - POC concentration as a function of total suspended load, 183
 - POC signatures, 184–185
 - POC sources, 175–182
 - regolith, geomorphic processes, 180–181
 - river POC ¹⁴C-contents (Fmod). ¹⁴C-contents, 186–188
 - riverine exported POC $\delta^{13}\text{C}$ values, 175–178
 - riverine POC ¹⁴C-content (Fmod)
 - as function of %OC in suspended load, 186–189
 - vs. watershed area for global rivers, 187
 - soil POC, $\delta^{13}\text{C}$ (top) and $\delta^{14}\text{C}$ (bottom) values, 178–180
 - terrestrial organic carbon (OC) indicators, 176–177
 - vegetation, 175–178
- Siene River and estuary, fate of nutrients in aquatic continuum
- agricultural soils, aquifers, surface waters, 607–610
 - agriculture, 606
 - annual nitrogen, silica, phosphorus and organic carbon fluxes, model, 620–622
 - discharge and specific nitrogen, silica, phosphorus fluxes, 612–614
 - drainage network, land use, 607
 - drinking water protection areas (DWPAs), 619–620
 - N, P, Si inputs and fluxes, 606, 612–614
 - N content of crop harvest and total N fertilization, long-term variation, 607–608
 - nitrate concentration, seasonal variations, 608–609
 - N:P, Si:P Redfield ratios, 620
 - overview, 606–607, 623
 - phosphorus retention, 609–610
 - population density, 606
 - relationships between percentages of total nitrogen, phosphorus and carbon retention, associated discharges, 619
 - total nitrogen, phosphorus and silica urban inputs, 610–611
 - total nitrogen (TN), silica (TSi), phosphorus (TP) and organic carbon (TOC) fluxes, seasonal variations for two dry years, 618
 - transfer in drainage network, RIVE model, 611–614
 - transfer in drainage network, SENEQUE/RIVERSTRAHLER model, 611–614, 623
 - transfer through estuarine zone, 614–619
 - transfer through estuarine zone, LIFT model, 616–618
 - transfer through estuarine zone, SiAM-1D/RIVE model, 614–619
 - urban sources, wastewater treatment plants (WWTP), 610–611
 - wastewater treatment (WWT), 610
 - watershed characteristics, 606–607
 - watershed management, nutrient deliveries to coastal zone, 619–622
 - yearly average of total nitrogen, silica, phosphorus and organic carbon fluxes, 616
- Sundarbans, 379–380
- Teignmouth, plume dispersals, 55
- tropical river-coastal boundaries, carbon (CO) and nutrient fluxes
- Amazon, 378–379
 - anthropomorphic disturbances, global change and, 386–388
 - Brantas River delta, 377

- tropical river-coastal boundaries, carbon (CO) and nutrient fluxes (*cont.*)
- cross-boundary exchanges, 375–385
 - Everglades, 385
 - Fly Delta, 380–381
 - Fly Delta, carbon budget, 380–381
 - Gazi Bay, 382–385
 - global discharges, 375
 - global map, 374, 375
 - Godavari River, 379–380
 - Hinchinbrook Channel, carbon budget, 377
 - Hooghly River, 380
 - Irrawaddy Delta, 387
 - large river deltas, 378–382
 - litter and POC export, DOC flux, 382
 - Matang estuary, 385
 - Merbok estuary, 385
 - overview, 373–374
 - Paña River, 386
 - Red River delta, 377
 - Sawi Bay, 385
 - Segara Anakan Lagoon, 387
 - small to medium river deltas, 375–378
 - Sundarbans, 379–380, 387
 - Tana River delta, Masinga Dam, 386
 - Tapi River delta, 377
 - tidally-dominated creeks, estuaries, lagoons, 382–385
 - tropical coasts, types and diversity, 374
 - Tsengwen delta, 377
- water and sediment dynamics, wetlands and coastal water bodies of large river plains
- avulsion, distributary channel evolution, 27–28
 - climate processes, 38–42
 - controlling physical process, 31–33
 - delta formation, land-building models, 35
 - delta plain components, schematic diagram, 26
 - deltaic plain, lowland floodplain, 21
 - deltaic plain morphology classifications, 25
 - ecomorphology of deltaic plain wetlands, 36–38
 - estuaries, 32
 - groundwater, rainwater input, 29–30
 - high water events, 25
 - Holocene channel belts, paleogeographic maps, 28
 - hydrodynamic and sediment transport conditions, 31
 - ice raft formation, conceptual model, 39
 - large river wetlands, particle sources, sinks, loci of particle transformation, 42–43
 - large-river delta front estuary (LDE), 21
 - numerical modeling tools, 34–36
 - one-dimensional models, 35
 - overbank, crevasses processes, 28–29
 - Regional FVCOM model grid, 35–36
 - relative sea level rise (RSLR), 26, 41
 - stem height relative to water depth, schematic diagram, 37
 - stem properties, 36–38
 - strong winds, storm surge, 26
 - submarine groundwater discharge (SGD), 30
 - subsurface fluids withdrawal, 24–25
 - system differences, 43
 - three-dimensional circulation models, 34
 - tides, winds, 32–33
 - two-, three-dimensional storm surge models, 34–35
 - two-dimensional shallow-water circulation models, 34
 - water, sediment circulation in deltaic plains and estuaries, processes/simulation, 31–34
 - water, sediment input to delta plain mechanisms, 26–30
 - water, sediment process simulation, 33–34
 - water, sediment transport physics, 21
 - wetland deltaic development controls, 38–42
 - wetlands role, 36–43
- Yukon River basin, carbon and nutrient species, composition and fluxes
- average daily stream flow variations, 505, 521
 - C, N, and P partitioning, chemical speciation variations, 519–521
 - carbon species, concentrations and fluxes, 513
 - carbon species of river waters, variations, 505–512
 - chemical, isotopic composition of organic matter, 513
 - chemical composition and isotopic signatures of dissolved (DOM), colloidal (COM), and particulate organic matter (POM), 513, 516
 - chromophoric and fluorescent-DOM, 511–512
 - $\Delta^{14}\text{C}$ values (%) of POM, COM and LMW-DOM, 518
 - elemental, molecular composition, 513–516
 - hydrological regime, environmental change effects, 524–525
 - inorganic carbon species, variations, 509–511
 - isotopic composition, 516–518
 - isotopic composition, heterogeneity, 517–518
 - map, major tributaries, 503–505
 - N, P, and Si abundance and speciation variations, 518–521
 - nutrients export fluxes, 521–523
 - overview, 503–505
 - partitioning of carbon between DIC, DOC and POC pool, 509
 - recent studies on carbon, nutrients and trace elements, 507–508
 - riverine export fluxes of carbon species, quantification, 512–513
 - season and spatial variations, organic carbon species, 507–508
 - total organic carbon and total fatty acids in dissolved organic matter (DOM), colloidal organic matter (COM), and particulate organic matter (POM) pools, 513
 - trace elements transportation, organic matter role, 523–524