

Index

- σ -algebra, 8
- ABO blood group model, 472, 479, 489
- accessible (knowledge structure), 279
- accumulative prediction error, 571
- adapted QUERY routine, 311
- adaptive design optimization, 581
- additive Cauchy equation, 153
- additive representation, 152
- adjacency matrix, 198
- adjusted QUERY routine, 312
- AIC, 527
- Akaike information criterion, 569
- almost hanging (state), 310
- antimatroid, 279
- antisymmetric relation, 283
- assortativity, r , 262
- asymptotically stable, 344
- atom, 284
- attractor, 345
- attribution function, 285

- base, 283
- basin of attraction, 345
- Bayes factor, 525, 573
- Bayes rule, 510
- Bayesian hierarchical modeling, 504, 545
- Bayesian information criterion, 569
- Bayesian probability, 509
- Bernoulli
 - graph model, 252
 - model, 249
- BIC, 527
- big-O notation, $O(g(n))$, 201
- binary (preference) relations, 386
 - lexicographic, 388
 - orders
 - interval, 387
 - semi-, 387
 - strict linear, 387
 - strict partial, 387
 - strict weak, 387
 - weak, 387
- binary multinomial processing tree, 466, 469, 486
- binocular vision, 173
- bipolar coordinates, 174
- birth–death process, 364
- Bishop–Cannings theorem, 329
- Bonini’s paradox, 588
- BUGS, 517

- careless error probability, 300
- child, 299
- choice
 - binary, 383
 - Luce model, 431
 - multinomial logit (MNL) model, 431
 - probabilities
 - attraction effect, 440
 - best, 383
 - best choice induced by rankings, 399
 - best-worst, 383
 - best-worst choice induced by rankings, 400
 - binary, 383
 - binary choice induced by rankings, 397
 - compromise effect, 441
 - constant error model, 390
 - context effects, 439
 - distance-based specification, 393
 - independence of irrelevant alternatives, 431
 - regularity of, 429
 - similarity effect, 441
 - tremble model, 390
 - worst, 383
 - probability distributions, 382
- clause, 285
- clique, 206, 247
- cognitive modeling, 553
- competing risks model
 - hazard based, 30
 - latent-failure times, 32
 - Weibull, 30
- complementation, 246
- completely monotone function, 61
- computational complexity, 199
- concatenation (of a word and an item), 289
- conditional posterior distributions, 515

- conditionally
 - independent, 244
 - uniform models, 259
 - conjugate prior, 513
 - connected, 205
 - component, 205
 - constant ratio condition, 155
 - context-free grammar, 490
 - convex
 - hull, 395
 - polytope, 396
 - set, 395
 - convex dimension, 294
 - convolution, 20
 - copula, 56
 - Archimedean
 - generator, 61
 - Archimedean, 60
 - bivariate extreme value, 57
 - Clayton, 62
 - co-, 60
 - dual of, 60
 - n -dimensional, 57
 - density, 58
 - Fréchet bound, 60
 - independence, 58
 - survival, 59
 - vine, 58
- coupling, 44, 108, 119
 - event, 48
 - inequality, 48
- i.i.d., 46
- identity, 119
- independent, 108, 147
- maximal, 49
- quantile, 46
- reduced, 117–119, 122, 124, 129, 136
- self-, 46
- covered (state covered by), 278
- covering diagram, 278
- cross-validation, 571
-
- d'Alembert's functional equation, 169
- degree
 - digraph, 202
 - graph, 203
 - sequence, 203
- dependence, 63
 - Kendall's tau, 54
 - negative, 66
 - negative lower orthant (*NLOD*), 66
 - negative upper orthant (*NUOD*), 66
 - positive, 63
 - positive lower orthant (*PLOD*), 64
 - positive quadrant, 77
 - positive upper orthant (*PUOD*), 64
 - Spearman's rho, 67
- dependency graph, 247
- derivable operation, 62
- design optimization, 579
- determines (a word determines), 289
- Deviance Information Criterion (DIC), 524, 527
- diameter, 204
- digraph, 469, 485
- discrepancy function, 567
- discriminative (knowledge structure), 276
- discriminative (surmise system), 286
- distribution for degrees
 - β -negative binomial, 218
 - negative binomial, 219
 - Poisson, 218
 - power law, 218
 - Zipf's law, 222
- distribution function, 11
 - Beta, 75
 - bivariate Marshall–Olkin, 15
 - exponential, 12
 - extreme value, 37
 - Fréchet type, 40
 - Gumbel type, 40
 - Weibull type, 40
 - inverse Gauss (or Wald), 20
 - Kumaraswamy, 75
 - normal, 20
 - of the same type, 38
 - sub-, 30
 - uniform, 27
 - Weibull, 12
- domain, 276
- downgradable (knowledge structure), 279
- dyad independence model, 254
-
- eigenvalue, 226
 - dominant, 232
 - second smallest Laplacian, 238
- embedding, 207
- empty word, 289
- encoded (learning space), 294
- entropies of degree α , 170
- equality
 - in distribution, 10
 - point-wise, 10
- equilibrium evolutionarily stable set, 335
- error function, 171
- evolutionarily stable set, 334
- evolutionarily stable state, 348
- evolutionarily stable strategy
 - Maynard Smith and Price definition, 326
 - Taylor–Jonker definition, 327
- exchangeability (of random variables), 24
- explanatory adequacy, 563

- exponential random graph models (ERGMs),
 244, 258
 extended distribution, 300
 extra problem, 314

 facet
 defining, 396
 factor, 111
 failure rate, *see* hazard (rate) function
 faithfulness, 563
 falsifiable, 562
 Fechner's theory, 159
 finite (knowledge structure), 276
 Fisher information approximation, 565, 569
 fixed point
 definition, 344
 hyperbolic, 353
 Fréchet–Hoeffding bound, 50
 comonotonicity, 51
 countermonotonicity, 51
 frequentist probability, 508
 full binary tree, 469, 493
 functional equation, 151
 functional form, 565
 fundamental equation of information measures,
 153
 fundamental theorem of natural selection, 358

 gambles, 183
 branches of, 408
 first-order
 of gains, 408
 idempotence, 409
 mixed, 415
 risky, 409
 uncertain, 409
 utility representations
 configural weighted, 411
 cumulative prospect theory, 410
 ranked additive, 407
 ranked weighted, 407
 TAX, 411
 game
 Battle of the sexes, 342, 360
 coordination, 342
 doubly symmetric, 358
 hawk–dove, 343
 Prisoners' Dilemma, 341, 367
 Rock–Paper–Scissors, 328, 344
 generalizability, 567
 Gibbs sampler, 515, 531
 global identification, 468, 478
 goodness-of-fit, 563
 gradation, 312
 granular (knowledge space), 284
 graph, 202
 bipartite, 204
 connected, 205
 connectivity, κ , 206
 cycle, 202, C_N , 204
 path, 202, P_N , 204
 star, S_N , 204
 subgraph, 205, induced —, $\langle V \rangle$, 205
 groundedness, 57

 half-split rule, 305
 Hammersley–Clifford (H-C) theorem, 244, 248
 hanging (state), 310
 hazard (rate) function, 27
 cause-specific, 30
 cumulative, 28
 sub-, 30
 hidden Markov chain, 455, 458, 460
 hierarchical dual-process models, 540
 hierarchical linear models, 507
 hierarchical model, 499
 hierarchical modeling, 506
 hierarchical nonlinear models, 507
 hierarchical prior, 520
 hierarchical shrinkage, 520
 hierarchical single-process models, 544
 horse race model
 context-dependent, 430
 context-free, 429
 distribution-free, 428
 independent random utility, 429
 linear ballistic accumulator model
 additive, 438
 drift rate variability, 438
 multiattribute, 441
 multiplicative, 434
 of choice and response time, 439
 parallel, 439
 start point variability, 434
 of choice, 427
 hyperbolae of Hillebrand, 173

 identity
 Hoeffding's, 19
 imitate-the-best, 367
 improper prior, 513
 inner fringe, 288
 input, 85, 110, 111
 instance, 275
 interaction neighborhood, 367
 interpretability, 563
 item, 276

 JAGS, 516, 545
 joint posterior, 514
 joint prior, 514
 joint receipt, 183

- knowledge space, 279
 knowledge state, 276
 knowledge structure, 276
- Laplace approximation, 526
 Laplace transform, 61
 Laplacian
 - eigenvalue, algebraic connectivity, 236, 238, 262
 - eigenvector, Fiedler, 238
 - matrix, 236
- latent parameters, 244
 latent state, 306
 learning diagram, 292
 learning path, 312
 learning sequence, 289
 learning space, 278
 learning word, 289
 least squares estimation, 509, 560
 leave-one-out-cross-validation, 572
 length (of a word), 289
 likelihood function, 497, 511, 512, 561
 linear ballistic accumulator model, 433
 linear ordering
 - model
 - of best choice probabilities, 399
 - of best-worst choice probabilities, 400
 - polytope, 398
 - of best choice probabilities, 399
 - of best-worst choice probabilities, 402
 - of worst choice probabilities, 399
- local identification, 468
 local interaction model, 366
 lucky guess probability, 300
 Lyapunov stability, 344
 Lyapunov stability theorem, 347
 Lyapunov's indirect method, 353
- m*-dimensional fundamental equation of
 - information, 171
- marginal likelihood, 525, 573
 marginal posterior, 514
 marginal selectivity, 125
 Markov
 - chain, 249
 - chain Monte Carlo (MCMC), 256
 - graph, 255
- Markov chain, 458
 Markov chain Monte Carlo (MCMC), 515
 mass-at-chance (MAC) link, 530
 maximized likelihood, 563
 maximum likelihood distribution, 567, 570
 maximum likelihood estimation, 509, 560
 measurable
 - function, 9
 - space, 8
- measurable set, *see* sigma-algebra
 minimum description length, 570
 model complexity, 565
 model evaluation, 562
 model revision, 587
 model selection, 569
 model validity, 478
 morphism
 - auto—, 207
 - homeomorphic, 208
 - iso—, 207
- multinomial distribution, 496
 multiplicative updating rule, 306
- Nash equilibrium
 - components of, 335
 - definition, 324
 - existence of, 328
 - inadequate for evolutionary stability, 325
 - relation between strict and ESS, 326
 - rest point of continuous replicator dynamics, 350
 - solution concept, 324
 - strict, 326
- neutrally stable, 344
 normalized maximum likelihood, 565, 569
 normalizing constant, 258
 nuisance variation, 506
 numerosity decision task, 528
- Occam's razor, 567
 OpenBUGS, 516, 517
 optimal design, 577
 order
 - convex, 71
 - dilation, 71
 - dispersive, 72
 - hazard rate, 69
 - likelihood ratio, 70
 - Lorenz, 72
 - lower orthant, 76
 - positive dependence, 76
 - quantile spread, 73
 - stochastic, 68
 - univariate, 68
 - strong stochastic, 75
 - upper orthant, 76
 - variability, 71
- order statistics, 34
 ordinal (space), 282
 outer fringe, 288
 output, 85, 110, 112
 - dependence on inputs, 110
 - selectivity in, *see* selective influence
- overfitting, 564

- pair-clustering model, 474
- parameter estimation, 560
- parametric order constraint, 481
- parametrized (probabilistic knowledge structure), 300
- parent (structure), 299
- partial order, 282
- partially ordinal (space), 282
- payoff function, 324
- penalized-likelihoods, 569
- percent variance accounted for, 563
- Pexider equation, 152
- Pexiderized version, 159
- Plateau's experiment, 164
- plausibility, 563
- Poisson process, 22
 - non-homogeneous, 23
- polytope
 - face of a , 396
 - facet description of, 396
 - facet of a , 396
 - facet-defining inequalities of a , 396
 - full-dimensional, 396
 - hypercube, 384
 - minimal description of, 396
 - vertex description, 396
- posterior, 510
- posterior beliefs, 511
- posterior distribution, 513
- power set, 278
- preference order, 183
- preferences
 - mixture of, 394
 - random, 395
 - revealed, 376
 - stated, 376
- preferential attachment, 219
- prefix (of a word), 289
- preorder traversal, 493
- prior, 510
- prior beliefs, 511
- prior distribution, 512
- probabilistic (knowledge structure), 300
- probabilistic digraph model (PDM), 241, 252
- probabilistic graph model (PGM), 241, 252
- probabilistic projection (of a probabilistic knowledge structure), 300
- probability
 - measure, 8
 - space, 8
- probability density function, 559
- probability redistribution vector, 332
- probability simplex, 467
- probit transformation, 529
- product multinomial, 497
- product space, 19
- projected distribution, 300
- projection (of a knowledge structure), 297
- psychological measurement, 152
- psychophysical invariances, 176
- quantile function, 26
 - density, 26
 - hazard, 29
 - quantile spread, 73
- quasi order, 282
- quasi ordinal (space), 282
- quasiarithmetic mean, 181
- query, 308, 382
- race model, 51
 - inequality, 51
- radius, rad, 204
- random
 - advantage model, 444
 - element, 46
 - function model, 422
 - preference, 395
 - relation model, 395
 - utility model, 420
 - distribution-free, 420
 - noncoincident, 420
 - unidimensional, 420
 - utility representation
 - EBA, 430
 - of best choice, 424
 - of best-worst choice, 424
 - of interval orders, 421
 - of semiorders, 421
 - of strict linear orders, 421
 - of strict weak orders, 421
 - of weak orders, 421
- variable, 10
- vector, 13
- vector
 - associated random vectors, 64
 - negatively associated random vectors (NA), 66
- random variable, 92, 100
 - as measurable function, 105
 - continuous, 101
 - discrete, 100
 - distribution of, 92
 - equality, 109
 - functions of, 102
 - identity of, 94, 98, 109
 - jointly distributed, 95
 - marginal, 98, 100, 105
 - sameness, *see* identity of
 - stochastically independent, 97
 - stochastically unrelated, 107
- rank, 387

- rank-dependent utility representation, 184
 receiver operating characteristic (ROC), 538, 543
 recognition memory, 456, 476, 479, 495, 538, 540
 record, 40
 counting process, 41, 43
 inter-times, 41
 time, 41
 value sequence, 41
 relational structure, 198, 199
 replicator dynamics
 continuous, 337
 discrete, 360
 two populations, 361
 replicator–mutator dynamics, 362
 representation
 configural weighted, 411
 cumulative prospect theory, 410
 full-dimensional, 391
 order-preserving, 408
 ranked additive, 407
 ranked weighted, 407
 separable, 416
 stakes sensitive, 415
 TAX, 411
 root mean square error, 560
- Savage–Dickey ratio, 526
 selective influence, 116
 canonical form, 114
 diagram of, 112
 invariance to transformations, 139
 nestedness, 124
 test, *see* tests of selective influence
 semiorder, 387
 simple lexicographic, 389
 sequence, 289
 Shannon's entropy, 153
 sigma-algebra, 92
 Borel, 93, 103
 discrete, 93, 102
 Lebesgue, 94, 98, 100, 101, 103
 product, 95–97, 99, 100
 signal detection models, 537
 software
 Mathematica[®], 209
 nauty, 267
 Pajek, 209
 RSIENNA, 267
 Traces, 267
 source monitoring, 476, 501
 span, 283
 split-half CV, 571
 splitting representation, 49
 Stan, 545
 star, 204, 255
 state, 276
 straight (parametrized probabilistic knowledge structure), 300
 strategic form game, 324
 strategy
 completely mixed, 330
 locally superior, 331
 mixed, 324
 neutrally stable, 334
 pure, 323
 strictly dominated, 327
 support of, 329
 weakly dominated, 327
 strategy profile, 324
 strict weak order model
 of ternary paired-comparison probabilities, 405
 string language, 486
 structurally unstable point, 345
 subliminal priming, 507, 527, 528, 537
 surmise function, 285
 surmise system, 286
 survival function, 27
 sub-, 30
 symmetric difference, 279
- take-the-last heuristic, 340
 ternary paired-comparison probabilities, 385
 testability, 562
 tests of selective influence, 86, 124, 128
 cosphericity test, 139, 141, 143
 distance test, 135
 joint distribution criterion, 117, 118
 linear feasibility test, 116, 129, 132
 marginal selectivity test, 129
 theorem
 de Finetti, 25
 Sklar, 57
 Strassen, 47
 Tsiatis, 33
 theory of signal detection, 538
 total positivity, 65
 multivariate (MTP_2), 65
 of order 2 (TP_2), 65
 total variation distance, 50
 trace, 297
 transitive closure, 309
 treatment, 111
 allowable, 111, 115
 trial, 301
 trial number, 305
 triangle inequalities, 397
 trivial (child), 299
- uniform extension (of a probabilistic knowledge structure), 300

-
- uniform invasion barrier, 330
 - uniform random graph, 260
 - union-closed (knowledge structure), 279
 - uniqueness theorem, 153
 - unstable point, 345
 - update neighborhood, 367
 - utility
 - configural weighted, 411
 - cumulative prospect theory, 410
 - multiplicative separability of, 418
 - of gambling, 409
 - order-preserving, 408
 - parametric forms, 416
 - rank-dependent, 410
 - ranked weighted, 407
 - representation
 - of gambles, 409
 - of interval orders, 406
 - of semiorders, 406
 - of strict linear orders, 406
 - of strict weak orders, 406
 - TAX, 411
 - weak, 394
 - variability across items, 506
 - variability across participants, 506
 - vector, 224
 - eigen—, 226
 - eigen— centrality, 232
 - inner product of, 224
 - normal, 224
 - orthogonal, 228
 - Perron, 232
 - vertex, 384
 - representation, 391
 - Vieth–Müller circles, 173
 - Weber’s law, 160
 - well-graded (knowledge structure), 279
 - WinBUGS, 516, 545
 - word, 289