Contents

Preface

1

2

| 1.1 | Carle | n structures and hubridizations | |
|------|--|--|--|
| 1.1 | | n structures and hybridizations | |
| | | Carbon nanostructures Guide to the book | |
| | | r reading | |
| Elec | tronic p | roperties of carbon-based nanostructures | |
| 2.1 | Introd | uction | |
| 2.2 | Electro | onic properties of graphene | |
| | 2.2.1 | Tight-binding description of graphene | |
| | 2.2.2 | Effective description close to the Dirac point and massless Dirac fermions | |
| | 2.2.3 | Electronic properties of graphene beyond the linear approximation | |
| 2.3 | Electro | onic properties of few-layer graphene | |
| 2.4 | Electronic properties of graphene nanoribbons | | |
| | 2.4.1 | Electronic properties of armchair nanoribbons (aGNRs) | |
| | 2.4.2 | Electronic properties of zigzag nanoribbons (zGNRs) | |
| 2.5 | Electronic properties of carbon nanotubes | | |
| | 2.5.1 | Structural parameters of CNTs | |
| | 2.5.2 | Electronic structure of CNTs within the zone-folding | |
| | | approximation | |
| | | Curvature effects: beyond the zone-folding model | |
| | | Small-diameter nanotubes: beyond the <i>tight-binding</i> approach | |
| | | Nanotubes in bundles | |
| | | Multiwall nanotubes | |
| 2.6 | - | orbit coupling in graphene | |
| 2.7 | Magnetic field effects in low-dimensional graphene-related materials | | |
| | 2.7.1 | Short historical perspective | |
| | 2.7.2 | Peierls substitution | |

page xi

| | | 2.7.3 Parallel field, Aharonov–Bohm gap opening and orbital | |
|---|-------|---|-----|
| | | degeneracy splitting | 58 |
| | | 2.7.4 Perpendicular field and Landau levels | 62 |
| | | 2.7.5 Landau levels in graphene | 65 |
| | 2.8 | Defects and disorder in graphene-based nanostructures | 67 |
| | | 2.8.1 Structural point defects in graphene | 68 |
| | | 2.8.2 Grain boundaries and extended defects in graphene | 71 |
| | | 2.8.3 Structural defects at graphene edges | 76 |
| | | 2.8.4 Defects in carbon nanotubes | 82 |
| | 2.9 | Further reading and problems | 85 |
| 3 | Qua | ntum transport: general concepts | 91 |
| | 3.1 | Introduction | 91 |
| | | 3.1.1 Relevant time and length scales | 91 |
| | | 3.1.2 Coherent versus sequential transport | 92 |
| | 3.2 | Landauer–Büttiker theory | 94 |
| | | 3.2.1 Heuristic derivation of Landauer's formula | 97 |
| | 3.3 | Boltzmann semiclassical transport | 98 |
| | | 3.3.1 The relaxation time approximation and the Boltzmann | |
| | | conductivity | 99 |
| | 3.4 | Kubo formula for the electronic conductivity | 101 |
| | | 3.4.1 Illustrations for ballistic and diffusive regimes | 105 |
| | | 3.4.2 Kubo versus Landauer | 107 |
| | | 3.4.3 Validity limit of Ohm's law in the quantum regime | 108 |
| | | 3.4.4 The Kubo formalism in real space | 108 |
| | | 3.4.5 Scaling theory of localization | 111 |
| | 3.5 | Quantum transport beyond the fully coherent or decoherent | |
| | | limits | 115 |
| | 3.6 | Further reading and problems | 116 |
| 1 | Kleiı | n tunneling and ballistic transport in graphene and related materials | 118 |
| | 4.1 | The Klein tunneling mechanism | 118 |
| | | 4.1.1 Klein tunneling through monolayer graphene with a single | |
| | | (impurity) potential barrier | 119 |
| | | 4.1.2 Klein tunneling through bilayer graphene with a single (impurity) | |
| | | potential barrier | 124 |
| | 4.2 | Ballistic transport in carbon nanotubes and graphene | 126 |
| | | 4.2.1 Ballistic motion and conductance quantization | 127 |
| | | 4.2.2 Mode decomposition in real space | 128 |
| | | 4.2.3 Fabry-Pérot conductance oscillations | 132 |
| | | 4.2.4 Contact effects: SWNT-based heterojunctions and the role of | |
| | | contacts between metals and carbon-based devices | 135 |

| | | Contents | vii |
|---|-----|--|-----|
| | | | |
| | 4.3 | Ballistic motion through a graphene constriction: the 2D limit and the | 140 |
| | 4.4 | minimum conductivity | 140 |
| | 4.4 | Further reading and problems | 141 |
| 5 | Qua | ntum transport in disordered graphene-based materials | 143 |
| | 5.1 | Elastic mean free path | 143 |
| | | 5.1.1 Temperature dependence of the mean free path | 146 |
| | | 5.1.2 Inelastic mean free path in the high-bias regime | 148 |
| | | 5.1.3 Quantum interference effects and localization phenomena in | |
| | | disordered graphene-based materials | 150 |
| | | 5.1.4 Edge disorder and transport gaps in graphene nanoribbons | 152 |
| | 5.2 | Transport properties in disordered two-dimensional graphene | 154 |
| | | 5.2.1 Two-dimensional disordered graphene: experimental and | |
| | | theoretical overview | 154 |
| | | 5.2.2 Metallic versus insulating state and minimum conductivity | 158 |
| | | 5.2.3 Boltzmann transport in two-dimensional graphene | 158 |
| | | 5.2.4 Kubo transport: graphene with Anderson disorder | 166 |
| | | 5.2.5 Kubo transport: graphene with Gaussian impurities | 169 |
| | | 5.2.6 Weak localization phenonema in disordered graphene | 173 |
| | | 5.2.7 Strong localization in disordered graphene | 181 |
| | 5.3 | Quantum Hall effect in graphene | 182 |
| | | 5.3.1 Hall quantization in graphene | 183 |
| | | 5.3.2 The mystery of the zero-energy Landau level splitting | 184 |
| | | 5.3.3 Universal longitudinal conductivity at the Dirac point | 185 |
| | 5.4 | Graphene with monovacancies | 187 |
| | | 5.4.1 Electronic structure of graphene with monovacancies | 190 |
| | | 5.4.2 Transport features of graphene with monovacancies | 191 |
| | 5.5 | Polycrystalline graphene | 195 |
| | | 5.5.1 Motivation and structural models | 195 |
| | | 5.5.2 Electronic properties of polycrystalline graphene | 199 |
| | | 5.5.3 Mean free path, conductivity and charge mobility | 201 |
| | 5.6 | Amorphous graphene | 202 |
| | 5.0 | 5.6.1 Structural models | 202 |
| | | 5.6.2 Electronic properties of amorphous graphene | 202 |
| | | 5.6.3 Mean free path, conductivity and localization | 203 |
| | 5.7 | Phonon transport in graphene-related materials | 204 |
| | 5.7 | 5.7.1 Computational phonon propagation methodology | 200 |
| | | 5.7.2 Disordered carbon nanotubes with isotope impurities | 200 |
| | | 5.7.2 Disordered graphene nanoribbons with edge disorder | 208 |
| | 5.8 | | 209 |
| | 3.8 | Graphene quantum dots 5.8.1 Generalities on Coulomb blockade | |
| | | | 212 |
| | 5.0 | 5.8.2 Confining charges in graphene devices | 214 |
| | 5.9 | Further reading and problems | 217 |

| viii | Cont | ents | |
|------|--------------|--|------------|
| 6 | Quar | ntum transport beyond DC | 219 |
| U | | | 219 |
| | 6.1 | Introduction: why AC fields? | 219 |
| | | Adiabatic approximation Floquet theory | 220 |
| | 0.5 | 6.3.1 Average current and density of states | 221 |
| | | 6.3.2 Homogeneous driving and the Tien–Gordon model | 222 |
| | | 6.3.3 Time-evolution operator | 224 |
| | 6.4 | | 225 |
| | 6.5 | AC transport and laser-induced effects on the electronic properties | 220 |
| | 010 | of graphene | 227 |
| | 6.6 | Further reading and problems | 230 |
| 7 | Ab ii | nitio and multiscale quantum transport in graphene-based materials | 232 |
| | 7.1 | Introduction | 232 |
| | 7.2 | Chemically doped nanotubes | 233 |
| | | 7.2.1 Tight-binding Hamiltonian of the pristine carbon nanotube | 233 |
| | | 7.2.2 Boron-doped metallic carbon nanotubes | 233 |
| | | 7.2.3 Nitrogen-doped metallic carbon nanotubes | 236 |
| | 7.3 | Two-dimensional disordered graphene with adatoms defects | 242 |
| | | 7.3.1 Monatomic oxygen defects | 242 |
| | | 7.3.2 Atomic hydrogen defects | 246 |
| | | 7.3.3 Scattering times | 248 |
| | 7.4 | | 249 |
| | 7.5 | Ab initio quantum transport in 1D carbon nanostructures | 251 |
| | | 7.5.1 Introduction | 251 |
| | | 7.5.2 Carbon nanotubes | 253 |
| | | 7.5.3 Defective carbon nanotubes | 255 |
| | | 7.5.4 Doped carbon nanotubes | 259 |
| | | 7.5.5 Functionalized carbon nanotubes | 261 |
| | | 7.5.6 Carbon nanotubes decorated with metal clusters | 269 |
| | | 7.5.7 Graphene nanoribbons7.5.8 Graphene nanoribbons with point defects | 278 |
| | | | 280 281 |
| | | 7.5.9 Graphene nanoribbons with edge reconstruction7.5.10 Graphene nanoribbons with edge disorder | 281 |
| | | 7.5.11 Doped graphene nanoribbons | 282 |
| | | 7.5.12 GNR-based networks | 200 293 |
| | 7.6 | Conclusion | 293 298 |
| 8 | Applications | | 300 |
| | 8.1 | Introduction | 300 |
| | 8.2 | Flexible electronics | 301 |
| | 8.3 | High-frequency electronics | 302 |

| | Contents | b |
|------------|---|------------|
| | | |
| 8.4 | Optoelectronics-photonics-plasmonics | 304 |
| | 8.4.1 Opacity of graphene and fine structure constant | 304 |
| | 8.4.2 Harnessing graphene with light: a new dimension of possibilities | 305 |
| 8.5 | 8 | 306 |
| 8.6 | | 307 |
| 8.7 | 1 | 308 |
| 8.8 | 1 | 309 |
| 8.9 | Further reading | 313 |
| Appendix A | Electronic structure calculations: the density functional theory (DFT) | 314 |
| A.1 | | 314 |
| A.2 | Overview of the approximations | 314 |
| | A.2.1 The Schrödinger equation | 314 |
| | A.2.2 The Born–Oppenheimer approximation | 315 |
| | A.2.3 The Hartree approximation | 316 |
| A 2 | A.2.4 The Hartree–Fock approximation | 317 318 |
| A.3 | Density functional theory A.3.1 The Thomas–Fermi model | 318 |
| | A.3.2 The Hohenberg–Kohn theorem | 319 |
| | A.3.3 The Kohn–Sham equations | 320 |
| | A.3.4 The exchange–correlation functionals | 320 |
| A.4 | Practical calculations | 324 |
| | A.4.1 Crystal lattice and reciprocal space | 324 |
| | A.4.2 The plane wave representation | 325 |
| | A.4.3 k-point grids and band structures | 326 |
| | A.4.4 The pseudopotential approximation | 326 |
| | A.4.5 Available DFT codes | 330 |
| Appendix B | Electronic structure calculations: the many-body perturbation | |
| | theory (MBPT) | 332 |
| B.1 | Introduction | 332 |
| B.2 | Many-body perturbation theory (MBPT) | 333 |
| | B.2.1 Hedin's equations | 333 |
| | B.2.2 <i>GW</i> approximation | 334 |
| B.3 | 1 0 0 | 335 |
| | B.3.1 Perturbative approach | 335 |
| | B.3.2 Plasmon pole | 336 |
| Appendix C | Green's functions and <i>ab initio</i> quantum transport in the Landauer–Büttiker formalism | 338 |
| | | |
| C.1 | The second | 338 |
| C.2 | Self-energy corrections and recursive Green's functions techniques | 344 |
| | | |

| Cambridge University Press |
|---|
| 978-1-107-03083-1 - Introduction to Graphene-Based Nanomaterials: From Electronic Structure |
| to Quantum Transport |
| Luis E. F. Foa Torres, Stephan Roche and Jean-Christophe Charlier |
| Table of Contents |
| More information |
| |

| X | Contents | | | |
|----------|----------|--|-----|--|
| | C.3 | Dyson's equation and an application to treatment of disordered systems | 347 | |
| | C.4 | Computing transport properties within ab initio simulations | 351 | |
| Appendix | D | Recursion methods for computing the DOS and wavepacket dynamics | 358 | |
| | D.1 | Lanczos method for the density of states | 358 | |
| | | D.1.1 Termination of the continued fraction | 361 | |
| | D.2 | Wavepacket propagation method | 363 | |
| | D.3 | Lanczos method for computing off-diagonal Green's functions | 367 | |
| | Refe | rences | 370 | |
| | Inde. | x | 405 | |