

Index

Locators in **bold** refer to figures and tables

- ABM (attention bias modification), 268
- absolute ceiling effect, assessment, 235
- abuse, childhood, 5. *see also* childhood trauma
- academic functioning. *see* educational functioning
- action-based cognitive remediation (ABCR), 315, 316, 317
- activities of daily living (ADL), 16, 161
- activity scheduling, cognitive remediation, 315, 316
- acute clinical stage, inflammation, 163, 164
- AD. *see* Alzheimer's disease
- ADHD. *see* attention deficit/hyperactivity disorder
- adherence to treatment, 161
- clinician–patient relationship, 42, 43
- and cognitive dysfunction, 32–33
- exercise, 333
- functional outcomes, 140
- adolescent depression. *see* pediatric depression
- Advanced Finances Task (AFT), 139
- aerobic exercise. *see* exercise
- affective neuroscience, 257
- age of onset, 1, 113, 117, 119. *see also* late-onset MDD
- agitation. *see* psychomotor speed
- AIC (anti-inflammatory cytokines), 162
- alcohol intoxication, 18, 19, 20
- algorithms, assessment, 37
- alternative medicine, 275, 286
- Alzheimer's disease (AD), 16, 17, 18, 24, 237
- insulin resistance, 214, 215, 216
- American Academy of Child and Adolescent Psychiatry (AACAP), 47
- amphetamine, 23
- amygdala
- anatomy, 63
- biological mechanisms of MDD development, 5–6
- biomarkers of illness, 149, 153, 154
- emotional processing, 83, 202, 263, 264
- inflammation, 168, 169
- negative cognitive bias, 202
- neural circuits, 197
- post-recovery deficits, 111
- progressive cognitive deficits, 117
- rumination, 202–203
- social emotional processing, 83, 93, 96
- social exclusion, 88
- anatomy of brain, 60–66, 116–117
- anhedonia, 83, 84
- animal studies, erythropoietin, 292, 293
- ANT (Attention Network Task), 49–50, 51
- antecedents to illness. *see* predictions of future depression; premorbid cognitive function
- anterior cingulate cortex. *see* cingulate cortex
- anticholinergic effects, medication, 283. *see also* tricyclic antidepressants
- antidepressant medication, 275–280, 278, 283
- amygdala, 197
- baseline cognitive function, 274
- biomarkers of illness, 74, 151
- combined with CBT, 274
- effects on cognitive dysfunction, 160, 275
- emotional processing, 264
- erythropoietin, 291
- functional outcomes, 126
- hot cognition, 72, 73
- insulin resistance, 213
- neurotoxicity hypothesis, 119, 120
- novel/fast-acting, 74–76
- response to treatment, 169, 260
- social cognition, 106
- see also* SNRIs; SSRIs; tricyclic antidepressants
- antiglucocorticoid medication, 188
- anti-inflammatory cytokines (AIC), 162
- anti-inflammatory treatments, 162
- erythropoietin, 293
- insulin treatment, 282–283
- anxiety disorders, 39, 64–65, 83
- apoptosis, 117, 293
- Applied Cognition-General Concerns (ACGC) Scale, 250
- arithmetic, 261, 309–310
- ARTIST study, pain comorbidity, 39
- ASRS (Attention Deficit Hyperactivity Disorder Self-Report Scale), 280
- assessment, indications for, 35–37. *see also* measurement
- astrocytes, 61–62
- attention, 1
- age of illness onset, 113
- biomarkers of illness, 147, 147, 149, 150–151
- clinician–patient relationship, 42
- and functional outcomes, 129–134, 258–260
- measures, 229–230, 231
- negative cognitive bias, 201–202
- neural circuits, 199, 201–202
- number of lifetime depressive episodes, 116
- objective measures, 36
- pediatric depression, 48–52
- post-recovery deficits, 111
- predictive validity of deficits, 15–17, 17
- psychotherapy, 119–120
- and rumination, 264–265
- three-component model, 6
- treatment interventions, 275–276
- see also* cold cognition
- attention bias modification (ABM), 268
- Attention Control Training Intervention, 309
- attention-deficit/hyperactivity disorder (ADHD)
- antecedents to illness, 3–5, 8, 8–11
- white matter integrity, 65
- working memory, 71
- Attention/Deficit Hyperactivity Disorder Self-Report Scale (ASRS), 280
- Attention Network Task (ANT), 49–50, 51

- Auditory Verbal Learning Test (AVLT), 55
- Australian study, alcohol intoxication, 18–20, **19**
- autism, social cognition, 92
- autonomic nervous system, and insulin resistance, 210
- BAC (blood alcohol concentration), 18–19, **19**
- baseline cognitive function and antidepressant medication, 274
- measures, 237
- primary care, 37, 41
- basolateral nucleus (BLA), amygdala, 168
- BBB (blood–brain barrier), 61–62, 289, 291
- BDNF. *see* brain-derived neurotrophic factor
- BDQ (Brief Disability Questionnaire), 128, 135
- Beck Depression Inventory (BDI), 32–33, **33**
- Beck model. *see* cognitive model of depression
- Behavior and Symptom Identification Scale (BASIS-32), 138
- behavioral inhibition. *see* response inhibition
- behavioral treatment interventions. *see* cognitive behavioral therapy; cognitive remediation therapy; diet; exercise
- benzodiazepines, 40, 118
- bias, memory, 265, 268–269. *see also* negative cognitive bias
- biofeedback, 266
- biological models. *see* neurobiology of depression
- biomarkers of illness, 145, **147**, **150**, 151–154
- antidepressant medication, 151
- cytokines, 169
- default mode network, 217
- dehydroepiandrosterone, 183
- early morning cortisol, 74
- endophenotypes, 145–146, 147–148, 149
- epiphenomena, 148, 151–154
- fractional anisotropy, 64–65, 66
- glial cells, 61–62
- hippocampal volume, 63
- hot vs. cold cognition, 70
- insulin resistance, 213
- neuroimaging/genetic, 146
- oxytocin, 88
- pediatric depression, 55, 56
- symptomatic and remitted stages, **147**, **150**, **160**
- white matter neurobiology, 203
- bipolar disorder
- antecedents to illness, 5
- attention and processing speed, 260
- biomarkers of illness, 153
- cellular anatomy, 62
- erythropoietin, **300**
- HPA axis, 184–185, **186**
- insulin treatment, 282–283
- progressive cognitive deficits, 116
- white matter integrity, 65
- blood alcohol concentration (BAC), 18–19, **19**
- blood–brain barrier (BBB), 61, 289, 291
- blood flow to brain. *see* cerebral blood flow
- blood lipids, and insulin resistance, 210, 213, 215
- blood oxygen level detection (BOLD), 61
- body language, 92, 102, 106. *see also* social cognition
- body mass index (BMI), 330
- bottom-up process, hot cognition, 70, 73
- BPI (Brief Pain Index), 38
- brain anatomy, 62–64, 116–117
- brain connectivity. *see* neural connectivity
- brain-derived neurotrophic factor (BDNF)
- biomarkers of illness, 146
- erythropoietin, 289
- exercise effects, 329, 332–333
- HPA axis, 187
- progressive cognitive deficits, 117
- brain function, 5–6, 60
- biomarkers of illness, 153
- cognitive remediation, **314**
- and insulin resistance, 211, 215–217
- pediatric depression, 52–53
- reward processing, 54–55
- self-focus, 84–85
- social and emotional brain, 82
- see also* neural connectivity; neurobiology of depression; neuroscience of functional outcomes
- brain injury, cognitive remediation, 306, 319
- brain size, and cognitive dysfunction, 63–64
- brain structure, chronic exercise effects, 329, 330
- bridging activities, cognitive remediation, 315
- BRIEF-A self-report scale, 281
- Brief Disability Questionnaire (BDQ), 128, 135
- Brief Pain Index (BPI), 38
- British Columbia Cognitive Complaints Inventory (BC-CCI), 246–247, **247**, **251**
- Brixton Spatial Anticipation Test, 232
- bupropion, 276
- calcium homeostasis, 210
- California Verbal Learning Test (CVLT), 152, 276
- CAM. *see* complementary alternative medicines
- Cambridge Cognitive Examination-Revised (CAMCOR), 2, **9**
- cancer risks, erythropoietin, 301
- CANTAB (Cambridge Neuropsychological Test Automated Battery), 70, 71, 117, 232
- functional outcomes, **131**, **133**, **138**
- quality of life, 137
- cardiovascular disease, 113, 209
- case finding approach, primary care, 37
- CAT (computerized adaptive testing), 250
- catastrophic response, to negative feedback, 72
- catastrophizing
- cognitive dysfunction, 245
- insomnia comorbidity, 40
- see also* negative cognitive bias
- catecholamines, 328. *see also* dopamine; norepinephrine; serotonin
- caudate, 263
- causation, cognitive dysfunction, 153
- CBB (CogState Brief Battery), 21
- CBT. *see* cognitive behavioral therapy
- CCN. *see* cognitive control network
- CDQ (Cognitive Dysfunction Questionnaire), 250
- cellular anatomy, brain, 61–62, 65
- central nucleus (CeA), amygdala, 168
- central tendency, cognitive dysfunction, 237
- cerebral autosomal
- dominant arteriopathy, subcortical infarcts, and leukoencephalopathy (CADASIL), 200
- cerebral blood flow
- endophenotypes, 147
- insulin resistance, 210
- changes over time, 25
- self-report instruments, 245
- significance of, 23
- childhood trauma
- biological mechanisms of MDD
- development, 5–6
- biomarkers of illness, 152
- HPA axis stress response, 185
- physical and sexual abuse, 6
- see also* neurodevelopmental abnormalities; stress

- children. *see* pediatric depression
- Choice Reaction Time (CRT) task, 36, 231
- chronic fatigue syndrome (CFS), 182
- cingulate cortex, 147
- anatomy, 63
 - attentional bias, 201–202
 - biological mechanisms of MDD development, 5–6
 - deep brain stimulation, 76
 - emotion perception, 263
 - neural circuits, 199
 - neuroscience of functional outcomes, 259
 - pain, 38
 - pediatric depression, 53
 - progressive cognitive deficits, 117
 - self-focus, 85
 - social cognition, 96–99
 - social exclusion, 86, 87, 88
 - working memory and executive function, 262
- citalopram, 106
- clinical outcomes, 73, 140. *see also* functional outcomes and recovery
- clinician–patient relationship, primary care, 41–43
- coding tasks, 230
- cognition
- cytokine effects, 165–166
 - definitions, 60
 - emotion linkage, 149, 152, 153
 - HPA axis, 180–182, 185–187, 186
 - insulin resistance, 211
 - see also* social cognition
 - cognitive activation, 307–308
 - cognitive remission. *see* remission
- Cognitive and Physical Functioning Questionnaire (CPFQ), 35, 247–248, 251–252
- cognitive behavioral therapy (CBT)
- combined with antidepressant medication, 274
 - HPA axis, 187
 - neuroscience of functional outcomes, 268
 - progressive cognitive deficits, 120
 - response to treatment, 182
- cognitive control network (CCN), 258, 263, 268
- attention and processing speed, 259
 - emotion perception, 263
 - and treatment targets/outcomes, 266
 - working memory and executive function, 261
- cognitive distortions. *see* negative cognitive bias
- Cognitive Drug Research (CDR)
- computerized assessment system, 277
 - cognitive dysfunction in MDD, 64–65, 242–244, 243
 - biomarkers of illness, 147–154, 147, 150
 - and brain size, 63, 64
 - clinical significance, 16
 - efficacy of treatment interventions, 284–285
 - and functional outcomes, 128, 129–134
 - measures, 233, 234
 - neural correlates, 60, 64
 - pediatric studies, 49–50
 - self-report instruments, 34. *see also* Perceived Deficits Questionnaire
- Cognitive Dysfunction Questionnaire (CDQ), 250
- cognitive efficiency. *see* processing speed
- cognitive flexibility, 48, 52, 232, 308. *see also* executive function; neuroplasticity
- Cognitive Impairment Associated with Schizophrenia (CIAS), 237
- cognitive-interpersonal vulnerability to depression models, 265
- cognitive model of depression, 72, 149, 201
- and emotion distortion and biases, 258, 264
- cognitive neuropsychological model of depression, 149, 150
- cognitive neuroscience, distinction from affective neuroscience, 257
- cognitive reappraisal, 265
- cognitive remediation (CR), 266, 306–307
- brain function model, 314
 - evidence-based studies, 313, 314
 - future directions for study, 318–319
 - HPA axis, 188
 - medication interactions, 318
 - motivational issues, 317–318
 - pillars of treatment, 307–309
 - pseudospecificity of treatment response, 318
 - psychotherapy model, 314, 315–316, 315, 317
 - rationale for using with MDD, 309
 - symptom profiles, 317
 - treatment interventions, 282, 284–286
- cognitive reserve, 3. *see also* IQ
- cognitive therapies, 197. *see also* cognitive behavioral therapy
- CogState Brief Battery (CBB), 21
- Cohen's *d* values, 16, 17, 20, 21, 71
- cold cognition, 257
- antecedents to illness, 3, 4, 5, 6, 8–11
- clinician–patient relationship, 41, 42
- cognitive alterations during course of illness, 69–76
- combination treatments, 274
- neural circuits, 70
- neuroanatomy, 60
- novel/fast-acting medication, 75
- pediatric depression, 48, 51, 52
- psychological interventions, 154
- screening tools in primary care, 32–34, 33
- as trait biomarkers, 70, 71, 74
- comorbidity
- addressing in primary care, 31, 32
 - assessment in primary care, 37–40
 - cardiovascular, 113
 - cognitive remediation, 317
 - functional outcomes, 140
 - pediatric depression, 56
- complementary alternative medicines (CAM), 275, 284–286
- compliance. *see* adherence to treatment
- computer-assisted cognitive remediation, 307–308, 311, 312
- computerized adaptive testing (CAT), 250
- computerized assessment instruments, 126
- concentration, 15, 131, 147
- confounding factors
- erythropoietin, 294
 - measures of cognitive dysfunction and change, 236, 237
 - medication, 63, 117
 - pediatric illness, 56
 - progressive cognitive deficits, 117
- connectivity, brain. *see* neural connectivity
- Continuous Performance Test (CPT), 49, 51
- Controlled Oral Word Association Test, 232
- cortico-limbic circuits, 153
- corticosteroids, 183, 184. *see also* hypocortisolism
- cortico-striatal–pallidal–thalamic circuits, 198
- cortisol levels
- biomarkers of illness, 55, 74
 - salivary, 55, 180
 - social exclusion, 86
- CPFQ (Cognitive and Physical Functioning Questionnaire), 35, 247–278, 251–252
- CPT. *see* Continuous Performance Test

- CR. *see* cognitive remediation
 CRH-R1 (CRH receptor 1 gene), 183–184
 CRT (Choice Reaction Time) task, 36, 231
 cumulative number of episodes. *see* number of lifetime depressive episodes
 cyberball paradigm, social exclusion, 86, 88
 cytokine models of depression and cognition, 162
 amygdala, 165, 168–169
 molecular mechanisms, 165–166
 neurodegenerative effects, 166–167
see also inflammation
- data analysis, measures of cognitive dysfunction, 237–239
 DDS (Denver Development Screening Test), 5
 decision-making, DSM criteria, 15
 declarative memory, 112, 181
 deep brain stimulation (DBS), 76, 281
 default mode network (DMN), 82, 87
 insulin resistance, 217–219, 219
 neuroscience of functional outcomes, 258, 267, 268
 Deficits in Executive Function Scale (DEFS), 4
 definitions
 cognition, 60
 emotions, 81
 epiphenomena, 151
 executive function, 261
 exercise, 321–322
 social cognition, 102, 106
 social processes, 82
 working memory, 261, 325
 dehydroepiandrosterone (DHEA), 183, 188
 dementia
 prodromal phase, 119, 235
 and progressive cognitive deficits, 110, 118–120
 social cognition, 96–99
see also Alzheimer's disease
 DEMO trial, exercise, 332, 333
 demyelination, 195
 Denver Development Screening Test (DDS), 5
 developmental abnormalities. *see* neurodevelopmental abnormalities
 dexamethasone suppression test (DST), 183, 184, 186
 DHEA (dehydroepiandrosterone), 183, 188
 diabetes, 209. *see also* insulin resistance
Diagnostic and Statistical Manual of Mental Disorders (DSM-5), 1, 15, 69, 147
 diary techniques, 86
 diet, treatment interventions, 284–285
 diffusion-tensor imaging (DTI), 61, 64
 Digit Span Test, 334
 Digit Symbol Substitution Test (DSST), 36, 230, 238, 277, 278, 279
 distortions, cognitive. *see* negative cognitive bias
 diurnal patterns of cognitive dysfunction, 236
 divided attention, 258, 260. *see also* attention
 DLPFC. *see* dorsolateral prefrontal cortex
 DMN. *see* default mode network
 dmPFC (dorsomedial prefrontal cortex), 267
 donepezil, 23
 dopamine, 328
 dopamine reuptake inhibitors, 75
 dorsal anterior cingulate cortex. *see* cingulate cortex
 dorsolateral prefrontal cortex (DLPFC), 147
 attentional bias, 201–202
 biological mechanisms of MDD development, 5–6
 biomarkers of illness, 149, 153
 emotional processing, 202
 neuroscience of functional outcomes, 259
 self-focus, 84, 85
 social exclusion, 87
 dorsomedial prefrontal cortex (dmPFC), 267
 drug treatment. *see* medication
 DSST. *see* Digit Symbol Substitution Test
 DTI (diffusion-tensor imaging), 61, 64
 duloxetine, 275, 277, 279, 280, 283, 284, 285, 318
 duration of illness
 anatomy of brain, 63
 biomarkers of illness, 152, 153
 and clinical outcome, 73
 and rumination, 265
see also number of lifetime depressive episodes
 dyslipidemia, 210, 213, 215
 early intervention
 biomarkers of illness, 73–74
 pediatric depression, 56
 predicting future depression, 8
 and progressive cognitive deficits, 119
see also treatment interventions
 education, patient, 32, 43
 educational functioning, 48, 70
 ADHD, 3
 and cognitive dysfunction, 17–20, 18
 working memory, 71
 effect sizes, measures of cognitive dysfunction, 229, 278
 effective ceiling effect, assessment, 235
 effort-automatic hypothesis, 152
 electroconvulsive therapy (ECT), 281, 285, 291, 310
 emotion network (EN), 258, 267, 268
 emotion(s)/emotional processing
 antidepressant medication, 264
 bias, 1, 83, 202, 264. *see also* negative cognitive bias
 brain function, 82
 and cognition, 149, 152, 153
 cytokine model, 165, 166, 168, 170
 definitions, 81
 erythropoietin, 294, 295–296, 297, 298
 neuroscience of functional outcomes, 263–264
 regulation, 82, 87, 257, 258. *see also* response inhibition
 top-down processes, 197
see also hot cognition
 employment status, 70, 161
 and cognitive function, 16, 17, 18, 20, 21, 130, 131, 133
 functional outcomes, 125, 126, 127, 135–137
 primary care clinician–patient relationship, 43
 and progressive cognitive deficits, 119
 working memory and executive function, 262
see also psychosocial functioning
 EN (emotion network), 258, 267, 268
 Endicott Work Productivity Scale (EWPS), 280
 endophenotypes, 110
 antecedents to illness, 112
 biomarkers of illness, 145–146, 147, 149
 enjoyment of life, psychosocial functioning, 138–139
 environmental factors
 biomarkers of illness, 153
 measures of cognitive dysfunction and change, 236
 epidemiology
 cognitive dysfunction in MDD, 242, 243
 insomnia comorbidity, 39
 major depressive disorder, 1, 15
 pain comorbidity, 37, 38
 recurrence risk, 257
 youth depression, 47
 epigenetics, 146, 185

- epiphenomena
 cognitive deficit biomarkers of illness, 148, 151–154
 definitions, 151
see also state vs. trait cognitive dysfunction
- episodic memory
 and functional outcomes, **129**
 measures, 229, 231
 predictive validity, 2, 15
 Epworth Sleepiness Scale, 40
 erythropoietin (EPO), 289, 290–291, **302**
 animal studies, **292**, 293
 clinical limitations, 299–302
 clinical trials, 293–299, **295–296**, **298**, **300**
 confounding factors, 294
 future perspectives, 302
 memory, 290, 291, 293, 294, **295–296**, **298**, 299, **301**
- escitalopram, 275, 312, 318
 etiology, cognitive dysfunction, 153
 European Study of the Epidemiology of Mental Disorders, 135
 evidence-based studies
 cognitive remediation, 309–315, **313**
 exercise, 323, 333
see also meta-analyses
- EWPS (Endicott Work Productivity Scale), 280
 exclusion, social, 81, 86, 87, **87**, 88
 executive function, 1
 age of illness onset, 113
 biomarkers of illness, **147**, 149
 definitions, 261
 erythropoietin, **295–296**, 299
 and functional outcomes, **129–134**, 261
 insulin resistance, 209, 211
 measures, 230, 232
 neural circuits, 200
 number of lifetime depressive episodes, 116
 objective measures, **36**
 pediatric depression, 48, 51, 52
 post-recovery deficits, 111
 predictive validity, 3, 4, 15
 primary care clinician–patient relationship, **42**
 progressive cognitive deficits, 119
 three-component model, 6
 treatment interventions, 275
 tricyclic antidepressants, 117
see also cold cognition
- exercise, **286**, 321–323, 334
 acute exercise effects, 323–325, 328–329
 chronic exercise effects, 325–327, 329–331
 evidence-based studies, 323, 333
 functional mechanisms, 327–331
 pediatric depression, 56
 treatment interventions, 281, 282, **284–285**
 WHO recommendations, 322
- experiment effects, measures of cognitive dysfunction, 236
 explicit (declarative) memory, 112, 181
 eye-tracking
 pediatric depression, 52
 social and emotional processing, 83
- FA (fractional anisotropy), 64, 65, 66
 Faces Pain-Rating Scale (FPRS), 38
 facial processing, 92, 93–95, 265
 case-control studies, **96–99**, **101**
 emotion perception, 262–264
 negative bias, 83, 154
 novel/fast-acting medication, 75
 pediatric depression, 53
 and severity of depression, **103–105**
see also social cognition
- false positive classification, cognitive dysfunction, 20, 21
 familial studies, 149. *see also* twin studies
- far transfer, cognitive remediation, 266
 fast-acting medication, 75
 fatigue, and cognitive dysfunction, 18, 19, **19**
- fear conditioning, amygdala, 168
 feedback mechanisms, HPA axis, **180**. *see also* homeostasis
- fight or flight response, 72, 214
 first episodes of MDD, biomarkers, 148
 FKBP5 gene, 183
 flexibility. *see* cognitive flexibility; neuroplasticity
- fMRI. *see* functional magnetic resonance imaging
 FOCUS study, 238, **238**
 forced choice paradigm, 231
 FPRS (Faces Pain-Rating Scale), 38
 fractional anisotropy (FA), 64, 65, 66
 frontal cortex, 63. *see also* prefrontal cortex
 frontal lobes, white matter tracts, 200
 fronto-cingulate cortex, pediatric depression, 52
 fronto-striato-limbic white matter tracts, 200
 functional connectivity analysis, default mode network, 218, **219**
 functional magnetic resonance imaging (fMRI)
 neural circuits, 197
 pediatric depression, 52
 social exclusion, 88
- functional outcomes and recovery, 125, 126, 128, 140, 161
 assessment, 128, 135
 attention and processing speed, 260
 clinical implications, 139–141
 cognitive dysfunction, 70, 110, 128, **129–134**
 emotion distortion and biases, 265–266
 emotion perception, 263
 neuroscience of, 257–258
 occupational functioning, 135–137. *see also* employment status
 primary care, 30, 31
 and progressive cognitive deficits, 119
 quality of life, 137–138
 social cognition, 93, 95–100, 106
 subjective measures, **243**
 working memory and executive function, 262
see also psychosocial functioning; remission
- Gage, Phineas, 200
 gamma aminobutyric acid (GABA), 18, 195
 gaze behavior. *see* eye-tracking
 gender effects, cognitive deficits, 117
 genetics
 antecedents to illness, 112
 biomarkers of illness, 146, 147, 153, 169
 familial studies, 149
 HPA axis stress response, 183–184, 185
 predicting future depression, 6–8
 serotonin transporter gene, 74
 white matter neurobiology, 195–196
see also endophenotypes
- glial cells, 61, 62, 65
 cytokine model, 167
 pathology of white matter, 195
 progressive cognitive deficits, 117
 Global Severity Index, psychological functioning, **5**, **10**
 glucocorticoids/glucocorticoid receptors, 162, 180, 181, 183, 185
 glucose metabolism, 61, 63. *see also* insulin resistance
 glutamate neurotransmission, 61, 195
 glycogen synthase kinase 3 beta (GSK3 β), 293
 goal-setting, cognitive remediation, 317, 318
 gold standard measures
 cognitive remediation, 306
 neuropsychological assessment as, 244
 and subjective measures, 250

- Go/NoGo task, 232
 exercise, 333
 pediatric depression, **49–50**,
 50, 51
- gray matter, 62, 63, 66
- group therapy, 100
- Hamilton Depression Rating Scale (HDRS), **9**
- “hate circuit” (white matter tracts), 197
- health-related quality of life. *see* quality of life
- heritability, 7, 153. *see also* genetics
- HIF (hypoxia inducible factor), 290
- hippocampus/hippocampal volume, 60
 anatomy, 62–63, 63
 antidepressant medication, 119
 biological mechanisms of MDD development, 5–6
 biomarker for response to treatment, 63
 biomarkers of illness, 148, 152
 erythropoietin, 290, 291, 294, **295–296**, 297, **298**
 exercise effects, 329–331, 332–333
 insulin resistance, 209, 211, 215–217, **217**, **218**
 neural connectivity, 220. *see also* default mode network
 pathology of white matter, 195
 progressive cognitive deficits, 116–117
 rumination, 202–203
 stress effects, 153, 180–182
see also long-term potentiation
- holistic treatment interventions, 76. *see also* multi-modal approaches
- HOMA-IR, 216–217, **217**, **218**. *see also* insulin resistance
- homeostasis
 calcium, 210
 endocrine, 209
 HPA axis, **180**, 182
 and threat of social rejection, 87
- homework, cognitive remediation, 312, **313**
- hospitalizations, number of lifetime depressive episodes, 113–118
- hot cognition, 1, 257
 biological mechanisms of MDD development, 6
 biomarkers of illness, 154
 clinician–patient relationship, 41, **41**
 cognitive alterations during course of illness, 69–73
 combination treatments, 274
 deep brain stimulation, 76
 erythropoietin, 297
- insomnia comorbidity, 40
- mindfulness-based meditation processes, 283
- neural circuits, 70
- neuroanatomy, 60
- novel/fast-acting medication, 76
- objective measures, 35
- pediatric depression, 48
- screening tools in primary care, 32, **33**
- as state biomarkers, 70
see also emotion(s)/emotional processing
- HPA axis. *see* hypothalamic–pituitary–adrenal (HPA) axis
- hypercortisolism, 179, 182
 antecedents to illness, 112
 biological mechanisms of MDD development, 5–6
 insulin resistance, 214
 post-recovery deficits, 111
 progressive cognitive deficits, 117
- hyperreactivity, neural networks, 268
 default mode network, 267
 emotion perception, 263
 prefrontal cortex, 259
 working memory and executive function, 261–262
- hypocortisolism, bipolar disorder, 184
- hyporeactivity, neural networks, 268
 default mode network, 267
 emotion perception, 263
 prefrontal cortex, 259
- hypothalamic–pituitary–adrenal (HPA) axis, 179, 181, 188
 biological mechanisms of MDD development, 5
 bipolar disorder, 184–185, **186**
 cognition, 180, 185, **186**
 cytokine model, 162, 165
 future directions for study, 187
 homeostasis, **180**, 182
 insulin resistance, 210, 211, 213
 measures of activity, 180
 neurobiological effects, 187
 progressive cognitive deficits, 117, 162
 receptors. *see* glucocorticoid receptor; mineralocorticoid receptor
 social exclusion, 86, **87**
 stress response, 153, 179, 182, 183
- hypoxia, erythropoietin, 290
- hypoxia inducible factor (HIF), 290
- IAPS (International Affective Picture System), 297
- IGT (Iowa Gambling Task), **49–50**, 50, 51
- IMDCP (International Mood Disorders Collaborative Project), 280
- immune dysregulation, 86, 161, **170**. *see also* inflammation
- incidence. *see* epidemiology
- indications for cognitive assessment, 35–37
- individual-level measures of variation, cognitive dysfunction and change, 20, **21**, **22**, 24–26
- inflammation, 160, 169–170
 amygdala, 165, 168–169
 chronic exercise effects, 330
 emotional regulation, 165, 168, **170**
 HPA axis, 187
 molecular mechanisms, 165
 neurobiology, 161–163
 neurodegenerative effects, 166–168
 pathology of white matter, 195
 staging model, 163–165, **163**, **164**
- infliximab, 162
- inhibition, emotional/behavioral. *see* response inhibition
- insomnia, 39–40, 185, 265
- Insomnia Severity Index, 40
- insula, 86, 263
- insulin resistance, 209–211, 220
 and brain function, 211, 215–217
 cognitive impairment, 214–215
 default mode network, 217–219, **219**, **220**
 etiology, 213–214
 hippocampal volume, 215–217, **217**, **218**
 and major depressive disorder, 211–213
- insulin treatment, 282–283, **284–286**
- insulin-like growth factor 1 (IGF-1), 329, 333
- integrated models, 265
- intelligence quotient. *see* IQ
- interferon, 165, 166–167
- interleukins, 165, 166–167, 168–169
- International Affective Picture System (IAPS), 297
- International Mood Disorders Collaborative Project (IMDCP), 280
- interpersonal interactions, and MDD, 92. *see also* social cognition
- interventions. *see* treatment interventions
- introspection, measures of cognitive dysfunction, 235
- Iowa Gambling Task (IGT), **49–50**, 50, 51
- IQ (intelligence quotient)
 cognitive remediation, 310
 cognitive reserve, 3
 predicting future depression, **9–11**, 71
- irritability, in youth depression, 47

- ketamine, 24, 75–76
ketoconazole, 188
kindling effect, progressive cognitive deficits, 120
- late-onset MDD
biomarkers of illness, 151
neurotoxicity hypothesis, 113
vascular hypothesis, 194–195
see also age of onset
- learning, 1
cytokine effects, 165–166
objective measures, 36
see also cold cognition; memory
- length of the depressive episodes.
see duration of illness
- life satisfaction. *see* quality of life
- limbic cortex
anatomy, 63
insulin resistance, 209
progressive cognitive deficits, 116–117
- lipids, blood, 210, 213, 214–215
- lisdexamfetamine, 284–285
- Longitudinal Interval Follow-up
Evaluation–Range of Impaired Functioning Tool (LIFE–RIFT), 136, 138–139
- longitudinal studies
biomarkers of illness, 150–151
cognitive dysfunction in MDD, 242
functional outcomes, 129
neurotoxicity hypothesis, 114–115
pediatric depression, 54–55
predicting future depression, 2–3, 8–11
white matter integrity, 65
- long-term potentiation (LTP), 289
chronic exercise effects, 329–330
insulin resistance, 211
cytokine effects, 166
see also brain-derived neurotrophic factor
- low positive affect, 83. *see also* negative cognitive bias
- magnetic resonance imaging (MRI), 61, 116, 147–148
- maintenance treatment, youth depression, 47–48
- major depressive disorder (MDD)
age of onset, 1
cognitive dysfunction in, 243, 244
DSM criteria, 15
incidence and prevalence, 1, 15
interpersonal interactions, 92.
see also social cognition
protective factors, 3
- maladaptive strategies, pruning, 308.
see also cognitive remediation
- maladaptive thought processes, 154.
see also negative cognitive bias
managing symptoms
- primary care, 30–32, 41–43
progressive cognitive deficits, 118–120
see also treatment interventions
- MAPK (mitogen-activated protein kinase) pathway, 169
- Massachusetts General Hospital Cognitive and Physical Functioning Questionnaire, 35, 247–248, 251–252
- MATRICES Consensus Cognitive Battery (MCCB), 237
- McGill Pain Questionnaire, 38–39
- MDD. *see* major depressive disorder
- measurement of cognitive dysfunction and change, 15–16, 17, 70, 229, 239–240
cognitive deficits in MDD, 232–233, 234
cognitive domains, 232
computerized, 126
data analysis, 237–239
FOCUS study, 238
functional outcomes, 135
indications, primary care, 35
individual-level measures of variation, 20–22, 21, 25, 26
practice effects, 24–25
predictive validity, 15, 17
primary care, 32–35, 33
repeated assessment, 235–237
responses to treatment interventions, 22–24
screening, 233–235
social cognition, 102–106
socially relevant benchmarks, 17–20, 17, 18
subjective and objective ratings, 245–247, 246
- medial prefrontal cortex (MPFC)
rumination, 202–203
self-focus, 84–85
- medical comorbidity. *see* comorbidity
- Medical Outcomes Study, 127, 250
- medication, 278, 281
anatomy of brain, 63–64
benzodiazepines, 40, 118
cognitive remediation interactions, 318
confounding factors, 62–63, 117–118
cortisol lowering, 182–183
efficacy prediction, 257
novel, 75, 76. *see also* erythropoietin
progressive cognitive deficits, 117
psychostimulants, 280–281, 283.
see also modafinil
see also antidepressant medication; *specific drugs by name*
- meditation, mindfulness-based, 283
- memory, 1, 6
- bias, 202–203, 265, 268–269.
see also negative cognitive bias
clinician–patient relationship, 42
cytokine effects, 166
erythropoietin, 290, 291–293, 294–295, 298, 299, 301
and functional outcomes, 129–134, 135
HPA axis stress response, 180–182
insulin resistance, 209, 211
measures, 231, 231–232
negative bias, 32
neural circuits, 200–201, 202
pediatric depression, 55
predictive validity, 17
spatial, 147
see also cold cognition;
declarative memory; episodic memory; working memory
- mental processing speed. *see* processing speed
- mentalizing network, 82
- meta-analyses, 15, 20, 23
age of illness onset, 113
anatomy of brain, 63
biomarkers of illness, 148
cytokine model, 167
emotion perception, 263–264
exercise, 323–327, 331–334
eye-tracking, 83
neural circuits, 198
novel/fast-acting medication, 281
post-recovery deficits, 111
salience network, 268
working memory and executive function, 261
- metabolic syndrome, 210. *see also* insulin resistance
- metacognition, 308, 309
- methylphenidate, 75
- mirtazapine, 182–183
- microglia, 61–62, 167
- mifepristone, 185, 188
- mild cognitive impairment, 120, 235
- mindfulness-based meditation processes, 283
- mineralocorticoid receptor (MR), 179–180, 183, 184–185
- Mini Mental State Exam (MMSE), 2, 8–11, 35, 233, 244
- MiniCog, 35
- mirror neural system (MNS), 82
- mirtazapine, 213, 276, 284–285
- mitogen-activated protein kinase (MAPK) pathway, 169
- modafinil (Provigil), 75, 280–281, 283
- molecular mechanisms, inflammation, 165
- monitoring of cognitive processes (metacognition), 308, 309
- monitoring technology, biomarkers of illness, 74

- monoamines, HPA axis, 187
- Montreal Cognitive Assessment (MoCA), 35, 233
- motivational issues, cognitive remediation, 317–318
- MR. *see* mineralocorticoid receptor
- MRI. *see* magnetic resonance imaging
- Multidimensional Scale of Independent Functioning (MSIF), 128
- multi-modal approaches, 40–41, 75, 76. *see also* treatment interventions
- multiple sclerosis, 34, 201, 245, 293, 301
- National Adult Reading Test (NART), 234–235
- National Comorbidity Survey Replication (NCS-R), 127
- n-back test, 36, 276
- NEAR (Neuropsychological and Educational Approach to Remediation), 310
- near transfer, cognitive remediation, 266
- needs, fundamental, 86
- negative automatic thoughts, 154
- negative cognitive bias, 264
 - biological mechanisms of MDD development, 5–6
 - biomarkers of illness, 74, 149, 153
 - clinician–patient relationship, 41, 42
 - cognitive model of depression, 258
 - cognitive remediation, 308, 315–316
 - emotion perception, 262–263 and emotional bias, 1, 83–84, 202, 264
 - erythropoietin, 294
 - hot cognition, 72
 - memory, 202, 265, 268–269
 - neural circuits, 196–197, 203 and self-focus, 84 *see also* catastrophizing
- negative feedback, catastrophic response to, 72
- neural circuits (white matter tracts), 198
 - attention, 199, 201–202
 - executive function, 199–200
 - hot vs. cold cognition, 70
 - memory, 200–201, 202
 - negative cognitive bias, 203
 - processing speed, 198–199
- neural connectivity
 - antecedents to illness, 112
 - biomarkers of illness, 153
 - exercise effects on cognition, 329–331
 - psychotherapy, 120
 - stress effects, 153
- neural modulation, 266
- neural networks. *see* cognitive control network; default mode network; emotion network; hyperreactivity; hyporeactivity; neural connectivity
- neuroanatomy, 62–63, 63–64, 66, 116–117
- neurobiology of depression, 81, 161–163, 187
- neurocognitive model of depression, 73
- neurodegenerative disorders, 65, 166–167, 214
- neurodevelopment, and erythropoietin, 290
- neurodevelopmental abnormalities
 - antecedents to illness, 3–5
 - biological mechanisms of MDD development, 5–6
 - biomarkers of illness, 152
 - predicting future depression, 8–11
 - white matter integrity, 64–65
- neuroendocrine malfunction, 161–163
- neurogenesis
 - cytokine models, 166
 - erythropoietin, 290
 - HPA axis, 187
 - insulin treatment, 282–283
- neuroimaging
 - antecedents to illness, 110
 - biomarkers of illness, 146, 147–148
 - post-recovery deficits, 111
 - social and emotional processing, 83
 - white matter neurobiology, 196 *see also specific modalities*
- neuromodulatory treatment
 - interventions, 274. *see also* neurostimulation
- neuroplasticity
 - antecedents to illness, 112
 - cytokine model, 161–163, 166
 - erythropoietin, 289
 - exercise effects, 329–331
 - inducing, 306, 307. *see also* cognitive remediation
 - insulin treatment, 282–283
 - progressive cognitive deficits, 121 *see also* brain-derived neurotrophic factor; cognitive flexibility
- neuroprogression models, 52, 56, 162. *see also* neurotoxicity hypothesis; progressive cognitive deficits
- neuroprotection, 165, 166–167
 - erythropoietin, 289, 290, 292
 - glial cells, 167–168
- neuropsychological assessment
 - as gold standard, 244
 - snapshot assessments, 245
- Neuropsychological and Educational Approach to Remediation (NEAR), 310
- neuropsychological model of depression, 149, 150
- Neuropsychological Test Battery (NTB), 238
- neuroscience of functional outcomes, 260, 268–269
 - attention and processing speed, 260
 - emotion distortion and biases, 264
 - emotion network, 258, 267–268
 - emotion perception, 264
 - future perspectives, 269
 - intervention targets, 268
 - saliency network, 258, 268
 - working memory and executive function, 260–261 *see also* cognitive control network; default mode network
- neurostimulation, 281. *see also* deep brain stimulation; neuromodulatory treatment interventions; repetitive transcranial magnetic stimulation; transcranial direct current stimulation
- neurotoxicity hypothesis, 110, 118, 119
 - biomarkers of illness, 151
 - longitudinal studies, 114–115 *see also* neuroprogression models; number of lifetime depressive episodes
- neurotransmitter uptake, cellular anatomy, 61–62
- neurotrophic factors, 289. *see also* brain-derived neurotrophic factor
- nitrosative stress, 161–163, 293
- NMDA (*N*-methyl-D-aspartate) receptor antagonists, 75, 290
- norepinephrine (noradrenaline)
 - exercise effects, 329–331
 - hippocampal formation, 119
 - insulin resistance, 210
- novel medication, 75. *see also* erythropoietin
- NR3C1 (glucocorticoid receptor gene), 184, 185
- NTB (Neuropsychological Test Battery), 238
- nucleus accumbens, 54–55
- number of lifetime depressive episodes
 - biomarkers of illness, 152
 - clinical outcomes, 73–74, 140
 - exercise effects, 330–331
 - neurotoxicity hypothesis, 114–115, 118
 - post-recovery deficits, 111 *see also* duration of illness

- obesity, 210. *see also* insulin resistance
- objective cognitive measures, 233, 235
- primary care, **36**
- vs. subjective measures, 244–245
- occupational functioning, *see* employment status
- off-line social processes, social cognition, 82
- oligodendrocytes, 61, 195
- on-line social processes, social cognition, 82
- orbitofrontal cortex (OFC), 111, 117, 263
- other processing, theory of mind, 85
- outcomes, clinical, 73, 139. *see also* functional outcomes and recovery
- over-general memories, 265, 268–269
- oxidative stress, 161, 293
- oxytocin, biomarkers of illness, 88
- Paced Auditory Serial Addition Test, 309
- pain comorbidity, 39
- Pain Disability Index, 38
- parental depression, children and adolescents, 55
- paroxetine, 213, 277
- patient education, 32, 42–43
- patient engagement, cognitive remediation, 317–318
- Patient Health Questionnaire (PHQ-9), 135
- Patient Reported Outcome Measurement Information System (PROMIS), 250
- patient indications for cognitive assessment, 35
- pediatric depression, 47
- cognitive dysfunction studies, **49–50**
- executive function/attention, 48–52
- facial processing, 52–53
- future directions for study, 56
- memory, 55
- parental depression, 55
- processing speed, 53–54
- reward processing, 54–55
- social exclusion, 88
- treatment interventions, 47–48, 56 *see also* educational functioning
- Perceived Deficits Questionnaire (PDQ), 239, **243**, 248–250, **249**, 252–253
- occupational functioning, 135
- primary care use, 30, 32, **33**, 34
- persistent cognitive dysfunction. *see* residual cognitive symptoms
- pessimism. *see* negative cognitive bias
- PFC. *see* prefrontal cortex
- pharmacological treatment. *see* medication
- phase-specific neuroimmune model of clinical depression, **163**. *see also* staging model of inflammation
- Phineas Gage, case of, 199–200
- physical abuse, 5–6. *see also* childhood trauma
- physical activity, 321. *see also* exercise
- PICs. *see* pro-inflammatory cytokines
- planning, mental, 232. *see also* executive function
- polymorphisms, gene biomarkers of illness, 146
- HPA axis stress response, 183–184, 185
- serotonin transporter gene, 74
- white matter pathology, 195–196 *see also* genetics
- positron emission tomography (PET), 63
- post-acute stage of inflammation, 163–165, **163**
- post-recovery deficits. *see* residual cognitive symptoms
- practice effects, assessment tests, 25, 236–237
- precuneus, 52–53
- predictions of future depression, 3
- cognitive dysfunction, 17
- prevention strategies, 6–8 *see also* premorbid cognitive function
- predictions of treatment response, 260, 261, 262
- prednisolone suppression test (PST), 183
- prefrontal cortex (PFC), 60
- anatomy, 62–63
- biological mechanisms of MDD development, 6
- hyper or hyporeactivity, 259
- insomnia comorbidity, 40
- neural circuits, 198, 199
- pediatric depression, 52
- reward processing, 54–55
- social and emotional processing, 83 *see also* dorsolateral PFC; medial PFC; ventrolateral PFC
- premorbid cognitive function, 110
- biomarkers of illness, 147–148, 149
- cold cognitive deficits, 1, **8–11**
- neurotoxicity hypothesis, 118
- progressive cognitive deficits, 112 *see also* predictions of future depression
- prenatal stress, and MDD development, 5–6
- prevalence. *see* epidemiology
- prevention strategies, 6–8
- primary care, 30, 43
- assessment of cognitive impairment, **33**, 37
- comorbidities, 31, 32, 40
- management strategies for impairment, 32
- objective measures, **36**
- treatment interventions/management, **42**, 43
- PRMQ (Prospective and Retrospective Memory Questionnaire), 250
- problem-solving, 52. *see also* executive function
- problem-solving strategies, cognitive remediation, 308
- procedural learning, 180–182
- processing speed, 1
- age of onset, 113
- biomarkers of illness, **147**
- clinician–patient relationship, **42**
- cytokine effects, 166
- erythropoietin, 290, 299
- and functional outcomes, **129–134**, 135, 260
- neural circuits, 198–199
- objective measures, **36**
- pediatric depression, 53–54
- post-recovery deficits, 111 *see also* cold cognition; psychomotor speed
- prodromal phase of dementia, 120, 235
- progressive cognitive deficits, 110, 120–121
- antecedents to illness, 112
- dementia risk, 120
- neurotoxicity hypothesis, **114**, 118
- post-recovery deficits, 110–112 *see also* neuroprogression models
- pro-inflammatory cytokines (PICs), 163. *see also* cytokine models of depression and cognition; inflammation
- PROMIS (Patient Reported Outcome Measurement Information System), 250
- prosody, 92, 93, 102, 106. *see also* social cognition
- Prospective and Retrospective Memory Questionnaire (PRMQ), 250
- protective factors, MDD, 3
- proton MRI, 61
- Provigil (modafinil), 75, 280–281, 283
- pruning maladaptive strategies, cognitive remediation, 308
- pseudospecificity of treatment response, cognitive remediation, 318

- PST (prednisolone suppression test), 183
- psychoanalytic-interactive group therapy, 100
- psychological interventions. *see* psychotherapy
- psychomotor speed
clinician-patient relationship, 42
diagnostic criteria, 15
measures, 230-231
see also processing speed
- psychosocial functioning, 70, 139, 161
attention and processing speed, 260
clinician-patient relationship, 43
emotion perception, 264
functional outcomes, 125, 126-127
pediatric depression, 47, 52
predicting future depression, 1, 4, 8-11
and progressive cognitive deficits, 119
subjective measures, 242, 243
see also employment status; social cognition
- psychostimulants, 281, 283. *see also* modafinil
- psychotherapy, 286
cognitive remediation, 314, 315, 315, 317
cold cognition, 154
efficacy, 257
progressive cognitive deficits, 120
and social cognition, 103
see also cognitive behavioral therapy
- psychotic features in MDD, 100, 103, 118
- QIDS-SR (Quick Inventory of Depressive Symptomatology), 32, 33
- quality of life, 125, 126, 128, 130, 130, 137. *see also* functional outcomes and recovery; psychosocial functioning
- Quality of Life Enjoyment and Satisfaction Questionnaire (Q-LES-Q), 138
- RAVLT. *see* Rey Auditory Verbal Learning Test
- RBANS (Repeatable Battery for the Assessment of Neuropsychological Status), 136
- reappraisal, cognitive, 265
- reassurance seeking, 265
- reboxetine, 277
- recovery. *see* functional outcomes and recovery; remission
- recurrence risk
and cognitive dysfunction, 70
exercise effects, 331
progressive cognitive deficits, 118
see also number of lifetime depressive episodes; risk factors for MDD
- red blood cell mass, erythropoietin effects, 294, 299
- regional cerebral blood flow. *see* cerebral blood flow
- regulation of emotion, 82, 87, 257, 258. *see also* response inhibition
- rejection, social, 81, 86, 87, 197
- relapse rates, 70, 110. *see also* number of lifetime depressive episodes; recurrence risk
- relational functioning, 161. *see also* social cognition
- reliable change index (RCI), 25
- remission
biomarkers of illness, 148
post-recovery deficits, 110, 259.
see also residual cognitive symptoms
- primary care, 31
- working memory, 72
see also functional outcomes and recovery; state vs. trait cognitive dysfunction
- Repeatable Battery for the Assessment of Neuropsychological Status (RBANS), 136
- repeated assessment, measures of
cognitive dysfunction, 235, 236
- repetitive transcranial magnetic stimulation (rTMS), 266
- repression, memory, 200
- Research Domain Criteria (RDoC), 69
- reserve, cognitive, 3. *see also* IQ
- residual cognitive symptoms
biomarkers of illness, 151
HPA axis, 187
neuroscience of functional outcomes, 259
number of lifetime depressive episodes, 113, 114
progressive, 110-112. *see also* neuroprogression models
subjective measures, 243
see also state vs. trait cognitive dysfunction
- RESPECT trial, pain comorbidity, 39
- response inhibition
attention deficit/hyperactivity disorder, 3
measures of cognitive dysfunction and change, 232
objective measures, 36
pediatric depression, 52, 54
see also executive function; regulation of emotion
- response to treatment, 169, 185, 260
- attention and processing speed, 260
emotion distortion and biases, 265
emotion perception, 263
late-onset MDD, 194
number of lifetime depressive episodes, 113
and rumination, 265
working memory and executive function, 262
- resting state network. *see* default mode network
- retardation. *see* psychomotor speed
- reversible/functional changes, 93, 95, 106. *see also* functional outcomes and functional recovery
- reward processing, 54, 72, 84
- Rey Auditory Verbal Learning Test (RAVLT), 36, 135, 138, 238, 277
- Rey Complex Figure Test, 138
- risk factors for MDD
ADHD, 3
antecedents to illness, 113
biomarkers of illness, 149, 153
exercise effects, 331
predicting future depression, 6
progressive cognitive deficits, 118
see also number of lifetime depressive episodes; recurrence risk
- risk taking, and reward, 72
- rumination, 265, 268
biological mechanisms of MDD development, 5
biomarkers of illness, 154
clinician-patient relationship, 41
cognitive remediation, 309
default mode network, 267
emotion distortion and biases, 264
insomnia comorbidity, 40
neural circuits, 202
and self-focus, 84
and social cognition, 100, 103
and suicidal behavior, 265
- salience network (SN), 258, 268
- salivary cortisol levels, 55, 180, 182
- satisfaction with life. *see* quality of life
- scar effect, 153, 259. *see also* neurotoxicity hypothesis
- SCG. *see* subgenual cingulate cortex
- Schedule for Affective Disorders and Schizophrenia for School-Age Children, 9
- schizophrenia
biomarkers of illness, 153
cellular anatomy, 61-62
cognitive dysfunction, 16, 17, 17, 18
cognitive remediation, 266, 306, 319

- erythropoietin, 293, **300**
 social cognition, 92, **96**
 working memory, 71
- scientific brain training program, 312
- screening, 233. *see also* measurement of cognitive dysfunction and change
- screening tools
 predicting future depression, 6
 primary care, 30, 32, **33**
see also specific instruments by name
- SDS. *see* Sheehan Disability Scale
- selective attention, 258, 259. *see also* attention
- selective serotonin reuptake inhibitors. *see* SSRIs
- self-control. *see* response inhibition
- self-efficacy targeting, cognitive remediation, 315
- self-esteem, and social exclusion, 88
- self-focus, 84
- self monitoring (metacognition), 308, 309
- self-reference effect (SRE), word recall, 84
- self-referential appraisal, rumination, 203
- self-report instruments, 126, 235
 functional outcomes, 135
 primary care, 34
 subjective measures, 244
see also specific instruments by name
- SENSO framework, threat of social rejection, 87
- Sequenced Treatment Alternatives to Relieve Depression (STAR*D) study, 147, 243–244
- serotonin
 exercise effects, 328
 hippocampal formation, 119
 HPA axis, 187
 insulin resistance, 210, 211, 213
 serotonin reuptake inhibitors. *see* SNRIs; SSRIs
- serotonin transporter gene, 74, 146, 169
- sertraline, 276, 322
- severity of illness, 118
 biomarkers of illness, 152
 functional outcomes, 126, 127
 and rumination, 265
 and social cognition, 100, 102, **103**
- sexual abuse in childhood, 5.
see also childhood trauma
- SF-12 (12-item Short Form Health Survey), 137
- SGC. *see* subgenual cingulate cortex
- Sheehan Disability Scale (SDS), 31, 34, 127, 242, **243**
- Shipleigh Institute of Living Scale, 4
- short-term memory, 6
- Simple Reaction Time (SRT), **36**, 230, 231
- simulations, cognitive remediation, **315**
- single-nucleotide polymorphisms (SNPs), 146, 195. *see also* polymorphisms
- sleeping problems. *see* insomnia
- smartphones, monitoring technology, 74
- SMILE study (Standard Medical Interventions and Longterm Exercise), 322
- snapshot assessments, 245
- SNRIs (serotonin norepinephrine reuptake inhibitors), 119, **284**
- social cognition, 81, **87**, 92, 102, 106
 case-control studies, **96**, **101**
 definitions, 102
 facial processing, 93
 future directions for study, 106
 methodological issues, 102
 remitted and symptomatic stages, 95, **96**
 and severity of depression, 100, 102, **103**
see also psychosocial functioning
- social engagement, and cognitive dysfunction, 17, **18**, **19**
- social functioning. *see* psychosocial functioning
- social perception, 102
- social processes, definitions, 82
- social rejection, 81, 86, **87**, 197
- Social Skills Performance Assessment (SSPA), 139
- somatic disorders, 182
- spatial memory, **147**
- speed vs. accuracy effects, 236.
see also processing speed
- speed of memory scores, 231
- SSRIs (selective serotonin reuptake inhibitors), **284**
 biomarkers of illness, 151
 insulin resistance, 213
 novel/fast-acting medication, 75
 and progressive cognitive deficits, 119
 response to treatment, 260
 youth depression, 47
- staging model of inflammation, 163
- Standard Medical Interventions and Longterm Exercise (SMILE) study, 322
- STAR*D (Sequenced Treatment Alternatives to Relieve Depression) trial, 147, 243–244
- state vs. trait cognitive dysfunction
 biomarkers, 70, **147**, **150**, 151, 152, 154, 160
 cold cognition, 70, **71**
 endophenotypes, 145, 147
- executive function and attention, 111
- functional outcomes, 140
- post-recovery deficits, 110
- primary care, 40
- progressive cognitive deficits, 110
- reward dysfunction, 54
- social cognition, 95, **101**
see also remission; residual cognitive symptoms
- Sternberg memory store task, 261
- strategy planning, 232. *see also* executive function
- stress
 antecedents to illness, 112
 biological mechanisms of MDD development, 5–6
 biomarkers of illness, 152, 153
 HPA axis response, 179, 182, 185
 neurobiology, 161
 progressive cognitive deficits, 117, 119, 120
 reward processing, 54
- striatum, 196, 263
- Stroop Test, **9**
 antidepressant medication, 276
 exercise, 333
 objective measures, **36**
 pediatric depression, **49**, 51
 predicting future depression, 2
- subgenual cingulate cortex (SCG), 53, 63, 149, 197, 263, 267
- subjective measures, 242
 functional impairment, **243**
 incidence and prevalence of dysfunction, **243**
 vs. objective measures, 244
 psychometrically validated scales, **246**
see also British Columbia Cognitive Complaints Inventory; Massachusetts General Hospital Cognitive and Physical Functioning Questionnaire; Perceived Deficits Questionnaire
- sub-syndromal stage, staging model of inflammation, **163**
- suicidal ideation, 100, **103**, 161, 257, 265
- surveys, occupational functioning, 136
- sustained attention, 258, 259, 299.
see also attention
- Symptom Checklist
 90-revised, 4, **9**
- symptom profiling, 257, 317
- symptomatic and remitted stages.
see state vs. trait cognitive dysfunction

- symptoms, 32
 changes during course of illness, 69
 primary care, 30, 41
 progressive cognitive deficits, 118
 severity. *see* severity of illness
 youth depression, 47
 see also managing symptoms
- synaptic development, cellular
 anatomy, 61
- Syndrome X, 210. *see also* insulin
 resistance
- synergistic effects, catecholamines,
 119
- technology, monitoring, 74
- temporal lobe, 117
- thalamus, 149, 202
- theory of mind, 85, 92, 93, 95. *see*
 also social cognition
- therapeutic relationship, primary
 care, 41
- THINC group, 35, 239
- threat of social rejection (TSR), 87
- three-component model of
 executive function, 6
- TNF α (tumor necrosis factor), 165,
 166, 169
- “to-be-remembered-items” (TBRI),
 231
- top-down processes
 biological mechanisms of MDD
 development, 5
 biomarkers of illness, 154
 cold cognition as, 70, 73
 emotional regulation, 197
- total lifetime illness. *see* duration
 of illness; number of lifetime
 depressive episodes
- Tower of Hanoi task, 232
- Tower of London planning task, 73
- Trail Making Test (TMT), 9, 232, 239
- antidepressant medication, 276
 functional outcomes, 135
 number of lifetime depressive
 episodes, 116
 objective measures, 36
 predicting future depression, 2
- trait deficits. *see* state vs. trait
 cognitive dysfunction
- transcranial direct current
 stimulation (tDCS), 266
- transfer, cognitive remediation,
 307, 308
- transnosographic scar, cognitive
 dysfunction as, 153, 259.
 see also neurotoxicity
 hypothesis
- trauma. *see* childhood trauma;
 stress
- TRD. *see* treatment-resistant
 depression
- treatment interventions, 23, 274,
 283
 cognitive remediation, 282, 284
- compliance. *see* adherence to
 treatment
- dietary factors, 284
- efficacy, 119, 257, 282, 284
- exercise, 281, 284
- functional outcomes, 139, 268
 guiding principles, 275
- holistic approaches, 76
- HPA axis, 187
- insomnia comorbidity, 40
- insulin treatment, 282, 284
- multi-modal approaches, 40
- neuromodulatory, 274
- neurostimulation, 281. *see also*
 deep brain stimulation
- pain comorbidity, 39
- pediatric depression, 47, 56
- pharmacotherapy, 275, 278.
 see also antidepressant
 medication
- primary care, 32, 42
 see also behavioral treatment
 interventions; early
 intervention; erythropoietin;
 managing symptoms;
 multi-modal approaches;
 psychotherapy; response to
 treatment
- treatment-resistant depression
 (TRD)
 cognitive remediation, 312, 313
 erythropoietin, 299, 300
- tricyclic antidepressants, 117, 283
- tryptophan, 328. *see also* serotonin
- TSR (threat of social rejection), 87
- tumor necrosis factor (TNF α), 165,
 166, 169
- 12-item Short Form Health Survey
 (SF-12), 137
- twin studies, 9
 antecedents to illness, 112
 cognitive deficit biomarkers of
 illness, 147
 predicting future depression, 2
 see also genetics
- unipolar depression, HPA axis,
 182–184. *see also* major
 depressive disorder
- University of California San Diego
 Performance-Based Skills
 Assessment (UPSA), 279
- updating of new information, three-
 component model, 6
- vascular dementia, insulin
 resistance, 214
- vascular endothelial growth factor
 (VEGF), 329–330, 333
- vascular hypothesis
 late-onset MDD, 113
- white matter neurobiology,
 194–195
- venlafaxine, 47
- ventrolateral prefrontal cortex
 (VLPFC), 86, 87, 111, 201
- ventromedial frontal cortex, 93,
 263–264
- verbal impairments, 259, 261
 erythropoietin, 294, 297, 299
 and functional outcomes,
 129–134, 138
 pediatric depression, 49–50,
 51, 55
 word recall, 2, 84
- vigilance, biomarkers of illness,
 150–151
- visual memory, biomarkers of
 illness, 147
- vortioxetine, 75, 277–280, 278,
 283, 284
- biomarkers of illness, 151
 cognitive dysfunction and
 change, 23
 cognitive remediation
 interactions, 318
- vulnerability to MDD, 110,
 147–148, 149. *see also*
 premorbid cognitive
 functioning
- WCST. *see* Wisconsin Card Sorting
 Test
- Wechsler Adult Intelligence Scale
 (WAIS), 230, 261
- Wechsler Intelligence Scale for
 Children (WISC), 4, 53
- Wechsler Memory Scale (WMS),
 116, 135, 276
- weight gain, antidepressant
 medication, 213
- white matter hyperintensities
 (WMH), 195
- white matter (WM) neurobiology,
 195, 203
 attention, 199, 201–202
 executive function, 199–200
 genetics, 195–196
 memory, 200–201, 202
 negative cognitive bias, 201–203
 neural circuits, 196–198
 neuroanatomy, 64–65, 66
 neuroimaging, 196
 pathology, 195
 processing speed, 198–199
 vascular depression hypothesis,
 194–195
 see also neural circuits (white
 matter tracts)
- WHO Disability Assessment
 Schedule, 135
- WHO World Health Survey (WHS)
 study, 127
- Wide Range Achievement Test-
 Revised, 4
- Widespread Pain Index, 39
- WISC. *see* Wechsler Intelligence
 Scale for Children

Cambridge University Press

978-1-107-07458-3 - Cognitive Impairment in Major Depressive Disorder: Clinical Relevance,
Biological Substrates and Treatment Opportunities

Edited by Roger S. McIntyre

Index

[More information](#)

- Wisconsin Card Sorting Test (WCST), 148, 232, 276
- WMS (Wechsler Memory Scale), 116, 135, 276
- word recall, 2, 84. *see also* verbal impairments
- Work and Social Adjustment Scale (WSAS), 35
- work functioning. *see* employment status
- Work Limitations Questionnaire (WLQ), 135, 280
- Work Productivity and Activity Impairment (WPAI), 243, 243
- working memory, 1
 cold cognition, 70–72
 definitions, 260–261
 exercise, 325
 functional outcomes, 260–262
 measures, 229–230, 231–232
 novel/fast-acting medication, 75
 number of lifetime depressive episodes, 116
 objective measures, 36
 pediatric depression, 48, 51, 55
 predicting future depression, 3–5, 17
- three-component model of executive function, 6
 treatment interventions, 275–276
see also cold cognition
- World Health Organization, 128, 135, 322
- WSAS (Work and Social Adjustment Scale), 35
- years of education (YoE), 234
- youth depression. *see* pediatric depression