

Index

- AAGA. *See* accidental awareness under general anesthesia
- abdominal pain, liver disease causing, 102
- abnormal placentation. *See also* placenta accreta; placenta previa
- case study and key points, 107
- definitions, 107, 108
- delivery management for, 111
- anesthetic management, 111–12
- interventional radiology and urological management, 111
- delivery planning for, 109–11
- diagnosis of, 108–9, 110
- emergency hysterectomy for (*See* emergency hysterectomy)
- emergency hysterectomy risk and, 193
- epidemiology, risk factors, and pathophysiology of, 107–8
- interventional radiology for (*See* interventional radiology)
- learning points, 112–13
- accidental awareness under general anesthesia (AAGA)
- case study and key points, 156, 157
- classification of, 156–57
- during elective cesarean delivery, 123
- incidence of, 157
- intraoperative management of, 161
- learning points, 161–62
- postoperative management of, 161
- prevention of, 159–61
- psychological sequelae of, 157
- risk factors for, 157–58, 159
- accidental dural puncture (ADP)
- case study and key points, 53
- complications after, 55–56
- incidence of, 53
- learning points, 56
- management of, 54
- intrathecal catheter, 54–55
- second epidural attempt, 54
- post-dural puncture headache after (*See* post-dural puncture headache)
- risk factors for, 53–54
- signs of, 54
- ACHD. *See* adult congenital heart disease
- acidemia, fetal, 59
- classification and management
- system for, 59–61, 62
- heart rate patterns related to, 59, 61
- scalp sampling for detection of, 61–62
- spectrum of, 59, 61
- ACS. *See* acute coronary syndrome
- activated partial thromboplastin time (aPTT), 89
- acupuncture, 36–37
- acute coronary syndrome (ACS), 249.
- See also* myocardial infarction
- acute fatty liver of pregnancy (AFLP), 98, 101
- acute pain, during cesarean delivery, 221
- acute respiratory distress syndrome (ARDS), 268, 269
- extra corporeal membrane oxygenation for, 271
- ventilation for, 270–71
- acute respiratory failure (ARF), 94
- ADP. *See* accidental dural puncture
- adrenaline, 85
- adult congenital heart disease (ACHD)
- cardiac anatomy, physiology, and surgical history in patients with, 82–84
- case study and key points, 81–82
- labor and delivery in patients with, 84–85
- learning points, 85
- advance directives, 52. *See also* birth plans; Do Not Resuscitate orders
- AEDs. *See* antiepileptic drugs
- AFE. *See* amniotic fluid embolism
- AFI. *See* awake fiberoptic intubation
- AFLP. *See* acute fatty liver of pregnancy
- air loss of resistance technique, 11–12, 20
- airway, difficult. *See* difficult airway
- alloimmunization, cell salvage and, 188–89
- Alphanate. *See* factor VIII-vWF concentrates
- alteplase, 250
- American Society of Anesthesiologists (ASA), guidelines for anesthetic management of postpartum tubal sterilizations, 140–41
- amniotic fluid embolism (AFE)
- case study and key points, 206
- cell salvage and, 188
- clinical presentation of, 207, 208
- diagnosis of, 207–9
- incidence of, 207
- learning points, 209
- management of, 209
- outcome of, 209
- pathophysiology of, 207
- risk factors for, 207
- anemia, postpartum hemorrhage risk and, 181
- anesthetic agents. *See also* local anesthetic; *specific agents*
- intraoperative awareness and, 157, 158, 159–60
- resistance to, 157
- safety of, 148, 240, 241
- aneurysm, intracranial, 255
- angina, 249
- antacids, 3–4, 121
- antibiotics
- for elective cesarean delivery, 121–22, 258
- for sepsis, 260–61
- anticholinesterases, 104
- anticoagulation
- anesthesia considerations and, 265–66
- epidural analgesia and, 40
- in parturients with complex adult congenital heart disease, 84–85
- for venous thromboembolism, 264–65, 266
- antiepileptic drugs (AEDs), 105
- anti-RhD, 188–89, 245
- antivirals, for influenza, 269–70
- anxiety, intraoperative awareness and, 157
- aorto-caval compression, 244
- apixaban, 264–65

Index

- appendectomy
 case study and key points, 239
 clinical challenges/controversies of, 239–42
 learning points, 242
 aPTT. *See* activated partial thromboplastin time
 ARDS. *See* acute respiratory distress syndrome
 ARF. *See* acute respiratory failure
 Arnold-Chiari malformation, 255
 arrhythmia
 complex adult congenital heart disease with, 85
 local anesthetic toxicity with, 212, 213–14
 arteriovenous malformation, intracranial, 255
 ASA. *See* American Society of Anesthesiologists
 aspiration
 eating as risk for, 2, 3, 4
 prophylaxis for, 121
 aspirin, 250
 asthma, 95, 96
 atony, uterine. *See* uterine atony
 awake fiberoptic intubation (AFI), 173–74
 awareness, intraoperative. *See* accidental awareness under general anesthesia
 axonotmesis, 231

 backache, after accidental dural puncture, 56
 background infusion, 7–8
 balloon occlusion
 for placental abnormalities, 111
 for post-partum hemorrhage, 204, 205
 benzodiazepines, for seizure, 105–6
 beta-2 agonists, for asthma, 95
 birth plans, 48
 bleeding, post-partum. *See* post-partum hemorrhage
 bleeding disorders. *See* hematological disorders
 blood patch. *See* epidural blood patch
 blood products, refusal of, 111
 blood transfusion
 allogeneic compared with autologous, 186–87
 cell salvage in mothers refusing, 190
 for emergency hysterectomy, 196–97
 in trauma, 244–45
 Body Mass Index (BMI), 76
 bradycardia, fetal. *See* fetal heart rate
 brain aneurysms, 255
 brain injury, maternal fever causing, 16–17

 brain tumors. *See* intracranial lesions
 breach of duty, 217
 breakthrough pain, during labor epidural analgesia, 7
 case study and key points, 10
 learning points, 13
 management of, 12–13, 21–22
 prevention of, 11–12
 rate of occurrence of, 10–11
 risk factors for, 11
 Brice-modified questionnaire, 157
 Bromage motor scale, 122
 bupivacaine
 for elective cesarean delivery, 122
 hypotension occurrence based on dose of, 32
 for maintenance of labor epidural analgesia, 6–7
 volume versus dose of, 22

 cesarean delivery
 antacids prior to, 3–4, 121
 cell salvage for hemorrhage in (*See* cell salvage)
 combined spinal epidural for, 221–22, 223
 elective, 122–23
 hypotension concerns and, 32
 quality of anesthesia concerns and, 32–33
 elective (*See* elective cesarean delivery)
 emergency (*See* emergency delivery)
 epidural top-up for (*See* epidural top-up)
 intracranial lesions and, 253
 ischemic heart disease and, 250–51
 neuraxial anesthesia for
 clonidine for, 222–23
 elective, 122
 hypotension after, 166–69
 pain during and after
 anesthetic options for reducing, 221–24
 case study and key points, 220
 characteristics of, 220–21
 learning points, 224
 predicting women at risk for, 221, 222
 platelet count recommendations for, 91, 121
 sepsis risk and, 258
 spinal anesthesia for, 122, 221–22, 223
 neurologic deficits following (*See* neurologic deficits)
 suspension of DNR orders during, 52
 timing of, 195
 urgency classification for, 129

 caffeine, for post-dural puncture headache, 235–36
 CAM. *See* complementary or alternative medicine
 capacity, consent and, 46–48, 49–52
 carbetocin, 135–36, 137–38
 carboprost
 ischemic heart disease and, 252
 for uterine atony, 136–37
 cardiac arrest, local anesthetic toxicity causing, 213–14
 cardiac disease. *See* adult congenital heart disease; myocardial infarction
 cardiac failure. *See* heart failure
 cardiac output
 in parturients with complex adult congenital heart disease, 84
 pregnancy effects on, 239–40
 cardiomyopathy, peripartum, 169
 cardiocography (CTG)
 case study and key points, 58
 classification system for interpretation and management of heart rate patterns, 59–61, 62
 fetal scalp sampling and, 61–62
 heart rate pattern nomenclature used in, 59, 60
 learning points, 62–63
 for nonobstetric surgery, 241
 relationship between heart rate patterns and acidemia, 59, 61
 cardiovascular system (CVS)
 local anesthetic toxicity of, 212
 pregnancy effects on, 239–40
 catheter placement, 11–12
 anatomic considerations for, 12, 24
 combined spinal epidural versus epidural, 12, 20–21, 30–31
 fixation of catheter after, 21
 loss of resistance technique for, 11–12, 20
 maternal positioning for, 20
 recommended catheter length in epidural space, 21
 type of catheter for, 20
 ultrasound guidance of (*See* ultrasound)
 catheter reliability, in combined spinal epidural versus epidural, 31
 causation, 217
 cell salvage
 allogeneic compared with autologous blood transfusion, 186–87
 benefits of
 in nonobstetric surgery, 188
 in obstetrics, 189

- in placenta accreta, 189–90
- case study and key points, 186
- contraindications to, 189
- controversies regarding
 - alloimmunization, 188–89
 - amniotic fluid contamination, 188
- cost effectiveness of, 190
- for emergency hysterectomy, 197
- LDF problems in clinical practice, 189
- learning points, 190
- mechanism of, 187
- in mothers who refuse blood transfusion, 190
- central nervous system (CNS)
 - infections of, combined spinal epidural and, 31
 - local anesthetic toxicity of, 212
- cerebral palsy, maternal fever and, 16–17
- cerebrospinal fluid (CSF), saline compared with, 54
- cervical cerclage
 - anesthetic considerations for, 148
 - case study and key points, 147
 - indications for, 147–48
 - learning points, 148
 - safety of, 148
- CEWS. *See* Community Early Warning Score
- chain of prevention, 72
- childbirth. *See* labor
- cholestasis, obstetric, 100
- cholinergic crisis, 104
- chronic pain after pregnancy
 - anesthetic options to minimize risk for, 221–22
 - intravenous ketamine, 223–24
 - neuraxial clonidine, 222–23
 - oral gabapentin, 223, 224
 - transversus abdominis plane block, 223, 224
 - case study and key points, 220
 - characteristics of, 220–21
 - learning points, 224
 - predicting women at risk for, 221, 222
- clonidine, 222–23
- clopidogrel, for myocardial infarction, 250
- CNS. *See* central nervous system
- coagulation measurement, in obstetric hemorrhage, 183–84
- coagulopathy. *See also* hematological disorders
 - amniotic fluid embolism with, 207, 208, 209
 - in trauma, 244–45
- cocaine, 173
- codeine, 41
- postoperative, 124
- coloadung, 167
- combined spinal epidural (CSE), 30
 - for cesarean delivery, 221–22, 223
 - elective, 122–23
 - hypotension concerns and, 32
 - quality of anesthesia concerns and, 32–33
 - case study and key points, 30
 - epidural technique versus, 12, 20–21
 - catheter reliability, 31
 - onset time of analgesia, 30–31
 - quality of pain relief, 31
- ischemic heart disease and, 250–51
- in labor
 - advantages of, 30–31
 - potential risks of, 31–32
 - learning points, 33
 - for postpartum tubal sterilizations, 143
- communication, in multidisciplinary team management, 65, 66
- Community Early Warning Score (CEWS), 73
- complementary or alternative medicine (CAM), for childbirth, 35. *See also* non-pharmacologic analgesia
- computed tomography (CT), during pregnancy, 240–41
- computerized tomography pulmonary angiography (CTPA), for pulmonary embolism, 264
- conflict of interest, maternal fetal, 52
- congenital heart disease. *See* adult congenital heart disease
- consent
 - medicolegal issues and, 217
 - UK perspective on
 - advance directives, 52
 - capacity, 49–52
 - case study and key points, 49
 - learning points, 52
 - maternal fetal conflict of interest, 52
 - preoperative discussion of difficult situations, 52
 - suspension of DNR orders during cesarean delivery and under anesthesia, 52
 - US perspective on
 - birth plans, 48
 - capacity, 46–48
 - case study and key points, 46
 - disagreement among woman and her partner, 48
 - learning points, 48
 - risks of neuraxial anesthesia, 48
 - standards of disclosure, 48
 - written consent, 48
- conus medullaris, 229–30
- corticosteroids
 - for asthma, 95
 - for idiopathic thrombocytopenic purpura, 91
 - for preterm labor, 153–54
 - for sepsis in influenza, 271
- CSE. *See* combined spinal epidural
- CSF. *See* cerebrospinal fluid
- CT. *See* computed tomography
- CTG. *See* cardiotocography
- CTPA. *See* computerized tomography pulmonary angiography
- CVS. *See* cardiovascular system
- dabigatran, 264–65
- dalteparin, 264, 265
- DDAVP, 90
- decelerations, in fetal heart rate, 59, 60
 - acidemia associated with, 59–61, 62
- deep vein thrombosis (DVT), 263, 264, 265
- delivery
 - cesarean (*See* cesarean delivery)
 - emergency (*See* emergency delivery)
 - ischemic heart disease and, 250–51
 - location of, 202–3
 - placental abnormalities and, 111
 - anesthetic management of, 111–12
 - interventional radiology and urological management of, 111
 - planning for, 109–11
 - timing of, 195, 270
 - depth of anesthesia (DOA), 158, 160
 - desflurane, uterine relaxation with, 150
 - desmopressin. *See* DDAVP
 - diamorphine, 41, 123–24
 - DIC. *See* disseminated intravascular coagulation
 - diclofenac, 124
 - difficult airway
 - failed intubations and, 171
 - intraoperative awareness and, 157, 160–61
 - management of anticipated
 - awake fiberoptic intubation, 173–74
 - case study and key points, 171–72
 - learning points, 174
 - regional anesthesia considerations, 172–73
 - management of un-anticipated
 - case study and key points, 175
 - guidelines for, 175–78
 - learning points, 178
 - in obese parturients, 76–78, 79
 - disclosure, standards of, 48

Index

- disseminated intravascular coagulation (DIC), amniotic fluid embolism with, 207, 208
- Do Not Resuscitate (DNR) orders, 52
- DOA. *See* depth of anesthesia
- dose, local anesthetic, 22
- drinking during labor. *See* eating during labor
- dural puncture. *See* accidental dural puncture
- duty of care, 217
- DVT. *See* deep vein thrombosis
- dyspnea. *See* respiratory disorders
- early warning scores (EWS), 71
- CEWS, Community Early Warning Score, 73
 - IMEWS, Irish Maternity Early Warning System, 73
 - MEOWS, Modified Early Obstetric warning system, 71–72
 - NEWS, National Early Warning Score, 72–73
- obstetric, 71
- case study and key points, 69–71
 - chain of prevention and, 72
 - current problems with, 71–72
 - future of, 72–73
 - learning points, 73
 - practicalities of, 71
- eating during labor
- benefits of, 2–3
 - case study and key points, 1
 - current recommendations on, 3–4
 - historical recommendations on, 2
 - informing patients on, 4
 - learning points, 5
 - risks of, 3
 - summary, 4
- EBP. *See* epidural blood patch
- echocardiography, transthoracic. *See* transthoracic echocardiography
- eclampsia, 100
- ECMO. *See* extra corporeal membrane oxygenation
- elective cesarean delivery
- case study and key points, 120
 - enhanced recovery after, 125
 - indications and assessment for, 120, 121
- intraoperative anesthesia for
- antibiotic administration, 121–22, 258
 - combined spinal-epidural anesthesia, 122–23
 - general anesthesia, 123
 - neuraxial techniques, 122
 - spinal anesthesia, 122
 - learning points, 125
- neurologic deficits following (*See* neurologic deficits)
- postoperative analgesia for, 123
- neuraxial opioids, 123–24
- nonneuraxial regional techniques, 124
- nonopioid systemic analgesia, 124
- opioid systemic analgesia, 124
- preoperative care for
- aspiration prophylaxis, 121
 - investigations, 120–21
- embolism. *See* amniotic fluid embolism; pulmonary embolism
- embolization
- for placental abnormalities, 111
 - for post-partum hemorrhage, 204
- emergency delivery
- case study and key points, 128
 - established epidural in, 128–29
 - initial assessment of, 129
 - presurgical assessment of, 129–30
 - rapid top-up of, 130–31
 - regional anesthesia in absence of, 131
 - learning points, 131–32
 - after trauma, 245
- emergency hysterectomy, 193, 198
- anesthetic considerations for, 195–96
- case study and key points, 192–93
 - hemorrhage management in, 198
 - learning points, 199
 - management considerations for, 194–95, 196
 - post-operative considerations for, 198
 - reasons and risk factors for, 193–94
 - surgical considerations for, 197–98
 - transfusion considerations for, 196–97
- emotional support, during labor, 38
- enhanced recovery (ER), after elective cesarean delivery, 125
- enoxaparin
- epidural analgesia and, 40
 - for venous thromboembolism, 264, 265
- Entonox, 40
- ephedrine, for spinal hypotension, 167
- Epidrum, 20
- epidural anesthesia. *See also* combined spinal epidural; epidural top-up
- benefits and risks of, 163
 - in emergency delivery, 128–29
 - initial assessment of, 129
 - presurgical assessment of, 129–30
 - rapid top-up of, 130–31
 - regional anesthesia in absence of, 131
- inadequate
- case study and key points, 163
 - learning points, 165
 - management of, 164–65
 - risk factors for, 163–64
- intracranial lesions and, 255
- ischemic heart disease and, 250–51
- for postpartum tubal sterilizations, 141–43
- in preterm labor, 153
- epidural blood patch (EBP), 55–56, 235, 236–37
- epidural hematoma, thrombocytopenia and, 92
- epidural labor analgesia
- breakthrough pain during (*See* breakthrough pain)
 - catheter placement for (*See* catheter placement)
 - combined spinal epidural in (*See* combined spinal epidural)
 - contraindications to, 40
 - fever associated with
 - case study and key points, 15
 - discussion, 15–17
 - learning points, 17
 - maintenance of
 - breakthrough pain during (*See* breakthrough pain)
 - bupivacaine versus ropivacaine for, 6–7
 - case study and key points, 6
 - learning points, 9
 - local anesthetic solutions for, 6
 - lower extremity motor block during, 7
 - novel concepts in, 7–8
 - patient controlled epidural analgesia for, 7
 - programmed intermittent epidural bolus for, 8
 - variable-frequency automated mandatory boluses for, 8–9
 - medicolegal issues involving (*See* medicolegal issues)
 - multiple sclerosis and, 103–4
 - myasthenia gravis and, 104–5
 - in obese parturients, 76, 78
 - in parturients with complex adult congenital heart disease, 85
 - procedure tips and tricks for
 - case study and key points, 19
 - catheter length in epidural space, 21
 - catheter type, 20
 - combined spinal epidural versus epidural technique, 20–21

- evaluating inadequate pain relief/
unilateral block/patchy block,
21–22
- fixation of catheter, 21
- learning points, 22
- loss of resistance technique, 20
- maternal positioning, 20
- predicting suboptimal or failed
epidurals, 19–20, 25, 28
- ultrasound guidance, 22
- volume versus dose, 22
- risks of, 48
- epidural top-up
benefits and risks of, 163
- for emergency delivery, 130–31
- failure of
case study and key points, 163
- learning points, 165
- management after delivery,
164–65
- management before surgery, 164
- management during surgery,
164–65
- risk factors for, 163–64
- medicolegal issues involving, 217
- epilepsy, 105–6
- epinephrine
for epidural top-up, 130–31
- for spinal hypotension, 168
- Episure, 20
- ER. *See* enhanced recovery
- ergometrine, ischemic heart disease
and, 252
- ergonovine maleate, for uterine atony,
136, 137
- ergot alkaloids, for uterine atony, 136,
137
- ESBLs. *See* Extended-Spectrum Beta-
Lactamases
- EWS. *See* early warning scores
- Extended-Spectrum Beta-Lactamases
(ESBLs), 259
- extra corporeal membrane
oxygenation (ECMO), 271
- factor VIII-vWF concentrates
(Humate P, Alphanate), 90
- fasting during labor. *See* eating during
labor
- fatty liver. *See* acute fatty liver of
pregnancy
- feeding during labor. *See* eating during
labor
- fentanyl
for cesarean delivery, 123–24
- for epidural top-up, 130–31
- for maintenance of labor epidural
analgesia, 6
- fetal distress, emergency delivery for.
See emergency delivery
- fetal heart rate (FHR)
combined spinal epidural and,
31–32
- decelerations of, 59, 60
- acidemia associated with (*See*
acidemia)
- intrauterine resuscitation of, 129
- fetal heart rate (FHR) monitoring
case study and key points, 58
- classification system for
interpretation and
management of heart rate
patterns, 59–61, 62
- fetal scalp sampling and, 61–62
- heart rate pattern nomenclature
used in, 59, 60
- learning points, 62–63
- for nonobstetric surgery, 241
- relationship between heart rate
patterns and acidemia, 59, 61
- spectrum of acidemia, 59, 61
- fetal heart rate variability (FHRV), 59,
60
- acidemia associated with (*See*
acidemia)
- fetal resuscitation, intrauterine, 129
- fetal scalp sampling, 61–62
- fetus
legal personality of, 48
- maternal conflict of interest with, 52
- fever. *See* intrapartum fever
- FHR. *See* fetal heart rate
- FHRV. *See* fetal heart rate variability
- FIBTEM, 184
- fixation, catheter, 21
- flu. *See* influenza
- fluid resuscitation
for post-partum hemorrhage, 183
- for sepsis, 260
- Fontan procedure, pregnancy in
patients with. *See* adult
congenital heart disease
- functional residual capacity (FRC),
pregnancy effects on, 94–95,
239–40
- gabapentin, 223, 224
- gallstones, 99
- gastric emptying, during labor, 3–4
- gastrointestinal system, pregnancy
effects on, 240
- general anesthesia
accidental awareness under (*See*
accidental awareness under
general anesthesia)
- airway management in (*See* difficult
airway)
- for cervical cerclage, 148
- complex adult congenital heart
disease and, 85
- for elective cesarean delivery, 123
- for emergency hysterectomy, 195
- intracranial lesions and, 255
- ischemic heart disease and, 250–51
- for manual removal of placenta, 150
- multiple sclerosis and, 104
- myasthenia gravis and, 105
- for nonobstetric surgery in
pregnancy, 241
- obesity and, 76–78, 79
- placental abnormalities and, 111–12
- for postpartum tubal sterilizations,
141, 143–44
- in preterm labor, 153
- respiratory disorders and, 96
- venous thromboembolism and,
265–66
- gestational thrombocytopenia, 90–92
- glycopyrrolate, 173
- Group A Streptococcus, 259
- H1N1 virus. *See* influenza
- hemoglobin measurement, 183
- halothane, uterine relaxation with, 150
- harm, 217
- headache. *See* postdural puncture
headache
- heart disease. *See* adult congenital
heart disease; myocardial
infarction
- heart failure
preeclampsia and, 115, 167–69
- transthoracic echocardiography in
management of (*See*
transthoracic
echocardiography)
- heart rate, fetal. *See* fetal heart rate
- HELLP syndrome. *See* hemolysis,
elevated liver enzymes and low
platelet count syndrome
- hematological disorders
case study and key points, 87
- learning points, 92
- post-partum hemorrhage and, 182
- thrombocytopenia, 90–92
- von Willebrand disease
hemostasis and, 88–89
- obstetric anesthesia implications
of, 89–90
- hematoma, epidural. *See* epidural
hematoma
- hemolysis, elevated liver enzymes and
low platelet count (HELLP)
syndrome
liver disease with, 100–1
- thrombocytopenia with, 91–92
- hemorrhage
post-partum (*See* post-partum
hemorrhage)
- in trauma, 244–45

Index

- hemostasis, in patients with
 hematological disorders, 88–89
- heparin. *See* low molecular weight heparin
- hepatitis, 99
- hepatitis-like features, of liver disease in pregnancy, 101–2
- HFOV. *See* high-frequency oscillation ventilation
- high spinal, 55
- high-frequency oscillation ventilation (HFOV), 271
- high-reliability organizations (HROs), 65
- histamine-2 receptor antagonists, pre-operative, 3–4, 121
- HIV/AIDS, 259
- HROs. *See* high-reliability organizations
- Humate P. *See* factor VIII-vWF concentrates
- hydralazine, for preeclampsia, 115
- hydration therapy, for post-dural puncture headache, 56
- hydrotherapy, 35–36
- hyperemesis gravidarum, 99–100
- hypertensive disorders of pregnancy
 eclampsia, 100
 preeclampsia (*See* preeclampsia)
- hypnosis, 37
- hypotension
 amniotic fluid embolism with, 207, 208
 spinal (*See* spinal hypotension)
 in trauma, 244
- hypoxia
 amniotic fluid embolism with, 207, 208, 209
 fetal, detection of, 59 (*See also* fetal heart rate monitoring)
- hysterectomy
 emergency (*See* emergency hysterectomy)
 for placental abnormalities, 111
- ICP. *See* intracranial pressure;
 intrahepatic cholestasis of pregnancy
- idiopathic thrombocytopenic purpura (ITP), 90–92
- IMEWS. *See* Irish Maternity Early Warning System
- induction agents
 intraoperative awareness and, 158, 159–60
 safety of, 148, 241
- infection. *See also* sepsis
 central nervous system, 31
 control guidance for, 270
 influenza (*See* influenza)
- influenza
 case study and key points, 268
 clinical challenges of, 268–69
 diagnosis of, 269
 extra corporeal membrane oxygenation for, 271
 high-frequency oscillation ventilation for, 271
 infection control guidance for, 270
 intensive care management for, 270–71
 learning points, 272
 management of, 269
 maternal outcomes of, 271–72
 perinatal outcomes of, 272
 pneumonia with, 95–96, 258–59
 presentation of, 269
 sepsis management in, 271
 thromboembolic prophylaxis for, 271
 timing of delivery in, 270
 treatment of, 269–70
- informed consent. *See* consent
- inotropes
 in parturients with complex adult congenital heart disease, 85
 for spinal hypotension, 168
- Intensive Therapy Unit (ITU),
 emergency hysterectomy recovery in, 198
- intermittent bolus
 for maintenance of labor epidural analgesia, 7–9
 systemic opioid administration with, 41
- interventional radiology (IR)
 case study and key points, 202
 delivery location for patients at high risk of hemorrhage, 202–3
 emergency hysterectomy and, 198
 after failure of other hemorrhage treatments, 204, 205
 hemorrhage management in units without, 203–4
 learning points, 205
 patient transfer to units with
 anesthetic considerations in, 204–5
 cardiovascular stability considerations in, 203
 for placental attachment disorder, 111
- intracranial lesions
 case study and key points, 253
 general anesthesia in, 255
 learning points, 256
 neuraxial anesthesia in, 253
 intracranial pressure considerations, 254–55
 lesion size, location, and growth rate considerations, 253–54
 preferred technique for, 255
 vascular pathology, 255
- intracranial pressure (ICP), 254–55
- intradural space, catheter placement in, 12
- intrahepatic cholestasis of pregnancy (ICP), 100
- Intralipid, 211
- intraoperative awareness. *See*
 accidental awareness under general anesthesia
- intraoperative cell salvage (IOCS), 187.
See also cell salvage
- intrapartum fever, epidural-associated
 case study and key points, 15
 discussion, 15–17
 learning points, 17
- intrathecal catheter (ITC)
 after accidental dural puncture, 54–55
 for post-dural puncture headache, 56
- intrauterine fetal resuscitation, 129
- intravenous induction agents
 intraoperative awareness and, 158, 159–60
 safety of, 148, 241
- intubation, failed, 171. *See also* difficult airway
- IOCS. *See* intraoperative cell salvage
- ipratropium bromide, for asthma, 95
- IR. *See* interventional radiology
- Irish Maternity Early Warning System (IMEWS), 73
- ischemic heart disease. *See* myocardial infarction
- ITC. *See* intrathecal catheter
- ITP. *See* idiopathic thrombocytopenic purpura
- ITU. *See* Intensive Therapy Unit
- IV IgG, for idiopathic thrombocytopenic purpura, 91
- jaundice, 102
- Jehovah's witnesses, 50
- ketamine, 223–24
- KFS. *See* Klippel-Feil Syndrome
- Kirkpatrick, Donald, model for training evaluation, 67
- Klippel-Feil Syndrome (KFS), airway management in
 awake fiberoptic intubation, 173–74
 case study and key points, 171–72
 learning points, 174
 regional anesthesia considerations, 172–73

- LA. *See* local anesthetic
- labor
- eating and drinking in
 - benefits of, 2–3
 - case study and key points, 1
 - current recommendations on, 3–4
 - historical recommendations on, 2
 - informing patients on, 4
 - learning points, 5
 - risks of, 3
 - summary, 4
 - epidural analgesia during (*See* epidural labor analgesia)
 - ischemic heart disease and, 250–51
 - maternal capacity during, 46–48
 - nonpharmacologic analgesia for (*See* nonpharmacologic analgesia)
 - in patients with complex adult congenital heart disease, 84–85
 - preterm (*See* preterm labor)
 - laparoscopic surgery, 241
 - LAST. *See* local anesthetic toxicity
 - Lasting Power of Attorney (LPA), 48
 - lateral position, for epidural procedures, 20
 - LCHAD deficiency. *See* long-chain 3-hydroxyacyl-coenzyme A dehydrogenase deficiency
 - LDF. *See* leucocyte depletion filter
 - legal claims. *See* medicolegal issues
 - leucocyte depletion filter (LDF), 188, 189
 - Levobupivacaine, for epidural top-up, 130–31
 - lidocaine
 - for epidural top-up, 130–31
 - intrathecal, 55
 - ligation, for post-partum hemorrhage, 204
 - lipid emulsion, for local anesthetic toxicity, 211, 214
 - liver disorders
 - anesthesia and, 102
 - case study and key points, 98
 - clinical challenges/controversies with, 98–99
 - disease specific to pregnancy, 99–102
 - learning points, 102
 - unrelated or coincidental disease, 99
 - LMWH. *See* low molecular weight heparin
 - local anesthetic (LA)
 - consumption of, in combined spinal epidural versus epidural, 31
 - hypotension occurrence based on dose of, 32
 - mechanism of action of, 211–12
 - safety of, 148
 - solutions for maintaining labor epidural analgesia, 6
 - volume versus dose of, 22
 - local anesthetic toxicity (LAST), 211
 - case study and key points, 211
 - features of, 212
 - cardiovascular system toxicity, 212
 - central nervous system toxicity, 212
 - learning points, 214–15
 - mechanism of action of, 211–12
 - in obstetrics, 213
 - prevention and detection of, 213
 - treatment of, 213–14
 - LockIt Plus device, 21
 - long-chain 3-hydroxyacyl-coenzyme A dehydrogenase (LCHAD) deficiency, 101
 - loss of resistance (LOR) technique, 11–12, 20
 - low dose epidural solutions, 6
 - low molecular weight heparin (LMWH)
 - anesthesia considerations in parturient on, 265–66
 - epidural analgesia and, 40
 - for myocardial infarction, 250
 - in parturients with complex adult congenital heart disease, 84–85
 - for venous thromboembolism, 264–65, 266
 - lower extremity motor block, during labor epidural analgesia, 7
 - LPA. *See* Lasting Power of Attorney
 - lumbar spine surgery. *See* spine surgery
 - Mackenzie technique, 174
 - MAD. *See* Mucosal Atomising Device
 - magnesium sulfate
 - for preeclampsia, 115, 116, 118
 - for preterm labor, 153–54
 - magnetic resonance imaging (MRI) of placental abnormalities, 108–9
 - during pregnancy, 241
 - major obstetric hemorrhage (MOH)
 - case study and key points, 180–81
 - learning points, 184
 - management of, 182–83
 - point of care testing in, 183–84
 - recognition and causes of, 182
 - manual removal of placenta (MROP), 150
 - anesthetic considerations for, 150
 - case study and key points, 149
 - massive transfusion protocol (MTP), 197
 - mastitis, 258
 - maternal fetal conflict of interest, 52
 - MCA 2005. *See* Mental Capacity Act 2005
 - MDT management. *See* multidisciplinary team management
 - medicolegal issues, claim of negligence
 - considerations, 216–17, 219
 - appropriateness of epidural, 217
 - block assessment before surgery, 218
 - case study and key points, 216
 - consent, 217
 - follow-up and headache management, 218–19
 - intra-operative pain experience and management, 218
 - learning points, 219
 - standard of epidural insertion and subsequent management, 217
 - subsequent events, 219
 - Mental Capacity Act (MCA) 2005, 48
 - MEOWS. *See* Modified Early Obstetric warning system
 - meperidine. *See* pethidine
 - methylergonovine, 136, 137
 - methylprednisolone, epidural-associated fever and, 16
 - metoclopramide, preoperative, 121
 - MG. *See* myasthenia gravis
 - MI. *See* myocardial infarction
 - Michigan Awareness Classification Instrument, 156
 - midazolam, for awake fiberoptic intubation, 173
 - minimum local anesthetic doses (MLAD), 22
 - minimum local anesthetic volumes (MLAV), 22
 - misoprostol
 - ischemic heart disease and, 252
 - for uterine atony, 136–37
 - MLAD. *See* minimum local anesthetic doses
 - MLAV. *See* minimum local anesthetic volumes
 - modified Bromage score, 7
 - Modified Early Obstetric warning system (MEOWS), 71–72
 - MOH. *See* major obstetric hemorrhage
 - monitoring, fetal heart rate. *See* fetal heart rate monitoring
 - morphine, 41
 - for cesarean delivery, 123–24
 - for post-dural puncture headache, 55
 - after postpartum tubal sterilizations, 144
 - motor block, during labor epidural analgesia

Index

- bupivacaine versus ropivacaine in
 causing, 6–7
 lower extremity, 7
- MRI. *See* magnetic resonance imaging
- MROP. *See* manual removal of placenta
- MS. *See* multiple sclerosis
- MTP. *See* massive transfusion protocol
- Mucosal Atomising Device (MAD), 174
- multidisciplinary team (MDT)
 management
 barriers to effective, 66
 case study and key points, 64
 for emergency hysterectomy, 194–95
 in high-reliability organizations, 65
 importance of, 64–65
 keys to effective, 65–66
 learning points, 68
 simulation training in, 66–68
- multi-orifice catheters, 7, 12, 20
- multiple sclerosis (MS), 103–4
- myasthenia gravis (MG), 104–5
- myocardial infarction (MI)
 case study and key points, 247, 248
 diagnosis of, 249
 incidence and outcomes of, 248
 learning points, 252
 management of, 249
 atonic postpartum hemorrhage, 251–52
 labor and delivery, 250–51
 medical, 250
 postnatal care, 252
 pregnancy care, 250
 primary percutaneous coronary intervention, 249–50
 thrombolysis, 250
 pathophysiology of, 249
 risk factors for, 248–49
- naloxone, 41
- National Early Warning Score (NEWS), 72–73
- National Institute of Child Health and Human Development (NICHD), fetal heart rate pattern definitions and classification of, 59, 60, 61
- nausea, 31, 102
- near-side blood testing, 182
- negligence claims. *See* medicolegal issues
- neurapraxia, 231
- neuraxial anesthesia. *See also* epidural anesthesia; spinal anesthesia
 for cesarean delivery
 clonidine for, 222–23
 elective, 122
 hypotension after, 166–69
 for cervical cerclage, 148
 for emergency hysterectomy, 195
 intracranial lesions and, 253
 intracranial pressure
 considerations, 254–55
 lesion size, location, and growth rate considerations, 253–54
 preferred technique for, 255
 for manual removal of placenta, 150
 neurologic deficits following (*See* neurologic deficits)
 for nonobstetric surgery in pregnancy, 241
 for postpartum tubal sterilizations, 141–43
 in preterm labor, 153
 venous thromboembolism and, 265–66
- neuraxial analgesia, 10. *See also*
 epidural labor analgesia
 complex adult congenital heart disease and, 85
 hematological disorders and thrombocytopenia, 90–92
 von Willebrand disease, 89–90
 medicolegal issues involving (*See* medicolegal issues)
 multiple sclerosis and, 103–4
 myasthenia gravis and, 104–5
 obesity and, 76, 78
 opioid, 123–24
 placental abnormalities and, 111–12
 platelet count recommendations for, 121
 respiratory disorders and, 96
 risks of, 48
 neurologic complications, combined
 spinal epidural and, 31
 neurologic deficits, after spinal anesthesia
 case study and key points, 228
 conus medullaris injury, 230
 incidence of, 228–29
 learning points, 232
 management and prognosis of, 230–32
 prevention of, 232
 spinal cord anatomic and clinical considerations in, 229–30
- neurologic disorders
 case study and key points, 103
 epilepsy, 105–6
 learning points, 106
 multiple sclerosis, 103–4
 myasthenia gravis, 104–5
- neuromuscular blocking drugs
 neurological disorders and, 104, 105
 for postpartum tubal sterilizations, 144
- neuropathic pain, 221
- NEWS. *See* National Early Warning Score
- NICHD. *See* National Institute of Child Health and Human Development
- nil by mouth, during labor, 2
- nitroglycerin, uterine relaxation with, 150
- nitrous oxide, 40
 intraoperative awareness and, 158, 159, 160
 safety of, 241
- nonobstetric surgery
 case study and key points, 239
 clinical challenges/controversies of, 239–42
 learning points, 242
- nonopioid analgesics, 40–41, 124
- nonpharmacologic analgesia
 acupuncture and acupressure, 36–37
 adjuvant therapies, 38
 case study and key points, 35
 hypnosis, 37
 learning points, 38
 patient desire for, 35
 sterile water papules, 37
 TENS, 37
 water immersion, 35–36
- nonsteroidal anti-inflammatory drugs (NSAIDs), 41, 124
- OAA. *See* Obstetric Anesthetists' Association
- obesity
 case study and key points, 75
 classification of, 76
 complications associated with, 75–76
 general anesthesia and, 76–78, 79
 inadequate labor epidural analgesia and, 11
 intraoperative awareness and, 157
 learning points, 79
 multisystem effects of, 76, 77
 neuraxial procedures in patients with, 76, 78
 ultrasound-guided placement in, 25, 27–28
- Obstetric Anesthetists' Association (OAA), epidural information card for patients, 48
- obstetric cholestasis, 100
- onset time of analgesia, combined
 spinal epidural versus epidural in, 30–31
- opioids
 intraoperative awareness and, 158, 160
 ischemic heart disease and, 251

- for maintenance of labor epidural analgesia, 6
- for postoperative neuraxial analgesia, 123–24
- after postpartum tubal sterilizations, 144
- safety of, 148, 241
- for systemic analgesia bolus, 41
- patient controlled, 42–43
- postoperative, 124
- oral intake during labor. *See* eating during labor
- Oseltamivir (Tamiflu), 269–70
- oxygenation, fetal, 59. *See also* fetal heart rate monitoring
- oxytocin
- ischemic heart disease and, 251
 - for post-partum hemorrhage, 96
 - for uterine atony, 135–36, 137–38
- PAD. *See* placental attachment disorder
- pain. *See* breakthrough pain; chronic pain after pregnancy
- pandemic flu. *See* influenza
- paracetamol, 40–41, 124
- patchy block, 21–22
- patient
- positioning of, 20
 - satisfaction of, combined spinal epidural versus epidural in, 31
 - as team member, 65
- patient controlled analgesia (PCA), systemic opioid administration with, 42–43
- patient controlled epidural analgesia (PCEA)
- background infusion with, 7–8
 - for maintenance of labor epidural analgesia, 7
 - programmed intermittent epidural bolus with, 8
 - variable-frequency automated mandatory boluses with, 8–9
- PCA. *See* patient controlled analgesia
- PCEA. *See* patient controlled epidural analgesia
- PCI. *See* percutaneous coronary intervention
- PDPH. *See* post-dural puncture headache
- PE. *See* pulmonary embolism
- PEBP. *See* prophylactic epidural blood patch
- percutaneous coronary intervention (PCI), 249–50
- perioperative surgical home. *See* enhanced recovery
- peripartum cardiomyopathy (PPCM), 169
- persistent post-operative pain, after cesarean delivery, 221. *See also* chronic pain after pregnancy
- pethidine (meperidine), 41
- phenylephrine
- complex adult congenital heart disease and, 85
 - for spinal hypotension, 167, 168
- PIEB. *See* programmed intermittent epidural bolus
- placement, catheter. *See* catheter placement
- placenta, retained. *See* retained placenta
- placenta accreta, 107, 108
- cell salvage in, 189–90
 - conservative treatment for, 197–98
 - diagnosis of, 194
 - emergency hysterectomy for (*See* emergency hysterectomy)
 - epidemiology of, 194
 - pathophysiology of, 193–94
 - risk factors for, 194
- placenta previa, 107, 108
- placental attachment disorder (PAD)
- case study and key points, 107
 - definitions, 107, 108
 - delivery management for, 111
 - anesthetic management, 111–12
 - interventional radiology and urological management, 111
 - delivery planning for, 109–11
 - diagnosis of, 108–9, 110
 - epidemiology, risk factors, and pathophysiology of, 107–8
 - learning points, 112–13
- plans, birth, 48
- platelet count
- low, 90–92
 - recommended levels of, 91, 121
- pneumonia, 95–96, 258–59
- point of care (POC) testing, in obstetric hemorrhage, 182–83
- case study and key points, 180–81
 - coagulation measurement, 183–84
 - hemoglobin measurement, 183
 - learning points, 184
- positioning, patient, 20
- post-dural puncture headache (PDPH)
- after accidental dural puncture, 53–54, 55–56
 - case study and key points, 234–35
 - conservative management of, 235–36
 - diagnosis of, 235, 236
 - epidural blood patch for, 55–56, 235, 236–37
 - follow up after, 237
- learning points, 237–38
 - combined spinal epidural and, 31
 - medicolegal issues and, 218–19
- postoperative analgesia
- after elective cesarean delivery, 123
 - neuraxial opioids, 123–24
 - nonneuraxial regional techniques, 124
 - nonopioid systemic analgesia, 124
 - opioid systemic analgesia, 124
 - after postpartum tubal sterilizations, 144
- postoperative anticoagulation, 266
- postoperative pain, after cesarean delivery, 221. *See also* chronic pain after pregnancy
- postpartum hemorrhage (PPH)
- asthma and, 96
 - cell salvage for (*See* cell salvage)
 - emergency hysterectomy for (*See* emergency hysterectomy)
 - interventional radiology for (*See* interventional radiology)
 - ischemic heart disease and, 251–52
 - major (*See* major obstetric hemorrhage)
 - management of, 182–83, 198
 - in parturients with hematological disorders, 89–90
 - placental abnormalities causing, 111
 - recognition and causes of, 182
 - retained placenta causing, 150
 - uterine atony and, 133, 134–82
 - uterotonics for (*See* uterotonics)
- postpartum sterilization
- advantages and disadvantages of, 140, 141
 - anesthetic management of, 140–44
 - case study and key points, 140
 - learning points, 144
- posture therapy, for post-dural puncture headache, 56
- PPCM. *See* peripartum cardiomyopathy
- PPH. *See* post-partum hemorrhage
- preeclampsia
- heart failure in, 167–69
 - liver disease with, 100
 - spinal hypotension in parturient with
 - case study and key points, 166–67
 - learning points, 169
 - management of, 167–69
 - thrombocytopenia with, 91–92
 - transthoracic echocardiography role in (*See* transthoracic echocardiography)
- pregnancy

Index

- consent considerations in (*See* consent)
- infection control guidance for, 270
- physiological changes of
 cardiovascular, 239–40
 gastrointestinal, 240
 hemorrhage and, 244
 intraoperative awareness due to, 158
 liver function, 98–99
 respiratory, 94–95, 239–40
 radiation exposure during, 240–41
 safety of anesthetic drugs in, 148, 240, 241
- preoperative discussion, 52
- preterm labor
 anesthetic management and neonate considerations, 153
 parturient considerations, 153–54
 case study and key points, 152
 definition and complications of, 152
 etiology of, 152–53
 learning points, 154
- programmed intermittent epidural bolus (PIEB), 8
- prophylactic epidural blood patch (PEBP), 55–56
- propofol
 intraoperative awareness and, 158, 159–60
 for postpartum tubal sterilizations, 144
- prostaglandins
 ischemic heart disease and, 252
 for uterine atony, 136–37
- proton pump inhibitors, 3–4, 121
- pruritus, combined spinal epidural and, 31
- pulmonary aspiration. *See* aspiration
- pulmonary disease. *See* respiratory disorders
- pulmonary embolism (PE), 263, 264, 265
- pulmonary function, pregnancy effects on, 94–95
- pulmonary edema, preeclampsia with, 115–17
- radiation exposure, during pregnancy, 240–41
- rapid sequence induction (RSI),
 intraoperative awareness and, 158
- regional anesthesia. *See also* neuraxial anesthesia
 for cesarean delivery, 122
 postoperative, 124
 difficult airway and, 172–73
 for emergency delivery in absence of established epidural, 131
 ischemic heart disease and, 250–51
 for nonobstetric surgery in pregnancy, 241
 venous thromboembolism and, 265–66
- relaxation, uterine, for retained placenta, 150
- Relenza. *See* Zanamivir
- remifentanyl, 42–43
 for awake fiberoptic intubation, 173
- respiratory disorders. *See also* acute respiratory distress syndrome
 acute respiratory failure, 94
 anesthetic considerations in, 96
 asthma, 95, 96
 case study and key points, 94
 learning points, 96
 pneumonia, 95–96, 258–59
 pregnancy effects on respiratory physiology, 94–95
- respiratory system, pregnancy effects on, 94–95, 239–40
- resuscitation
 intrauterine, 129
 of post-partum hemorrhage, 182–83
 of pregnant patients, 52
 of sepsis, 260–61
- retained placenta
 anesthetic considerations for, 150
 case study and key points, 149
 learning points, 150–51
 management of, 150
 risk factors and complications of, 149–50
 third stage of labor and, 149
 uterine relaxation for, 150
- reteplase, for myocardial infarction, 250
- risk, disclosure of, 48
- rivaroxaban, 264–65
- rocuronium, 144
- ropivacaine, 6–7
- Rowland, Melissa, 52
- RSI. *See* rapid sequence induction
- rtPA, for myocardial infarction, 250
- safety climate, 66
- sagittal paramedian approach, to epidural placement, 25
- saline
 cerebrospinal fluid compared with, 54
 for post-dural puncture headache, 56
- saline loss of resistance technique, 11–12, 20
- SAYGO technique, 174
- scalp sampling, fetal, 61–62
- scoliosis
 inadequate labor epidural analgesia and, 11
 ultrasound-guided placement for, 25
- seizure
 epilepsy causing, 105–6
 local anesthetic toxicity causing, 212, 213
- sepsis
 case study and key points, 257
 causes of, 258–59
 clinical presentation and diagnosis of, 259–60
 early warning scores for detecting (*See* early warning scores)
 influenza with, 271
 learning points, 261
 management of, 260–61
 microbiology of, 259
 mortality and morbidity rates of, 257
 risk factors for, 257–58
- sevoflurane
 seizure and, 105
 uterine relaxation with, 150
- Sickle cell disease, 189
- simulation training, 66–68
- sitting position, for epidural procedures, 20
- specialist care management, for emergency hysterectomy, 195, 196
- spinal anesthesia. *See also* combined spinal epidural
 for cesarean delivery, 122, 221–22, 223
 neurologic deficits following (*See* neurologic deficits)
 coloadung with, 167
 for emergency delivery in absence of established epidural, 131
 multiple sclerosis and, 103–4
 for postpartum tubal sterilizations, 143
 in preterm labor, 153
- spinal cord
 anatomic and clinical considerations of, 229–30
 injury of (*See* neurologic deficits)
- spinal hypotension
 case study and key points, 166–67
 combined spinal epidural and, 31, 32
 learning points, 169
 management of, 167–69
- spinal ultrasound. *See* ultrasound spine surgery
 inadequate labor epidural analgesia and, 11
 ultrasound-guided placement for, 25
 standard of care, 217
 block assessment before surgery, 218

- epidural insertion and management, 217
 epidural top-up, 217
 standards of disclosure, 48
 sterile water papules, 37
 sterilization, postpartum. *See* postpartum sterilization
 Streptococcus, 259
 streptokinase, for myocardial infarction, 250
 subarachnoid space, catheter placement in, 12
 subdural space, catheter placement in, 12
 succinylcholine
 neurological disorders and, 104, 105
 for postpartum tubal sterilizations, 144
 sufentanil, 6
 Swansea criteria, 101
 syntocinon, 251
 syntometrine, 137
 systemic pharmacological analgesia
 case study and key points, 40
 indications for, 40
 inhaled nitrous oxide, 40
 learning points, 43
 nonopioid analgesics, 40–41, 124
 opioids
 bolus, 41
 patient controlled, 42–43
 postoperative, 124
 Tamiflu. *See* Oseltamivir
 TAP block. *See* transversus abdominis plane block
 team management. *See* multidisciplinary team management
 TEG. *See* thromboelastography
 TENS. *See* transcutaneous electrical nerve stimulation
 teratogens
 anesthetics, 148, 240, 241
 antiepileptic drugs, 105
 thiopental
 intraoperative awareness and, 158, 159–60
 for postpartum tubal sterilizations, 144
 thrombocytopenia, 90–92
 thromboelastography (TEG), 92, 184
 thromboelastometry, 184
 thromboembolic prophylaxis, for
 influenza, 271
 thromboembolism. *See* venous thromboembolism
 thrombolysis, for myocardial infarction, 250
 tinzaparin, 264
 tocolytic agents, 153–54
 top-up. *See* epidural top-up
 toxicity, local anesthetic. *See* local anesthetic toxicity
 tramadol, 41
 tranexamic acid
 for post-partum hemorrhage, 183, 198
 in trauma, 245
 transcutaneous electrical nerve stimulation (TENS), 37
 transfusion. *See* blood transfusion
 transthoracic echocardiography (TTE)
 case study and key points, 114–15
 key strategies for managing severe preeclampsia, 115–18
 learning points, 118
 transverse midline approach, to
 epidural placement, 25–28
 transversus abdominis plane (TAP) block, 124
 for cesarean delivery, 223, 224
 trauma
 case study and key points, 243
 coagulopathy and, 244–45
 hemorrhage in, 244–45
 incidence and epidemiology of, 243–44
 initial approach to, 244
 learning points, 245
 post-partum hemorrhage and, 182
 pregnancy physiology and, 244
 transport to major trauma center after, 244
 TTE. *See* transthoracic echocardiography
 tubal ligation, postpartum. *See* postpartum sterilization
 tubs, labor, 35–36
 Tuffier's line, 232
 UDCA. *See* ursodeoxycholic acid
 UFH. *See* unfractionated heparin
 ultrasound
 epidural placement using, 11, 22
 advantages of, 24–25
 case study and key points, 24
 conventional placement and, 24, 25
 current controversies and future directions of, 28–29
 learning points, 29
 sagittal paramedian approach to, 25
 technique of, 25–28
 transverse midline approach to, 25–28
 of placental abnormalities, 108–9, 110
 during pregnancy, 241
 unfractionated heparin (UFH), for
 venous thromboembolism, 264–65, 266
 unilateral block, 21–22
 United Kingdom, consent in. *See* consent
 United States, consent in. *See* consent
 urinary tract infection, 258
 urological management, for placental abnormalities, 111
 ursodeoxycholic acid (UDCA), 100
 uterine artery balloons, for placental abnormalities, 111
 uterine artery embolization, 111
 uterine atony
 emergency hysterectomy risk and, 193
 post-partum hemorrhage and, 133, 134–82
 uterotonics for (*See* uterotonics)
 uterine relaxation, for retained placenta, 150
 uterotonics, 133–34
 administration guidelines for, 137
 case study and key points, 133
 current controversies and future directions of, 137–38
 ergot alkaloids, 136, 137
 learning points, 138
 oxytocin and analogues, 135–36, 137–38
 post-partum hemorrhage and, 133, 134
 prostaglandins, 136–37
 vaccine, influenza, 269
 vaginal delivery, platelet count recommendations for, 121
 variability, fetal heart rate. *See* fetal heart rate variability
 variable-frequency automated mandatory boluses, 8–9
 VAS scores, combined spinal epidural versus epidural in, 31
 vascular access, for emergency hysterectomy, 195
 vascular pathology, intracranial, 255
 vasopressors, complex adult congenital heart disease and, 85
 vecuronium, for postpartum tubal sterilizations, 144
 venous thromboembolism (VTE)
 anesthesia considerations in, 265–66
 case study and key points, 263
 diagnosis of, 263–64
 learning points, 266–67
 management of, 264–65
 postoperative anticoagulation for, 266
 risk factors for, 263, 264

Index

- ventilation, for influenza, 270–71
ventilation-perfusion (VQ) scan, 264
ventricular septal defect. *See* adult
congenital heart disease
volatile anesthetic agents
intraoperative awareness and, 158,
159–60
safety of, 148, 241
volume, of local anesthetic, 22
- vomiting, 102
von Willebrand disease (vWD)
hemostasis and, 88–89
obstetric anesthesia implications of,
89–90
VQ scan. *See* ventilation-perfusion
scan
VTE. *See* venous thromboembolism
vWD. *See* von Willebrand disease
- warfarin, 264–65
water immersion, 35–36
whirlpools, labor, 35–36
wire-reinforced catheters, 20
written consent, 48
- x-ray, during pregnancy, 240–41
Zanamivir (Relenza), 269–70