

Index

- 2 × 2 table, 87, 89
- acceleration, 360
- actuarial assumption, 315
- additive bivariate model, 120
- additive linear regression model, 118
- additive model, 119, 141, 150, 177, 178, 180, 181, 186
- additive pairwise interaction, 196
- additive proportional hazard model, 366
- additive regression model, 114, 116, 122
- additivity, 114, 162, 365
- admixed, 458
- age-adjusted rate ratio, 145
- alternative distribution, 225
- analysis of variance, 182
- approximate average rate, 316
- Arbuthnott, 291
- ascertainment, 450
- assortative mating, 482
- attributable risk, 387
- attributable risk percentage, 92
- average survival probability, 341
- Bailey method, 515
- balanced polymorphism, 445, 448
- bandwidth, 42, 43
- Bartlett test of variance, 271, 276, 403
- Bartlett test statistic, 272, 275
- Bayes theorem, 109
- Bernoulli variable, 51, 52, 54, 56
- between group variability, 265, 266, 268
- bilinear regression model, 113
- binomial coefficient, 49, 50
- binomial distribution, 52, 53, 55, 56, 60, 90, 170, 263, 289, 440
- binomial mean value, 53
- binomial probability distribution, 23, 24, 49, 51, 57, 67, 183, 326
- binomial theorem, 498, 499
- binomial variance, 51, 53
- bivariate linear regression model, 112, 123, 124
- Bonferroni inequality, 229
- bootstrap estimate, 204, 207, 225
- boxplot, 227
- calibration, 250, 379
- case-load estimate, 395
- categorical variable, 186
- censored, 311
- central limit theorem, 5–7
- chi-square based measures of association, 81
- chi-square analysis, 134, 137
- childhood leukemia mortality data, 39, 40
- chi-square comparison, 105, 147
- chi-square distribution, 9, 10, 12, 16, 82, 106, 108–110, 130, 135, 138, 142, 144
- collinear, 123
- comparison of the proportions, 89, 90
- complete independence, 190, 193
- computer software, 151
- conditional independence, 191, 194, 195, 201
- conditional likelihood estimation, 346
- confidence band, 19, 20, 21, 23–27, 35, 53, 71, 72, 227
- confounder bias, 121, 122
- contingency coefficient, 70, 82
- Cook distance, 255
- correlation coefficient, 70
- Cox, 346
- Cramer coefficient, 82
- critical value, 4, 5
- crossover, 435
- crude ratio, 160
- cumulative distribution function, 230, 232, 240, 255, 419
- cumulative hazard function, 333
- cumulative normal probability, 5
- cumulative probability distribution, 235, 324
- degrees of freedom, 8, 10
- deMoivre, 3
- deMoivre-Laplace theorem, 53
- direct adjustment, 38
- dose-response relationship, 148
- effective sample size, 288
- Einstein, 52
- Eisenhart, 300
- exact confidence interval, 22
- expected value, 47, 49, 51, 52

- exponential cumulative probability function, 239, 248
- exponential distribution, 242
- exponential survival distribution, 238, 319, 320, 322, 353
- f*-ratio test statistic, 270
- false positive, 108
- family-wise error rate, 229
- family laws, 434
- Fisher, 3, 9, 71, 98
- Fisher exact test, 101
- Fisher information function, 508
- Friedman rank test, 302
- Friedman test statistic, 303
- Galton, 69
- gamma coefficient, 77
- Gauss, 3
- Gauss-Laplacian distribution, 3
- genetic selection, 442
- geometric mean, 283, 286
- geometric probability distribution, 57, 58
- goodness-of-fit, 11, 64, 65, 253, 257
- Graunt, 291, 316, 322
- Greenwood, 330
- Greenwood estimated variance, 331
- Greenwood variance, 330
- Halley, 316
- Hardy-Weinberg equilibrium, 434, 465
- harmonic mean value, 212
- harmonic series, 481
- hazard rate, 339
- hazard ratio, 339, 340, 343
- heterogeneity, 264, 265
- Hodgkins disease mortality, 95
- Hodgkins disease mortality rate, 44
- homogeneity, 145, 146, 148, 162, 169, 264, 265, 267
- hypergeometric probability distribution, 98, 99, 100–102, 338
- identically distributed binary variable, 102, 105
- infant mortality rate, 38
- interaction, 111, 119, 120, 150, 165, 188, 189, 192, 349, 365
- inverse function, 233
- joint independence, 191, 194, 197
- Kaplan-Meier estimate, 324, 326
- kappa statistic, 216
- keno, 100
- Knox analysis table, 400
- Knox method, 402
- Kolmogorov, 241, 242
- Kolmogorov-Smirnov test, 243
- Kruskal-Wallis analysis, 295–297, 302, 303
- least squares estimation, 464
- left-censored data, 283
- Levene test of variance, 273, 276, 387
- life table, 317
- likelihood statistic, 170, 172, 173, 356, 371, 372
- likelihood value, 171, 501
- limit of detection, 284
- linear bivariate regression model, 111
- linear relationship, 70
- log-cumulative hazard function, 362
- log-hazard ratio, 345
- log-likelihood value, 151, 171–174, 176, 189
- log-normal distribution, 278, 281
- log-rank test, 335, 339
- log-rate, 265
- logarithmic transformation, 71
- logistic model, 257
- logistic regression model, 371
- loglinear Poisson model, 141, 187
- loglinear model, 148
- Maclaurin series, 497
- Mann-Whitney, 128, 132, 133
- Mann-Whitney estimation, 81, 130
- Mann-Whitney test, 127, 129, 131, 417
- Mantel-Haenszel odds ratio, 36
- Mantel statistic, 404, 405
- matched pair, 156, 182–184, 287
- maximum likelihood estimate, 505, 510
- McNemar test, 184
- mean fitness, 448
- mean polish, 179, 181
- mean survival time, 318, 321, 327, 331
- measures of association, 92
- median value, 321
- median polish, 188
- median survival time, 334, 335, 360
- median value, 23, 24, 283
- Mendel, 450, 465
- Mendelian inheritance pattern, 441
- method of moments estimation, 51, 104
- Miller, 271
- model prediction, 371
- mortality rate, 33, 34, 39, 309
- mortality rate ratio, 46
- mother-child assortment, 439
- mother-child pair, 440
- multiple correlation coefficient, 117
- multivariable Weibull model, 363
- nested, 189
- net reclassification index (nri), 375
- Newman, 19
- noninformative, 330
- noninformative censoring, 311, 312

- nonparametric confidence interval, 207
- nonparametric measure of association, 77
- normal distribution, 4, 6, 7, 9, 16, 17, 42, 53, 55, 114, 129, 170, 278
- normal distribution approximation, 3, 57, 60, 96, 98, 134
- null distribution, 225
- null hypothesis, 222, 300
- odds ratio, 35–37, 39, 82, 92, 94–96, 190–192, 202
- one-way analysis of variance, 266, 295, 276, 294
- ordinal data, 300
- ordinal variable, 126, 129
- ordinary least squares estimate, 31, 383, 386
- partial correlation coefficient, 117
- Pascal, 50
- Pascal triangle, 50
- Pearson, 3, 9
- Pearson chi-square statistic, 102, 106, 126, 170
- Pearson product-moment correlation coefficient, 103
- percentage decrease, 209, 210
- performance table, 407
- perinatal mortality, 158, 161
- perinatal mortality ratio, 163
- perpendicular least squares, 75, 76, 382, 386
- phi-coefficient, 82
- Poisson, 58
- Poisson additive model, 146, 396
- Poisson distribution, 63, 64, 296
- Poisson loglinear regression, 390
- Poisson probability distribution, 47, 58, 60–62, 142, 263, 264, 313, 402
- population stratification, 458
- positive assortative, 483
- power residue method, 245
- power of the sign test, 291
- product-limit estimated survival probability, 324, 326, 330, 339
- product-moment correlation coefficient, 69, 466
- proportional hazard analysis, 345–346
- proportional reduction in error criterion, 83
- pseudo-random number, 245, 246, 249, 396
- quantile, 282
- quantile-quantile plot, 235, 238
- quasi-independence, 156
- Quetelet, 3
- quick test, 300
- random genetic drift, 473, 474
- randomization response survey technique, 56
- randomization test, 220
- randomized response interview technique, 516
- randomized test statistic, 225
- rate ratio, 34, 35, 143, 144–147, 161, 162
- ratio, 29–31
- ratio estimate, 33
- ratio of rates, 46
- reclassification technique, 372
- recombination, 435
- regression diagnostics, 250
- regression model, 297
- relative hazard ratio, 92, 94, 95, 345, 389, 390
- replicate estimation, 215
- replicate sample, 205
- residual value, 113–115, 179–182, 253
- ridit value, 81, 132
- right censored, 328
- risk ratio, 148
- robust, 271, 297
- Rutherford, 263
- sample covariance, 490
- sample mean value, 3
- sample variance, 3, 260
- sampling with replacement, 204
- saturated model, 175
- Scott, 289
- seed-value, 245
- segregation probability, 450, 453
- segregation ratio, 450
- sensitivity, 108, 380, 407
- shift model, 409
- Siegel-Tukey nonparametric evaluation, 274
- signed rank test, 291
- significance probability, 15
- sign test, 289, 306
- simple linear regression model, 31
- simulated data, 235, 248
- smoothed curve, 46
- smoothing technique, 40
- Somer coefficient, 80, 81
- spatial pattern, 62
- Spearman rank correlation coefficient, 73, 74
- specificity, 108, 407
- standard error, 20
- standard mortality ratio, 39, 40
- standard normal distribution, 4, 5, 9
- statistical transformation, 71
- Stein, 20
- stochastically independent, 88
- structural zero, 106, 154
- Student (Gosset), 352
- Student t-test statistic, 76
- survival probability, 319, 324
- Synder ratio, 441
- t*-distribution, 7
- Taylor series, 496, 500, 516
- test for linear trend, 134
- test of variance, 262, 268
- Bayes, 109
- three-group regression analysis, 297
- three-way table, 189

560

Index

- tricubic relationship, 44
- truncated data, 450, 451
- truncated Poisson distribution, 65, 66
- Tshuprow coefficient, 82
- Tukey, 179, 299, 300
- two-sample comparison, 98
- two-sample rank test, 417
- two-sample test statistic, 139
- uniform probability distribution, 314
- US presidents, 67
- Wahlund model, 471, 472
- Wald test, 511
- Weibull, 352
- Weibull parametric survival model, 352, 360, 366, 471, 472
- Weibull two-sample model, 359
- weighted average, 29, 30, 35, 39, 41, 42, 44, 48, 49, 136, 160, 163, 164, 259, 390, 394, 395
- Wilcoxon, 417
- Wilcoxon rank sum test, 126–128, 131, 132, 291, 293
- Wilcoxon two-sample nonparametric analysis, 297
- Wilcoxon two-sample rank test, 243, 274, 297
- within group variability, 266, 268
- Yule coefficient, 82, 83