

Index

- abnormality, psychological, 6–7
 Aboriginal and Torres Strait Islander populations
 see Indigenous population
 abuse see adult abuse; child abuse
 acceptance and commitment therapy (ACT),
 120–1, 124
 acquired diversity, 10, 34
 acute stress disorders (ASD), 143, 145–7, 148–9
 addictions, behavioural, 229
 See *also* substance-related and addictive
 disorders
 adjustment disorders, 143
 adult abuse, 278
 age–sex structures, of mental health
 professionals, 28, 48–50
 age structures
 of Australian Indigenous population, 17–18
 birth rates and, 23, 42
 birthplace and, 39–41
 employment rates and, 28, 47
 family/household compositions and, 25, 43,
 44
 income and, 48
 of migrant populations, 20–1
 mortality rates and, 24, 43
 of New Zealand's ethnic groups, 37–8, 39
 qualification levels and, 26, 27, 46
 spatial diversity and, 15–16
 suicidal behaviours related to, 292
 agoraphobia, 117, 122
 alarm training, 191
 Alzheimer's disease
 assessment, 246
 prevalence and mortality rates, 245, 248
 research, 252
 treatment and prevention, 247–8
 See *also* neurocognitive disorders (NCDs)
 ancestry (statistics), 21
 anger management, 221–2
 anorexia nervosa (AN)
 clinical description, 175
 obsessive compulsive disorder and, 135
 prevalence, 176–7
 treatment and assessment, 177–9, 183
 antisocial personality disorder, 224
 anxiety disorders
 in Australia, 117–22
 case study, 124
 clinical description, 116–17
 comorbidity with, 118, 122
 in New Zealand, 122–3
 prevalence, 117–20, 122–3
 treatment and assessment, 119–22, 123
 assisted suicide, 293
 attention deficit hyperactivity disorder
 (ADHD)
 diagnosis, 65
 prevalence, 66, 69
 treatment, 68, 70
 attention processes, 161–2
 Australia
 diversity in, 10, 11
 socio-economic characteristics, 26–8
 See *also* demographic factors; socio-
 demographic characteristics
 Australian Imaging, Biomarker and
 Lifestyle (AIBL) Flagship Studying of
 Aging, 245–6
 Australian Schizophrenia Research Bank (ASRB),
 88
 autistic spectrum disorder (ASD)
 case study, 71
 diagnosis, 65
 in the Māori population, 70
 prevalence, 66, 69
 research, 67
 support services, 69
 treatment and assessment, 68–9, 70
 avoidant/restrictive food intake disorder
 (ARFID), 176–7
 Balance NZ, 112
 behavioural addictions, 229
 See *also* substance-related and addictive
 disorders
 behavioural therapy see psychological
 treatment
 binge eating disorder (BED)

- clinical description, 176
- consequences, 177
- prevalence, 176–7
- treatment and assessment, 177–9
- bipolar and related disorders
 - in Australia, 107–9
 - case study, 111–12
 - clinical description, 106
 - comorbidity with, 106, 110
 - diagnostic criteria, 106–7
 - education and research on, 109
 - genetics and, 76
 - history, 76
 - in New Zealand, 109–11
 - prevalence and impact, 106, 107, 109–11
 - research, 108
 - support services, 112
 - treatment and assessment, 108–9, 111
 - in young people, 108
- bipolar I disorder, 107
- bipolar II disorder, 107
- birth rates, 23, 42
- birthplace data, 20–1, 39–44
- Bleuler, Eugen, 76
- body dysmorphic disorder (BDD)
 - aetiology, 131
 - clinical description, 130
 - treatment, 133, 136
- borderline personality disorder (BPD)
 - case study, 262–3
 - treatment, 260, 261, 263
- bowel program, 191
- brief psychotic disorder, 78
- bulimia nervosa (BN)
 - case study, 181
 - clinical description, 175
 - consequences, 177
 - obsessive compulsive disorder and, 135
 - prevalence, 176–7
 - treatment and assessment, 177–9
- Cade, John, 108–9
- cannabis use, psychosis and, 85
- CAPS-5 (diagnostic interview), 146–7
- case studies
 - anxiety disorders, 124
 - bipolar disorders, 111–12
 - compulsory treatment, 306–7
 - conversion disorder, 167
 - depression, 100–1
 - disruptive, impulse-control, and conduct disorders, 224
 - elimination disorders, 196
 - feeding and eating disorders, 181
 - hoarding disorder, 137–8
 - neurocognitive disorders, 250–1
 - neurodevelopmental disorders, 71
 - paraphilic disorders, 271–2
 - personality disorders, 262–3
 - posttraumatic stress disorder, 149–51
 - schizophrenia, 87–8
 - sleep-wake disorders, 209–10
 - substance-related and addictive disorders, 235–6
 - suicidal behaviours, 297–8
- child abuse, 277–9, 280–1
- childhood-onset fluency disorder (stuttering), 65, 67
- children, mental illness in, 86
- Christchurch earthquakes (2010, 2011), 35–6, 148, 282
- Christchurch Health and Development Study (CHDS), 85, 148, 284
- circadian rhythm sleep-wake disorder, 206–7
- clozapine, 80
- cognitive appraisal processes, 135
- cognitive behaviour therapy (CBT)
 - for anxiety disorders, 120–1, 124
 - for bipolar disorders, 109
 - for depression, 97, 99
 - for feeding and eating disorders, 182
 - for obsessive compulsive disorder, 133, 138
 - for schizophrenia, 80–1
 - for somatic symptom and related disorders, 166
- cognitive impairments, compulsory treatment for, 303–4
- cognitive inhibition, 161–2
- cognitive remediation therapy (CRT), 81, 88–9
- communication disorders, neurodevelopmental, 65, 67, 68
- community treatment orders (CTOs), 303
- compulsions see obsessive compulsive disorder (OCD)

- compulsory treatment
 - appointed decision makers, 303–4
 - case study, 306–7
 - in emergency situations, 305
 - for high risk offenders, 306
 - legislation regarding, 301–3
 - standard safeguards against, 301
 - of substance-use disorders, 304–5
- conduct disorder (CD)
 - assessment and diagnosis, 219–20
 - clinical description, 216–17, 219
 - comorbidity with, 219, 220
 - prevalence and impact, 219–20
 - treatment, 220–2
- Convention of the Rights of Persons with Disabilities*, 302
- conversion disorder, 167
- counsellors, 7, 8
- courtship disorder model, 268
- cyclothymic disorder, 107

- danger ideation reduction therapy (DIRT), 132
- data, diversity, 10–11, 34
- death rates *see* mortality rates
- delusional disorder, 78
- dementia *see* neurocognitive disorders (NCDs)
- demographic factors
 - Australia, 10, 11–18
 - New Zealand, 35–6
 - See also* socio-demographic characteristics; socio-economic characteristics
- depersonalisation/derealisation disorder (DRD), 155
- depression
 - aetiology, 96
 - assessment, 96
 - in Australia, 95–8
 - case study, 100–1
 - clinical description, 94–5
 - genetics and, 99
 - in New Zealand, 98–9
 - prevalence and impacts, 95, 98–9
 - support services, 95, 101, 112
 - treatment, 96–8, 99, 111
- depressive disorders, 94–5
 - See also* depression; major depressive disorder (MDD)
- developmental coordination disorder, 66

- Diagnostic and statistical manual of mental disorders* (5th ed.) (DSM-5), 6–7
- disordered eating, 182
- 'disparate impact' discrimination, 19, 45–6
- disruptive, impulse-control, and conduct disorders
 - case study, 224
 - clinical description, 216–17
 - support services for, 225
- dissociative amnesia (DA), 154–5, 156–7
- dissociative disorders
 - in Australia, 156–62
 - clinical description, 154–6
 - conflicting positions on, 156–7
 - in New Zealand, 162–3
 - research, 159–62
 - treatment, 158–9
- dissociative fugue, 155, 159
- dissociative identity disorder (DID)
 - clinical description, 155–6
 - conflicting positions on, 156–7
 - research, 159–62
 - treatment, 158–9
- diversion measures, 305
- diversity data, 10–11, 34
- divorce rates, 277, 280
- drug therapy *see* pharmacological treatment
- DSM-5 *see* *Diagnostic and statistical manual of mental disorders*
- Dunedin Multidisciplinary Health and Development Study (DMHDS), 85–6
- dyscalculia, 66
- dyslexia, 66, 67
- dysthymia, 111

- earthquakes, community impact of, 35–6, 148, 282
- eating disorders *see* feeding and eating disorders
- education (statistics), 26–8, 46–7
- electroconvulsive therapy (ECT), 109, 303
- electroencephalogram (EEG) coherence, 159–60
- elimination disorders
 - in Australia, 190
 - case study, 196
 - clinical definition, 189–90
 - in New Zealand, 194–6
 - prevalence, 194

- support services, 193–4, 195, 197
- treatment and assessment, 190–4, 195–6
- emergency situations, compulsory treatment in, 305
- employment rates, 27–8, 47
- encopresis (faecal soiling)
 - case study, 196
 - clinical definition, 189–90
 - prevalence, 190, 194
 - support services, 193–4, 195
 - treatment and assessment, 190–4, 195–6
- engagement, clinical, 82–3
- enuresis (wetting)
 - clinical definition, 189–90
 - prevalence, 190, 194
 - support services, 193–4, 195
 - treatment and assessment, 190–4, 195–6
- environmental risks, schizophrenia and, 79
- ethnic groups (New Zealand)
 - age structures of, 37–8, 39
 - anxiety disorders in, 123
 - bipolar disorders in, 110
 - neurocognitive disorders in, 248, 250
 - obsessive compulsive disorder in, 134, 135, 136–7
 - resident population, 36–9
 - schizophrenia in, 85
 - socio-demographic characteristics of, 39–44
 - socio-economic characteristics of, 46–8, 281
 - spatial distribution, 35
 - substance-related and addictive disorders in, 233, 234
 - See also Māori population
- excoriation (skin picking), 130, 133
- exhibitionistic disorder, 267–8, 271–2
- exposure and response prevention (ERP), 132, 133, 134
- externalising disorders see disruptive, impulse-control, and conduct disorders
- family/household compositions, 24–6, 43–4
- feeding and eating disorders
 - in Australia, 176–9
 - case study, 181
 - clinical description, 175–6
 - disordered eating and, 182
 - in New Zealand, 179–81, 183
 - obsessive compulsive disorder and, 135
 - prevalence, 176–7, 179–80
 - risk factors and consequences, 177, 180
 - support services, 179, 180–1
 - treatment and assessment, 177–9
- fertility rates see birth rates
- fetishistic disorder, 271
- gambling disorder, 229–30, 231–3, 234–5
- gender differentials see sex differentials
- generalised anxiety disorder
 - age of onset, 118, 122
 - case study, 282–3
 - clinical description, 117
 - prevalence, 117, 122
 - treatment, 119, 121
- generations (population waves), 19, 21
- genetics
 - bipolar disorders and, 106
 - depression and, 99
 - obsessive-compulsive and related disorders and, 131
 - schizophrenia and, 76, 79, 88
 - screening, 68
- global developmental delay (GDD), 64, 68
- guardianship acts, 303–4
- hallucinations, 160–1
- heritability see genetics
- hoarding disorder, 130, 131, 133, 137–8
- homelessness, 279, 281
- household/family compositions, 24–6, 43–4
- hypersomnolence disorder, 205–6
- hypomanic episode, 107
- illness anxiety disorder (health anxiety), 165
- income (statistics), 16, 35, 47–8
- Indigenous Network Suicide Intervention Skills Training (INSIST), 101
- Indigenous population (Australia)
 - age structures in, 16, 17–18
 - anxiety disorders in, 119–20
 - clinical approaches and considerations, 57–60
 - conduct disorders in, 222
 - culture and identity, 57, 59–60
 - depression in, 95
 - disparate impact on, 19
 - diversion programs for, 305
 - feeding and eating disorders in, 176, 178
 - mental illness in, 107

- Indigenous population (Australia) (*cont.*)
 narrative therapy and, 60, 97–8
 neurocognitive disorders in, 251–2
 neurodevelopmental disorders in, 68
 obsessive-compulsive and related disorders in, 133–4
 parenting intervention programs in, 277
 population composition, 17–18, 39
 posttraumatic stress disorder in, 145
 socio-demographic characteristics of, 22, 23–6
 socio-economic characteristics of, 26–7, 28, 279
 substance-related and addictive disorders in, 230, 231, 232
 suicidal behaviours in, 291
 suicide risk in, 95
 support services for, 54–5, 101, 283
 violence in, 278
 See *also* Māori population
- inherent diversity, 10, 34
 insomnia disorder, 204–5
 intellectual development disorder, 64–5
 intellectual disability, 64–5
 intentional self-harm *see* suicidal behaviours
 intermittent explosive disorder, 222–4
 internet gaming disorder (IGD), 230, 233, 235, 236
- jet lag, 207
- Kimberly Indigenous Cognitive Assessment (KICA-Cog), 251–2
 Kraeplin, Emil, 76
- language disorder, 65
 languages, dominant, 22, 41–4
 learning disorders *see* specific learning disorders
 legislation, mental health, 301–3
 life expectancy, 24, 43
 lithium, 108–9
- major depressive disorder (MDD)
 aetiology, 96
 clinical description and diagnosis, 94–5
 prevalence, 98
 treatment, 97
 See *also* depression
 manic depression *see* bipolar and related disorders
- manic episode, 107
- Māori population
 age structure and, 41
 anxiety disorders in, 122–3
 autism spectrum disorder in, 70
 bipolar disorders in, 110
 disparate impact on, 19, 45–6
 dissociation in, 163
 diversion programs for, 305
 elimination disorders in, 195
 feeding and eating disorders in, 180
 mental health needs in, 99
 neurocognitive disorders in, 248, 250
 obsessive compulsive disorder in, 134, 135, 136–7, 138
 posttraumatic stress disorder in, 148
 resident population, 36–9
 schizophrenia in, 85, 87–8
 socio-demographic characteristics of, 41–4, 282
 socio-economic characteristics of, 46–8, 281
 spatial distribution, 35
 substance-related and addictive disorders in, 233, 234
 suicidal behaviours in, 291
 See *also* ethnic groups
- marital status (statistics), 43–4
 masochism, 268–9
 mathematics, impairment in (dyscalculia), 66
 medications *see* pharmacological treatment
 memory processes, 161–2
 mental health legislation, 301–3
 mental health professionals, 7–8, 28, 48–50
 mental illness, compulsory treatment of, 301–3
 metacognitive therapy, 99, 132
- migrant populations (Australia)
 depression in, 95
 socio-demographic characteristics of, 19–26
 socio-economic characteristics of, 15–16, 27–8
 spatial distribution, 12
- migrant populations (New Zealand) *see* ethnic groups
- mood disorders *see* bipolar and related disorders;
 depressive disorders
- mortality rates, 24, 43, 77
- motor disorders, neurodevelopmental, 66, 67
- ‘multiple count’ method, 36, 37
- mutism, selective, 116

- narrative therapy, 60, 97–8
- National Disability Insurance Scheme, 69
- natural disasters, psychological effects of, 148, 279–80, 282
- neurocognitive disorders (NCDs)
 - assessment, 246–7, 249–50
 - in Australia, 245–8, 251–2
 - case study, 250–1
 - clinical description, 243–4
 - DSM-5 coding for, 244–5
 - in New Zealand, 248–50, 252
 - prevalence and mortality rates, 245, 248
 - research, 245–6, 252
 - treatment and prevention, 247–9
- neurodevelopmental disorders
 - in Australia, 66–9
 - case study, 71
 - clinical description, 64–6
 - incidence and prevalence, 66–7, 69–70
 - in the Māori population, 70
 - in New Zealand, 69–70
 - related factors, 70
 - research, 67
 - support services, 69
 - treatment and assessment, 67–9, 70
- neurosurgical interventions (psychosurgery), 303
- New Zealand
 - diversity in, 10, 34, 36–9
 - socio-economic characteristics, 35, 46–8, 281
 - See also demographic factors; socio-demographic characteristics
- nightmares, 208
- obsessive-compulsive and related disorders
 - in Australia, 130–4
 - case study, 137–8
 - clinical description, 129–30
 - in New Zealand, 134–7
- obsessive compulsive disorder (OCD)
 - aetiology, 131–2, 135–6
 - clinical description, 129–30
 - comorbidity with, 135
 - prevalence and impact, 130–1, 134–5
 - support services, 138
 - treatment and assessment, 132–4, 136–7
- offenders (high risk), compulsory treatment for, 306
- online therapies
 - for anxiety disorders, 121
 - for depression, 97
 - for obsessive compulsive disorder, 132, 134, 138
- oppositional defiant disorder (ODD)
 - assessment and diagnosis, 218
 - case study, 224
 - clinical description, 216–17
 - comorbidity with, 217, 218
 - prevalence and impact, 218
 - treatment, 218–19
- oral tradition, 60
- other specific dissociative disorder (OSDD), 156
- other specified bipolar disorder, 107
- other specified feeding or eating disorder (OSFED), 176
- paedophilic disorder, 270
- Pākehā populations see ethnic groups
- panic disorder
 - clinical description, 117
 - prevalence, 117, 122
 - treatment, 119, 121, 123
- paraphilic disorders
 - case study, 271–2
 - clinical description and diagnosis, 266–7
 - support services, 272–3
- parasomnias, 207–8
- partner abuse, 280–1
- personality disorders
 - in Australia, 260–1
 - case study, 262–3
 - clinical description, 258–9
 - cultural perspectives on, 259–60
 - in New Zealand, 261
 - prevalence, 260, 261
 - resembling schizophrenia, 79
 - treatment and assessment, 260–1, 263
- pervasive developmental disorders see autistic spectrum disorder (ASD)
- pharmacological treatment
 - for bipolar disorders, 108–9
 - for exhibitionistic disorder, 268
 - for neurodevelopmental disorders, 68
 - for schizophrenia, 79, 80
 - for somatic symptom and related disorders, 166

- pica, 176–7
- population share, 12
- posttraumatic stress disorder (PTSD)
 - in Australia, 144–7
 - case study, 149–51
 - clinical description and diagnosis, 143–4
 - comorbidity with, 145, 146, 148
 - prevalence and risk factors, 144–5, 148
 - support services, 147
 - treatment and assessment, 145–7, 148–9
- poverty, 279, 281
- professionals *see* mental health professionals
- psychiatric nurses, 49–50
- psychiatrists, 7
- psychological disorders, 6–7
- psychological treatment
 - for anxiety disorders, 119–22
 - of Australian Indigenous population, 57–60
 - for bipolar disorders, 109
 - for dissociative disorders, 158–9
 - for elimination disorders, 191–3, 196
 - for exhibitionistic disorder, 268
 - for neurodevelopmental disorders, 68–9
 - for obsessive compulsive disorder (OCD), 132–4
 - for personality disorders, 260, 261, 263
 - for posttraumatic stress disorder, 145–7
 - for schizophrenia, 79, 80–1
 - for somatic symptom and related disorders, 166
- See also* compulsory treatment
- psychologists, 7, 8, 28, 48
- psychopathology, 6–7
- psychosis, 78, 81–2, 84
 - See also* schizophrenia
- psychosurgery, 303
- psychotic disorders *see* schizophrenia
- qualification levels (statistics), 26–8, 46–7
- reading, impairment in (dyslexia), 66, 67
- regulation, of mental health professionals, 7–8
- rehabilitation, vocational, 81
- relational problems, 277, 280, 282–3
- religious affiliations, 22–3, 42, 279, 282
- residential distribution *see* spatial distribution
- rumination disorder, 176–7
- rural populations, 15, 119, 292
- sadism, 269–70
- schizoaffective disorder, 76, 78, 83–4
- schizophrenia
 - in Australia, 83–4
 - case study, 87–8
 - causes, 76, 79, 85–6
 - clinical features and diagnosis, 77–8, 81–2
 - history of, 76
 - in New Zealand, 85–7
 - obsessive compulsive disorder and, 136
 - prevalence and impact, 77, 83–4, 85
 - research, 84, 88
 - spectrum and related disorders, 78–9
 - support services, 83
 - treatment and assessment, 79–83, 84, 86–7, 88–9
- schizophreniform disorders, 78, 85
- selective mutism, 116
- self-harm, intentional *see* suicidal behaviours
- separation anxiety disorder, 116
- sex differentials (statistics)
 - in anxiety disorders, 118
 - in disruptive, impulse-control, and conduct disorders, 219, 220
 - in employment rates, 28
 - of mental health professionals, 28, 29, 48, 50
 - in qualification levels, 26
 - in substance-related and addictive disorders, 232
 - in suicidal behaviours, 291
- sexual masochism disorder, 268–9
- sexual sadism disorder, 269–70
- sleep terrors, 208
- sleep-wake disorders
 - case study, 209–10
 - clinical description, 202–3
 - prevalence and impact, 203
 - research, 210–11
- sleepwalking, 207–8
- social anxiety disorder (social phobia)
 - case study, 124
 - clinical description, 116
 - prevalence, 117, 122
 - treatment, 119, 121, 123
- Social Cognition and Interaction Training (SCIT)
 - tool, 88–9
- social (pragmatic) communication disorder, 65
- social phobia *see* social anxiety disorder

- social workers, 7, 8
- socio-demographic characteristics (Australia)
 - ancestry, 21
 - birth rates, 23
 - birthplace data, 20–1
 - dominant languages, 22
 - family and household compositions, 24–6
 - methodological implications, 17
 - mortality rates, 24
 - religious affiliations, 22–3, 279
 - See *also* demographic factors; socio-economic characteristics
- socio-demographic characteristics (New Zealand)
 - birth rates, 42
 - birthplace data, 39–44
 - dominant languages, 41–4
 - ethnic diversity measurement, 36–9
 - family and household compositions, 43–4
 - methodological implications, 45
 - mortality rates, 43
 - religious affiliations, 42, 282
 - See *also* demographic factors; socio-economic characteristics
- socio-economic characteristics
 - Australia, 10, 15–16, 26–8, 279
 - New Zealand, 35, 46–8, 281
- somatic symptom and related disorders
 - aetiology, 165
 - case study, 167
 - clinical description, 163–4
 - management, 166
 - risk factors, 165
- somatic symptom disorder (somatisation disorder), 164–5
- somatoform disorders *see* somatic symptom and related disorders
- somatoform dissociation, 165
- spatial distribution, 11–18, 20, 35–6
- specific learning disorders, 66, 67
- specific phobia, 116, 123
- speech sound disorder, 65, 67
- standardisation technique (data), 17, 45, 46
- stereotypic movement disorder, 66
- structural ageing (Australia), 16
- stuttering (childhood-onset fluency disorder), 65, 67
- substance-induced disorders, 229
- substance/medication-induced anxiety disorder, 117
- substance-related and addictive disorders
 - in Australia, 230–3, 236
 - case study, 235–6
 - clinical description, 229–30
 - in New Zealand, 233–5, 236
- substance-use disorders (SUDs)
 - assessment and treatment, 231, 233–4, 236, 304–5
 - case study, 235–6
 - clinical description, 229
 - comorbidity with, 233
 - prevalence and associated risks, 230–1, 233
- suicidal behaviours
 - in Australian Indigenous population, 95
 - in bipolar disorders, 106
 - case study, 297–8
 - causes and risk factors, 292–3
 - depression and, 95, 98–9
 - in obsessive compulsive disorder, 135
 - presentation, 290
 - prevalence and impact, 290–2
 - reduction, 293–5
 - research, 298
 - risk assessment, 295–6
 - in schizophrenia, 84
 - services addressing, 101
 - stigmatisation of, 290, 298
 - treatment and prevention, 296–7
- The Black Dog Institute, 112
- tic disorders, 66, 68
- Torres Strait Islander population *see* Indigenous population (Australia)
- Tourette's syndrome, 67
- trauma (as aetiological factor), 160–1, 165
- Trauma and Dissociation Unit (TDU) (Brisbane), 158–9
- trauma- and stressor-related disorders (TSRD)
 - in Australia, 144–7
 - clinical description, 143
 - in New Zealand, 148–9
- trauma-focused psychological therapy, 145–6
- treatment, psychological *see* psychological treatment
- trichotillomania (hair pulling), 130, 133
- unspecified communication disorder, 65
- unspecified dissociative disorder (USDD), 156

- unspecified feeding or eating disorders (USFED),
176

urbanisation, 15, 35

urotherapy, 191

vascular dementia, 250–1
See also neurocognitive disorders (NCDs)
- violence, 85–6
See also adult abuse; child abuse

vocational rehabilitation, 81

voice disorders, 67

written expression, impairment in,
66