

## Index

- Abbreviations used in the index:
- ARDS adult respiratory distress syndrome
- DIC disseminated intravascular coagulation
- ED emergency department
- HHS hyperglycemic hyperosmolar state
- HUS hemolytic uremic syndrome
- IAP intra-abdominal pressure
- TCA tricyclic antidepressants
- TTP thrombotic thrombocytopenic purpura
- ABCs (airway, breathing, circulation), 41–3  
 acute spinal cord compression, 136  
 adrenal crisis, 406  
 burns, 108–9  
 diabetic ketoacidosis, 392  
 drug overdose, 442, 445  
 neck trauma, 67–9  
 pulmonary embolism, 226  
 shock management, 6, 9  
 traumatic brain injury, 48, 52  
*see also* airway assessment; breathing; circulation
- abdomen, acute/surgical, *see* surgical abdomen
- abdominal aortic aneurysm (AAA), 237–43  
 critical management, 241–3  
 diagnosis and evaluation, 239–41  
 nonemergent treatment, 242  
 presentation, 238–9  
 risk factors, and rupture risk, 238  
 ruptured, 237–41, 243  
 screening, 237, 240  
 sudden deterioration, 238, 243
- abdominal compartment syndrome (ACS), 274–7
- abdominal examination, 270, 286
- abdominal guarding, 270
- abdominal organs, arterial supply, 262
- abdominal pain, 268–70  
 abdominal aortic aneurysm, 238–9  
 acute pancreatitis, 246  
 location and differential diagnosis, 268–9  
 referred, 269  
 sickle cell emergencies, 349
- abdominal trauma (solid organ), 82–9  
 blunt, 83, 86, 88  
 critical management, 87–8  
 diagnosis and evaluation, 83–5, 88  
 penetrating, 82, 86  
 presentation, 83  
 sudden deterioration, 89
- abdominal radiographs, 84, 240, 265, 271, 286–7
- absolute neutrophil count (ANC), 339
- ACE inhibitors, 143, 168–9
- acetaminophen, 401, 432
- acetaminophen overdose/toxicity, 431–5  
 critical management, 256–8, 434  
 fulminant hepatic failure in, 253, 256–8, 432  
 presentation, and diagnosis, 432–3  
 sudden deterioration, 434–5
- acetone, serum, 391
- N*-acetylcysteine (NAC), 257–8, 433–4
- acetylsalicylic acid, *see* aspirin overdose
- acid–base disturbances, 289–97  
 aspirin overdose, 436  
 compensation, 290  
 heatstroke, 419  
 shock, 4  
 sudden deterioration, 297  
*see also* metabolic acidosis; metabolic alkalosis; respiratory acidosis; respiratory alkalosis
- acidemia, 289, 291
- acidosis, 289  
*see also* metabolic acidosis; respiratory acidosis

**458 Index**

- acquired immunodeficiency syndrome,  
*see* AIDS
- ACTH, 403
- ACTH stimulation test, 406
- activated charcoal, 434, 436, 440, 442,  
445, 448
- activated partial thromboplastin time  
(aPTT), 327
- acute abdomen, *see* surgical abdomen
- acute chest syndrome (ACS), 350, 353
- acute coronary syndrome (ACS), 139–45
- critical management, 142–3, 188
  - diagnosis and evaluation, 140–1
  - etiology, 139
  - pharmacological agents, 142–3
  - presentation, 140
  - sudden deterioration, 143–5
  - see also* myocardial infarction, acute
- acute decompensated heart failure  
(ADHF), 164–70
- critical management, 168–9
  - diagnosis and evaluation, 166–8
  - differential diagnosis, 165–6
  - etiology, 164–5
  - presentation, 165–6
  - sudden deterioration, 169–70
- acute interstitial nephritis (AIN), 318
- acute kidney injury (AKI), 314–19
- critical management, 316–18
  - definition and classes, 314, 345
  - etiologies, 315
  - presentation and diagnosis, 314–16
  - rhabdomyolysis and, 318–19
  - severity score (KDIGO), 316
  - special circumstances, 318–19
  - sudden deterioration, 318, 347
  - tumor lysis syndrome and, 345–7
  - types, 314–15
- acute lung injury (ALI), prevention, 31
- acute mesenteric ischemia, 261–6
- critical management, 266–7
  - diagnosis and evaluation, 264–5
  - etiology and pathogenesis, 261–3
  - presentation, 264
  - risk factors, 263
- acute pancreatitis, 245–52
- critical management, 250–2
  - diagnosis and evaluation, 248–50, 391
  - etiologies, 245–6
  - gallstone, *see* gallstone pancreatitis
  - hemorrhagic, 251–2
  - mild, management, 250
  - presentation, 246–8
  - prognostic scores, 247–9
  - severe, 245, 247–8
  - SIRS with, 245, 251–2
  - sudden deterioration, 251–2
- acute respiratory distress syndrome  
(ARDS), 213–17, 251, 362
- critical management, 215–17, 435
  - diagnosis and evaluation, 215
  - etiology, 213–14
  - presentation, 213–14
  - sudden deterioration, 214, 217
- acute spinal cord compression,  
*see* spinal cord
- Addison disease, 403, 405
- adenosine, 150–1
- adrenal crisis, 403–7
- critical management, 406–7
  - diagnosis and evaluation, 405–6
  - presentation, 404–5
  - special circumstances, 407
- adrenal hemorrhage, 407
- adrenal insufficiency, 403
- acute, 404–5
  - causes, 404
  - chronic, 404, 407
  - presentation, 403–4
  - see also* adrenal crisis
- advance directives, 452
- afterload, 37, 168
- agitation, control in thyroid storm, 401
- AIDS, 381–7
- criteria, 381–2
  - opportunistic infections, 382–3
  - pneumonia, 364–5
  - sudden deterioration/  
complications, 382–7
- airway
- collapse, warning signs, 198
  - foreign bodies, 201
  - injuries, 66
  - upper, emergencies, *see* upper airway  
emergencies
- airway assessment
- massive hemoptysis, 220–1
  - primary survey, 41–2, 198
  - burns, and injuries, 48, 60, 108–9
- airway management, 17–26
- alternative techniques, 26
  - difficulty, assessment, 25–6
  - double setup, 25, 68
  - failure, recognition, 17–18

- principles, 17–19
- specific conditions
  - abdominal trauma, 87
  - acute kidney injury, 317
  - aspirin overdose, 437
  - asthma and COPD, 206–7, 211
  - heatstroke deterioration, 421
  - intracranial hemorrhage, 126, 128
  - massive hemoptysis, 220–1
  - neck trauma, 67–9
  - spinal cord compression/injury, 61–2, 136
  - status epilepticus, 132–3
- see also* intubation
- albumin, calcium binding, 308
- albuterol, 205–6, 211, 346
- aldosterone deficiency, adrenal crisis, 405
- alkalemia, 294
- alkalosis, 289
  - see also* metabolic alkalosis; respiratory alkalosis
- allergic syndrome, life-threatening,
  - see* anaphylaxis
- allopurinol, 345–6
- alpha-1 receptor antagonists, 410
- aluminum hydroxide, 346
- alveolar edema, 213
- American Spinal Injury Association (ASIA) scale, 58
- amiodarone, 151
- ammonia levels, 258
- ammonium, renal excretion, 292
- amylase, serum, 248–9
- analgesia, *see* opioid analgesics; pain control
- anaphylactoid reaction, 413
- anaphylaxis, 199–200, 413–17
  - causes, 413–14
  - critical management, 415–17
  - diagnosis and evaluation, 415
  - differential diagnosis, 414
  - patients on beta-blockers, 416–17
  - presentation, 413–15
  - rebound reaction, 416
  - sudden deterioration, 416–17
- anemia, in sickle cell disease, 352
- anesthetic agents, malignant hyperthermia and, 421
- aneurysm, aortic, *see* abdominal aortic aneurysm (AAA)
- angina, intestinal/mesenteric, 262
- angina pectoris, 140
  - unstable, 139–40, 143
- angioedema, 199–200
- angiography
  - abdominal, 85, 240–1, 265
  - CT-angiography, 65–7, 103, 265
- anion gap, 291–2
  - see also* metabolic acidosis
- ankle-brachial index (ABI), 103
- anorexia, forced, acute pancreatitis, 250
- anterior cerebral artery syndrome, 116
- anterior cord syndrome, 58–9
- antidiuretic hormone (ADH), 7, 360
  - deficiency/insensitivity, 299
  - inappropriate secretion (SIADH), 301–2
  - increased secretion, 299–300
- antiarrhythmics, 144
- antibiotics
  - COPD, 210
  - febrile neutropenia, 340
  - infective endocarditis, 373–4
  - meningitis and encephalitis, 369
  - multi-resistant infections, 360
  - necrotizing soft tissue skin infections, 377–9
  - open fractures, 105–6
  - penetrating abdominal trauma, 88
  - pneumonia, 365
  - severe sepsis and septic shock, 359
  - small bowel obstruction, 288
  - tube thoracostomy, pneumothorax, 73
  - variceal bleeding, 234
- anticholinergics, inhaled, 205, 210
- anticoagulants, 142, 227–8, 325
  - mechanism of action, 326
- anticoagulation reversal, 128, 325–8
  - in brain/head trauma, 53–4, 326
  - critical management, 327–8
  - diagnosis and evaluation, 326–7
  - in intracranial hemorrhage, 127–8, 325–7
  - presentation (prior to), 325–6
  - sudden deterioration, 328
- anticonvulsants, 132
- antiemetics, acute pancreatitis, 250
- antihistamines, anaphylaxis management, 416
- antiplatelet agents, 142–3
  - reversal, 54, 328
- antipsychotics, 422–3
- antiretroviral therapy (ART), 386
- antithrombin, 143

**460 Index**

- aortic aneurysm, *see* abdominal aortic aneurysm (AAA)
- aortic dilatation, 237
- aortic dissection, 171–6, 237
  - classification/types, 173, 175
  - critical management, 173–6, 188–9
  - diagnosis and evaluation, 172–3, 186
  - hypertensive emergency, 189
  - presentation, 171–2
  - risk factors, 171–2
  - sudden deterioration, 175
- aortic injury, 76–8
- aortic regurgitation, acute, 193, 195–6
  - critical management, 195–6
  - diagnosis and evaluation, 194–5
  - presentation, 191–3
- aortic rupture, traumatic, 76
- aortic stenosis
  - critical management, 195–6
  - diagnosis and evaluation, 194–5
  - presentation, 191, 193
- aortocaval fistula, 243
- aortoenteric fistula, 231, 235, 239, 242
- aortovenous fistula, 239
- APACHE II score, acute pancreatitis, 248
- aplastic crisis, sickle cell disease, 349, 352
- ARDSNet Protocol, 31
- arrhythmia, *see* cardiac arrhythmia
- arterial blood gas (ABG), 270, 289, 391
  - asthma, or COPD, 204, 209
  - interpretation approach, 289–90
  - normal values, 289
- arterial injuries, 100–1
  - critical management, 105
  - diagnosis and evaluation, 103
  - hard and soft findings, 103–4
- arterial line, shock, 5
- arterial supply, abdominal organs, 262
- arteriography, arterial injuries, 103
- aspergillosis, 383
- aspiration pneumonia, 362, 365
- aspirin, 182, 326, 435
- aspirin overdose, 435–8
- assist volume control (assist control: AC), 30
- asthma, 203–7
  - critical management, 205–7
  - diagnosis and evaluation, 204
  - presentation, 203–4
  - sudden deterioration, 207
- ATACH trial, 127
- Atlanta classification, 247
- atrial fibrillation, 148, 150
- atrial tachycardia, multifocal, 149–50
- atrioventricular (AV) block 155–6
- atrioventricular (AV) node dysfunction, 153, 155–6
- atropine, 156–7, 443, 445
- auto-PEEP, 38–9
- autonomic dysreflexia, 59, 62
- autonomic nerves, 268
- awake intubation, 25–6, 68, 199–200
- awake visualization, upper airway, 199–200
- azithromycin, 211
- back pain, 134
- bag-mask ventilation (BMV), 21
- barotrauma, 28
- bartonellosis, 384
- basilar artery syndrome, 116
- Beck's triad, 75, 172
- bedside index of severity in acute pancreatitis (BISAP), 248–9
- benzodiazepines, 132, 425
- beta-agonists, 205, 210, 416
- beta-blockers, 441
  - anaphylaxis management in patients on, 416–17
  - aortic dissection, 175
  - beta1 and beta2, 441
  - indications and mechanism, 441
  - overdose, 441–4
  - pheochromocytoma, 411
  - STEMI, 142–3
  - thyroid storm, 401–2
- bicarbonate, normal, 289
- Bi-PAP (biphasic positive airway pressure), 206, 296
- bleeding/blood loss
  - abdominal aortic aneurysm, 239
  - control, 43–5, 221–2
  - DIC, 331–3, 335–6
  - massive hemoptysis, 218, 221–2
  - shock, classes, 43
  - trauma, 43–6
  - see also* hemorrhage
- blood count, complete, 4, 232
- blood cultures, 339, 341, 359, 363
- blood flow, shock, 1
- blood pressure
  - aortic dissection management, 174–5
  - control in intracranial hemorrhage, 126–7
  - endpoint of resuscitation and, 36
  - hypertension definition, 185

- hypertensive crisis, 185
- ischemic stroke management, 121
- monitoring, 10–11
- noninvasive measurement, 11
- reduction, hypertensive emergency, 187, 189
- systolic, *see* systolic blood pressure (SBP)
- trauma, 42
- blood products, in management of
  - DIC, 334–5
  - shock, 6–8, 44
  - trauma, 44
- blood supply, abdominal organs, 262
- blood transfusion
  - acute abdomen, 272
  - acute chest syndrome, 353
  - gastrointestinal bleeding, 232, 234–5
  - Jehovah's Witnesses, 235
  - massive, 44, 234
  - massive hemoptysis, 221–2
  - trauma, 44, 88
- bougie, 23
- bowel dilatation, 284
- bowel distension, 284–5
- bowel obstruction, *see* small bowel obstruction
- bowel perforation, 285
- bradyarrhythmias, 153–7
  - critical management, 156–7
  - diagnosis and evaluation, 154–6
  - digoxin overdose, management, 448–9
  - etiology, 153–4
  - presentation, 153–4
  - sudden deterioration, 157
- brain injury, 159
  - traumatic, *see* traumatic brain injury (TBI)
- brain (B-type) natriuretic peptide (BNP), 167, 225, 227
- breath stacking, 32, 34, 38–9
- breathing
  - assessment, primary survey, 42
  - burns, 109
  - spinal cord injury, 60
  - traumatic brain injury, 52
- critical management
  - abdominal trauma, 87–8
  - COPD, 210
  - neck trauma, 69
  - spinal cord injury, 61–2, 136
  - see also* ABCs (airway, breathing, circulation)
- bright red blood per rectum (BRBPR), 231
- bromocriptine, 423
- bronchial artery embolization, 221
- bronchial artery injury, hemoptysis, 218
- bronchodilators, 353
- bronchoscopy, massive hemoptysis, 221
- Brown–Séquard syndrome, 59
- BUN/creatinine ratio, 232
- Burch and Wartofsky scoring system, 399
- burns, 108–12, 198
  - initial assessment, 108–9
  - initial management, 110–11
  - initial resuscitation failure, 111
  - secondary assessment, 111
  - size estimation, 108, 110
  - triage, Burn Center, 112
- calcium
  - administration, in drug overdose, 443, 445, 448
  - ionized (iCa), 308, 345
  - serum, 308
- calcium channel blockers, 151, 411, 444
  - overdose, 444–6
- calcium chloride, 45, 307, 346, 443, 445
- calcium gluconate, 307, 309, 346, 443, 445
- calcium metabolism disorders, 307–10
  - see also* hypercalcemia; hypocalcemia
- Candida albicans*, 384
- candidal esophagitis, 383
- capnography/capnometry, 12, 23
  - asthma/COPD, 204, 209
  - colorimetric, 12, 23
  - quantitative, 12
  - waveform, 23
- carbon dioxide, arterial PaCO<sub>2</sub> levels, 13, 160, 289
- carbon monoxide, measurement, 14
- cardiac arrest, 147, 158
  - in cardiac tamponade, 75–6
  - care after, *see* post-cardiac arrest care
  - complications, 159
  - soft tissue injury deterioration, 106–7
- cardiac arrhythmia
  - acute coronary syndrome, 143–4
  - digoxin overdose, 447–9
  - hypothermia, 162, 429
  - management, 429, 440, 448–9
  - soft tissue injury deterioration, 106–7
  - tricyclic antidepressant overdose, 440
  - tumor lysis syndrome, 346–7
- cardiac biomarkers, 141

**462 Index**

- cardiac biopsy, 181
- cardiac enzymes, 233
- cardiac index (CI), shock, 5
- cardiac output, reduced, 164–7, 275
- cardiac pacing, 443, 446, 449
- cardiac resynchronization therapy (CRT), 169
- cardiac tamponade, 74–6, 183
- cardiogenic shock, 1–5, 7
  - acute coronary syndrome deterioration, 143–4
  - aortic dissection, 172
  - management, 144–5, 162–3, 183, 360
  - presentation, 140, 358
- cardiopulmonary resuscitation (CPR), 13, 75
- cardiovascular emergencies, 139–96
- cardiovascular system
  - HIV/AIDS complications, 386–7
  - injury screening, 64
  - spinal cord injury management, 62
  - thyrotoxicosis complications, 397
- cardioversion, 150, 152, 449
  - biphasic, guidelines, 152
- catecholamines, 398, 408, 443, 446
- cauda equina syndrome, 59, 138
- ceftazidime, 211
- celiac artery, 262
- Center to Advance Palliative Care, 456
- central cooling, 420
- central cord syndrome, 58
- central nervous system (CNS), AIDS complications, 386
- central pontine myelinosis, 302
- central venous catheter (CVC), 5, 15
- central venous oxygen saturation (ScvO<sub>2</sub>), 5, 8, 15
  - endpoint of resuscitation, 38
- central venous pressure (CVP), 15
  - endpoint of resuscitation, 37–8
  - factors affecting, 15
  - increased IAP, 276
  - invasive measurement, 15
  - noninvasive measurement, 14
  - shock, 5, 8, 37
- cerebellar infarction, 116
- cerebral blood flow, 123
- cerebral edema, 258–9, 304, 395, 435
- cerebral embolization, 115
- cerebral hypoperfusion, 367
- cerebral perfusion pressure (CPP), 53, 123, 189, 258
  - acute ischemic stroke, 189
  - decreased, increased IAP and, 275
  - traumatic brain injury, 52
- cerebral thrombosis, 115
- cerebrospinal fluid (CSF) analysis, 339, 368–9
- cerebrospinal fluid (CSF) pressure, subdural hemorrhage, 124
- cerebrovascular accident (CVA), 115
- cervical spine injury, 56, 61, 134
  - NEXUS criteria for clearance, 60
  - penetrating neck trauma, 69
  - primary survey and, 45, 48, 60
- chemical pneumonitis, 109
- chest, flail, 77, 79–80
- chest pain, 140, 171, 178, 278
- chest trauma, *see* thoracic trauma
- chest tube, 73–4
- chest radiography
  - acute abdominal pain, 271
  - acute decompensated heart failure, 167
  - aortic dissection, 173–4
  - aortic injury, 78
  - ARDS, 215–16
  - asthma, 204
  - COPD, 209
  - esophageal perforation, 280
  - hemothorax detection, 74
  - myocarditis, 180
  - neutropenic fever, 339
  - pericarditis, 179
  - pneumonia, 364
  - post-cardiac arrest, 159
  - pulmonary embolism, 224–5
  - shock, 4
  - thoracic trauma, 72
  - thyroid storm, 400
  - valvular diseases, 195
- children
  - abdominal trauma, 88–9
  - acute abdomen, 272
  - coping with death, 455
  - hemolytic uremic syndrome, 332–3, 335
  - shock, 8
  - sickle cell emergencies, 349–50
  - tumor lysis syndrome, 343, 346
- chloride, urine, 294
- cholecystectomy, 250–1
- chronic obstructive pulmonary disease (COPD), 208–12
  - critical management, 210–11
  - presentation and diagnosis, 208–9

- sudden deterioration, 212
- Chvostek's sign, 309
- circulation
  - assessment, primary survey, 42–3
  - burns, 109
  - spinal cord injury, 60
  - traumatic brain injury, 52
- critical management, in trauma, 62, 69, 88
  - see also* ABCs (airway, breathing, circulation)
- cisatracurium, 216
- clevipidine, 188
- clindamycin, 378
- clopidogrel, 326
- coagulation profile, 232, 271
- coagulopathy, 161–2, 233–4, 259, 419
  - see also* disseminated intravascular coagulation (DIC)
- coccidioidomycosis, 385
- cold diuresis, 162
- colorimetric capnometry, 12, 23
- compartment syndrome, 96–8
  - abdominal, *see* abdominal compartment syndrome
  - critical management, 98
  - presentation and diagnosis, 96–8
  - in rhabdomyolysis, 321, 324
- computed tomography (CT)
  - abdominal aortic aneurysm, 240
  - abdominal trauma, 84–5
  - acute pancreatitis, 249–50
  - ARDS, 215
  - esophageal perforation, 280
  - hemoptysis, massive, 220
  - intracranial hemorrhage, 125
  - ischemic stroke, 121
  - lumbar puncture, before, 368
  - pelvic trauma, 93–4
  - post-cardiac arrest, 159
  - shock, 5
  - small bowel obstruction, 287
  - traumatic brain injury, 47
  - traumatic spinal cord injury, 61
- computed tomography angiography (CTA), 65–7, 103, 265
- congestive heart failure, 186, 195, 372
- consciousness, loss, 47, 201–2
- continuous positive airway pressure, asthma, 206
- continuous veno-venous hemofiltration (CVVH), 317, 324
- contrast-induced nephropathy, 318
- controlled mandatory ventilation (CMV), 30
- cooling measures, in heatstroke, 420
- corticosteroids
  - anaphylaxis, 416
  - asthma, 205
  - COPD, 210
  - traumatic spinal cord injury, 62
  - see also* steroid therapy
- cortisol, 403, 406
- creatine kinase (CK),
  - rhabdomyolysis, 321–2
- creatinine clearance (CrCl), 316
- creatinine phosphokinase (CPK), 102, 141
- creatinine, serum, increased, 314
- crescent sign, 240
- cricothyrotomy, 200–1
- critical care (CC) rotations, ED-based, 40
- crush injury, 100–1
  - critical management, 105
  - diagnosis and evaluation, 102
  - sudden deterioration, 106–7
- crush syndrome, 101
- cryptococcal infection, 383
- CT Severity Index, pancreatitis, 248
- Cullen's sign, 83
- CURB-65 rule, 364–5
- Cushing reflex, 52
- Cushing's response, 128
- cyproheptadine, 425
- cytomegalovirus (CMV) retinitis, 383
- D-dimers, 173, 225
- dabigatran (Pradaxa), 54, 326–7
  - reversal, critical management, 327–8
- dantrolene, 422
- decision-making, for dying patients, 452
- decompressive laparotomy, 276
- deep venous thrombosis (DVT), 39, 223–5
- dehydration, signs, diabetic ketoacidosis, 390
- delayed sequence intubation (DSI), 26
- delirium, terminal, 455
- delta-delta gap ( $\Delta\Delta$ ), 291
- desaturation, 19
- desmopressin (DDAVP), 54, 303, 328
- dexamethasone, 369, 401, 406
- diabetes insipidus, 299, 303
  - critical management, 304



**464 Index**

- diabetic ketoacidosis (DKA), 292, 389–93
  - critical management, 391–3
  - diagnosis and evaluation, 390–1
  - differential diagnosis, 390–1
  - etiologies, 389–90
  - presentation, 390
  - sudden deterioration, 395
- diagnostic peritoneal lavage (DPL), 85
- dialysis, emergent, 317–18, 323–4, 347
  - see also* hemodialysis
- diaphragm, thoracic/abdominal trauma, 71
- diastolic heart failure, 164
- diastolic pressure, 11
- diazepam, 347
- digoxin, 151, 446–7
- Digoxin Immune Fab, 448–9
- digoxin overdose, 446–50
  - acute vs. chronic, 447
- dihydropyridines, 444
- diltiazem, 151, 411
- disability, 45, 88
- disseminated intravascular coagulation (DIC), 98, 330–6
  - characteristics, mechanism, 330–2
  - conditions associated, 322, 331, 419
  - critical management, 334–6
  - diagnosis and evaluation, 333–4
  - presentation, 332
  - sudden deterioration, 335–6
- diuretics, 169, 317, 323, 345
- dobutamine, 7, 145, 162–3, 196
  - acute decompensated heart failure, 7, 169–70
  - cardiac tamponade, pericarditis, 184
  - contraindication, 144
- dopamine, 7, 144–5, 157, 162–3, 196
  - acute decompensated heart failure, 7, 169–70
  - cardiac tamponade, pericarditis, 184
- DOPES mnemonic, 33–4
- double setup, airway management, 25, 68, 200
- Duke Criteria, 372–3
- duplex ultrasound, arterial injuries, 103
- dying patient, care of, 451–6
  - critical management, 454–5
  - family experience, improving, 455
  - goal-setting conference, 452–3
  - presentation, and evaluation, 452–3
  - special circumstances, 455–6
  - symptom-based therapy, 452
- e-FAST, 42, 46, 94
- ear-to-sternal notch, 20
- early goal-directed therapy (EGDT), 357, 359–60, 366, 380
- echocardiography
  - acute decompensated heart failure, 167–8
  - cardiac tamponade, 183
  - myocarditis, 180–1
  - pericarditis, 179, 183
  - post-cardiac arrest, 159
  - shock, 4–5
  - thyroid storm, 400
- eclampsia, 133, 188, 190
- ED-centric model, 40
- elderly patients
  - acute abdomen, 272
  - chest trauma, 141, 169, 76, 80
  - gastrointestinal bleeding, 231
  - hyperglycemic hyperosmolar state, 394
  - neutropenic fever, 338
  - pneumonia, 362
  - shock, 9
  - subdural hemorrhage, 124
- electrocardiography (ECG)
  - acute coronary syndrome, 141
  - acute decompensated heart failure, 167
  - aortic dissection, 172
  - bradyarrhythmias, 154–6
  - COPD, 209
  - diabetic ketoacidosis diagnosis, 391
  - digoxin overdose, 448
  - hyperkalemia, 102–3, 107, 306
  - hypokalemia, 305
  - hypothermia, 428
  - monitoring, 10
  - myocarditis, 180
  - pericarditis, 179–80, 182–3
  - post-cardiac arrest care, 159
  - pulmonary embolism, 224, 227
  - QRS complex, 146
    - narrow complex, 146–8
    - TCA overdose, 439–40
    - wide complex, 146–7, 149
  - rhabdomyolysis, 322
  - shock, 4
  - ST segment
    - depression, digoxin overdose, 448
    - elevation, 139, 141, 172, 179, 182
  - tachyarrhythmias, 148–9
  - thyroid storm, 400
  - tricyclic antidepressant overdose, 439–40
  - tumor lysis syndrome, 346



- valvular diseases, 194–5
- electrolyte abnormalities, 299–312
  - diabetic ketoacidosis, 391, 393, 395
  - digoxin overdose, 447–8
  - hypothermia complication, 162
- electrolyte management,
  - rhabdomyolysis, 323
- emboli, valve vegetation fragments, 372
- emergency department (ED), ICU boarder/
  - boarding, 35–40
  - intubated patients, 38–9
  - ongoing assessment, 37
  - sepsis, 38
  - specific management strategies, 40
- emergency department thoracotomy (EDT), 46
- enalaprilat, 169
- encephalitis, 367–70
  - see also* meningitis, and encephalitis
- end of life, *see* dying patient, care of
- end-tidal carbon dioxide (EtCO<sub>2</sub>), 12–13
  - post–cardiac arrest, 160
  - rising, malignant hyperthermia, 421–2
- endobronchial blocker, 221
- endocarditis, *see* infective endocarditis
- endocrine emergencies, 389–411
- endograft, abdominal aortic aneurysm, 242
- endoscopic retrograde
  - cholangiopancreatography (ERCP), 251
- endoscopy, flexible upper, 280
- endotracheal tube (ET), 160, 199, 221
  - placement confirmation, 12–13, 23
  - see also* intubation
- endpoints of resuscitation (EOR), 36–8
- enoxaparin, 227–8, 326–7
  - reversal, critical management, 328
- environmental emergencies, 413–50
- epidural hemorrhage, 49, 123–9
- epinephrine, 7, 157, 205–6, 360, 415–16
- Epstein–Barr virus (EBV), 385
- escharotomy, 109
- esmolol, 151, 175, 401
- esophageal injuries, 66–7
- esophageal perforation (rupture), 278–83
  - causes, and sites, 278–9
  - critical management, 282–3
  - differential diagnosis, 281
  - presentation, and diagnosis, 278–80
  - prognostic factors, and mortality, 278, 283
- esophageal tears, 278
- esophageal varices, bleeding, 229, 231, 234
- esophagram, esophageal perforation, 280, 282
- etomidate, 24, 52, 126
- evaporative cooling, 420
- external laryngeal manipulation (ELM), 22–3
- extraglottic airway (EGA), 18, 21
- eye trauma, sickle cell emergencies, 351, 354
- fasciotomy, 98
- fentanyl, 48
- fever, 339, 398–9
  - neutropenic, *see* neutropenic fever
- FEVER mnemonic, 423
- fibrinolytics, *see* thrombolysis/
  - thrombolytics
- flail chest, 77, 79–80
- flatus, loss, 284
- fluid
  - deficit, hyperglycemic hyperosmolar state, 394
  - overload, 438
  - requirements, burns, 110–11
- fluid replacement
  - diabetic ketoacidosis, 392
  - hyperglycemic hyperosmolar state, 395
  - hypovolemic hypotonic
    - hyponatremia, 301–2
  - rhabdomyolysis, 322–3
  - tumor lysis syndrome, 345
  - vaso-occlusive crisis in sickle cell disease, 351–2
- fluid resuscitation, 233
  - abdominal compartment syndrome, 276–7
  - abdominal trauma, 88
  - acute abdominal pain, 271–2
  - acute pancreatitis, 250
  - anaphylaxis deterioration, 416–17
  - aortic dissection, 173, 175–6
  - brain injury, 52
  - crush injury, 105
  - gastrointestinal bleeding, 233
  - heatstroke, 421
  - rhabdomyolysis, 322–3
  - septic shock, 359
  - shock, 6, 8, 44, 328
  - thyroid storm, 400–1
  - trauma, 43–5, 52
- Focused Assessment with Sonography in Trauma (FAST), 4, 84
  - extended (eFAST), 42, 46, 94
- Foley catheter, 271

**466 Index**

- fondaparinux, 227–8
- foreign bodies, airway, 201
- four-vessel cerebrovascular angiography (FVCA), 67
- fractional excretion of sodium (FENA), 316
- fractional inspired percentage of oxygen (FiO<sub>2</sub>), 29, 39, 160, 215
- fractures, open, 100–1
  - classification schemes, 101, 104
  - critical management, 105–6
- fresh frozen plasma (FFP), 44, 327, 335
- fulminant hepatic failure, 253–60
  - acetaminophen overdose, 253, 256–8, 432
  - complications, management, 258–9
  - critical management, 256–60
  - etiologies, 254
  - presentation, and diagnosis, 253–6
  - prognosis, 256
  - sudden deterioration, 259–60
- fungal infections, AIDS, 383
- furosemide, 317, 346
- gallstone pancreatitis, 250
  - critical management, 250–1
- gastric lavage, 232, 434, 436, 440, 448
- Gastrografin study, 280
- gastrointestinal bleeding, 229–35, 326
  - critical management, 233–5
  - diagnosis and evaluation, 231–3
  - differential diagnosis, 229–30
  - etiologies, 229–30
  - lower tract, 229–31, 233–4
  - presentation, 231
  - upper tract, 229–31, 234
  - upper tract vs. lower tract, 233
- gastrointestinal complications, of AIDS, 387
- gastrointestinal emergencies, 229–88
- gastrointestinal tract decontamination, 442, 445, 448
- generalized convulsive status epilepticus (GCSE), 130
- geriatrics, *see* elderly patients
- Glasgow–Blatchford Score, 233
- Glasgow Coma Scale, 47–8
- glucagon, 266, 416, 442–3, 445
- glucometry, 391, 394
- glucose (blood), 121, 391
  - elevated, *see* hyperglycemia
  - reduced (hypoglycemia), 259, 395, 405
- glutathione, 432
- glyceryl trinitrate, 175, 188
- glycopyrrolate, 454
- “goals of care,” dying patients, 452–3
- GP IIb/IIIa receptor antagonists, 143
- Grey–Turner’s sign, 83
- guaiac test, 230
- gunshot wounds, abdominal, 82
- Gustilo classification system, 104
- hairy leukoplakia, oral, 385
- Hamman’s sign, 79, 279
- “Hampton’s hump,” 224
- harmless acute pancreatitis score, 248
- head, elevation, for intubation, 20
- head trauma, 47–54, 326
- heart block, complete, 155–6
  - see also* atrioventricular (AV) block
- heart failure, 164
  - acute decompensated, *see* acute decompensated heart failure (ADHF)
  - congestive, 186, 195
  - left-sided, presentation, 165
  - right-sided, presentation, 165
- heart rate, 10, 189
- heatstroke, 418–21
- heliox, asthma management, 206
- hematemesis, 231
- hematochezia, 231
- hematocrit, in trauma, 44
- hematology-oncology emergencies, 325–55
- hemodialysis, 317
  - aspirin overdose, 437–8
  - beta-blocker overdose, 444
- hemodynamic instability, arrhythmias, 147, 154
- hemodynamic monitoring
  - invasive, 5–6, 14–15
  - noninvasive, 13–14
- hemodynamic optimization/management, 24, 160, 183, 317
- hemofiltration, 317
- hemolytic uremic syndrome (HUS), 330–6
  - characteristics, 330
  - critical management, 335
  - diagnosis and evaluation, 334
  - presentation, 332–3
  - sudden deterioration, 335–6
- hemoptysis, massive, 218–22
- hemorrhage
  - acute pancreatitis, 251–2
  - intracranial, *see* intracranial hemorrhage

- pelvic trauma, 90, 95
  - see also* bleeding/blood loss
- hemorrhagic shock, 6, 8, 43, 62, 74
  - abdominal trauma, 87, 89
- hemothorax, 73–4, 172
- heparin, 326
  - DIC, 335–6
  - reversal, critical management, 328
  - unfractionated, pulmonary embolism, 227
- hepatic encephalopathy, 255, 258
- hepatic failure, *see* liver failure
- herpes simplex virus (HSV) encephalitis, 367, 369
- high-output heart failure, 164, 402
- histoplasmosis, 384
- HIV infection, 381–7
  - presentation, diagnosis, management, 382
  - see also* AIDS
- hospice, 455
- Hunter criteria, 424–5
- hydralazine, 188, 190
- hydrocephalus, acute
  - noncommunicating, 123
- hydrocortisone, 406–7
- hypercalcemia, 309–10
- hypercapnia, 295
  - permissive, 32, 216
- hypercarbia, 295
- hypercatabolic state, malignant
  - hyperthermia, 421
- hypercoagulable state, 332
- hyperglycemia, 389, 394
  - hypothermia complication, 162
  - pancreatitis and, 391
  - presentation, 390–1
  - see also* diabetic ketoacidosis (DKA)
- hyperglycemic hyperosmolar state (HHS), 394–5
- hyperinflation, 38–9
- hyperkalemia, 304, 306–7, 345
  - critical management, deterioration, 307, 346–7, 448
  - diagnosis and evaluation, 306
  - digoxin overdose, 447–8
  - electrocardiogram, 102–3, 107, 306
  - etiologies, 307
  - exertional heatstroke, 421
  - metabolic acidosis, 291–2
  - presentation, 306–7
  - rhabdomyolysis, 322–3
  - succinylcholine and, 24
  - tumor lysis syndrome, 344
- hypermagnesemia, 312
- hypermetabolic state, 398, 400–1
- hypernatremia, 299, 302–4
- hyperphosphatemia, 345
- hyperpigmentation, 405
- hypertension, 185
  - chronic, management, 187
  - effects, 185
  - management in intracranial hemorrhage, 126–7
  - pheochromocytoma, 409–10
- hypertensive crisis, 185
  - pheochromocytoma, 409–10
- hypertensive emergencies, 185–90
  - complications, management, 188
  - critical management, 187–8
  - diagnosis and evaluation, 186–7
  - presentation, 186
  - specific conditions causing, 189–90
- hypertensive encephalopathy, 188
- hypertensive urgency, 185, 187
- hyperthermia, 418–25
  - thyroid storm management, 398, 401
  - see also* heatstroke; malignant hyperthermia
- hypertonic saline, 53, 127, 132, 259
- hyperuricemia, 343, 345
  - rhabdomyolysis, 322–3
  - tumor lysis syndrome, 343–4
- hyperventilation, 53, 122, 258, 397
- hyphema, 351, 354
- hypocalcemia, 308–9, 345
  - rhabdomyolysis, 322–3
  - tumor lysis syndrome, 344
- hypocapnia, 296
- hypocarbia, 160
- hypoglycemia, 259, 395, 405
- hypokalemia, 304–6
  - aspirin overdose and, 436
  - critical management, deterioration, 306, 393
  - diabetic ketoacidosis, 389, 391
  - diagnosis and evaluation, 305
  - digoxin overdose, 447–8
  - etiologies, 305
  - metabolic alkalosis, 295
  - presentation, 304–5
- hypomagnesemia, 311
- hyponatremia, 299–302
  - adrenal insufficiency, 405
  - status epilepticus due to, treatment, 132

**468 Index**

- hypotension
  - abdominal aortic aneurysm, 238
  - acute coronary syndrome, 144
  - acute decompensated heart failure, 166–9
  - anaphylaxis, 415–17
  - aortic dissection, 172, 175–6
  - DIC deterioration, 336
  - management in intracranial hemorrhage, 127
  - management, TCA overdose, 440
  - massive pulmonary embolism, 227–8
  - meningitis and encephalitis, 370
  - permissive, 44
  - refractory, adrenal crisis, 404–5
  - sepsis and septic shock, 358, 360, 366
  - shock, 2–3
  - sickle cell emergency deterioration, 354
  - in trauma, 43, 60
- hypothermia, 427–30
  - avoidance/prevention, 95, 111
  - critical management, 429
  - presentation, and diagnosis, 427–9
  - sudden decompensation, 430
- hypothermia, therapeutic
  - after cardiac arrest, 160–1
  - fulminant hepatic failure, 259
  - methods and monitoring, 161
  - rewarming, 161
- hypovolemic hypotonic
  - hyponatremia, 300–2
- hypovolemic shock, 1–6
  - pelvic trauma, 91
  - soft tissue injury, 106
- hypoxemia, refractory, in ARDS, 217
- hypoxia, 336, 415
  
- I:E ratio, 29
- ICU-centric model, 40
- imaging
  - abdominal trauma, 84
  - intracranial hemorrhage, 125–6
  - post-cardiac arrest, 159
  - secondary survey in trauma, 46
  - shock, 4–5
  - spinal/spinal cord injuries, 60–1, 136
  - upper airway emergencies, 199
  - see also individual modalities*
- immediate hypersensitivity, 413
- immune reconstitution inflammatory syndrome (IRIS), 386
- implantable cardioverter-defibrillators (ICDs), 169
  
- induction agents, rapid sequence
  - intubation, 52, 126
- infections
  - acute spinal cord compression, 137
  - graft, abdominal aortic aneurysm, 242
  - hypothermia complication, 162
  - multi-resistant, 360
  - neutropenic fever, 337–8
  - pneumonia, *see* pneumonia
  - severe sepsis and septic shock, 359
  - sickle cell emergencies, 349, 352–3
  - status epilepticus, 132–3
- infectious disease emergencies, 357–87
- infective endocarditis, 371–4
- inferior mesenteric artery, 262
- inferior vena cava (IVC),
  - ultrasound, 13–14
- inflammation reduction, pericarditis, 182
- inhalation injury, 109
- inotropic agents, 7, 157
- inspiratory flow rate (IFR), 29
- inspiratory pressure, 29
- inspiratory time ( $T_i$ ), 29
- insulin, 346–7
  - diabetic ketoacidosis, 392–3
  - high-dose, drug overdoses, 443, 445–6
  - hyperglycemic hyperosmolar state, 395
- intensive care unit (ICU), 35
  - ICU boarders in ED, *see* emergency department (ED)
- INTERACT trial, 127
- intestinal angina, 262
- intra-abdominal pressure (IAP), 276
- intraosseous access, 110
- intracerebral hemorrhage, 123–9
  - acute, hypertensive emergency, 189–90
  - warfarin-related risk, 325–6
- Intracerebral Hemorrhage Score, 327
- intracranial hemorrhage, 123–9
  - anticoagulant use and, 325–7
  - critical management, 126–8
  - diagnosis and evaluation, 125–6, 326–7
  - ischemic stroke deterioration, 121
  - presentation, 125
  - sudden deterioration, 128–9, 328
  - types, 123–5
- intracranial pressure (ICP), 52, 123
  - invasive monitoring, 54, 128
- intracranial pressure (ICP), increased
  - intracranial hemorrhage, 123, 127–8
  - ischemic stroke, 121–2
  - laryngeal manipulation effect, 126

- management, 53, 126–8, 258–9
  - goal, 258
  - trauma, 52–3
- intrafacility transfers, upper airway emergencies, 201
- intraparenchymal hemorrhage (IPH), 50
- intravenous immunoglobulin (IVIg), 380
- intraventricular septum rupture, 145
- intubation
  - awake, 25–6, 68, 199–200
  - decision-making, 18
  - delayed sequence (DSI), 26
  - difficult, assessment, 25–6
  - EtCO<sub>2</sub> application, 12–13
  - failure, 17
  - ICU boarder in ED, 38–9
  - indications, 19, 41–2, 48
    - acute spinal cord compression, 136
    - asthma, 206–7
    - COPD, 211
    - massive hemoptysis, 220
    - post–cardiac arrest, 160
    - spinal cord injury, 61
    - upper airway emergency, 199–200
  - nasal route, upper airway emergency, 200
  - placement confirmation, 12–13, 23
  - preparation and positioning, 20
  - process, 20–1
  - rapid sequence, *see* rapid sequence intubation (RSI)
  - sedative-only, 26
- iodine, 401
- ipratropium bromide, 205–6, 210–11
- ischemic heart disease, *see* acute coronary syndrome (ACS)
- ischemic stroke, 115–22
  - acute, hypertensive emergency, 189
  - critical management, 120–1, 189
  - diagnosis and evaluation, 116–20
  - NIH scale, 118
  - presentation, 115–16
  - sudden deterioration, 121–2
  - thrombolytics, 117, 121
- isoproterenol, 151, 157
- IV access, burns, 110
- Janeway lesions, 373
- jaw thrust, 20
- JC virus infection, 385
- Jehovah's Witnesses, 235
- ketamine, 24, 26, 52
- ketones, 389, 391–2
- Kidney Disease Improving Global Outcomes (KDIGO) group, 314, 316
- King's College Criteria, 256, 434–5
- Kussmaul respiration, 291, 390
- labetalol, 175, 188
- Laboratory Risk Indicator for Necrotizing Fasciitis Score (LRINEC), 377, 379
- lactate
  - abdominal trauma, 84
  - elevation, causes, 36
  - endpoint of resuscitation and, 36–7
  - hyperglycemic hyperosmolar state, 394
  - shock, 4, 8, 44
  - trauma, 44, 84
- lactic acidosis, 292–3, 419
- lacunar syndromes, 116
- laparotomy, 88, 276
- laryngeal mask, 21
- laryngoscopy, 20–1, 199, 201
  - apneic oxygenation, 21
  - bimanual, 22–3
  - positioning, 20, 22
  - procedure, 22
- laryngotracheal injuries, 66
- “lead pipe rigidity,” 423
- levofloxacin, 211
- lidocaine, 48
- lipase, serum, 248–9
- lipid emulsion therapy, 443, 446
- liver disease, gastrointestinal bleeding and, 235
- liver enzymes, acetaminophen overdose, 432–3
- liver failure, 54, 258–9
  - acute, *see* fulminant hepatic failure
- liver injury, grading scale, 86
- liver transplant, indications, 434–5
- loop diuretics, 169, 323, 345
- lorazepam, 347
- low–molecular weight heparin, 228
- lower gastrointestinal bleed (LGIB), 229–31, 233–4
- lower leg compartment syndrome, 97
- “lucid interval,” 124–5
- Lugol solution, 401
- lumbar puncture, 368–9
- Lund and Browder chart, 111
- lung injury, ventilator-induced, 28, 31
- lung mechanics, 33
- lung-protective ventilation, 31–2
  - ARDS, 216

Cambridge University Press

978-1-107-62685-0 - Practical Emergency Resuscitation and Critical Care

Edited by Kaushal Shah, Jarone Lee, Kamal Medlej and Scott D. Weingart

Index

[More information](#)**470 Index**

- MAD LoVE mnemonic, 37, 40
- magnesium, 150–1, 449
- magnesium metabolism disorders, 311–12
- magnesium sulfate, asthma, 206
- magnetic resonance imaging (MRI)
- abdominal aortic aneurysm, 241
  - abdominal trauma, 85
  - acute spinal cord compression, 136
  - intracranial hemorrhage, 125
  - myocarditis, 181
  - traumatic spinal cord injury, 61
- malignant hyperthermia, 421–2
- mangled extremity severity score (MESS), 101–2
- mannitol, 53, 127, 258, 323
- MARS (molecular adsorbants recirculation system), 260
- mean arterial pressure (MAP), 11, 123, 189
- calculation/measurement, 11
  - endpoint of resuscitation, 38
  - hypertensive emergency, 187
  - increasing, trauma/ICP management, 53, 127
  - in intracranial hemorrhage, 127
  - post-cardiac arrest, 160
  - trauma resuscitation, 44
- mechanical ventilation, 28–34
- acute decompensated heart failure, 168
  - ARDS, 216
  - aspirin overdose, 437
  - asthma, 206–7
  - complications, 28–9
  - COPD, 211
  - definitions related to, 29
  - ICU boarder in ED, 38–9
  - IVC diameter, 14
  - lung mechanics, 33
  - modes, 30–1
  - pneumonia, 364
  - strategies, 31–3
  - sudden deterioration, 33–4
  - ventilator settings, 29
  - withdrawal, dying patient, 454–5
- Meckler's triad, 279
- mediastinitis, 278–83
- see also* esophageal perforation (rupture)
- melena, 231
- meningitis, and encephalitis, 367–70, 386
- critical management, 369–70
  - diagnosis and evaluation, 368–9
  - presentation, 368–9
  - sudden deterioration, 370
- mental status (altered), 19, 43, 47, 130, 335, 386
- aspirin overdose, 438
  - diabetic ketoacidosis and HHS, 395
  - shock (classes), 43
  - sickle cell emergency, 354–5
- mesenteric angina, 262
- mesenteric arterial embolism (MAE), 262, 264, 267
- mesenteric arterial thrombosis (MAT), 262, 264, 267
- mesenteric ischemia, *see* acute mesenteric ischemia
- mesenteric venous thrombosis (MVT), 263–4, 267
- metabolic acidosis, 21, 289–93, 389
- anion-gap, 290–1, 389, 394, 436
  - etiologies, 292
  - critical management, 292–3
  - etiologies, 292–3
  - non-anion gap, 292–3
  - presentation, and diagnosis, 290–2
  - rhabdomyolysis deterioration, 324
- metabolic alkalosis, 290, 294–5
- critical management, 295
  - etiologies, 294
  - presentation, and diagnosis, 294–5
- metanephrines, 409
- methimazole, 401
- methylprednisolone, 206, 211
- metoprolol, 151
- metyrosine, 410
- microsporidial diarrhea, 384
- midazolam, 24, 161
- middle cerebral artery syndrome, 115
- milrinone, 7, 169, 402
- minute ventilation ( $V_E$ ), 29
- mitral regurgitation
- acute, 194, 196
  - chronic, 192–3
  - critical management, 196
  - diagnosis and evaluation, 194–5
  - presentation, 192–4
- mitral stenosis
- critical management, 196
  - ECG and chest radiograph, 195
  - presentation, 192–3
- mixed model (ICU boarder in ED), 40
- Mobitz type I/II block, 155–6
- monitoring, 10–16
- basic, 10–11

- multi-organ dysfunction syndrome (MODS), 1–2
- multi-organ failure, acetaminophen overdose, 434–5
- multifocal atrial tachycardia, 149–50
- muscle weakness, diabetic ketoacidosis and HHS, 395
- musculoskeletal dysfunction, increased IAP, 275
- Mycobacterium avium* complex, 383
- Mycobacterium tuberculosis*, 385
- myelography, CT, 136
- myocardial infarction, acute, 139–40, 144, 159
  - aortic dissection causing, 172, 175
  - NSTEMI, 141
    - critical management, 143
  - STEMI, 141
    - critical management, 142–3
- myocardial ischemia, 139, 141
- myocarditis, 177–8
  - critical management, 182, 184
  - diagnosis and evaluation, 179–81
  - presentation, 178–9
  - sudden deterioration, 183
- myoglobin, 102, 141, 322
- NAPQI, 432
- narcotic analgesia, *see* opioid analgesics
- nasogastric aspiration, GI bleeding, 232
- nasogastric decompression, 271
- National Acute Spinal Cord Injury Studies (NASCIS II and III), 62
- National Emergency X-Radiography Utilization Study (NEXUS), 60
- neck
  - pain, 134
  - zones, 64–5
- neck trauma, 64–9
  - blunt, 64
    - critical management, 67–9
    - diagnosis and evaluation, 65–7
    - laryngotracheal injuries, 66
    - penetrating, 64, 69
    - pharyngoesophageal injuries, 66–7
    - presentation (hard/soft signs), 64–5, 199
    - vascular injuries, 67
- necrotizing soft tissue skin infections (NSTI), 375–80
  - causes and risk factors, 375–6
  - classification, 376
  - critical management, 377–80
  - presentation, 376–7
  - sudden deterioration, 380
- nephritis, acute interstitial, 318
- nephropathy, contrast-induced, 318
- neurogenic shock, 59, 62
- neuroleptic malignant syndrome (NMS), 422–3, 425
  - diagnosis and management, 423
- neurological emergencies, 115–38
- neurological examination, spinal cord injury, 60
- neurological protection, post-cardiac arrest, 160
- neuromuscular blockers, 126, 216, 423, 425
- neurosurgery, 49, 54, 122, 128–9
- neutropenia, 337–8
  - diagnosis and levels, 339
- neutropenic fever, 337–41
  - critical management, 339–41
  - diagnosis and evaluation, 339
  - presentation, 338
  - sudden deterioration, 341
  - timing and causes, 337–8
- nicardipine, 175, 188
- nitrates, 142–3, 168
- nitroglycerin, *see* glyceryl trinitrate
- nitroprusside, 188
- nonconvulsive status epilepticus (NCSE), 130
- noninvasive positive-pressure ventilation (NPPV)
  - acute decompensated heart failure, 168–9
  - asthma, or COPD, 206, 211
  - respiratory acidosis, 296
  - respiratory distress, 354
- noninvasive ventilation (NIV), 18–19, 21
- nonocclusive mesenteric ischemia (NOMI), 262–4, 267
- nonrebreather mask (NRB), 21
- norepinephrine, 7
  - acute decompensated heart failure, 170
  - cardiac tamponade, pericarditis, 184
  - cardiogenic shock, 145
  - DIC and TTP/HUS, 336
  - gastrointestinal bleeding, 235
  - HIV/AIDS deterioration, 387
  - infective endocarditis, 374
  - post-cardiac arrest care, 162–3
  - septic shock, 7, 360
  - valvular diseases, 196
- obesity, acute abdomen, 272



**472 Index**

- obstructive strategy, ventilation, 32–3
- ocular ultrasound, 53
- oliguria, 275
- opioid analgesics, 143
  - acute abdominal pain, 272
  - acute pancreatitis, 250
  - acute spinal cord compression, 137
  - dying patients, 454–5
- opportunistic infections, AIDS, 382–3
- organ donation, 456
- Osborn wave, 428
- oscillometric blood pressure, 11
- Osler's nodes, 373
- osmolal gap (OG), 291
- osmolality, serum, 300, 394
- osmotic demyelination syndrome, 132
- osmotic therapy, 53, 122, 127, 258–9
- overdoses, 431–50
- oxygen
  - demand decrease, shock management, 8
  - PO<sub>2</sub>, 289
  - therapy, 352, 364
    - asthma, 205–6
    - ischemic stroke, 120
- oxygen delivery (DO<sub>2</sub>)
  - consumption (VO<sub>2</sub>) imbalance, 35
  - ICU boarders in ED, 35–6
  - shock management, 6–8
- oxygen saturation (SpO<sub>2</sub>), 19, 21, 39, 160, 216
- oxygenation, 19–20, 168
  - apneic, during laryngoscopy, 21
- pain control, 272, 455
  - ICU boarder in ED, 39
- palliative care, *see* dying patient, care of
- pancreas injury, grading scale, 87
- pancreatitis, *see* acute pancreatitis
- papaverine, 266
- papillary muscle rupture, 144–5
- paradoxical breathing, 61
- paraganglioma, 408
- paralytic agents, 161, 422
  - rapid sequence intubation, 24, 52, 126
- Parkland formula, 110–11
- paroxysmal supraventricular tachycardia, 149–50
- peak inspiratory pressure (PIP), 33, 38, 275
- pediatrics, *see* children
- pelvic binding, 94
- pelvic fractures, 90–1
- pelvic trauma, severe, 90–5
  - critical management, 94–5
  - presentation, and diagnosis, 91–4
  - sudden deterioration, 95
- pelvic radiograph, 53, 93–4
- pelvis stability, assessment, 43
- peptic ulcer disease, 232
- percutaneous coronary intervention (PCI), 142
- pericardial effusion, 179, 181, 183
- pericardial window, 183
- pericardiocentesis, 175, 183–4
- pericardiotomy, 75
- pericarditis, 177–84
  - critical management, 182, 184
  - diagnosis and evaluation, 179–81
  - etiology, 178
  - high risk, 182
  - presentation, 178
  - sudden deterioration, 183
- peritonitis, 285
- permissive hypercapnia, 32, 216
- permissive hypotension, 44
- petechiae, 373, 405
- pH, normal, 289
- pharyngoesophageal injuries, 66–7
- phenoxybenzamine, 410
- phentolamine, 188, 410
- phenylephrine, 7, 336
- pheochromocytoma, 408–11
- pheochromocytoma multisystem crisis (PMC), 409
- phosphodiesterase inhibitors, 443, 446
- plateau pressure (Pplat), 33, 39, 216
- platelet transfusion, 334–5, 341
- pleurocentesis, 280
- Pneumocystis jirovecii* (PCP)
  - pneumonia, 384
- pneumonia, 362–6
  - AIDS, 364–5
  - aspiration, 362, 365
  - atypical, and typical, 363
  - causative organisms, 362–4, 385
  - community-acquired (CAP), 362–5
  - critical management, 364–6
  - diagnosis and evaluation, 363–5
  - healthcare-associated (HCAP), 362–3, 365
  - presentation, 363
  - sudden deterioration, 365–6
  - ventilator associated, 28–9, 39
- pneumothorax, 72–3

- deterioration on ventilator due to,
  - 34, 73
  - tension, *see* tension pneumothorax
- point-of-care ultrasonography (US), 4–5
- poisoning/overdoses, 131, 133
  - see also specific overdoses*
- popliteal artery injury, 103
- positioning of patient, massive
  - hemoptysis, 221
- positive airway pressure (PAP), 168
- positive end-expiratory pressure (PEEP),
  - 29, 122, 168, 216
  - auto-PEEP (intrinsic PEEP), 38–9
- positive pressure ventilation, 80, 380
- post–cardiac arrest care, 158–63
  - critical management, 159–63
  - diagnosis and evaluation, 159
  - special circumstances, 162
- posterior cerebral artery syndrome, 116
- potassium iodide, 401
- potassium metabolism disorders, 304–7
  - see also hyperkalemia; hypokalemia*
- potassium repletion, 393, 437
- preeclampsia, 190
  - critical management, 188
- prednisone, 206, 211
- pregnancy
  - acute abdomen, 272
  - HIV/AIDS, 386
  - shock, 8–9
- preload, 168
  - estimation, 15, 37
- preoxygenation, 21, 26
- pressor agents, *see* vasopressors
- pressure control (PC), 30–1
- pressure support (PS), 30–1
- priapism, 349, 352–3
- primary survey, trauma, 41–5, 59–61
- procainamide, 151
- progressive multifocal
  - leukoencephalopathy (PML), 385
- propofol, 24, 126
- propranolol, 401, 441
- propylthiouracil (PTU), 401
- protective equipment, 108
- prothrombin complex concentrate (PCC),
  - 44, 327, 335
- proximal airway pressure (PAP), 38–9
- P's, Five, 101
- pseudoaneurysm, aortic, 237, 243
- pulmonary arteries, massive hemoptysis,
  - 218–19
- pulmonary artery catheter (PAC), 5–6,
  - 14–15, 276
  - measurements taken, 14
- pulmonary capillary wedge pressure
 (PCWP), 14, 276
- pulmonary compliance, increased IAP
  - and, 275
- pulmonary complications
  - AIDS deterioration, 386
  - thyrotoxicosis, 397
- pulmonary edema, 157, 317
  - ARDS, 214
  - critical management, 188
  - presentation/diagnosis, 167
- pulmonary embolism, 223–8, 336
  - clinical prediction scoring, 225
  - critical management, 226–8
  - diagnosis and evaluation, 224–5
  - massive and submassive, 227
  - presentation, 223–4
  - sudden deterioration, 228
- pulmonary imaging, pulmonary
  - embolism, 225–6
- pulmonary optimization, post–cardiac
  - arrest, 160
- pulmonary vascular congestion, 165
- pulse oximetry, 11–13
  - continuous, COPD, 209
  - SpO<sub>2</sub> lag, 19
- pulse pressure, 15
- pulse pressure variation (PPV),
  - 15–16, 37
- pulseless electrical activity (PEA),
  - 163, 224
- pulsus paradoxus, 75
- purpura, atraumatic systemic, 331
- quantitative capnography, 12
- radiographs
  - spinal cord injury, 60–1
  - see also specific radiographs (e.g. chest radiograph)*
- Ranson criteria/score, 247
- rapid sequence intubation (RSI), 23–5, 42,
  - 126, 392
  - infective endocarditis, 374
  - neck trauma, 68
  - traumatic brain injury, 48, 52
- rasburicase, 346
- rectal examination, 286
- red cell aplasia, transient, 349, 352

**474 Index**

- renal complications
  - HIV/AIDS, 387
  - thyrotoxicosis, 397
- renal emergencies, 289
- renal failure
  - critical management, 188
  - drug overdose, 435–6
  - fulminant hepatic failure, 259
  - septic shock deterioration, 361
  - see also* acute kidney injury (AKI)
- renal injury, grading scale, 87
- renal replacement therapy (RRT), 317, 347
  - types and indications, 317–18
- respiratory acidosis, 32, 289–90, 295–6
  - diagnosis, critical management, 295–6
  - etiologies, 296, 419
  - presentation, 295
- respiratory alkalosis, 290, 296–7
  - aspirin overdose, 436
  - critical management, 297
  - etiologies, 297
  - presentation, and diagnosis, 296–7
- respiratory distress, 168, 251, 354, 395
- respiratory emergencies, 197–228
- respiratory failure, 109, 198, 360–1
  - ARDS, 214–15
- respiratory monitoring, 11–12
- respiratory rate (RR), 11, 29, 126
- resuscitation
  - acute abdominal pain, 271
  - assessment in shock, 8
  - CPR, 13, 75
  - endpoints, 36–8
  - fluid/volume, *see* fluid resuscitation
  - goal, in trauma, 44
  - massive hemoptysis, 221
- return of spontaneous circulation (ROSC),
  - 13, 158, 163
- reversal of anticoagulation,
  - see* anticoagulation reversal
- rewarming, 429
- rhabdomyolysis, 101, 105, 318–19,
  - 320–4, 421
  - acute kidney injury due to, 318–19, 320
  - critical management,
    - deterioration, 322–4
  - diagnosis and evaluation, 321–2
  - etiologies, 320–1
  - presentation, 320–1
- rocuronium, 24, 52, 126
- Roth spots, 373
- Rule of 2s, 37
- Rumack–Matthew nomogram, 257, 433
- salicylate, 435–7
- secondary survey, trauma, 45–6
- sedation, 39, 54, 161, 200
  - EtCO<sub>2</sub> monitoring, 13
- sedative agents, rapid sequence intubation,
  - 24, 52, 126
- seizures, 130–1
  - hypothermia complication, 161
  - traumatic brain injury, 54
  - tricyclic antidepressant overdose, 440
  - tumor lysis syndrome deterioration, 347
- sepsis, 357
  - adrenal crisis and, 407
  - chronic adrenal insufficiency, 407
  - ICU boarder in ED, 38
  - risk factors, 357
  - severe, 357–61
- septic shock, 341, 357–61
  - adrenal crisis presentation, 404, 407
  - critical management, 359–60
  - diagnosis and evaluation, 358–9
  - necrotizing soft tissue skin
    - infections, 380
  - presentation, 358
  - sudden deterioration, 360–1
  - vasopressors in, 7, 360
- serotonin syndrome, 424–5
  - critical management, 424–5
  - presentation and diagnosis, 424–5
- shock, 1–9, 35, 43–5
  - cardiogenic, *see* cardiogenic shock
  - compensated/decompensated, 2
  - critical management, 1, 6–9, 35, 328
  - diagnosis and evaluation, 2–6, 87
  - distributive, 1–5
  - hemodynamic monitoring
    - (invasive), 5–6
  - hemorrhagic, *see* hemorrhagic shock
  - hypotensive, aortic dissection, 175
  - hypovolemic, *see* hypovolemic shock
  - neurogenic, 59, 62
  - obstructive, 1–5
  - presentation, 2–3, 231
  - refractory, 2
  - septic, *see* septic shock
  - special patient groups, 8–9
  - types/classes, 1–2, 43
- sickle cell disease, anemia causes, 352
- sickle cell emergencies, 348–55
  - critical management, 351–5
  - diagnosis and evaluation, 351–2
  - presentation, 348–51
  - sudden deterioration, 354–5

- sickle cells, 348  
 “silent chest,” 204  
 sinoatrial (SA) node dysfunction, 153, 155  
 sinus bradycardia, 154  
 sinus tachycardia, 148–9, 400  
 SIRS (systemic inflammatory response syndrome), 357, 375  
   pancreatitis with, 245, 251–2  
   presentation/criteria, 358  
 skin infections, necrotizing, *see* necrotizing soft tissue skin infections (NSTI)  
 small bowel obstruction, 284–8  
   critical management, 287–8  
   diagnosis and evaluation, 285–7  
   etiology, 285  
   presentation, 284–5  
 small bowel perforation, 285  
 sodium bicarbonate  
   aspirin overdose, 437  
   beta-blocker overdose, 444  
   diabetic ketoacidosis, 393  
   intravenous, 292–3  
   rhabdomyolysis, 323  
   tricyclic antidepressant overdose, 440  
 sodium, in diabetic ketoacidosis, 391  
 sodium metabolism disorders, 299–304  
   *see also* hyponatremia; hyponatremia  
 sodium nitroprusside, 175, 410  
 sodium polystyrene sulfonate (Kayexalate), 347, 393  
 sodium-potassium ATPase,  
   blockade, 446–7  
 soft tissue infections, *see* necrotizing soft tissue skin infections (NSTI)  
 soft tissue injury, 100–7  
   critical management, 105–6  
   diagnosis and evaluation, 102–4  
   presentation, and severity, 100–2  
   sudden deterioration, 106–7  
 somatic nerves, 268  
 sotalol, 444  
 spinal cord  
   acute compression, 138  
     causes, 134–5  
     critical management, 136–8  
     diagnosis and evaluation, 135–6  
     presentation, 134  
     sudden deterioration, 137–8  
   complete/incomplete injury, 58  
   incomplete injury syndromes, 58–9  
   trauma, *see* traumatic spinal cord injury (TSCI)  
 spinal shock, 59, 135, 137–8  
 spirometry, asthma, 204  
 splanchnic blood flow, 234, 261, 275  
 splenic injuries, 83, 86  
 splenic sequestration, sickle cell emergencies, 350, 353  
 splinter hemorrhages, 373  
 sputum features/culture, 363–4  
 stab wounds, abdominal, 82  
 status epilepticus, 130–3, 370  
   critical management, 131–2  
   diagnosis and evaluation, 131–2  
   presentation, 130  
 steroid therapy  
   acute spinal cord compression, 137  
   adrenal crisis and, 407  
   neutropenic fever and, 341  
   *see also* corticosteroids  
 stool, black/red, 229, 231  
 stroke  
   hemorrhagic, 124–5  
   ischemic, *see* ischemic stroke  
   sickle cell emergencies, 350, 353  
 stroke volume variation (SVV), 37  
 subglottic obstruction, 201–2  
 subarachnoid hemorrhage (SAH),  
   49, 123–9  
   frequency and causes, 124  
   presentation, 125  
   radiology/diagnosis, 125–6  
 subcutaneous emphysema, 279, 377  
 subdiaphragmatic thrusts, 201  
 subdural hemorrhage, 50, 123–9  
   causes, 124  
 succinylcholine, 24, 52, 126, 421  
 superior mesenteric artery, 262  
 supraventricular tachycardia (SVT), 148  
   paroxysmal, 149–50  
 surgery  
   abdominal aortic aneurysm, 241–3  
   abdominal compartment syndrome, 277  
   acute mesenteric ischemia, 266  
   esophageal perforation, 282–3  
   massive hemoptysis, 221–2  
   necrotizing soft tissue infections,  
     377, 379  
   small bowel obstruction, 288  
   traumatic spinal cord injury, 62–3  
 surgical abdomen, 268–72  
   critical management, 271–2  
   diagnosis and evaluation, 269–71  
   preparation for operating suite, 272  
   presentation, 268–9  
   special circumstances, 272

**476 Index**

- sustained low efficiency dialysis (SLED), 317–18
- Swan–Ganz catheter, *see* pulmonary artery catheter (PAC)
- sweating, impaired, 419
- sympathetic crisis, critical management, 188
- synchronized intermittent mandatory ventilation (SIMV), 30–1
- syncope, 154
- syndrome of inappropriate antidiuretic hormone secretion (SIADH), 301–2
- systemic inflammatory response syndrome, *see* SIRS
- systemic vascular resistance (SVP), shock, 5
- systolic blood pressure (SBP), 10, 15, 160  
traumatic brain injury, 52
- systolic heart failure, 164
- systolic pressure variation (SPV), 15–16
- tachyarrhythmias, 146–52  
critical management, 149–51, 449  
diagnosis and evaluation, 148–9  
digoxin overdose, 449  
presentation, 147  
sudden deterioration, 150–2
- tachycardia, 146  
atrial, 148–50  
reflexive, lack in adrenal crisis, 405  
shock, 3  
sinus, 148–9, 400  
thyroid storm, 399  
ventricular, 149, 150, 398, 447
- telemetry monitor, 10–11
- temperature, 11, 418  
core, 11  
elevated, 419–20  
low, 427  
rectal, 11, 420
- tension pneumothorax, 42, 72  
critical management, 73  
deterioration, 73  
presentation, 72
- terbutaline, subcutaneous, 205–6
- tetanus toxoid, 105
- THAM (tromethamine), 293
- thermoregulation, 418, 427
- thienopyridine antiplatelet agents, 143
- thoracic fluid content (TFC), 37
- thoracic trauma, 71–80, 87–8  
aortic injury, 76–8  
blunt, 75–6  
cardiac tamponade, 74–6  
evaluation, 71  
flail chest, 77, 79–80  
hemothorax, 73–4, 172  
penetrating, 71–2, 75  
pneumothorax, *see* pneumothorax  
tracheobronchial injury, 78–9  
thoracostomy, tube, *see* tube thoracostomy  
thoracotomy (emergency department), 46, 75
- thrombin inhibitors, direct, 228
- thrombocytopenia, 233–4, 333
- thromboembolism, 332  
life/limb threatening, 336  
venous, 223–4
- thrombolysis/thrombolytics, 142–3  
contraindications, 117  
ischemic stroke, 117, 121  
pulmonary embolism, 227
- thrombotic thrombocytopenic purpura (TTP), 330–6  
characteristics, 330  
critical management, 335  
diagnosis and evaluation, 334  
presentation, 332–3  
sudden deterioration, 335–6
- thyroid hormones  
excess, thyroid storm, 397  
inhibition of toxic effects, 401  
release inhibition, 401  
synthesis inhibition, 401
- thyroid storm, 397–402  
critical management, 400–2  
precipitants, 398  
presentation, and diagnosis, 398–400  
sudden deterioration, 402
- thyrotoxicosis, 397
- tidal volumes ( $V_t$ ), 28–9, 216
- tissue plasminogen activator (tPA), 121, 189
- torsades de pointes, 149–50
- total body surface area (TBSA), 108, 110
- total body water (TBW), 304
- toxins/overdoses, 131, 133
- toxoplasmosis, 383
- tracheal obstruction, 201–2
- tracheobronchial injury, 78–9
- tracheostomy, neck trauma, 69
- tranexamic acid (TXA), 45, 88
- transesophageal ultrasound, 173, 372
- trans-thoracic ultrasound, 173

- transient red cell aplasia, 349, 352
- transjugular intrahepatic portosystemic shunt (TIPS), 234
- transtubular potassium gradient (TTKG), 305–6
- “trauma triad of death,” 44
- traumatic brain injury (TBI), 47–54
  - anticoagulation reversal, 53–4
  - common patterns, 49
  - management, 47–53
  - neurosurgery, 49, 54
  - pitfalls to avoid, 54
- traumatic spinal cord injury (TSCI), 56–63, 272
  - critical management, 61–3
  - experimental treatment, 63
  - imaging, 60–1
  - initial evaluation, 59–61
  - pathophysiology, 56–7
  - presentation and grading, 58–9
  - primary, 57
  - secondary, 57–60
- tricyclic antidepressants, 438–9
  - overdose, 438–41
- triiodothyronine (T<sub>3</sub>) synthesis
  - inhibition, 401
- troponin, 141, 167, 179
  - pulmonary embolism, 225, 227
- Trousseau’s sign, 309
- Tscherne classification, fractures, 101
- tube thoracostomy
  - hemothorax, 74
  - tension pneumothorax, 73
- tuberculosis, 219, 385
- tumor lysis syndrome (TLS), 343–7
- ultrasonography
  - abdominal aortic aneurysm, 240
  - acute abdominal pain, 271
  - acute kidney injury, 316
  - acute mesenteric ischemia, 265
  - acute pancreatitis, 250
  - hemothorax detection, 74
  - inferior vena cava assessment, 13–14
  - necrotizing soft tissue infections, 377
  - pneumothorax detection, 72
  - pulmonary embolism, 224–5
  - small bowel obstruction, 287
- unconscious patient, 47, 201–2
- unilateral lung ventilation, 221
- upper airway emergencies, 197–202
- upper airway obstruction, 198–200
  - complete, or incomplete, 201
  - consciousness, loss, 201–2
- upper gastrointestinal bleed (UGIB), 229–31, 233–4
- uric acid, reduction therapy, 345–6
- urinalysis, 84, 391
- urinary retention, acute spinal cord compression, 137
- urine
  - alkalinization, 105, 346, 421, 437
  - diabetic ketoacidosis, 391
  - microscopy/sediment, acute kidney failure, 316
  - myoglobin, 322
  - osmolality, 301–3
  - sodium, 301, 303
  - toxicology screens, 431
- valvular diseases, 191–6, 374
  - critical management, deterioration, 194–6
  - diagnosis and evaluation, 194–5
  - presentation, 191–4
- vancomycin, 340, 379
- variceal bleeding, 229, 231, 234
- varicella zoster virus (VZV), 384
- vascular injuries, neck, 67
- vascular obstruction, aortic
  - dissection, 171–2
- vaso-occlusive crisis, sickle cell disease, 348–9, 351–2
- vasodilators, aortic dissection, 175
- vasopressin, *see* antidiuretic hormone (ADH)
- vasopressors
  - acute coronary syndrome, 145
  - acute decompensated heart failure, 169–70
  - acute kidney injury, 317
  - acute spinal cord compression, 138
  - aortic dissection, 175–6
  - bleeding, anticoagulation reversal, 328
  - cardiac tamponade, pericarditis, 184
  - DIC and TTP/HUS, 336
  - HIV/AIDS deterioration, 387
  - hyperthermia, 425
  - infective endocarditis, 374
  - intracranial hemorrhage, 129
  - meningitis and encephalitis, 370
  - pheochromocytoma, 411
  - pneumonia, 366
  - post-cardiac arrest care, 162–3

Cambridge University Press

978-1-107-62685-0 - Practical Emergency Resuscitation and Critical Care

Edited by Kaushal Shah, Jarone Lee, Kamal Medlej and Scott D. Weingart

Index

[More information](#)**478 Index**

- vasopressors (*cont.*)
  - septic shock deterioration, 360, 380
  - shock management, 6–7
  - sickle cell emergencies, 355
  - status epilepticus, 133
  - trauma, 45, 53
  - valvular diseases, 196
- vecuronium, 52
- venous blood gases (VBG), 289
- venous return, reduced, increased IAP, 275
- venous thromboembolism, 223–4
- ventilation
  - mechanical, *see* mechanical ventilation
  - noninvasive (NIV), 18–19, 21
  - withdrawal, dying patient, 454–5
- ventilator associated pneumonia (VAP), 28–9, 39
- ventilator-induced lung injury (VILI), 28, 31
- ventricular fibrillation, 429
- ventricular free wall rupture, 145
- ventricular tachycardia, 149–50, 398, 447
- verapamil, 151
- vertebrobasilar artery syndrome, 116
- video laryngoscopes/laryngoscopy, 20, 22, 26, 199
- viral infections, AIDS, 383
- vital signs, shock, 3
- vitamin K, 327
- vitiligo, 405
- vocal cords, 22
- volume resuscitation, *see* fluid resuscitation
- volutrauma, 28
- warfarin, 227, 325–6
  - reversal, 53, 327
- Wenckebach block, 155–6
- “Westermarck’s sign,” 225
- white blood cell count (WBC), 270, 351
- whole bowel irrigation, 442, 445
- withdrawal, of life-sustaining
  - treatment, 451–2
- withdrawal of ventilation, dying patient, 454–5
- X-rays, *see* radiographs
- yellow-green chromatopsia, 447
- Young–Burgess system, 90–1