

INDEX

- Abnormalities of *Helminthostachys*, 52 (Fig. 344)
Actinostachys, 165
 Age and area (Willis), 332
Alsophila, 287, 293, 310; scales on prothallus of, 303; sorus of, 300
A. aculeata, anatomy of, 306
A. atrovirens, 298 (Fig. 561); young sorus of, 301 (Fig. 564)
A. blechnoides, 289 (Fig. 552)
A. excelsa, ontogenetic history of, 297; vascular system of, 299 (Fig. 562)
A. pruinata, 283
A. Williamsii, 298 (Fig. 561)
 "Amphiphloic siphonostele," 137
 Anachoropterideae, 6, 10, 34
Anachoropteris, synangial fusion of, 39
 Anatomy of *Alsophila aculeata*, 306; of *Asteropteris*, 37; of Cyatheaceae, 296; of *Cyathea Imrayana*, 298 (Fig. 560); of Dipteridaceae, 315; of Fossil Osmundaceae (Table), 141; of *Gleichenia*, 196; of Hymenophyllaceae, 239; of *Lophosoria*, 285; of Loxsomaceae, 254; of Marattiaceae, 101; of Marsileaceae, 180; of Matoniaceae, 222; of *Metaxya*, 289; of Osmundaceae, 133; of *Plagiogyria*, 277; of *Platyzoma*, 200; of Schizaeaceae, 157; of *Stromatopteris*, 200
Anemia, 153, 157 (Fig. 441), 175; axis of a young plant of, 159 (Fig. 443); comparison of, 174; leaf of, 156 (Fig. 440); sporangia of, 163
A. § *Anemidictyon*, 5
A. elegans, 166
A. Phyllitidis, petiole of, 161 (Fig. 445); sporangia of, 167 (Fig. 451)
A. (*Trochopteris*), 165; fertile leaf of, 162 (Fig. 446)
Anemidictyon, 162
Anemiorrhiza, 160; solenostele of, 172
Angiopteris, 95, 124; apex of stem of, 101 (Fig. 395); spermatozoids of, 120; stem of, 108; suspensor of, 120, 122 (Fig. 414); wings of leaves of, 101 (Fig. 396)
A. execta, sections of stem of, 102 (Fig. 397); sporangium of, 111 (Fig. 404); stem of, 98 (Fig. 391)
A. Teysmanniana, 95 (Fig. 388)
Ankyropteris, 17; vegetative structure of, 18
A. corrugata, dichotomy of, 18 (Fig. 318)
A. Grayi, leaf-trace of, 21; stele of, 20 (Fig. 321); stem of, 19 (Fig. 319)
A. westphalensis, petiole of, 26 (Fig. 326)
 Annulus (oblique) of *Plagiogyria*, 275
 "Antennae," 26, 37
 Antheridia, 4; of *Botrychium virginianum*, 76 (Fig. 376); of *Gleichenia laevigata*, 212 (Fig. 494); of *G. polypodoides*, 212 (Fig. 494); of *Helminthostachys*, 77 (Fig. 378)
 Apex of stem, leaf, or root, 2; of *Trichomanes radicans*, 237 (Fig. 507); of stem of *Angiopteris*, 101 (Fig. 395); of *Osmunda regalis*, 132 (Fig. 422)
 Aphlebia of *Diplolabis*, 19
 Aphlebiae, 25; of Coenopterids, 97; of *Hemitelia*, 305; of *Zygopteris Grayi*, 97
 "Aphlebioid" pinnae, 306 (Fig. 567)
 Appendages, dermal, 2, 307
Archangiopteris, 95, 124
 Archegonia, 4
 Archegonium, fertilised, of *Botrychium Lunaria*, 83 (Fig. 383); in *B. virginianum*, 77 (Fig. 377); of *Helminthostachys*, 77 (Fig. 378); in *Ophioglossum pendulum*, 80 (Fig. 380); of *Osmunda cinnamomea*, 146 (Fig. 435)
 Archetype of Ferns, 5, 40
 "Armature," 296; of Cyatheaceae, 215, 307
Asterochlaena, 17; stele of, 36
A. laxa, stele of, 23 (Fig. 323); stem of, 22 (Fig. 322)
Asteropteris, 17, 34; anatomy of, 37; stele of, 36
A. noveboracensis, stem of, 24 (Fig. 324)
Asterotheca, 100 (Fig. 393), 114 (Fig. 406)
 Axillary branches in Hymenophyllaceae, 237; buds in *Botrychium*, 54; buds of *Helminthostachys*, 54
Balantium, 263, 274
B. Culcita, 264 (Fig. 530)
 Basipetal sequence of sporangia, 249; succession, deviations from, 271; succession of sporangia of Cyatheaceae, 302
 Bifurcation of *Plagiogyria semicordata*, 275
 Blechnoids, 333
 "Borsten," 267
 "Borstenförmige Zellflächen," 267
Botrychioxylon, 25, 139; dichotomy of, 18
Botrychium, 42, 93; axillary buds in, 54; genus, 92; pinna-trace of, 64; root of, 55
B. daucifolium, development of sporangium in, 71 (Fig. 370); sporangium of, 72 (Fig. 371)
B. jenmani, venation of, 63 (Fig. 358)
B. Lunaria, 45 (Fig. 337), 68 (Fig. 364); fertilised archegonium and embryos of, 83 (Fig. 383); medullated stele of, 57 (Fig. 349); old embryo of, 83 (Fig. 384); stelar structure of, 58 (Fig. 350); sterile lamina of, 46 (Fig. 338)

- Botrychium (Sceptridium) obliquum*, suspensor of, 80, 81 (Figs. 381, 382)
B. simplex, 44 (Fig. 336)
B. ternatum, fertile spike of, 64 (Fig. 359)
B. virginianum, antheridium of, 76 (Fig. 376); archegonium in, 77 (Fig. 377); cotyledon of, 63 (Fig. 357); pinna-trace of, 64 (Fig. 360); prothallus of, 75 (Fig. 375); strands which supply the fertile spike in, 65 (Fig. 360*B*); venation of, 63 (Fig. 358)
- Botryopterideae, 6, 10, 228; comparison of, 15
- Botryopteris*, 10, 34; "equisetoid" hairs of, 13; vascular structure of, 12
B. antiqua, 13 (Fig. 314); sporangia of, 15 (Fig. 317)
B. cylindrica, 10 (Fig. 312), 11; branching of, 16; comparison of, 16; two forms of, 12
B. forensis, sporangia of, 14, 16 (Fig. 316)
B. hirsuta, young petiole of, 12 (Fig. 313)
B. ramosa, 10 (Fig. 311), 11
- Boweria*, sporangia of, 6, 7 (Fig. 310), 151
- Branching of *Botryopteris cylindrica*, 16; of leaf of *Gleichenia*, 193; (scheme of) leaf of *Mertensia*, 195 (Fig. 474)
- "Brevicide" sporangia, 308
- Bristles (stiff) of *Gleichenia*, 215
- Campteridinae, 319
- Camptopteris*, 319
- Cardiomanes*, 248, 250
- Caulopteris*, 104; trunk of, 100 (Fig. 394)
- Cellular constitution, 2; construction of *Osmundaceae*, 132
- Chaffy scales, 295
- Cheiroglossa*, 93
- Cheiropleuria*, 5; two-rowed segmentation of Fern sporangia of, 291 (Fig. 555)
- Christensenia*, 5, 95, 125; rhizome of, 96 (Fig. 390)
- Cibotium*, 263, 274
C. Barometz, 264 (Fig. 530); solenostelic rhizome of, 264 (Fig. 531); vascular system of stem of, 264 (Fig. 532)
C. Menziesii, sporangia of, 266 (Fig. 535)
C. Scheideri, marginal receptacle of, 265 (Fig. 534); young sorus of, 266 (Fig. 535)
- Clathropteris*, 319, 323
- Clepsydroid leaf-trace, 24, 25, 37
- Clepsydroideae, 19, 25, 38
- Clepsydroopsis*, 25, 27 (Fig. 325)
C. antiqua, foliar trace of, 28 (Fig. 328)
C. australis, stem of, 24
- Coenopteridaceae, 6, 8, 10 *sgg.* (ch. XVIII); aplebiae of, 97; comparison of, 34, 40; known only as fossils, 228; leaves of, 38; petioles, 36; sporangia of, 39; stems of, 35
- Comparison of *Anemia*, 174; of Coenopteridaceae, 40; of Cyatheaceae, 304, 309; of Dipteridaceae, 318; of Gleicheniaceae, 214; of Hymenophyllaceae, 248; of *Lophosoria*, 287; of Loxsomaceae, 257; of *Lygodium*, 174; of Marsileaceae, 186; of Matoniaceae, 226; of *Metaxya*, 292; of *Mohria*, 174; of *Plagiogyria*, 280; of *Schizaea*, 174; of Schizaeaceae, 172
- Coniopteris*, 267
- Corynopteris*, fertile pinna of, 113 (Fig. 405); sporangia of, 39
C. Essenghi, fertile pinna of, 33 (Fig. 334)
- Cotyledon of *Botrychium virginianum*, 63 (Fig. 357)
- Criteria of comparison, 1
- Cyathea*, 293, 310; cup-like indusium of, 300; venation of, 295 (Fig. 557)
C. dealbata, young sorus of, 302 (Fig. 565)
C. Imrayana, anatomy of, 298 (Fig. 560)
C. medullaris, bifurcated trunk of, 294 (Fig. 556); Tree-Fern, 293; vascular system of, 297 (Fig. 559)
C. sinuata, habit of, 295 (Fig. 557); sori of, 300 (Fig. 557)
- Cyatheaceae, 7, 293 *sgg.* (ch. XXXIII); anatomy of, 296; "armature" of, 215, 307; basipetal succession of sporangia of, 302; broad scales of, 307; comparison of, 304, 309; dichotomy of, 305; distinct from Dicksoniaceae, 326; gametophyte of, 303; gradate sorus of, 307; habit of, 293; indusium of, 307; leaf-trace of, 297; leaves of, 294; scales of, 296; sori and sporangia of, 299; sporangia of, 300, 301 (Fig. 563); sporangial segmentation of, 302; spore-output of, 301
- Danaea*, 95, 125; suspensor of, 120
D. alata, 96 (Fig. 389); fertile pinna of, 110 (Fig. 403); stelar system of, 103 (Fig. 398)
D. elliptica, partial septations of, 117 (Fig. 409)
- Danaeopsis*, 115, 124; spore-counts of, 115
D. fecunda, 116 (Fig. 408)
- Davallioids, 333
- Dendroid habit, 309
- Dennstaedtia*, 268, 274
D. apiifolia, sorus and sporangia of, 271 (Fig. 539)
D. dissecta, polycyclus in, 269 (Fig. 537)
D. rubiginosa, mixed sorus of, 271
- Dennstaedtiinae, habit of, 268; intra-marginal sori of, 273 (Fig. 541); sorus of, 270; sporangia of, 271; stelar structure of, 269 (Fig. 536)
- Dermal appendages, 2, 307; of Gleicheniaceae, 195, 215; of Marsileaceae, 187; of Matoniaceae, 221; of Schizaeaceae, 163
- Diacalpe*, 303, 309

- Dichotomised steles of *Stromatopteris*, 202 (Fig. 485)
- Dichotomy of *Ankyropteris corrugata*, 18 (Fig. 318); of *Botrychioxylon*, 18; of Cyatheaceae, 305; of *Dineuron*, 18; (equal) of *Gleichenia*, 214; of *G. linearis*, 193 (Fig. 472); of *Platyzoma*, 214
- Dicksonia*, 263, 274
D. antarctica, section of stem of, 265 (Fig. 533)
D. arborescens, 264 (Fig. 530)
- Dicksoniaceae, 260 *sqq.* (ch. XXIX), 268 *sqq.* (ch. XXX); distinct from Cyatheaceae, 326; sori of, 264 (Fig. 530)
- Dicksonieae, habit of, 263; sorus and sporangia of, 265; prothalli of, 267
- Dicranopteris*, 219; (= *Mertensia*), 193; spermatocytes of, 213
- Dictyophyllum*, 319, 323
D. exile, 321
D. rugosum, 321
- Dictyoxylic cylinder of *Osmunda*, 136
- Dineuroideae, 19, 38; of Sahni, 27
- Dineuron*, 27; dichotomy of, 18
- Diplolabis*, 17; aplebia of, 19
D. Römeri, ramification of, 19 (Fig. 320); stele of, 21
- Dipteridaceae, 311 *sqq.* (ch. XXXIV), 323; anatomy of, 315; comparisons of, 318; Fossil, 321; habit of, 311; Mesozoic, 318; sori and sporangia of, 316; spore-counts of, 321
- Dipteridinae, 7
- Dipteris*, 7, 311, 323; leaves of, 312 (Fig. 568); solenostele of, 315; *venatio anaxeti* in, 311
D. conjugata, habit of, 322 (Fig. 580); juvenile leaves of, 313 (Fig. 570); leaf of, 315 (Fig. 573); prothallus of, 318 (Fig. 577); young sorus of, 317 (Fig. 575)
D. Lobbiana, fertile segment of, 314 (Fig. 571); juvenile leaves of, 313 (Fig. 569); leaf-trace of, 316 (Fig. 574); sporangia of, 317 (Fig. 576)
D. quinquefurcata, leaf-segments of, 314 (Fig. 572)
- Dipteroids, 333
- Distribution of Gleicheniaceae, 328; of Marattiaceae, 331; of *Matonia-Dipteris*, 329; of Schizaeaceae, 330
- Dryopteroids, 333
- Embryology, 4, 80 *sqq.*; exoscopic, 120; in the genus *Ophioglossum*, 85; of *Botrychium Lunaria*, 83 (Fig. 383); of *Eu-Botrychium*, 80; of *Helminthostachys*, 84 (Fig. 385), with a suspensor, 82; of *Macroglossum*, 121 (Fig. 413); of Marattiaceae, 120, 121 (Fig. 412); of *Marattia Douglasii*, 122 (Fig. 415); of *Marsilea vestita*, 186 (Fig. 468); of Ophioglossaceae, 80; of *Ophioglossum moluccanum*, 86; of *O. vulgatum*, 84 (Fig. 386); of Osmundaceae, 146; of *Trichomanes*, 246
- Endodermis of *Helminthostachys zeylanica*, 56 (Fig. 347)
- “Equisetoid” hairs of *Botryopteris*, 13
- Etaopteris Lacattai*, sporangia of, 32 (Fig. 333)
- E. Scotti*, rachis of, 27 (Fig. 327)
- Eu-Botrychium*, 93; embryology of, 80
- Eu-Dicranopteris*, 218, 219
- Eu-Gleichenia*, 5, 193, 218, 219; broad scales of, 195; spermatocytes of, 213
- Eu-Ophioglossum*, 93
- Eu-Schizaea*, 165
- Eu-Todea*, 152
- Eusporangiate type, 3; Ferns, antiquity of, 8
- Exoscopic embryogeny, 120
- Feea*, 250
- Ferns, antiquity of Eusporangiate, 8; arche-type of, 5, 40; “filmy,” 234; geographical distribution of, 327; of Mesozoic Period, 8, 325; of Mount Ophir, 322; (primitive) general review of, 325
- Fertile leaf of *Anemia (Trochopteris) elegans*, 162 (Fig. 446); leaves of *Schizaea*, 154 (Fig. 437); pinna of *Corynepteris*, 113 (Fig. 405); pinna of *Danaea alata*, 110 (Fig. 403); pinnule of *Klukia exilis*, 169 (Fig. 453); segment of *Dipteris Lobbiana*, 314 (Fig. 571); segments of *Lygodium*, 163; spike, morphological nature of, 87
- “Filmy” Ferns, 234; habit, 237; habit of *Leptopteris*, 151
- Foliar trace of *Clepsydropsis antiqua*, 28 (Fig. 328)
- Fossil Coenopteridaceae, 228; Dipteridaceae, 321; Gleicheniaceae, 211; Marattiaceous, sori of, 113; Osmundaceae, anatomy of (Table), 141; Palaeozoic, 228
- Fungus, mycorrhizic, 75
- Gametangia of Marattiaceae, 120
- Gametophyte, saprophytic, 75; of Cyatheaceae, 303; of Gleicheniaceae, 212; of *Helminthostachys*, 78; of Hymenophyllaceae, 244; of Loxsomaceae, 256; of Marattiaceae, 118; of *Marattia fraxinea*, 118 (Fig. 410); of Ophioglossaceae, 75; of Osmundaceae, 145
- Gemmae, 245; of *Trichomanes alatum*, 246 (Fig. 517); of *T. Kaulfussii*, 246 (Fig. 517)
- Genera of Marattiaceae, 95; of Ophioglossaceae, 93
- Geographical distribution of Ferns, 327
- Gleichenia*, 7, 193, 219; anatomy of, 196; branching of leaf of, 193; equal dichotomy in, 214; juvenile leaves of, 194 (Fig. 473); mechanical deadlock in, 233; petiole of, 198 (Fig. 479); pinna-

INDEX

337

- trace of, 198 (Fig. 480); protosteles of, 196, 197 (Fig. 477); protostelic structure of, 215; spermatocytes of, 213; spore-counts of, 208; stiff bristles of, 215
- Gleichenia circinata*, sporangia of, 206 (Fig. 489), 207 (Fig. 490)
- G. dichotoma*, sori of, 206 (Fig. 489), 207 (Fig. 490); spore-mother-cells of, 209 (Fig. 491)
- G. flabellata*, flattened scale of, 216 (Fig. 495); rhizome of, 197 (Fig. 478); sori of, 196 (Fig. 476), 203, 207 (Fig. 490); sporangia of, 207 (Fig. 490); spore-mother-cells of, 209 (Fig. 491)
- G. laevigata*, antheridia of, 212 (Fig. 494)
- G. linearis*, dichotomy of, 193 (Fig. 472); hairs of, 195; pinnule of, 204 (Fig. 486); soral construction of, 199; sorus of, 203, 204 (Fig. 487)
- G. pectinata*, 282; hairs of, 195, 196 (Fig. 475); pinnule of, 205 (Fig. 488); solenostelic structure of, 199 (Fig. 481); solenostely of, 199, 215; soral construction of, 199; sorus of, 205 (Fig. 488)
- G. polypodioides*, antheridia of, 212 (Fig. 494)
- Gleicheniaceae, 7, 8, 193 *sgg.* (ch. XXIV); comparison of, 214–17; dermal appendages of, 195, 215; distribution of, 328; Fossil, 211; gametophyte of, 212; habit of, 193; phyletic grouping of, 217; sori of, 196, 216; spore-producing members of, 203
- Gradatae, 325
- Gradate sorus of Cyatheaceae, 307; sorus, principle of, 244; state, origin of, 304; type, 233
- Grammatopteris*, 10; protostele of, 139; stems of, 13
- Gymnogrammoids, 333
- Habit, dendroid, 309; “filmy,” 237; of Cyatheaceae, 293; of *Cyathea sinuata*, 295 (Fig. 557); of Dennstaedtiinae, 268; of Dicksonieae, 263; of Dipteridaceae, 311; of *Dipteris conjugata*, 322 (Fig. 580); of Hymenophyllaceae, 234; of *Hymenophyllum*, 236 (Fig. 505); of *Lophosoria quadripinnata*, 283; of Loxsomaceae, 252; of *Marsilea*, 176; of *M. quadrifolia*, 177 (Fig. 459); of Matoniaceae, 220; of *Matonia pectinata*, 220 (Fig. 496); of *Metaxya rostrata*, 288; of Ophioglossaceae, 42–7; of *Ophioglossum vulgatum*, 47 (Fig. 340); of Plagiogyriaceae, 275; of *Plagiogyria*, 276 (Fig. 542); of *Platyzoma*, 199; of *P. microphylla*, 201 (Fig. 482); of Protocyatheaaceae, 282; of *Regnellidium*, 178 (Fig. 460); of Schizaeaceae, 152 *sgg.* (ch. XXII); of *Stromatopteris*, 200; of *S. moniliformis*, 202 (Fig. 484); of Thyrsopterideae, 260; of *Trichomanes*, 235 (Fig. 504)
- Hairs of *Gleichenia linearis*, 195; of *G. pectinata*, 195, 196 (Fig. 475); of *Lophosoria*, 283; of Marattiaceae, 97; of Osmundaceae, 143; of *Platyzoma*, 196
- Hausmannia*, 323; sp., Fig. 579
- H. dichotoma*, 320 (Fig. 578)
- H. Forchammeri*, 321
- Hawlea*, 114 (Fig. 406)
- Helminthostachys*, 42, 93, 230; abnormalities of, 52 (Fig. 344); antheridium of, 77 (Fig. 378); archegonium of, 77 (Fig. 378); axillary buds of, 54; embryogeny of, with a suspensor, 82; embryos of, 84 (Fig. 385); gametophyte of, 78; leaf-insertion of, 53 (Fig. 346); leaf-trace of, 60 (Fig. 352); origin of spike of, 72; rhizome of, 49; root of, 55; single species of, 91; spike of, 49, 65 (Fig. 361); sporangia of, 74 (Fig. 374); sporangiophores of, 73 (Fig. 373), 74 (Fig. 374)
- H. zeylanica*, 68 (Fig. 364); endodermis of, 56 (Fig. 347); prothalli and sexual organs of, 77 (Fig. 378); rhizome of, 46 (Fig. 339); stele of, 59 (Fig. 351); vascular skeleton of, 60 (Fig. 353); young spike of, 73 (Fig. 372)
- Hemiphlebium*, 234
- Hemitelia*, 293, 310; “aphlebiae” of, 305; indusium of, 300; vein-fusion in, 305
- H. capensis*, segmentation of sporangia of, 302 (Fig. 566)
- H. Karsteniana*, venation of, 296 (Fig. 558)
- Heterophylly of *Platyzoma microphylla*, 201 (Fig. 482), 214
- Heterosporous character of Hydropterideae, 176
- Hooker Lecture, by Seward, 327
- Hydroglossum*, 162
- Hydropterideae, heterosporous character of, 176
- Hymenophyllaceae, 7, 234 *sgg.* (ch. XXVII); anatomy of, 239; axillary branches in, 237; comparison of, 248; gametophyte of, 244; habit of, 234; indusium of, 242, 249; soral estimates of, 244; sori and sporangia of, 240; spore-counts of, 243
- Hymenophyllites quadridactylites*, 7, 234
- Hymenophyllum*, 234, 250; habit of, 236 (Fig. 505)
- H. australe*, habit of, 236 (Fig. 505)
- H. cruentum*, habit of, 236 (Fig. 505)
- H. dilatatum*, habit of, 236 (Fig. 505); sex-organs of, 247 (Fig. 519); sporangia of, 241 (Fig. 512); transverse section of, 240 (Fig. 510)
- H. Weissii*, 7
- H. Wilsoni*, 241 (Fig. 513)

- Hypoderris*, 309
Hypolepis, 268, 274; sorus of, 273
- Indusia of Schizaeaceae, 173
 Indusial protection, 3
 Indusium (cup-like) of *Cyathea*, 300; of Cyatheaceae, 307; of *Hemitelia*, 300; of Hymenophyllaceae, 242, 249
 "Inversicatenal" type, 38
- Juvenile leaf of *Leptopteris*, 128 (Fig. 417); of *Dipteris conjugata*, 313 (Fig. 570); of *D. Lobbiana*, 313 (Fig. 569); of *Gleichenia*, 194 (Fig. 473); of *Marsilea*, 179 (Fig. 461); of *Osmunda regalis*, 127 (Fig. 416)
- Kaulfussia*, 95, 125
Kidstonia, 153; sporangia of, 151
K. heracleiensis, sporangium of, 169 (Fig. 454)
Klukia, 153, 168
K. exilis, fertile pinnule of, 169 (Fig. 453)
- Lacopteris*, 7, 224, 226
L. Woodwardi, 226 (Fig. 503)
- Leaf, apices of, 2; morphology of, 2; meristele of, 2; of *Anemia*, 156 (Fig. 440); sterile, of *Botrychium Lunaria*, 46 (Fig. 338); of *Dipteris conjugata*, 315 (Fig. 573); of *Loxsomopsis notabilis*, 252 (Fig. 520); of *Mohria caffrorum*, 155 (Fig. 439); of *Ophioglossum vulgatum*, 67 (Fig. 363); of *Osmunda Presliana*, 130 (Fig. 419); of *Thyrsopteris elegans*, 261 (Fig. 527); venation of, 2
- Leaf-construction, *Sphenopteris*-type of, 248
- Leaf-insertion of *Helminthostachys*, 53 (Fig. 346)
- Leaf-segments of *Dipteris quinquefurcata*, 314 (Fig. 572)
- Leaf-trace, Clepsydroid, 37; in *Ophioglossum vulgatum*, 60; of *Ankyropteris Grayi*, 21; of Cyatheaceae, 297; of *Dipteris Lobbiana*, 316 (Fig. 574); of *Helminthostachys*, 60 (Fig. 352); of Ophioglossaceae, 62; (small) of *Platyzoa*, 201 (Fig. 483); (undivided) of Schizaeaceae, 160; of *Thamnopteris Schlechtendalii*, 142 (Fig. 432); of Zygopterideae, 25
- Leaves of Coenopterids, 38; of Cyatheaceae, 294; of *Dipteris*, 312 (Fig. 568); of *Lygodium palmatum*, 155 (Fig. 438); of Marattiaceae, 97; of Schizaeaceae, 153; segmentation at the margin of, 238 (Fig. 508)
- Leptolepia*, 268, 274
Leptopteris, 126, 152; filmy habit of, 151; juvenile leaf of, 128 (Fig. 417)
 Leptosporangiate type, 3
 "Longicide" sporangia, 308
- "Looking-glass" images of sporangia, 308
Lophidium, 165
Lophosoria, 292; anatomy of, 285; comparison of, 287; hairs of, 283; pinnule of, 284 (Fig. 548); sorus of, 285 (Fig. 549); sporangia of, 287 (Fig. 551); stem of, 286 (Fig. 550); stock of, 284 (Fig. 547)
L. quadripinnata, habit of, 283
Loxsona, 252; meristele of, 254 (Fig. 523); prothallus of, 257 (Fig. 525)
L. Cunninghami, solenostele of, 254 (Fig. 522); sorus and sporangium of, 253 (Fig. 521)
 Loxsomaceae, 252 *sgg.* (ch. XXVIII); anatomy of, 254; comparison of, 257; gametophyte of, 256; habit of, 252; sorus and sporangia of, 255
Loxsomopsis, 252, 257
L. notabilis, leaf of, 252 (Fig. 520); sporangia of, 256 (Fig. 524)
Lygodium, 153, 157 (Fig. 441), 175; comparison of, 174; fertile segments of, 163; "monangial" sorus of, 164 (Fig. 447); rhizome of, 158 (Fig. 442); spermatocytes of, 172; sporangia of, 163
L. circinatum, sporangium of, 170 (Fig. 456)
L. §Hydroglossum, 5
L. japonicum, sporangia of, 167 (Fig. 451)
L. lanceolatum, sporangium of, 169 (Fig. 454)
L. palmatum, leaves of, 155 (Fig. 438)
L. scandens, rhizome of, 153 (Fig. 436)
- Macroglossum*, 95, 124; embryos of, 121 (Fig. 413)
- Maps of Fern distribution, A 328, B 329, C 330, D 331
- Marattia*, 95, 125
M. Douglasii, embryo of, 122 (Fig. 415); sex-organs of, 119 (Fig. 411)
M. fraxinea, gametophyte of, 118 (Fig. 410)
- Marattiaceae, 6, 8, 95 *sgg.* (ch. XX), 230; anatomy of, 101; comparative treatment of, 123; distribution of, 331; embryology of, 120; embryos of, 121 (Fig. 412); external characters of, 95; gametangia of, 120; gametophyte of, 118; hairs and scales of, 97; leaves of, 97; pinnae of, 99 (Fig. 392); roots of, 98; seven genera of, 95; fossil sori of, 113; sori and sporangia of, 109 (Fig. 402); spore-output of, 112; spore-producing members of, 108; stipules of, 96
- Marginal receptacle of *Cibotium Scheidei*, 265 (Fig. 534)
- Marginales, 8, 231, 282, 304, 309, 326
Marsilea, 176, 192; comparison with *Schizaea*, 191; habit of, 176; juvenile leaves of, 179 (Fig. 461); morphology of sporo-

- carp of, 188; spermatozoids of, 185; sporocarp, orientation of, 190
Marsilea polycarpa, sporocarp-trace of, 180 (Fig. 462)
M. quadrifolia, development of sporocarp of, 183 (Fig. 466); habit of, 177 (Fig. 459); "raphe" of the sporocarp of, 189 (Fig. 469); sporocarp of, 182 (Fig. 464)
M. salvatrix, megasporangia of, 182 (Fig. 465); microsporangia of, 182 (Fig. 465); prothallus of, 185 (Fig. 467); ripe sporocarp of, 182 (Fig. 465)
M. vestita, embryos of, 186 (Fig. 468)
 Marsileaceae, 176 *sqq.* (ch. XXIII); anatomy of, 180; comparison of, 186; dermal appendages of, 187; morphology of, 177; sporangium of, 184; sporocarp of, 188
Matonia, 7, 220; meristele of, 223; polycyclus of, 222; spore-counts of, 225
M.-Dipteris, distribution of, 329
M. pectinata, habit of, 220 (Fig. 496); prothallus and young plant of, 226 (Fig. 502); sorus of, 224 (Fig. 500); sporangia of, 224 (Fig. 501); stelar system of, 222 (Fig. 498); venation of, 221 (Fig. 497)
M. sarmentosa, stelar structure of, 223 (Fig. 499); venation of, 221 (Fig. 497)
 Matoniaceae, 220 *sqq.* (ch. XXV); anatomy of, 222; comparison of, 226; dermal appendages of, 221; habit of, 220; sporangia of, 225; spore-producing members of, 223
Matonidium, 7, 222, 226
 Matonineae, 7
Matteuccia, 309
 Mechanical deadlock in *Gleichenia*, 233
 Medullary system of vascular strands, 306
Megaphyton, 100, 104
 Megasporangia of *Marsilea salvatrix*, 182 (Fig. 465)
 Megaspore of *Pilularia*, 186 (Fig. 468)
 Meristele of leaf, 2; of *Loxsona*, 254 (Fig. 523); of *Matonia*, 223; of *Osmundites Carnieri*, 142 (Fig. 431); of *Stromatopteris*, 202 (Fig. 485); of Zygopterideae, 21
Mertensia, scheme of branching of leaf of, 195 (Fig. 474)
 Mesozoic Dipteridaceae, 318
 Mesozoic Period, Ferns of, 8, 325
Metaclepsydropsis, 17, 25, 34; rhizome of, 22
M. duplex, stele of, 21
Metaxya, 292, 298 (Fig. 561); anatomy of, 289; comparison of, 292; sori of, 290; sporangial development of, 291 (Fig. 555)
M. rostrata, habit of, 288; pinna of, 289 (Fig. 552); sporangial stalks of, 291 (Fig. 554); sporangium of, 290 (Fig. 553)
Microlepia, 268, 274
M. hirta, sorus of, 272 (Fig. 540)
M. speluncae, sorus of, 272 (Fig. 540)
 Microsporangia of *Marsilea salvatrix*, 182 (Fig. 465)
 "Mixed pith," 21, 25
 Mixed type, 233
Mohria, 153, 157 (Fig. 441), 175; comparison of, 174; scales of, 163; sporangia of, 163
M. caffrorum, leaf of, 155 (Fig. 439); sporangia of, 167 (Fig. 451); sporangium of, 166 (Fig. 450)
 "Monangial sorus," 163; of *Lygodium*, 164 (Fig. 447); of Schizaeaceae, 233
 Mount Ophir, Ferns of, 322
 Mycorrhiza, of Ophioglossaceae, 86
 Mycorrhizic fungus, 75
Oligocarpia, 7, 210; sorus of, 211 (Fig. 493)
 Ontogenetic history of *Alsophila excelsa*, 297
Onychopsis mantelli, 8
Ophioderma, 93
 Ophioglossaceae, 6, 8, 42 *sqq.* (ch. XIX), 229; comparative treatment of, 87; embryology of, 80; fertile spikes of, 66; gametophyte of, 75; genera of, 93; habit of, 42-7; leaf-traces of, 62; meristems of, 54; mycorrhiza of, 86; phyletic arrangement of, 91; sporangia of, 66, 74; stem structure of, 57; vascular anatomy of, 55; venation of, 43 (Fig. 335)
Ophioglossum, 42, 93; development of fertile spike in, 66; embryogeny in the genus, 85; genus, 92; prothallus of, 78; relatively late and derivative, 89; root of, 55; spike, pleiomery of, 90; sporangium of, 67; synangial state of, 110; variety in detail of, 49
O. Bergianum, 50 (Fig. 342); root of, 56 (Fig. 348); stock of, 61 (Fig. 354)
O. intermedium, 50, 53; degradation of, 90
O. moluccanum, embryogeny of, 86
O. palmatum, 68 (Fig. 364); distended sappy stem of, 89 (Fig. 385); spikes of, 50, 51 (Fig. 343), 52, 65 (Fig. 362); vascular system of, 61 (Fig. 355)
O. pendulum, archegonium in, 80 (Fig. 380); sex-organs of, 119 (Fig. 411); spikes of, 52 (Fig. 344), 69 (Figs. 366, 367); stock of, 62 (Fig. 356)
O. reticulatum, sporangium of, 70 (Fig. 368)
O. simplex, 50, 53 (Fig. 345); degradation of, 90
O. vulgatum, embryogeny of, 84 (Fig. 386); habit of, 47 (Fig. 340); leaf-trace in, 60; prothallus of, 79 (Fig. 379); shoot of, 48 (Fig. 341); sporogenous tissue of, 70 (Fig. 369); very young spike of, 68 (Fig. 365); young leaf of, 67 (Fig. 363); young sporeling of, 85 (Fig. 387)
 Orientation of *Marsilea* sporocarp, 190

- Osmunda*, 126, 152, 231; dictyoxyle cylinder of, 136; sporangia of, 129; spore-output of, 145; stele of, 137
- O. cinnamomea*, apex of young leaf of, 132 (Fig. 422); archegonium of, 146 (Fig. 435); stele of, 136 (Fig. 427)
- O. Presliana*, leaf of, 130
- O. regalis*, abnormal sporelings of, 129 (Fig. 418); adult leaf of, 130 (Fig. 419); apical region of, 132 (Fig. 422); juvenile leaves of, 127 (Fig. 416); root-tip of, 133 (Fig. 423); scheme showing xylem-ring of, 137 (Fig. 428); sporangium of, 145 (Fig. 434); sporophyll of, 131 (Fig. 420); zylem-strands of, 135 (Fig. 426)
- Osmundaceae, 6, 8, 126 *sgg.* (ch. XXI), 230; anatomy of, 133; cellular construction of, 132; comparison of, 147–52; comparison with *Plagiogyria*, 278; embryo of, 146; external characters of, 126; gametophyte of, 145; hairs of, 143; sporangia of, 6; spore-numbers of, 150; spore-producing members of, 143; sporophylls of, 150; stelar state of, 149; stele of, 135; “stipular leaf-bases” of, 127
- Osmundites Carnieri*, 140
- O. Dowkeri*, 140
- O. Dunlopi*, 140
- O. Gibbeana*, 140
- O. Kidstoni*, 139
- O. Kolbei*, 140
- O. skidegatensis*, 140
- Osmundopteris*, 93
- Palaeontology, facts of, 6
- Palaeozoic fossils, 228
- Pecopteris pennaeformis*, 168
- Peranema*, 309
- Petiole of *Anemia Phyllitidis*, 161 (Fig. 445); of *Ankyropteris westphalensis*, 26 (Fig. 326); of *Gleichenia*, 198 (Fig. 479); of *Stauropteris*, 28; of *S. oldhamia*, 29 (Fig. 329); of Coenopterids, 36
- Phyletic arrangement of Ophioglossaceae, 91; grouping of Gleicheniaceae, 217; scheme, 333; seriation, 8; slide, 39
- Phyllopodium, 2
- “Phyllosiphonic” state, 215
- Pilularia*, 176, 192; megaspore of, 186 (Fig. 468); sporophyte of, 176
- P. americana* compare with *Schizaea*, 185
- P. globulifera*, rhizome of, 177 (Fig. 458); sporocarp of, 181 (Fig. 463)
- Pinna, fertile, of *Corynepteris Essenghi*, 33 (Fig. 334); of *Metaxya rostrata*, 289 (Fig. 552); of *Schizaea*, 165 (Fig. 449)
- Pinna-nature of spike, 88
- Pinna-origin of fertile spike, 66
- Pinna-trace, 2; of *Botrychium*, 64; of *B. virginianum*, 64 (Fig. 360); of *Gleichenia*, 198 (Fig. 480)
- Pinnae, “aphleboid,” 306 (Fig. 567); of Marattiaceae, 99 (Fig. 392)
- Pinnule of *Gleichenia linearis*, 204 (Fig. 486); of *G. pectinata*, 205 (Fig. 488); of *Lophosoria*, 284 (Fig. 548); of *Todea pellucida*, 131 (Fig. 420)
- “Pith, mixed,” 21, 25
- Plagiogyria*, 275, 281; anatomy of, 277; comparison of, 280; comparison with Osmundaceae, 278; habit of, 276 (Fig. 542); mixed sorus of, 281; oblique annulus of, 275; pneumatophores of, 277 (Fig. 543); sorus and sporangium of, 279; sporangia of, 280 (Fig. 546)
- P. pycnophylla*, stem of, 278 (Fig. 544); stolons of, 275, 279 (Fig. 545)
- P. semicordata*, bifurcation of, 275
- Plagiogyriaceae, 275 *sgg.* (ch. xxxi), 281, 326; habit of, 275
- “Plaque élastique,” 168
- Platyzoma*, 193, 219; anatomy of, 200; dichotomous branching of, 214; habit of, 199; hairs of, 196; heterophylly in, 214; incipient heterospory of, 210; small leaf-traces of, 201 (Fig. 483); sporangia of, 209 (Fig. 492); spore-counts of, 210; transverse section of rhizome of, 201 (Fig. 483)
- Platyzomamicrophylla*, habit of, 201; heterophylly of, 201
- Pleiomery of *Ophioglossum* spike, 90
- Pneumatodes, 295
- Pneumatophores of *Plagiogyria*, 277 (Fig. 543)
- Polycyclic axial system of *Saccoloma*, 270 (Fig. 538)
- Polycycly in *Dennstaedtia dissecta*, 269 (Fig. 537); of *Matonia*, 222
- Polypodiaceae, 8
- Primitive Filicales, general review of, 325; phyletic scheme of, 333
- Prothallus of *Botrychium virginianum*, 75 (Fig. 375); of Dicksonieae, 267; of *Dipteris conjugata*, 318 (Fig. 577); of *Helminthostachys zeylanica*, 77 (Fig. 378); of *Loxsona*, 257 (Fig. 525); of *Marsilea salvatrix*, 185 (Fig. 467); of *Matonia pectinata*, 226 (Fig. 502); of *Ophioglossum*, 78; of *O. vulgatum*, 79 (Fig. 379); of *Schizaea pusilla*, 171 (Fig. 457); of *Thyrsopteris*, 263; of *Trichomanes rigidum*, 245 (Fig. 516); scales on, 303; vegetative features of, 4
- Protocyatheaceae, 282 *sgg.* (ch. xxxii), 292, 326; habit of, 282
- Protomarattia*, 95, 125
- Protostele, 2; of *Gleichenia*, 196, 197 (Fig. 477); 215; of *Grammatopteris*, 139
- Psaronius*, 100; sections of stems of, 106 (Fig. 400); stems (“starling-stones”) of, 104

- Psaronius brasiliensis*, stem of, 105 (Fig. 399)
- P. Renaultii*, stem of, 107 (Fig. 401)
- Pteridotheca*, 151
- Pteroids, 333
- Ptychocarpus*, 125; synangial sori of, 110
- P. unites*, 6; fructification of, 115 (Fig. 407); synangia of, 115 (Fig. 407)
- Ptychopteris*, 104
- Rachis of *Etaopteris Scotti*, 27 (Fig. 327)
- Ramifications of *Diplolabis Römeri*, 19 (Fig. 320); of *Stauropteris*, 30 (Fig. 331)
- “Ramular gap,” 135
- “Raphe” of the sporocarp of *Marsilea quadrifolia*, 189 (Fig. 469)
- Regnellidium*, 176, 192; habit of, 178 (Fig. 460)
- Rhizome of *Christensenia*, 96 (Fig. 390); of *Gleichenia flabellata*, 197 (Fig. 478); of *Helminthostachys*, 49; of *H. zeylanica*, 46 (Fig. 339); of *Lygodium*, 158 (Fig. 442); of *L. scandens*, 153 (Fig. 436); of *Metaclepysdropsis*, 22; of *Pilularia globulifera*, 177 (Fig. 458); (transverse section) of *Platyzoma*, 201 (Fig. 483); of *Thyrsopteris elegans*, 262 (Fig. 528)
- Rhynie Flora, 5
- Root apices of, 2; of *Botrychium*, 55; of *Helminthostachys*, 55; of *Ophioglossum*, 55; of *O. Bergianum*, 56
- Root-tip of *Osmunda regalis*, 133 (Fig. 423)
- Roots of Marattiaceae, 98 (Fig. 391)
- Ruffordia*, 169
- Saccoloma*, 268, 274; polycyclic axial system of, 270 (Fig. 538)
- S. elegans*, stock of, 270 (Fig. 538)
- Sagenopteris*, 191
- Sahni, Clepsidroidae, 25; Dineuroideae, 27
- Saprophytic gametophyte, 75
- Scales, broad, of Cyatheaceae, 307; broad, of *Eu-Gleichenia*, 195; of *Gl. flabellata*, 216 (Fig. 495); chaffy, 295; of Cyatheaceae, 296; of Marattiaceae, 97; of *Mohria*, 163; on prothallus, 303; on prothallus of *Alsophila*, 303
- Sceptridium*, 93
- Schizaea*, 153, 175; antheridia of, 171 (Fig. 457); archegonia of, 171 (Fig. 457); compare with *Pilularia americana*, 185; comparison of, 174; comparison with *Marsilea*, 191; fertile leaves of, 154 (Fig. 437); sporangia of, 163
- S. dichotoma*, stelar construction of, 159 (Fig. 444)
- S. digitata*, pinna of, 165 (Fig. 449)
- S. pennula*, sporangia of, 167 (Fig. 451)
- S. pusilla*, prothallus of, 171 (Fig. 457)
- S. rupestris*, origin of the sporangia of, 164 (Fig. 448); young fertile leaves of, 190 (Fig. 471); young leaves of, 164 (Fig. 448), 189 (Fig. 470)
- Schizaeaceae, 7, 8; anatomy of, 157; comparison of, 172; dermal appendages of, 163; distribution of, 330; habit of, 153 *sqg.* (ch. XXII); indusia of, 173; leaves of, 153; “monangial sorus” of, 233; sporangia of, 157 (Fig. 441), 167 (Fig. 451); spore-output of, 170; spore-producing members of, 163; undivided leaf-trace of, 160; venation of, 160; young sporangium of, 170 (Fig. 455)
- Scolecopteris*, 114 (Fig. 406)
- S. elegans*, 100 (Fig. 393)
- Secondary wood of Zygoterids, 25
- Section of stem of *Angiopteris evecta*, 102 (Fig. 397); of stem of *Psaronius*, 106 (Fig. 400); of stem of *Todea Fraseri*, 134 (Fig. 425)
- Segmentation at the margin of leaves, 238 (Fig. 508); of young sporangium, 308
- Senftenbergia*, 114 (Fig. 406), 153; sporangium of, 168 (Fig. 452)
- Senftenbergia elegans*, 7
- Seriation, phyletic, 8
- Seward, Hooker Lecture by, 327
- Sex-organs of *Hymenophyllum dilatatum*, 247 (Fig. 519); of *Marattia Douglasii*, 119 (Fig. 411); of *Ophioglossum pendulum*, 119 (Fig. 411); of *Trichomanes pyxidiferum*, 247 (Fig. 518)
- Sexual organs, 4; of *Helminthostachys zeylanica*, 77 (Fig. 378)
- Sheathing stipule, 67
- Shoot, simple, 2; of *Ophioglossum vulgatum*, 48 (Fig. 341)
- Simplices, 233, 325; general review of, 228 *sqg.* (ch. XXVI)
- “Siphonostele, amphiphloic,” 137
- Solenostele of *Anemiorrhiza*, 172; of *Dipteris*, 315; of *Loxsona Cunninghami*, 254 (Fig. 522)
- Solenostelic rhizome of *Cibotium Barometz*, 264 (Fig. 531); structure of *Gleichenia pectinata*, 199 (Fig. 481)
- Solenostely of *Gleichenia pectinata*, 199, 215
- Soral construction of *Gleichenia linearis*, 199; *G. pectinata*, 199; estimates of Hymenophyllaceae, 244; position, constant feature of, 310
- Sori of *Cyathea sinuata*, 300 (Fig. 557); of Cyatheaceae, 299; (intra-marginal) of Dennstaedtiinae, 273 (Fig. 541); of Dicksoniaceae, 264 (Fig. 530); of Dipteridaceae, 316; of *Gleichenia dichotoma*, 206 (Fig. 489), 207 (Fig. 490); of *G. flabellata*, 196 (Fig. 476), 207 (Fig. 490); of Gleicheniaceae, 196; of Hymenophyllaceae, 240; of Marattiaceae, 109 (Fig. 402); of Marattiaceous fossils, 113; of *Metaxya*, 290
- Sorus, comparison of, 3; individuality of, 3;

- marginal position of, 249; monangial, 163; monangial, of *Lygodium*, 164 (Fig. 447); of *Alsophila*, 300; of *Dennstaedtia apiifolia*, 271 (Fig. 539); (mixed) of *D. rubiginosa*, 271; of Dennstaedtiinae, 270; of Dicksonieae, 265; of *Gleichenia flabellata*, 203; of *G. linearis*, 203, 204 (Fig. 487); of *G. pectinata*, 205 (Fig. 488); of Gleicheniaceae, 216; of *Hypolepis*, 273; of *Lophosoria*, 285 (Fig. 549); of *Loxsona Cunninghamsi*, 253 (Fig. 521); of Loxsomaceae, 255; of *Matonia pectinata*, 224 (Fig. 500); of *Microlepia speluncae*, 272 (Fig. 540); of *Oligocarpia*, 211 (Fig. 493); of *Plagiogyria*, 279, (mixed) 281; of *Stachypteris Hallei*, 258 (Fig. 526); of *Thyrsopteris elegans*, 143 (Fig. 527); of *Trichomanes tenerum*, 241 (Fig. 511); simple, gradate, and mixed types of, 3
- Spermatocytes of *Dicranopteris*, 213; of *Eu-Gleichenia*, 213; of *Gleichenia*, 213; of *Lygodium*, 172
- Spermatozoids of *Angiopteris*, 120; of *Marsilea*, 185
- Sphenopteris*-type, 236; of leaf-construction, 248
- Spike, development of, 69; Fern-pinna theory of, 88; fertile, in *Ophioglossum*, 66; fertile, of *Botrychium ternatum*, 64 (Fig. 359); in *B. virginianum* 65 (Fig. 360 B); of *Helminthostachys*, 49, 65 (Fig. 361), 72; of *H. zeylanica*, 73 (Fig. 372); of *Ophioglossum vulgatum*, 68 (Fig. 365); of *O. pendulum*, 69 (Figs. 366, 367); pinna-nature of, 88; pinna-origin of fertile, 66
- Spikes, fertile, of Ophioglossaceae, 66; of *Ophioglossum palmatum*, 50, 51 (Fig. 343), 52 (Fig. 344), 65 (Fig. 362); of *O. pendulum*, 52 (Fig. 344)
- Sporangia, basipetal sequence of, 249; "brevicide," 308; development of, 71; "longicide," 308; "looking-glass" images of, 308; of *Anemia*, 163; of *A. Phyllitidis*, 167 (Fig. 451); of *Botryopteris forensis*, 14 (Fig. 316), 16; *B. antiqua*, 15 (Fig. 317); of *Boweria*, 6, 151; of *Cibotium Menziesii*, 266 (Fig. 535); of Coenopteridaceae, 39; of *Corynepteris*, 39; of Cyatheaceae, 299, 300, 301 (Fig. 563), 302; of *Dennstaedtia apiifolia*, 271 (Fig. 539); of Dennstaedtiinae, 271; of Dicksonieae, 265; of Dipteridaceae, 316; of *Dipteris Lobbiana*, 317 (Fig. 576); of *Etapteris Lacatzei*, 32 (Fig. 333); of *Gleichenia circinata*, 206 (Fig. 489), 207 (Fig. 490); of *G. flabellata*, 207 (Fig. 490); of *Helminthostachys*, 74 (Fig. 374); of Hymenophyllaceae, 240; of *Hymenophyllum dilatatum*, 241 (Fig. 512); of *Kidstonia*, 151; of *Lophosoria*, 287 (Fig. 551); of Loxsomaceae, 255; of *Loxsomopsis notabilis*, 256 (Fig. 524); of *Lygodium*, 163; of *L. japonicum*, 167 (Fig. 451); of Marattiaceae, 109 (Fig. 402); of *Matonia pectinata*, 224 (Fig. 501); of Matoniaceae, 225; of *Mohria*, 163; of *M. caffrorum*, 167 (Fig. 451); of Ophioglossaceae, 66, 74; of *Osmunda*, 129; of Osmundaceae, 6; of *Plagiogyria*, 280 (Fig. 546); of *Platyzoma*, 209 (Fig. 492); of *Schizaea*, 163; of *S. pennula*, 167 (Fig. 451); of Schizaeaceae, 157 (Fig. 441), 167 (Fig. 451); of *Stauropteris*, 39; of *S. oldhamia*, 31 (Fig. 332); of *Thyrsopteris elegans*, 262 (Fig. 529); of *Todea*, 129; of *T. barbara*, 143; of Zygopterids, 30; (origin of) of *Schizaea rupestris*, 164 (Fig. 448); segmentation of *Hemitelia capensis*, 302 (Fig. 566)
- Sporangial development of *Metaxya*, 291 (Fig. 555); segmentation of Cyatheaceae, 302; stalks of *Metaxya rostrata*, 291 (Fig. 554)
- Sporangiogenic band, 67
- Sporangiophores, 72; of *Helminthostachys*, 73 (Fig. 373), 74 (Fig. 374)
- Sporangium, characters of, 3; "distal face" of, 168; in *Botrychium daucifolium*, 71 (Fig. 370); of *Angiopteris evecta*, 111 (Fig. 404); of *Botrychium daucifolium*, 72 (Fig. 371); of *Kidstonia heraclensis*, 169 (Fig. 454); of *Loxsona Cunninghamsi*, 253 (Fig. 521); of *Lygodium circinatum*, 170 (Fig. 456); of *L. lanceolatum*, 169 (Fig. 454); of Marsileaceae, 184; of *Metaxya rostrata*, 290 (Fig. 553); of *Mohria caffrorum*, 166 (Fig. 450); of *Ophioglossum*, 67; of *O. reticulatum*, 70 (Fig. 368); of *Osmunda regalis*, 145 (Fig. 434); of *Plagiogyria*, 279; of *Senftenbergia*, 168 (Fig. 452); of *Todea barbara*, 131 (Fig. 421)
- Spore-counts of *Danaeopsis*, 115; of Dipteridaceae, 321; of *Gleichenia*, 208; of Hymenophyllaceae, 243; of *Matonia*, 225; of *Platyzoma*, 210
- Sporelings, abnormal, of *Osmunda regalis*, 129 (Fig. 418)
- Spore-mother-cells of *Gleichenia dichotoma*, 209 (Fig. 491); of *G. flabellata*, 209 (Fig. 491)
- Spore-numbers of Osmundaceae, 150
- Spore-output, 3; of Cyatheaceae, 301; of Marattiaceae, 112; of *Osmunda*, 145; of Schizaeaceae, 170; of *Todea*, 145
- Spore-producing members of Gleicheniaceae, 203; of Marattiaceae, 108; of Matoniaceae, 223; of Osmundaceae, 143; of Schizaeaceae, 163

- Sporocarp of *Marsilea quadrifolia*, 182 (Fig. 464), 183 (Fig. 466); of *M. salva-trix*, 182 (Fig. 465); of *Pilularia globulifera*, 181 (Fig. 463)
- Sporocarp-trace of *Marsilea polycarpa*, 180 (Fig. 462)
- Sporogenous tissue of *Ophioglossum vulgatum*, 70 (Fig. 369)
- Sporophyll of *Osmunda regalis*, 131 (Fig. 420); of Osmundaceae, 150
- Stachypteris*, 258
S. Hallei, sorus of, 258 (Fig. 526)
- Stauropteris*, 17, 229, 232; petiole of, 28; ramifications of, 30 (Fig. 331); sporangia of, 39
S. oldhamia, 10, 28; petiole of, 29 (Fig. 329); sporangia of, 31 (Fig. 332); vascular bundle of, 29 (Fig. 330)
- Stelar construction of *Schizaea dichotoma*, 159 (Fig. 444); of Osmundaceae, 149; of *Botrychium Lunaria*, 58 (Fig. 350); of Dennstaedtiinae, 269 (Fig. 536); of *Matonia sarmentosa*, 223 (Fig. 499); system of *Danaea alata*, 103 (Fig. 398); system of *Matonia pectinata*, 222 (Fig. 498)
- Stele of *Ankyropteris Grayi*, 20 (Fig. 321); of *Asterochlaena*, 36; of *A. laxa*, 23 (Fig. 323); of *Asteropteris*, 36; of *Botrychium Lunaria*, 57 (Fig. 349); of *Diplolabis Römeri*, 21; of *Helminthostachys zeylanica*, 59 (Fig. 351); of *Metaclepsydropsis duplex*, 21; of *Osmunda*, 137; of *O. cinnamomea*, 136 (Fig. 427); of Osmundaceae, 135; of *Thamnopteris Schlechtendalii*, 139 (Fig. 430); of *Todea barbara*, 135 (Fig. 426); of *Trichomanes scandens*, 239 (Fig. 509)
- Stellate stele of Zygopterideae, 23
- Stem, apices of, 2; of *Angiopteris*, 108; of *A. evecta*, 98 (Fig. 391); of *Ankyropteris Grayi*, 19 (Fig. 319); of *Asterochlaena laxa*, 22 (Fig. 322); of *Asteropteris noveboracensis*, 24 (Fig. 324); of *Clepsydropsis australis*, 24; of *Lophosoria*, 286 (Fig. 550); of *Plagiogyria pycnophylla*, 278 (Fig. 544); of *Psaronius brasiliensis*, 105 (Fig. 399); of *P. Renaulti*, 107 (Fig. 401); of *Thamnopteris Schlechtendalii*, 138 (Fig. 429); (section) of *Dicksonia antarctica*, 265 (Fig. 533); structure of Ophioglossaceae, 57
- Stems of Coenopteridaceae, 35; of *Grammatopteris*, 13; of *Tubicaulis*, 13; ("starling-stones") of *Psaronius*, 104
- "Stipular leaf-bases" of Osmundaceae, 127
- Stipule, sheathing, 67; of Marattiaceae, 96
- Stock of *Lophosoria*, 284 (Fig. 547); of *Ophioglossum Bergianum*, 61 (Fig. 354); of *O. pendulum*, 62 (Fig. 356); of *Saccoloma elegans*, 270 (Fig. 538)
- Stolons of *Plagiogyria pycnophylla*, 275, 279 (Fig. 545)
- Stromatopteris*, 193, 219; anatomy of, 200; dichotomised steles of, 202 (Fig. 485); habit of, 200; meristele of, 202 (Fig. 485)
S. moniliformis, habit of, 202 (Fig. 484)
- Superficiales, 8, 231, 282, 304, 309, 326
- Suspensor, 4; of *Angiopteris*, 120, 122 (Fig. 414); of *Botrychium obliquum*, 80; of *Danaea*, 120
- Synangia of *Ptychocarpus unitus*, 115 (Fig. 407)
- Synangial fusion of *Anachoropteris*, 39; sori of *Ptychocarpus*, 110; of *Ophioglossum*, 110
- Taenitis*, 300
- Tempskya*, 106
T. Knowltoni, 169
- Thamnopteris*, 126
T. Schlechtendalii, leaf-trace of, 142 (Fig. 432); part of stele of, 139 (Fig. 430); stem of, 138 (Fig. 429)
- Thaumatopteris*, 319, 323
T. schenki, 321
- Thyrsopterideae, habit of, 260
- Thyrsopteris*, 257, 260, 274; prothallus of, 263
T. elegans, 260; leaf and sorus of, 261 (Fig. 527); rhizome of, 262 (Fig. 528); sporangia of, 262 (Fig. 529)
- Tietea singularis*, 106
- Todea*, 126, 152, 231; sporangia of, 129; spore-output of, 145
T. barbara, scheme showing xylem-ring of, 137 (Fig. 428); sporangium of, 131 (Fig. 421), 143; stele of, 135 (Fig. 426); young sporangia of, 133 (Fig. 424), 144 (Fig. 433)
T. Fraseri, section of stem of, 134 (Fig. 425)
T. pellucida, pinnule of, 131 (Fig. 420)
- Transverse section of *Hymenophyllum dilatatum*, 240 (Fig. 510)
- Tree-Fern, vascular ontogeny of, 299
- Tree-Fern-habit, 305
- Tree-Ferns, 263, 293
- Trichomanes*, 234, 250; embryo of, 246; habit of, 235 (Fig. 504)
T. alatum, gemmae of, 246 (Fig. 517)
T. Goebelianum, 237 (Fig. 506)
T. Kaulfussii, gemmae of, 246 (Fig. 517)
T. Lyallii, habit of, 235 (Fig. 504)
T. membranaceum, habit of, 235 (Fig. 504)
T. Motleyi, 239
T. muscoides, 239
T. pyxidiferum, sex-organs of, 247 (Fig. 518)
T. radicans, apex of, 237 (Fig. 507)
T. rigidum, prothallus of, 245 (Fig. 516)
T. scandens, stele of, 239 (Fig. 509)
T. speciosum, young receptacle of, 242 (Fig. 514)

- Trichomanes spicatum*, habit of, 255 (Fig. 504)
T. tenerum, sorus of, 241 (Fig. 511)
Trichopteris, 300
Trochopteris, 162
 Trunk of *Caulopteris*, 100 (Fig. 394); (bifurcated) of *Cyathea medullaris*, 294 (Fig. 556)
Tubicaulis, 10; stems of, 13
T. solenites, 14 (Fig. 315)
 Type, Gradate, 233; Mixed, 233
- Vascular anatomy of Ophioglossaceae, 55; bundle of *Stauropteris oldhamia*, 29 (Fig. 330); ontogeny of Tree-Fern, 299; skeleton of *Helminthostachys zeylanica*, 60 (Fig. 353); strands, medullary system of, 306; structure, 2; structure of *Botryopteris*, 12; system of *Alsophila excelsa*, 299 (Fig. 562); system of *Ophioglossum palmatum*, 61 (Fig. 355); system of, *Cyathea medullaris*, 297 (Fig. 559); system of stem of *Cibotium Barometz*, 264 (Fig. 532)
- Vegetative features of prothallus, 4; structure of *Ankyropteris*, 18
- Vein-fusion in *Hemitelia*, 305
- Venation of *Botrychium jenmani*, 63 (Fig. 358); *B. virginianum*, 63 (Fig. 358); of Cyathea, 295 (Fig. 557); of *Hemitelia Karsteniana*, 296 (Fig. 558); of leaf, 2; of *Matonia pectinata*, 221 (Fig. 497); of *M. sarmentosa*, 221 (Fig. 497); of Ophioglossaceae, 43 (Fig. 335); of Schizaeaceae, 160
- Wings of leaves of *Angiopteris*, 101 (Fig. 396)
Woodsia, 303
- Xylem-ring of *Osmunda regalis*, 137 (Fig. 428); of *Todea barbara*, 137 (Fig. 428)
 Xylem-strands of *Osmunda regalis*, 135 (Fig. 426)
- Young fertile leaves of *Schizaea rupestris*, 190 (Fig. 471); leaf of *Osmunda cinnamomea*, 132 (Fig. 422); leaves of *Schizaea rupestris*, 164 (Fig. 448), 189 (Fig. 470); receptacle of *Trichomanes speciosum*, 242 (Fig. 514); sorus of *Alsophila atrovirens*, 301 (Fig. 564); sorus of *Cibotium Scheidei*, 266 (Fig. 535); sorus of *Cyathea dealbata*, 302 (Fig. 565); sorus of *Dipteris conjugata*, 317 (Fig. 575); sorus of *Hymenophyllum Wilsoni*, 241 (Fig. 513); sporangia of *Todea barbara*, 133 (Fig. 424), 144 (Fig. 433); sporangium of Schizaeaceae, 170 (Fig. 455); sporangium, segmentation of, 308; sporeling of *Ophioglossum vulgatum*, 85 (Fig. 387)
- Zalesskya*, 126, 139
 "Zellflächen, borstenförmige," 267
- Zygopterideae, 6, 10, 17, 229; age of, 17; leaf-trace of, 25; meristele of, 21; stellate stele of, 23
- Zygopterids, secondary wood of, 25; sporangia of, 30
- Zygopteris Grayi*, apophyses of, 97