

## I N D E X.

## A.

## ACID.

- Acetic*. Composition; and relation to that of aldehyde, 279, 280.
- Benzoic*. Composition, and relation to that of oil of bitter almonds, 279, 280. Converted into hippuric acid in the human body, 150, 325.
- Carbonic*. Is the form in which the inspired oxygen and the carbon of the food are given out, 13. Its formation in the body the chief source of animal heat, 17—22. Occurs combined with potash and soda, in the serum of the blood, 41. Formed by the action of oxygen on the products of the metamorphosis of the tissues, 60. Its formation may also be connected with the production of fat from starch, 85—91. Generated by putrefaction of food in the stomach of animals, 115. Also by the fermentation of bad wine in man, when it causes death by penetrating into the lungs, 116. Escapes through both skin and lungs, *ib.* Produced, along with urea, by the oxidation of uric acid, 140. Produced, with several other compounds, by the oxidation of blood, *ib.* May be formed, along with choleic acid, from hippuric acid, starch, and oxygen, 152. Also, along with choleic acid, urea, and ammonia, by the action of water and oxygen on starch and proteine, *ib.* Produced, along with fat and urea, from proteine, by the action of water and oxygen, in the absence of soda, 154. Combines with the compound of

## ACID.

- iron present in venous blood, and is given off when oxygen is absorbed, 269. Is absorbed by the serum of blood in all states, 270.
- Cerebric*. Its composition, 184. Its properties, 186.
  - Choleic*. Represents the organic portion of the bile, 133. Its formula, 134. Its transformations, 125. Half its formula, added to that of urate of ammonia, is equal to the formula of blood + a little oxygen and water, 136. Produced in the oxidation of blood, 140. Views which may be taken of its composition, 148. May be formed by the action of oxygen and water on proteine and starch, 152. Products of its oxidation, 154. Various ways in which it may be supposed to be formed in the body, 160. Its composition, 315. Cannot be said to exist ready formed in the bile, 317.
  - Cholic*. Its composition, 318. Derived from choleic acid, 134, 135. Possible relation to choleic acid, 148.
  - Choloidic*. Its composition, 315. Derived from choleic acid, 135. Possible relation to choleic acid, 148. Possible relation to starch, 157. Possible relation to proteine, 141.
  - Cyanic*. Its formula, 281.
  - Cyanuric*. Its formula, 281.
  - Hippuric*. Its composition, 319. Appears in the urine of stall-fed animals, 82. Is destroyed by exercise, 82, 139. Is probably formed in the oxidation of blood, 140. Is found in the human urine after benzoic acid has been administered, 150, 325. May be derived from proteine when acted on by oxygen and uric acid, 151. With starch and oxygen, it may produce choleic and carbonic acids, 152. May be derived from the oxidation of choleic acid, 154.
  - Hydrocyanic* or *Prussic*. Its poisonous action explained, 274.
  - Lithofellic*. Its composition, 322. Probably derived from the oxidation of choleic acid: is the chief constituent of bezoar stones, 154.
  - Lactic*. Its composition, 309. Its origin, 111. Does not exist in the healthy gastric juice, 112.
  - Margaric*. Exists in bile, 317.
  - Muriatic*. Exists in the free state in the gastric juice, 109, 112. Is derived from common salt, 112, 161.
  - Oxaluric*. Analysis of, 321.

## ACID.

- Parabanic*. Analysis of, 321.
- Phosphoric*. Exists in the urine of the carnivora in considerable quantity, 78, 163. Its proportion very small in that of the graminivora, 79. Derived from the phosphorus of the tissues, 78. It is retained in the body to form bones and nervous matter, 80.
- Sulphuric*. Exists in the urine of the carnivora, 78, 163. Derived from the sulphur of the tissues, 78.
- Uric*. Its composition, 318. Products of its oxidation, alloxan, oxalic acid, carbonic acid, urea, &c., 137, 140. Is probably derived, along with choleic acid, by the action of oxygen and water on blood or muscle, 136. Disappears almost entirely in the system of man and of the higher animals, 55, 137. Appears as calculus, when there is a deficiency of oxygen, 137. Never occurs in pthysical cases, *ib*. Yields mulberry calculus when the quantity of oxygen is somewhat increased, but only urea and carbonic acid with a full supply of oxygen, *ib*. Uric acid calculus promoted by the use of fat and of certain wines, 139. Unknown on the Rhine, *ib*. Uric acid and urea, how related to allantoine, 141; to gelatine, 142. Forms the greater part of the urine of serpents, 54. Yields, with the elements of proteine and oxygen, hippuric acid and urea, 151. How related to taurine, 155, 156. Calculi of it never occur in wild carnivora, but often in men who use little animal food, 146.
- AFFINITY, Chemical.** Is the ultimate cause of the vital phenomena, 9, 10. Is active only in the case of contact, and depends much on the order in which the particles are arranged, 205. Its equilibrium renders a compound liable to transformations, 207. In producing the vital phenomena, it is modified by other forces, 209. It is not alone the vital force or vitality, but is exerted in subordination to that force, 232.
- AIR.** Introduced into the stomach during digestion with the saliva, 113. Effects of its temperature and density, dryness, &c., in respiration, 15, 16.
- ALBUMEN.** Animal and vegetable albumen identical, 47, 48. Their composition, 293, 294, 308, 309. Vegetable albumen, how obtained, 45. Is a compound of proteine, and in organic composition identical with fibrine and caseine, 47, 104, 106.

- Exists in the yolk as well as the white of eggs, 107. Also in the serum of the blood, 41. Is the true starting point of all the animal tissues, 107, 108.
- ALCOHOL.** Is hurtful to carnivorous savages, 179. Its mode of action: checks the change of matter, 239. In cold climates serves as an element of respiration, 22.
- ALDEHYDE.** Its composition; how related to that of acetic acid, 279, 280.
- ALKALIES.** Mineral alkalies essential both to vegetable and animal life, 164. Vegetable alkalies all contain nitrogen, all act on the nervous system, and are all poisonous in a moderate dose, 177, 182. Theory of their action: they take a share in the transformation or production of nervous matter, for which they are adapted by their composition, 182—189. Action of caustic alkalies on bile, or choleic acid, 134.
- ALLANTOINE.** Is found in the urine of the foetal calf. How derived from proteine. How related to uric acid and urea, 141. How related to choleic acid, 148. Its composition, 319.
- ALLEN and PEPYS.** Their calculation of the amount of inspired oxygen, 283.
- ALLOXAN.** Formed by the oxidation of uric acid, 137. Converted by oxidation into oxalic acid and urea, oxaluric and parabanic acids, or carbonic acid and urea, *ib.* How related to taurine, 156. Seems to act as a diuretic. Recommended for experiment in hepatic diseases, *ib. (note)*.
- ALMONDS, BITTER.** Oil of. Its composition; how related to benzoic acid, 280.
- AMMONIA.** Combined with uric acid it forms the urine of serpents, birds, &c., 54. Its relation to choleic, choloidic, and cholic acids, 135. Is one of the products which may be formed by the oxidation of blood, 140; or of proteine, 152. Its relation to uric acid, urea, and taurine, 155. To allantoin and taurine, 155, 156. To alloxan and taurine, 156. To choleic and choloidic acid and taurine, 158. To urea, water, and carbonic acid, 159. Is found in combination with acids in the urine of the carnivora, 163.
- ANALYSIS.** Of dry blood, 283, 314. Of dried flesh, 314. Of fæces, 285. Of black bread, *ib.* Of potatoes, *ib.* Of peas, *ib.* Of beans, *ib.* Of lentils, *ib.* Of fresh meat, *ib.* Of moist bread, *ib.* Of moist potatoes, *ib.* Of the fibrine and

albumen of blood, 293, 310, 311. Of vegetable fibrine and albumen, vegetable caseine and gluten, 294, 295. Of animal caseine, 295. Of starch, 296, 297. Of grape or starch sugar, 297. Of sugar of milk, 298. Of gum, *ib.* Of oats, *ib.* Of hay, 299. Of fat, 300. Of cane-sugar, *ib.* Of cholesterine, *ib.* Of wax, 307. Of cyanic acid, cyanuric acid, and cyamelide, 308. Of aldehyde, metaldehyde, and elaldehyde, 307. Of proteine, 308. Of albumen from the yolk and white of egg, *ib.* Of lactic acid, 309. Of gas from the stomach of cows after eating to excess, *ib.* Of gas from stomach and intestines of executed criminals, *ib.* Of gelatinous tissues, 311. Of tissues containing chondrine, 312. Of arterial membrane, *ib.* Of horny tissues, *ib.* Of the lining membrane of the egg, 313. Of feathers, *ib.* Of the pigmentum nigrum, *ib.* Of choleic acid, 315. Of taurine, *ib.* Of choloidic acid, *ib.* Of cholic acid, 318. Of uric acid, *ib.* Of alloxan, 319. Of urea, *ib.* Of hippuric acid, *ib.* Of allantoin, *ib.* Of xanthic oxide, 320. Of cystic oxide, *ib.* Of oxalic acid, 320. Of oxaluric acid, *ib.* Of parabanic acid, *ib.* Of roasted flesh, 322. Of lithofellic acid, *ib.* Of solanine, 323. Of picrotoxine, *ib.* Of quinine, *ib.* Of morphia, 324. Of caffeine, theine, or guaranine, *ib.* Of theobromine, *ib.* Of asparagine, 325.

**ANIMAL HEAT.** Derived from the combination of oxygen with the carbon and hydrogen of the metamorphosed tissues, which proceed ultimately from the food, 17, 18. Is highest in those animals whose respiration is most active, 19. Is the same in man in all climates, 19, 20. Is kept up by the food in proportion to amount of external cooling, 22. Is not produced either by any direct influence of the nerves, or by muscular contractions, 29—34. Its amount in man, 34. Chemical action the sole source of it, 38. The formation of fat from starch or sugar must produce heat, 91, 94. The elements of the bile, by combining with oxygen, serve chiefly to produce it, 61.

**ANIMAL LIFE.** Distinguished from vegetable life by the absorption of oxygen, and the production of carbonic acid, 2. Must not be confounded with consciousness, 6, 7. Conditions necessary to animal life, 9, 12. Depends on an equilibrium between waste and supply, 245, 254, 265.

**ANTISEPTICS.** They act by putting a stop to fermentation, putre-

faction, or other forms of metamorphosis, 170. Their action on wounds and ulcers, 121.

**ARTERIES.** Composition of their tunica media, 312. How derived from proteine, 126.

**ARTERIAL BLOOD.** Conveys oxygen to every part of the body, 60, 269. Contains a compound of iron, most probably peroxide, 269. Yields oxygen in passing through the capillaries, 60, 271. Contains carbonic acid dissolved or combined with soda, 272.

**ASPARAGINE.** Its composition, 325. Its relation to taurine and bile, 180. Theory of its action on the bile, 181.

**ASSIMILATION.** In animals it is independent of external influences, 3. Depends on the presence in the blood of compounds of proteine, such as fibrine, albumen, or caseine, 40, 106. Is more energetic in the young than in the adult animal, 67. Is also more energetic in the herbivora than in the carnivora, 81.

**ATMOSPHERE.** See **AIR**.

**AZOTISED PRODUCTS.** Of vegetable life, 45, 176—182. Of the metamorphosis of tissues. Necessary for the formation of bile in the herbivora, 158. In man, 166, 168. May be replaced by azotised vegetable compounds, 169—170. Theory of this, 177—182. Of the transformation of the bile, or of choleic acid; how related to the constituents of urine, 155.

## B.

**BEANS.** Composition of, 285.

**BEER.** Forms part of the diet of soldiers in Germany, 286, 288.

**BEEES.** Their power of forming wax from honey, 301—306.

**BENZOIC ACID.** See **ACID**, *Benzoic*.

**BERTHOLLET.** His analysis of oxalic acid, 321.

**BERZELIUS.** His analysis of potato starch, 297; of sugar of milk, 298; of gum, *ib.*; of cane sugar, 300.

**BEZOAR STONES.** See **ACID**, *Lithofellic*.

**BLANCHET.** His analysis of solanine, 323.

**BILE.** In the carnivora is a product of the metamorphosis of the tissues, along with urate of ammonia, 136. May be represented by choleate of soda, with which, however, it is not identical, 317. Products of its transformation, 135, 317. Remarks on these, 315—318. Origin of bile, 61, 144. Starch, &c., contribute to its formation in the herbivora, 146—

- 150, 159, 160, 166. Soda essential to it, 154, 162—164. Relation of bile to urine, 156. To starch, 157. To fibrine, 136. To caffeine, &c., asparagine, and theobromine, 180. For the acid substances derived from bile, choleic, choloidic, and cholic acids, see **Acid**, choleic, &c. Yields taurine, 135. Contains cholesterine, 85, 317. Also stearic and margaric acids, 317. Its function : to support respiration and produce animal heat by presenting carbon and hydrogen in a very soluble form to the oxygen of the arterial blood, 61—64. Amount secreted by the dog, the horse, and man, 64. It returns entirely into the circulation, and disappears completely, 60—66.
- BLOOD.** The fluid from which every part of the body is formed, 8. Its chief constituents, 40. How formed from vegetable food, 45. Can only be formed from compounds of proteine, 48. Is therefore entirely derived from vegetable products in the herbivora, and indirectly also by the carnivora, which feed on the flesh of the former, 48. Its composition identical with that of flesh, 133. Analysis of both, 314. The secretions contain all the elements of the blood, 132. Its relation to bile and urine, 136. Products of the oxidation of blood, 140. Excess of azotised food produces fulness of blood and disease, 145. Soda is present in the blood, 161—164. Important properties of the blood, 171—175. Venous blood contains iron, probably as protoxide; arterial blood, probably as peroxide, 271, 273. Theory of the poisonous action of sulphuretted hydrogen and Prussic acid : they decompose the compound of iron in the blood, 274. The blood, in analogous morbid states, ought to be chemically examined, 275.
- BLOOD-LETTING.** Theory of its mode of action, 258. It may produce opposite effects in different cases, 264.
- BÆCKMANN.** His analysis of black bread, 285; of potatoes, *ib.*; of dry beef, 314; of dry blood, *ib.*; of roasted flesh, 322.
- BONES.** Phosphoric acid of the food retained to assist in forming them, 80. Gelatine of bones digested by dogs, 97. See, further, **GELATINE.** Cause of brittleness in bones, 99.
- BOUSSINGAULT.** His analysis of potatoes, 285. His comparison of the food and excretions in the horse and cow, Table, 290. His analysis of gluten, 294; of vegetable albumen, *ib.*; of vegetable caseine, 295; of oats, 298; of hay, 299.

- BRACONNOT.** On the presence of lactic acid in gastric juice, 112 ;  
 of iron in the gastric juice of the dog, 113.
- BRAIN.** See **ACID**, *Cerebric*, and **NERVOUS MATTER**.
- BREAD.** Analysis of, 285.
- BRUNN.** His analysis of sugar of milk, 298.
- BUCKWHEAT.** Analysis of starch from, 296.
- BURDACH.** His statement of the amount of bile secreted by animals, 64.
- BUTTER.** Forms a part of the food of soldiers in Germany, 286, 288.
- BUZZARD.** Its excrements consist of urate of ammonia, 54.

## C.

- CAFFEINE.** Identical with theine, 179. Its relation to taurine and bile, 180. Theory of its mode of action, 181. Its composition, 324.
- CANE SUGAR.** Its composition, 300.
- CARBON.** Is accumulated in the bile, 61. Is given off as carbonic acid, 13. Excess of carbon causes hepatic diseases, 24. By combining with oxygen, it yields the greater part of the animal heat. See **ANIMAL HEAT**, **BILE**, and **ACID**, *Carbonic*. Amount of carbon oxidised daily in the body of a man, 14. Calculations on which this statement is founded, 284—289. Amount consumed by the horse and cow, 14. Different proportions of carbon in different kinds of food, 17. Carbon of flesh compared with that of starch, shewing the advantage of a mixed diet, 76. Calculation on which this statement is founded, 299. Amount of carbon in dry blood calculated, 283. Amount in the food of prisoners calculated, 293.
- CARBONIC ACID.** See **ACID**, *Carbonic*.
- CARBONATES.** They occur in the blood, 41.
- CALCULUS**, Mulberry. Derived from the imperfect oxidation of uric acid, 137. Uric acid calculus is formed in consequence of deficiency of inspired oxygen, or excess of carbon in the food, 137. See **ACID**, *Uric*. Bezoar stones composed of lithofellic acid, 154.
- CARNIVORA.** Their nutrition the most simple, 44. It is ultimately derived from vegetables, 48, 49. Their young, like graminivora, require non-azotised compounds in their food, 50. Their bile is formed from the metamorphosis of their tissues,



- 59, 61. The process of assimilation in adult and young carnivora compared, 67. Their urine, 78. The assimilative process in adult carnivora less energetic than in graminivora, 80. They are destitute of fat, 82. They swallow less air with their food than graminivora, 118. Concretions of uric acid are never found in them, 146. Both soda and ammonia found in their urine, 163.
- CASEINE.** One of the azotised nutritious products of vegetable life, 47. Abundant in leguminous plants, 47. Identical in organic composition with fibrine and albumen, 47, 48. Animal caseine found in milk and cheese: identical with vegetable caseine, 51. Furnishes blood to the young animal, 52. Is one of the plastic elements of nutrition, 96. Yields proteine, 105, 106. Its relation to proteine, 126. It contains sulphur, *ib.* Potash essential to its production, 164. Contains more of the earth of bones than blood does, 52. Its analysis, 295.
- CEREBRIC ACID.** See **ACID, Cerebric.**
- CHANGE OF MATTER.** See **METAMORPHOSIS OF TISSUES.**
- CHEMICAL ATTRACTION.** See **AFFINITY.**
- CHEVREUL.** His researches on fat, 84. His analysis of fat, 300; of cholesterine, *ib.*
- CHLORIDE OF SODIUM.** See **COMMON SALT.**
- CHOLEIC ACID.** See **ACID, Choleic.**
- CHOLESTERINE.** See **BILE.**
- CHOLIC ACID.** See **ACID, Cholic.**
- CHOLOIDIC ACID.** See **ACID, Choloidic.**
- CHONDRINE.** Its relation to proteine, 126. Analysis of tissues containing it, 312.
- CHRONIC DISEASES.** The action of inspired oxygen is the cause of death in them, 27, 28.
- CHYLE.** When it has reached the thoracic duct, it is alkaline, and contains albumen coagulable by heat, 145.
- CHYME.** It is formed independently of the vital force, by a chemical transformation, 108. The substance which causes this transformation is derived from the living membrane of the stomach, 109. Chyme is acid, 145.
- CLOTHING.** Warm clothing is a substitute for food to a certain extent, 22. Want of clothing accelerates the rate of cooling, and the respirations, and thus increases the appetite, *ib.*
- COLD.** Increases the appetite by accelerating the respiration, 22.

Is most judiciously employed as a remedy in cerebral inflammation, 261.

CONCRETIONS. See CALCULUS, and ACID, *Uric*; also ACID, *Lithofellic*.

CONSTITUENTS, *Azotised*. Of blood: see FIBRINE and ALBUMEN. Of vegetables: see FIBRINE, Vegetable; ALBUMEN, Vegetable; CASEINE, Vegetable; ALKALIES, Vegetable; and CAFFEINE. Of bile: see ACID, *Choleic*, *Cholic*, and *Choloidic*. Of urine: see ACID, *Uric*; UREA, and ALLANTOINE.

COOLING. See COLD and CLOTHING.

COURBE. His analysis of cholesterine, 300.

COW. Amount of carbon expired by the, 14. Comparison of the food with the excretions of the cow, 291.

CRUM. His analysis of cane sugar, 300.

CULTIVATION. Is the economy of force, 78.

CYAMELIDE. Its formula, 280.

CYANIC ACID. See ACID, *Cyanic*.

CYANIDE OF IRON. Its remarkable properties, 269.

CYANURIC ACID. See ACID, *Cyanuric*.

## D.

DAVY. Oxygen consumed by an adult man, 283.

DEATH. Cause of, in chronic diseases, 27, 28. Caused in old people by a slight depression of temperature, 255. Definition of it, 254.

DEMARÇAY. His analysis of choleic acid, choloidic acid, and taurine, 315. Remarks on his Researches on Bile, 316.

DENIS. His experiments on the conversion of fibrine into albumen, 42.

DESPREZ. His calculation of the heat developed in the combustion of carbon, 34.

DIABETES MELLITUS. The sugar found in the urine in this disease is grape sugar, and is derived from the starch of the food, 95.

DIASTASE. Analogy between its solvent action on starch, and that of the gastric juice on coagulated albumen, 111.

DIFFUSION OF GASES. Explains the fact that nitrogen is given out through the skin of animals, 118; and the poisonous action of feather-white wine, 116.

DIGESTION. Is effected without the aid of the vital force, by a

metamorphosis derived from the transformation of a substance proceeding from the lining membrane of the stomach, 109. The oxygen introduced with the saliva assists in the process, 113. Lactic acid has no share in it, 111, 112.

**DISEASE.** Theory of, 254 *et seq.* Cause of death in chronic disease, 27. Disease of liver caused by excess of carbon or deficiency of oxygen, 23. Prevails in hot weather, 24.

**DOG.** Amount of bile secreted by, 64. Digests the gelatine of bones, 97. His excrements contain only bone earth, 98. Concretion of urate of ammonia said to have been found by Lassaigne in a dog, doubtful, 146 (*note*).

**DUMAS.** His analysis of choleic acid, 315; of choloidic acid, *ib.*; of taurine, *ib.*; of cholic acid, 318; of hippuric acid, 319.

## E.

**EGGS.** Albumen of the white and of the yolk identical, 107. Analysis of both, 308; of lining membrane, 313. The fat of the yolk may contribute to the formation of nervous matter, 108. This fat contains iron, 107.

**ELALDEHYDE.** See ALDEHYDE.

**ELEMENTS.** Of nutrition, 96. Of respiration, *ib.*

**EMPYREUMATICS.** They check transformations, 170. Their action on ulcers, 121.

**EQUILIBRIUM.** Between waste and supply of matter is the abstract state of health, 245, 258. Transformations occur in compounds in which the chemical forces are in unstable equilibrium, 109.

**ETTLING.** His analysis of wax, 307. **ETTLING and WILL.** Their analysis of lithofellic acid, 322.

**EXCREMENTS.** Contain little or no bile in man and in the herbivora, none at all in the dog and other carnivora, 64. Those of the dog are phosphate of lime, 98. Those of serpents are urate of ammonia, 54. Those of birds also contain that salt, 54. Those of the horse and cow compared with their food, 290, 291.

**EXCRETIONS.** Contain, with the secretions, the elements of the blood or of the tissues, 132—136. Those of the horse and cow compared with their food, 290, 291. Bile is not an excretion, 63.

## F.

FÆCES. Analysis of, 285.

FAT. Theory of its production from starch, when oxygen is deficient, 83 *et seq.*; from other substances, 86. The formation of fat supplies a new source of oxygen, 89; and produces heat, 90 *et seq.* Maximum of fat, how obtained, 94. Carnivora have no fat, 82. Fat in stall-fed animals, 89. Occurs in some diseases in the blood, 95. Fat in the women of the East, 99. Composition compared with that of sugar, 84, 85. Analysis of fat, 300. Disappears in starvation, 25. Is an element of respiration, 96.

FATTENING OF ANIMALS. See FAT.

FEATHERWHITE WINE. Its poisonous action, 116.

FEBRILE PAROXYSM. Definition of, 256.

FEHLING. His analysis of metaldehyde and elaldehyde, 307.

FERMENTATION. May be produced by any azotised matter in a state of decomposition, 120. Is arrested by empyreumatics, *ib.* Is analogous to digestion, 119.

FEVER. Theory and definition of, 256.

FIBRE. Muscular. See Flesh.

FIBRINE. Is an element of nutrition, 96. Animal and vegetable fibrine are identical, 45. Is a compound of proteine, 105. Its relation to proteine, 126. Convertible into albumen, 42. Is derived from albumen during incubation, 107. Its analysis, 293, 294, 311. Vegetable fibrine, how obtained, 45, 46.

FISHES. Yield phosphuretted hydrogen, 191 (*note*).

FLESH. Consists chiefly of fibrine, but, from the mixture of fat and membrane, has the same formula as blood, 133. Analysis of flesh, 314, 322. Amount of carbon in flesh compared with that of starch, 77, 299.

FOOD. Must contain both elements of nutrition and elements of respiration, 96. Nutritious food, strictly speaking, is that alone which is capable of forming blood, 40. Whether derived from animals or from vegetables, nutritious food contains proteine, 44, 106 *et seq.* Changes which the food undergoes in the organism of the carnivora, 53 *et seq.* The food of the herbivora always contains starch, sugar, &c., 70. Food, how dissolved, 108 *et seq.* Azotised food has no direct influence on the formation of uric acid calculus, 138. Effects of super-

abundant azotised food, 145, 146. Non-azotised food contributes to the formation of bile, and thus to respiration, 147 *et seq.* Salt must be added to the food of herbivora, in order to yield soda for the bile, 162. Caffeine, &c., serve as food for the liver, 188. The vegetable alkalies may be viewed as food for the organs which form the nervous matter, 189. Amount of food consumed by soldiers in Germany, 286. Its analysis, 284. Food of the horse and cow compared with their excretions, 290, 291.

FORMULÆ. Explanation of their use, 280. How reduced to 100 parts, 281. Formulæ of albumen, fibrine, caseine, and animal tissues, 126. Formula of proteine, 121; of blood and flesh, 133; of fat, 85; of cholesterine, 85; of aldehyde, acetic acid, oil of bitter almonds, and benzoic acid, 280; of cyamelide, cyanic acid, and cyanuric acid, 280; of choleic acid, 134; of choloidic acid and cholic acid, 135; of gelatine, 142; of hippuric acid, 150; of lithofellic acid, 154; of taurine, 155; of alloxan, 156. See ANALYSIS.

FRANCIS. His analysis of picrotoxine, 323.

FREMY. Lameyran and Frémy. Their analysis of gas from the abdomen of cows after excess in fresh food, 309. His researches on the brain, 43, 184.

FREQUENCY of the pulse and respiration in different animals, 19, 292.

FRUITS. Contain very little carbon, and hence are adapted for food in hot climates, 17.

## G.

GAS. Analysis of gas from abdomen of cows after excess in fresh food, 115, 309. Analysis of gas from the stomach and intestines of executed criminals, 115, 309.

GASTRIC JUICE. Contains no solvent but a substance in a state of metamorphosis, by the presence of which the food is dissolved, 109. Contains free acid, *ib.* Contains no lactic acid, 112. In the dog has been found to contain iron, 113. See DIGESTION, CHYME, FOOD.

GAY-LUSSAC and THENARD. Their analysis of starch, 297; of sugar of milk, and of gum, 298; of cane sugar, 300; of wax, 307; of oxalic acid, 321.

GELATINE. Is derived from proteine, but is no longer a com-

- pound of proteine, and cannot form blood, 127 *et seq.* May serve as food for the gelatinous tissues, and thus spare the stomach of convalescents, 98, 130. In starvation the gelatinous tissues remain intact, 97. Its relation to proteine, 126. Its formula, 142. Its analysis, 311, 322.
- GOEBEL. His analysis of gum, 298.
- GLOBULES of the blood are the carriers of oxygen to all parts of the body, 171—175. They contain iron, 265 *et seq.*
- GLUTEN. Contains vegetable fibrine, 46. Analysis of it, 295.
- GMELIN. On the sugar of bile, 147.
- GOOSE. How fattened to the utmost, 94.
- GRAMINIVORA. See HERBIVORA.
- GRAPE-SUGAR. An element of respiration, 96. Is identical with starch sugar and diabetic sugar, 72. Its composition, 73. Its analysis, 297.
- GROWTH, or increase of mass, greater in graminivora than in carnivora, 80. Depends on the blood, 40; and on compounds of proteine, 106. See NUTRITION.
- GUM. An element of respiration, 96. Its composition, 73. Is related to sugar of milk, *ib.* Its analysis, 298.
- GUNDLACH. His researches on the formation of wax from honey by the bee, 301.

## H.

- HAIR. Analysis of, 312. Its relation to proteine, 126. Analysis of proteine from hair, 308.
- HAY. Analysis of, 299.
- HEPATIC DISEASES. Cause of, 23.
- HERBIVORA. Their blood derived from compounds of proteine in their food, 48. But they require also for their support non-azotised substances, 70. These last assist in the formation of their bile, 147 *et seq.* They retain the phosphoric acid of their food to form bone and nervous matter, 80. Their urine contains very little phosphoric acid, 79. The energy of vegetative life in them is very great, 81. They become fat when stall-fed, 82.
- HESS. His analysis of wax, 307.
- HYBERNATING ANIMALS. Their fat disappears during the winter sleep, 25. They secrete bile and urine during the same period, 61.

- HIPPURIC ACID.** See **ACID**, *Hippuric*.
- HORN.** Analysis of, 312. Contains proteine; its relation to proteine, 126. Analysis of proteine from horn, 308.
- HORSE.** Amount of carbon expired by, 14. Comparison of his food with his excretions, 290. Force exerted by a horse in mechanical motion compared to that exerted by a whale, 337.
- HYDROCYANIC ACID.** See **ACID**, *Hydrocyanic*.
- HYDROGEN.** By combining with oxygen contributes to produce the animal heat, 25.

## I.

- ICE.** Is judiciously employed as a remedy in cerebral inflammation, 261.
- INORGANIC constituents of albumen, fibrine, and caseine,** 41, 121, 126.
- JOBST.** His analysis of theine, 324.
- JONES, Dr. Bence.** His analysis of vegetable fibrine, 294; of vegetable albumen, *ib.*; of vegetable caseine, 295; of gluten, *ib.*; of the albumen of yolk of egg, 308, 310; of the albumen of brain, 310.
- IRON.** Is an essential constituent of the globules of the blood, 265 *et seq.* Is found in the fat of yolk of egg, 107. Also in the gastric juice of the dog, 113. Singular properties of its compounds, 268.
- ISOMERIC BODIES.** 103, 280.

## K.

- KELLER.** His researches on the conversion of benzoic acid into hippuric acid in the human body, 325.
- KIDNEYS.** They separate from the arterial blood the nitrogenised compounds destined for excretion.

## L.

- LACTIC ACID.** See **ACID**, *Lactic*.
- LAVOISIER.** His calculation of the amount of inspired oxygen, 12, 283.
- LEHMANN.** On the presence of lactic acid in gastric juice, 112.
- LIEBIG.** His analysis of sugar of milk, 298; of cane sugar, 300; of aldehyde, 307; of uric acid, 318; of hippuric acid, 319; of quinine, 323; of morphia, 324; of asparagine, 325.

His calculation of the carbon daily expired as carbonic acid, 14, 284. Table, 289. His remarks on Demarçay's researches on bile, 315—318.

LIEBIG AND PFAFF. Their analysis of caffeine, 324.

LIEBIG AND WÖHLER. Their analysis of alloxan, 319; of urea, *ib.*; of allantoine, *ib.*; of xanthic oxide, 320; of oxaluric acid, 321; of parabanic acid, *ib.*

LENTILS. Contain vegetable caseine, 47. Analysis of, 284, 285. Form part of the diet of soldiers in Germany, 287. Table, 289.

LIGHT. Its influence on vegetable life analogous to that of heat on animal life, 233.

LIME. Phosphate of. See BONES.

LIVER. It separates from the venous blood the carbonised constituents destined for respiration, 58. Diseases of the liver, how produced, 23. Accumulation of fat in the liver of the goose, 95.

## M.

MAIZE. Analysis of starch from, 297.

MARCHAND. On the amount of urea in the urine of the dog when fed on sugar, 61. His analysis of cholesterine, 300.

MARÇET. His analysis of gluten, 294.

MARTIUS. His analysis of guaranine, 324.

MECHANICAL EFFECTS. See MOTION.

MEDICINE. Definition of the objects of, 257 *et seq.* Action of medicinal agents, 170 *et seq.*

MENZIES. His calculation of the amount of inspired oxygen, 12, 283.

METALDEHYDE. See ALDEHYDE.

METAMORPHOSIS OF TISSUES. 103 *et seq.* In other parts of the volume, *passim.*

MILK. Is the only natural product perfectly fitted to sustain life, 51. Contains caseine, *ib.* Fat (butter), *ib.* Sugar of milk, *ib.* Earth of bones, 52. And potash, 164.

MORPHIA. Contains less nitrogen than quinine, 177. Its analysis, 324.

MITSCHERLICH. His analysis of uric acid, 318; of hippuric acid, 319.

MOMENTUM. Of force, 202. Of motion, *ib.*



- MOTION.** Phenomena of motion in the animal body, 196 *et seq.*  
Different sources of motion 199. Momentum of motion, 202.  
Motion propagated by nerves, 219. Voluntary and involuntary motions accompanied by a change of form and structure in living parts, 220. Motion derived from change of matter, 221 *et seq.* The cause of motion in the animal body is a peculiar force, 232. The sum of the effects of motion in the body proportional to the amount of nitrogen in the urine, 245.
- MULBERRY CALCULUS.** See **CALCULUS.**
- MULDER.** Discovered proteine, 105. His analysis of fibrine of blood, 293. Of animal caseine, 296. Of proteine, 308. Of fibrine, 311. Of gelatine, *ib.* Of chondrine, 312.
- MUSCLE.** See **FLESH.**
- MUSCULAR FIBRE.** Its transformation depends on the amount of force expended in producing motion, 220.

## N.

- NERVES.** Are the conductors of the vital force, and of mechanical effects, 219. Effects of the disturbance of their conducting power, 229. They are not the source of animal heat, 29.
- NERVOUS LIFE.** Distinguished from vegetative, 38.
- NERVOUS MATTER.** Contains albumen, and fatty matter of a peculiar kind, 43. Vegetables cannot produce it, 50. The fat of yolk of egg probably contributes to its formation, 108. The phosphoric acid and phosphates, formed in the metamorphosis of the tissues of the herbivora, are retained to assist in the formation of nervous matter, 80. The vegetable alkalies affect the nervous system, 182—184. Composition of cerebriic acid, 184. Theory of the action of the vegetable alkalies, 185.
- NITROGEN.** Essential to all organised structures, 42, 43. Substances in the body which are destitute of it not organised, 43. Abounds in nutritious vegetables, 45. Nutritious forms in which it occurs, *ib. et seq.* Occurs in all vegetable poisons, 177; also in a few substances which are neither nutritious nor poisonous, but have a peculiar effect on the system, such as caffeine, 177 *et seq.*
- NITROGENISED.** See **AZOTISED.**
- NON-AZOTISED.** Constituents of food. See **STARCH.**
- NUTRITION.** Depends on the blood, 40. On Albumen, fibrine, or caseine, 40 *et seq.* Elements of nutrition, 96. Compounds

of proteine alone are nutritious, 106. Occurs when the vital force is more powerful than the opposing chemical forces, 198. Theory of it, 210. Is almost unlimited in plants from the absence of nerves, 212. Depends on the momentum of force in each part, 227. Depends also on heat, 243.

## O.

- OATS.** Amount required to keep a horse in good condition, 74. Analysis of, 298.
- OIL OF BITTER ALMONDS.** Its composition. How related to benzoic acid, 279, 280.
- OLD AGE.** Characteristics of, 248 *et seq.*
- OPPERMANN.** His analysis of wax, 307.
- ORGANS.** The food of animals always consists of parts of organs, 2. All organs in the body contain nitrogen, 42, 43. There must exist organs for the production of nervous matter, 189; and the vegetable alkalies may be viewed as food for these organs, *ib.*
- ORGANISED TISSUES.** All contain nitrogen, 42, 43. All such as are destined for effecting the change of matter are full of small vessels, 223. Their composition, 126. The gelatinous and cellular tissues, and the uterus, not being destined for that purpose, are differently constructed, 224. Waste of organised tissues rapid in carnivora, 76.
- ORIGIN.** Of animal heat, 17, 31. Of fat, 81 *et seq.* Of the nitrogen exhaled from the lungs, 114 *et seq.* Of gelatine, 127 *et seq.*, 143. Of uric acid and urea, 135 *et seq.* Of bile, 135, 143, 146 *et seq.*, 159. Of hippuric acid, 150, 325. Of the chief secretions and excretions, 152. Of the soda of the bile, 161 *et seq.* Of the nitrogen in bile, 168. Of nervous matter, 183 *et seq.*
- ORTIGOSA.** His analysis of starch, 297.
- OXALIC ACID.** A product, along with urea, of the partial oxidation of uric acid, occurring in the form of mulberry calculus, 137. Its analysis, 321.
- OXYGEN.** Amount consumed by man daily, 12, 283. Amount consumed daily in oxidising carbon by the horse and cow, 14. The absorption of oxygen characterizes animal life, 2. The action of oxygen is the cause of death in starvation and in chronic diseases, 25—28. The amount of oxygen inspired

varies with the temperature, dryness, and density of the air, 16. Is carried by arterial blood to all parts of the body, 171. Fat differs from sugar and starch only in the amount of oxygen, 84. It also contains less oxygen than albumen, fibrine, &c., 86. The formation of fat depends on a deficiency of oxygen, 88 *et seq.*; and helps to supply this deficiency, 89. Oxygen essential to digestion, 113. Relation of oxygen to some of the tissues formed from proteine, 126. Oxygen and water, added to blood or to flesh, yield the elements of bile and of urine, 135. Action of oxygen on uric acid, 136, 139; on hippuric acid, 82, 139; on blood, 140; on proteine, with uric acid, 151; on proteine and starch, with water, 152; on choleic acid, 154; on proteine, with water, 154. By depriving starch of oxygen and water, choloidic acid may be formed, 157. Oxygen is essential to the change of matter, 173. Its action on the azotised constituents of plants when separated, 213. Its action on the muscular fibre essential to the production of force, 220—226. Oxygen is absorbed by hibernating animals, 241. Is the cause of the waste of matter, 243; and of animal heat, 244, 252. Blood-letting acts by diminishing the amount of oxygen which acts on the body, 258. Its absorption is the cause of the change of colour from venous to arterial blood, 265. The globules probably contain oxide of iron, protoxide in venous blood, peroxide in arterial, 267 *et seq.* All parts of the arterial blood contain oxygen, 173, 174, 266, 271.

## P.

- PEARS. Analysis of starch from unripe, 297.
- PEAS. Form part of the diet of soldiers in Germany, 287, 289. Abound in vegetable caseine, 47. Analysis of peas, 285; of starch from peas, 296.
- PEPYS and ALLEN. Their calculation of the amount of inspired oxygen, 283.
- PEROXIDE OF IRON. Probably exists in arterial blood, 267 *et seq.*
- PFLÜGER. His analysis of the gas obtained by puncture from the abdomen of cattle after excess in green food, 309.
- PHENOMENA of motion in the animal body, 195 *et seq.*
- PHOSPHATES. See BONES.
- PHOSPHORIC ACID. See ACID, *Phosphoric*.
- PHOSPHORUS. Exists in albumen and fibrine, 41, 48, 126. It is

not known in what form, 121 *et seq.* Is an essential constituent of nervous matter, 184, 190.

**PHOSPHURETTED HYDROGEN.** Occurs among the products of the putrefaction of fishes, 190, 191.

**PICROTOXINE.** Contains nitrogen, 177 (*note*). Its analysis, 323.

**PLANTS.** Distinguished from animals by fixing carbon and giving out oxygen, 2, 213; by the want of nerves and of locomotive powers, 3. Their capacity of growth almost unlimited, 212. Cause of death in plants, 214.

**PLAYFAIR, DR. L.** His formula for blood, 113. His analysis of fæces, of peas, of lentils, of beans, 285; of flesh and of blood, 314; of roasted flesh, 322.

**POISONS, VEGETABLE.** Always contain nitrogen; 176 *et seq.* Different kinds of poisons, 170. Theory of the action of Prussic acid and sulphuretted hydrogen, 274.

**POLYMERIC BODIES,** 103.

**POTASH.** Essential to the production of caseine or milk, 164.

**POTATOES.** Amount of carbon in, 287. They form part of the diet of soldiers in Germany, *ib.* Analysis of, 285; of starch from, 297; of solanine from the buds of germinating potatoes, 323.

**PREVOST AND DUMAS.** On the frequency of the pulse and respirations, 292.

**PRODUCTS.** Of the metamorphosis of tissues found in the bile and urine, 132. Of the action of muriatic acid on bile, 133. Of the action of potash on bile, 134. Of the action of water and oxygen on blood or fibre, 136. Of the oxidation of uric acid, 137. Of the oxidation of blood, 140. Of the action of water on proteine, 141. Of the action of urea on lactic and benzoic acids, 150. Of oxygen and uric acid on proteine, 151. Of oxygen on starch and hippuric acid, 152. Of oxygen and water on proteine and starch, 153. Of oxygen and water on proteine when soda is absent, 154. Of the separation of oxygen from starch, 157. Of the action of water on urea, 159. Of the action of water and oxygen on caffeine or theine, asparagine, and theobromine, 180.

**PROTEINE.** Discovered by Mulder, 105. Its composition, *ib.* Produced alone by vegetables, 106. Is the source of all the organic azotised constituents of the body, 107. Its formula, 121. Its relation to fibrine, albumen, caseine, and all the

animal tissues, 126. Gelatine no longer yields it, although formed from it, 129. Its relation to bile and urine, 136. Its relation to allantoine and choloidic acid, 141; to gelatine, 142; to hippuric acid, 151; to the chief secretions and excretions, 152, 153; to fat, 154. Analysis of proteine from the crystalline lens, from albumen, from fibrine, from hair, from horn, from vegetable albumen and fibrine, from cheese, 308.

PROUT. His analysis of starch, 297; of grape sugar from honey, *ib.*; of sugar of milk, 298; of cane sugar, 300; of urea, 319. His discovery of free muriatic acid in the gastric juice, 112. On the effect of fat food on the urine, 139.

PRUSSIC ACID. See ACID, *Hydrocyanic*.

PULMONARY DISEASES. Arise from excess of oxygen, 23. Prevail in winter, 24.

PULSE. Its frequency in different animals, 292.

PUTREFACTION. Is a process of transformation, 109. Membranes very liable to it, 110. Effects of the putrefaction of green food in the stomach of animals, 115. Is analogous to digestion, 119. Putrefying animal matters cause the fermentation of sugar, 120. Is checked by empyreumatics, 121, 170.

## Q.

QUININE. Contains nitrogen, 177. Its analysis, 323.

## R.

REGNAULT. His analysis of morphia, 324.

REPRODUCTION OF TISSUES. See NUTRITION.

REPRODUCTION OF THE SPECIES, 39.

RHENISH WINES. Contain so much tartar, that their use prevents the formation of uric acid calculus, 139.

RESPIRATION. Theory of, 265 *et seq.* Its connection with the food and with the animal heat, 12 *et seq.*

## S.

SALT, Common. Essential to the formation of bile in the herbivora, and to that of gastric juice, 161 *et seq.*

SAUSSURE, DE. His analysis of grape sugar and of starch sugar, 297; of wax, 307.

SCHERER, DR. JOS. His analysis of albumen from serum of blood, 293; of fibrine of blood, *ib.*; of vegetable fibrine, 294; of vegetable caseine, 295; of animal caseine, *ib.*; of proteine from

different sources, 308; of albumen from white of egg, *ib.*; of albumen from different sources, 310; of fibrine, 311; of gelatine from different sources, *ib.*; of tissues containing chondrine, 312; of the tunica media of arteries, *ib.*; of horny tissues, *ib.*; of the lining membrane of the egg, 313; of feathers, *ib.*; of the pigmentum nigrum oculi, *ib.* Results of his researches, 125, 126.

SECRETIONS. See BILE and URINE.

SEGUIN. His calculation of the amount of inspired oxygen, 283.

SERPENTS. Their excrements consist of urate of ammonia, 54.

The process of digestion in them, 53.

SLEEP, Theory of, 228. Amount of sleep necessary for the adult, the infant, and the old man, 247 *et seq.* Induced by alcohol or wine, 240.

SODA. Essential to blood and bile, and derived from common salt, 161 *et seq.*

SODIUM, Chloride of. See SALT.

SOLANINE. Contains nitrogen, 177. Its analysis, 323.

STARCH. Exists in the food of the herbivora, 70. Is convertible into sugar, 70, 71. Its relation to gum and sugar, 73. Its function in food, 74 *et seq.* Amount of carbon in starch compared with that in flesh, 76, 77. Its composition compared with that of fat, 84, 90. Is the source of diabetic sugar, 95. Is an element of respiration, 96. Dissolved by diastase, 111. Its relation to choleic acid, 152. Its relation to the principal secretions and excretions, 153; to choloidic acid, 157; to bile, 158, 162, 164, 166. Its analysis from fifteen different plants, 297.

STARVATION. Process of, 25. Cause of death in, 27.

STRECKER. His analysis of starch from 12 different plants, 297.

SUGAR. Analysis of grape-sugar, 597; of sugar of milk, 298; of cane-sugar, 300. Is an element of respiration, 96.

SULPHUR. Exists in albumen, fibrine, and caseine, 41, 126.

SULPHURETTED HYDROGEN. Theory of its poisonous action, 274.

SULPHURIC ACID. See ACID, *Sulphuric*.

SUPPLY of matter. See NUTRITION.

SUPPLY and WASTE. Equilibrium between them constitutes the abstract state of health, 254, 255. Effects of its disturbance, *ib. et seq.* Means for restoring the equilibrium, 248, 257 *et seq.*

## T.

- TABLES** of the food consumed by soldiers in Germany, 289. Of the food and excretions of the horse and cow, 290, 291.
- TAURINE.** How produced from bile, 133. Its relation to choleic acid, 135. Its relation to uric acid and urea, and to allantoine, 155; to uric acid 156; to alloxan, *ib.*; to choloidic and choleic acids, and ammonia, 158; to caffeine or theine, 180; to asparagine, *ib.*; to theobromine, *ib.*, 181.
- TEMPERATURE.** Its effects on the amount of inspired oxygen, 16, and on the appetite, 17 *et seq.* A slight depression of temperature causes death in aged people, 255. Temperature of the blood in different animals, 292. Temperature of the body constantly kept up by internal causes, 19—22.
- TENDONS.** Analysis of, 311.
- THAULOW.** His analysis of cystic oxide, 320.
- THEINE.** Identical with caffeine, 179. And with guaranine, 324. Theory of its action, 181 *et seq.* Its relation to bile, 180. Its analysis, 324.
- THEOBROMINE.** Analogous to theine, 179. Theory of its action, 181 *et seq.* Its relation to bile, 180, 181. Its analysis, 324.
- THEORY.** Of animal heat, 17 *et seq.* Of digestion, 108 *et seq.* Of respiration, 265 *et seq.* Of the motions in the animal organism, 195 *et seq.* Of disease, 254 *et seq.* Of the action of caffeine, &c., 181 *et seq.* Of the action of the vegetable alkalies, 182 *et seq.* Of health, 254, 255.
- TIEDEMANN and GMELIN.** Their attempt to support a goose upon albumen alone, unsuccessful, 106.
- TISSUES, METAMORPHOSIS OF:** see **METAMORPHOSIS.** Analysis of the animal tissues, 310, 313. Formulæ of, 126.
- TOBACCO.** Arrests or retards the change of matter, 179.
- TRANSFORMATION.** See **METAMORPHOSIS.**
- TURNIPS.** Juice of, contains vegetable fibrine and albumen, 45, 46.

## U.

- UREA.** Derived from uric acid, 137, 140. Also from the oxidation of blood, 140; from allantoine, 141. Its relation to choleic acid, 148; to hippuric acid, 150; to proteine, 151; to proteine and starch, 153; to proteine and fat, 154; to taurine, 155, 156; to carbonate of ammonia, 159; to theobromine, 180. Its analysis, 319. Occurs in the urine of those who have taken benzoic acid along with hippuric acid, 327.

URINARY CALCULI. See CALCULUS.

URIC ACID. See ACID, *Uric*.

## V.

VARRENTRAPP and WILL. Their analysis of vegetable albumen, 294. Of sulphate of potash and caseine, 295.

VEGETABLES. Alone produce compounds of proteine, 106. Azotised constituents of, nutritious, 45; medicinal or poisonous, 176. Analysis of those vegetables which are used for food, 285 *et seq.*

VEGETATIVE LIFE. Distinguished from nervous life, 38. Predominates in the early stages of life, *ib.* Also in the female, 39.

VENOUS BLOOD. See BLOOD.

VITAL FORCE, or vitality. Definition of, 1 *et seq.* Theory of, 195 *et seq.*

VOGEL. His analysis of gas from the abdomen of cattle after excess in green food, 309.

## W.

WATER. Is one of the two constituents of the body which contain no nitrogen, 43. Its use as a solvent, *ib.* Contributes to the greater part of the transformations in the body, 136, 140, 141, 142, 148, 153, 154, 155, 156, 157, 159, 180, 181.

WAX. On its production from honey by the bee, 301—306. Its analysis, 307.

WHEAT. Contains vegetable fibrine, 46. Analysis of fibrine, albumen, and gluten, from wheat, 294.

WILL and ETTLING. Their analysis of lithofellic acid, 322.

WINE. The wines of the south promote the formation of calculus, 139. But not Rhenish wines, *ib.* Theory of its action, 239, 240.

WOSKRESENSKY. His analysis of theobromine, 324.

## Y.

YAMS. Analysis of starch from, 297.