

Index

- active coaching, therapeutic meals and, 52
- adherence ratings
- CBT-AR and, 32
 - Stage 1 and
 - session 1 in, 140–141
 - session 2 in, 142–143
 - session 3 in, 144–145
 - session 4 in, 146–147
 - Stage 2 and
 - session 1 in, 148–149
 - session 2 in, 150–151
 - Stage 3, fear of aversive consequences module, and
 - session 1 in, 158–159
 - sessions, later, in, 159–160
 - Stage 3, lack of interest in eating or food module, and
 - session 1 in, 161–162
 - sessions, later, in, 164–165
 - Stage 3, sensory sensitivity module, and
 - session 1 in, 150–153
 - session 2 in, 154–155
 - sessions, later, in, 163–166
 - Stage 4 and
 - session 1 in, 166–167
 - session 2 in, 168–169
- age, 4, 26–27
- age-of-onset criterion, 2
- anorexia nervosa
- bone loss caused by, 4
 - delay discounting and, 4–5
 - fat phobia and, 4
 - FBT and, 10, 50
 - starvation syndrome and, 24
 - weight loss caused by, 4
- antecedent manipulation, pediatric feeding disorders and, 7
- anxiety, 13
- anxiety disorders. *See also* panic disorder; phobias, specific
- avoidance and, 11, 128
 - case examples of, 127–132
 - CBT for, 11–12
 - exposure and, 11–12
 - psychoeducation for, 12
 - treatment for, 11–13
- ARFID. *See specific topics*
- assessment, 14–19
- fear of aversive consequences and, 15–16
 - food avoidance and restriction and, 14–15
 - height and, 17
 - lack of interest in eating or food and, 16
 - medical sequelae of ARFID and, 16–18
 - medication and, 18
 - oral-motor difficulties and, 18
 - PARDI and, 14–16
 - psychosocial impairment and, 16
 - rule-outs, for ARFID diagnosis, and, 16
 - sensory sensitivity and, 15
 - specific psychopathology of ARFID and, 14–16
 - supplement drinks and, 17
 - treatment and, 18–19
 - tube feeding and, 17
 - vitamin deficiencies and, 17
 - vitamin supplements and, 17
 - weight and, 17
- at-home practice. *See also* self-monitoring
- CBT-AR and, 30, 43
 - goals and, 30–31
 - Stage 1 and
 - session 1 in, 43–46
 - session 2 in, 51
 - session 3 in, 52–53
 - session 4 in, 52–53
 - Stage 2 and, 58–68
 - Stage 3, fear of aversive consequences module, and, 92, 94
 - Stage 3, lack of interest in eating or food module, and, 100–101
 - Stage 3, sensory sensitivity module, and
 - session 1 in, 73–74
 - session 2 in, 75
 - sessions, later, in, 78–87
- attentional deficit/hyperactivity disorder, 120, 122
- autism spectrum disorder, 4, 6, 16
- avoidance
- anxiety disorders and, 11, 128
 - psychoeducation and, 88
- avoidant/restrictive food intake disorder (ARFID). *See specific topics*
- BED. *See* binge eating disorder
- behavior therapy
- antecedent manipulation and, 7
 - differential reinforcement and, 7
 - extinction-based techniques and, 7
 - pediatric feeding disorders and, 7, 9
- binge eating disorder (BED), case examples of, 123–127
- biological contributors, 5, 20, 25, 137
- brain derived neurotrophic factor and, 24
 - food variety and, 34
 - ghrelin and, 23–24
 - lack of interest in eating or food and, 23–24, 40, 97
 - sensory sensitivity and, 21
 - smell and, 21
 - supertasters and, 21, 34
- body shape concerns, 3, 10
- case examples and, 123–124, 127
 - classical eating disorders and, 3
- bone loss, 4
- brain derived neurotrophic factor (BDNF), 24
- cannabis, 16
- care team, 19
- case examples, 116–135
- anxiety disorders and, 127–132
 - attentional deficit/hyperactivity disorder and, 120, 122
 - BED and, 123–127
 - body shape concerns and, 123–124, 127
 - CBT-AR, family-supported, and, 116–123, 127–132
 - CBT-AR, individual, and, 132–135
 - comorbidity and, 119–127
 - daily schedules and, 128–129, 131
 - depression and, 120, 128
 - exposure and, 117–118, 125, 130, 133–134

- exposure hierarchy and, 129–130
 fear of aversive consequences and, 127–132
 Five Steps and, 118, 121, 125–126, 133–134
 formulation, individualized, and, 117, 121, 124–125, 132–133
 IBS and, 128
 incorporating and, 117–119, 126, 133–135
 interoceptive exposure and, 121
 lack of interest in eating or food and, 119–132
 medical evaluations and, 116, 119, 133
 nutrition deficiencies and, 121, 125, 133
 obesity and, 123–127
 oppositional behavior and, 120, 122
 outcomes and conclusions from, 118–119, 122–123, 127, 131–132, 135
 parent monitoring and, 128
 psychoeducation and, 117, 128, 132–133
 psychosocial impairment and, 116, 118–119, 123–124, 128, 130–131, 133
 relapse prevention plans and, 118, 122, 126–127, 135
 reward schedules and, 120, 122, 129
 self-monitoring and, 124–127, 133
 sensory sensitivity and, 116–127, 132–135
 Stage 1 and, 117, 120–121, 124–125, 128–129, 132–133
 Stage 2 and, 117, 121, 125, 129, 133
 Stage 3, fear of aversive consequences module, and, 129–131
 Stage 3, lack of interest in eating or food module, and, 121–122
 Stage 3, sensory sensitivity module, and, 117–118, 122, 125–127, 133–134
 Stage 3 and, 121–123
 Stage 4 and, 118, 122, 126–127, 131, 134–135
 supplement drinks and, 122
 therapeutic meals and, 121, 129
 treatment interruptions and, 134
 underweight patients and, 119–123, 127–132
 weight gain and, 119–122, 128, 130–132
 weight loss and, 130–131
 categorical model, of ARFID, 14–15
 CBT. *See* cognitive-behavioral therapy
 CBT-AR. *See* cognitive-behavioral therapy for avoidant/restrictive food intake disorder
 CBT-AR, family-supported
 case examples of, 116–123, 127–132
 CBT-AR, individual, *versus*, 26–27
 CBT-AR, individual
 case examples of, 132–135
 CBT-AR, family-supported, *versus*, 26–27
 CBT-E. *See* cognitive-behavioral therapy, enhanced version
 child-targeted food products, 5–6
 choking, fear of, exposure hierarchy and, 88, 91
 classical eating disorders. *See also* anorexia nervosa
 ARFID, diagnostic crossover with, 5
 ARFID *versus*, 3–5
 body shape concerns and, 3
 CBT-E for, 9–10
 preferred foods and, 3
 treatment for, 9–11
 weight and, 3
 clinical severity of ARFID, 2
 cognitive-behavioral model of ARFID, 20–25
 fear of aversive consequences and, 22–23
 lack of interest in eating or food and, 23–25
 sensory sensitivity and, 21
 cognitive-behavioral therapy, enhanced version (CBT-E), 29
 CBT-AR and, 10
 classical eating disorders and, 9–10
 cognitive-behavioral therapy (CBT), anxiety disorders and, 11–12
 cognitive-behavioral therapy for avoidant/restrictive food intake disorder (CBT-AR). *See also* Stage 1, of CBT-AR; Stage 2, of CBT-AR; Stage 3, of CBT-AR; Stage 3, of CBT-AR, fear of aversive consequences module; Stage 3, of CBT-AR, lack of interest in eating or food module; Stage 3, of CBT-AR, sensory sensitivity module; Stage 4, of CBT-AR; *specific topics*
 completing, 103–112, 115
 development of, 136
 first four weeks of treatment in, 31
 refinement of, 136–137
 typical session of, 30–31
 comorbidity, 2, 4–6
 case examples of, 119–127
 Stage 1 and, 54
 competence ratings, for CBT-AR, 32, 138–139
 contraindications, for CBT-AR, 16
 cyproheptadine, 18
 daily schedules, case examples and, 128–129, 131
 delay discounting, 4–5
 depression, case examples of, 120, 128
 dietitians, CBT-AR and, 54
 differential reinforcement, for pediatric feeding disorders, 7
 dismantling studies, for CBT-AR, 136–137
 DSM-5, ARFID in, 1–2, 25, 136
 DSM-IV, feeding disorder of infancy and early childhood in, 1
 eating, erratic, 40
 eating disorder not otherwise specified (EDNOS), 2
 EDNOS. *See* eating disorder not otherwise specified
 electronic application, for CBT-AR, 137
 etiology, of ARFID, 5–6
 exposure. *See also* exteroceptive exposure; interoceptive exposure
 anxiety disorders and, 11–12
 case examples of, 117–118, 125, 130, 133–134
 CBT-AR and, 12–13, 40–43
 Five Steps and, 71–73
 food variety and, 34–40
 limited opportunities for, 20, 22–23
 panic disorders and, 12
 phobias, specific, and, 11–12
 psychoeducation and, 71–72, 88
 Stage 3, fear of aversive consequences module, and, 92–94
 Stage 3, sensory sensitivity module, and,
 session 1 in, 73
 session 2 in, 74–75
 sessions, later, in, 78
 systematic desensitization and, 11
 exposure hierarchy
 case examples of, 129–130
 choking, fear of, and, 88, 91
 Stage 3, fear of aversive consequences module, and, 88–93
 exteroceptive exposure, 12
 panic disorders and, 12
 phobias, specific, and, 12
 extinction-based techniques, pediatric feeding disorders and, 7

Index

- family meal environment, 5–6
- family-based treatment (FBT), 10–11
- anorexia nervosa and, 10, 50
 - one more bite intervention and, 52
 - underweight patients and, 10–11
 - weight gain and, 10–11
- fat phobia, 3–4
- FBT. *See* family-based treatment
- fear of aversive consequences, 47.
- See also* Stage 3, of CBT-AR, fear of aversive consequences module
 - assessment and, 15–16
 - case examples of, 127–132
 - CBT-AR goals and, 31
 - cognitive-behavioral model, of ARFID, and, 22–23
 - exposure, limited opportunities for, and, 23
 - gastrointestinal symptoms and, 23
 - psychoeducation and, 34
 - Stage 2 and, 68
 - trauma, food-related, and, 22–23
- feeding, division in responsibility in, 5
- feeding disorder of infancy and early childhood, 1. *See also* pediatric feeding disorders
- feeding or eating difficulties, prevalence of, 2
- Five Steps
- case examples and, 118, 121, 125–126, 133–134
 - exposure and, 71–73
 - preferred foods and, 97–100
 - Stage 3, lack of interest in eating or food module, and, 97–101
 - Stage 3, sensory sensitivity module, and, 71–73
- food allergies, 6
- food enjoyment, CBT-AR and, 31
- food neophobia, 5
- food variety. *See also* healthy diet
- biological contributors and, 34
 - CBT-AR impact on, 31
 - exposure and, 34–40
 - lack of interest in eating or food and, 40
 - Stage 1 and
 - session 1 in, 43–46
 - session 2 in, 46
 - session 3 in, 52
 - session 4 in, 52–53
 - Stage 2 and, 55–58, 69
 - Stage 3, sensory sensitivity module, and
 - session 1 in, 71–74
 - session 2 in, 74–75
 - sessions, later, in, 78–87
- food volume
- Stage 1 and
 - session 1 in, 43
 - session 2 in, 46, 48–50
 - session 3 in, 51
 - session 4 in, 52–53
 - Stage 3 and, 70
- foods, non-preferred, nutrition deficiencies linked to, 55
- formulation, individualized, 9
- case examples of, 117, 121, 124–125, 132–133
 - lack of interest in eating or food and, 47
 - sensory sensitivity in, 47
 - Stage 1 and, 46–47
 - Stage 4 and, 103
- gastrointestinal symptoms, fear of
- aversive consequences and, 23
- gender, 3–4
- ghrelin, lack of interest in eating or food and, 23–24, 48
- goals
- at-home practice and, 30–31
 - of CBT-AR, 31
- handouts
- Stage 1 and, 34–35, 39–43
 - Stage 2 and, 55–57, 59
 - Stage 3, fear of aversive consequences module, and, 88
 - Stage 3, lack of interest in eating or food module, and, 95
 - Stage 3, sensory sensitivity module, and, 72–73, 75–78
- healthy diet
- MyPlate and, 58–59
 - Stage 2 and, 57–59
- height, assessment and, 17
- high-energy drinks. *See* supplement drinks
- homework. *See* at-home practice
- hunger and fullness ratings
- parent monitoring and, 97–98
 - self-monitoring of, 97–98, 100
- IBS. *See* irritable bowel syndrome
- incorporating
- case examples of, 117–119, 126, 133–135
 - number of presentations required for, 78
 - Stage 3, sensory sensitivity module, and, 78–87
 - tasting versus, 75–78
- interoceptive exposure, 12–13
- case examples of, 121
 - panic disorder and, 12
 - phobia, specific, and, 12
 - Stage 3, lack of interest in eating or food module, and, 94–97, 100–101
 - SUDS scale and, 96–97, 100–101
- irritable bowel syndrome (IBS), 128
- lack of interest in eating or food, 47.
- See also* Stage 3, of CBT-AR, lack of interest in eating or food module
 - assessment and, 16
 - biological contributors to, 23–24, 40, 97
 - brain derived neurotrophic factor and, 24
 - case examples of, 119–132
 - CBT-AR goals and, 31
 - cognitive-behavioral model, of ARFID, and, 23–25
 - eating, erratic, and, 40
 - food variety and, 40
 - formulation, individualized, and, 47
 - ghrelin and, 23–24, 48
 - schedule, regular eating, and, 48
 - sensory-specific satiety and, 40
 - social expectations and, 25
 - Stage 2 and, 68
 - weight and, 24–25, 40
- maintaining mechanisms, 27–30.
- See also* fear of aversive consequences; food, lack of interest in; sensory sensitivity
 - psychoeducation on, 34–39
 - Stage 3 and, 70
 - weight gain and, 70–71
- medical evaluations, case examples of, 116, 119, 133
- medical sequelae, of ARFID
- assessment of, 16–18
 - height and, 17
 - medication and, 18
 - oral-motor difficulties and, 18
 - vitamin deficiencies and, 17
 - vitamin supplements and, 17
 - weight and, 17
- medical stability, CBT-AR and, 16–17
- medication, 54
- assessment and, 18
 - stimulant, 18
- MyPlate, healthy diet and, 58–59.
- See also* Primary Food Group Building Blocks

- negative consequences, ARFID
 - associated with, 1
- nutrition deficiencies
 - case examples of, 121, 125, 133
 - foods, non-preferred, and, 55
 - psychoeducation and, 55–57
 - Stage 2 and, 55–57
- nutritional compromise, 20–21, 69
- obesity, case examples of, 123–127
- observation, of therapeutic meals,
 - 51–52
- one more bite intervention
 - FBT and, 52
 - therapeutic meals and, 52
- operant conditioning, 74
- oppositional behavior, case examples
 - of, 120, 122
- oral-motor difficulties, assessment and,
 - 18
- outcomes, of CBT-AR, 118–119,
 - 122–123, 127, 131–132,
 - 135–136
- overweight patients, Stage 1 and, 33
- panic disorder
 - exposure and, 12
 - exteroceptive exposure and, 12
 - interoceptive exposure and, 12
 - treatment for, 12
- PARDI. *See* Pica, ARFID, and
 - Rumination Disorder Interview
- parent monitoring
 - case examples of, 128
 - discontinuing, 112
 - hunger and fullness ratings and,
 - 97–98
 - Stage 1 and, 43
 - Stage 3, lack of interest in eating or
 - food module, and, 97–98
 - Stage 4 and, 112
- parents, 26–27, 137
 - ARFID, blamed for, 32
 - relapse prevention plans and, 115
 - schedule, regular eating, and, 48
 - Stage 2 and, 59, 68–69
 - Stage 3, sensory sensitivity module,
 - and, 73
 - Stage 3 and, 71
- pathophysiology, of ARFID, 137
- patient blaming, 31–32
- pediatric feeding disorders
 - antecedent manipulation and, 7
 - behavior therapy and, 7, 9
 - differential reinforcement and, 7
 - extinction-based techniques and, 7
 - SOS and, 7–8
 - systematic desensitization, play-
 - based, and, 7–8
 - treatment for, 7–9
- phobias, specific
 - exposure and, 11–12
 - exteroceptive exposure and, 12
 - interoceptive exposure and, 12
- Pica, ARFID, and Rumination Disorder
 - Interview (PARDI), 14–16
- preferred foods, 3
 - classical eating disorders and, 3
 - Five Steps and, 97–100
 - Stage 1 and, 50
- preterm births, 6
- prevalence
 - of ARFID, 2–3
 - of feeding or eating difficulties, 2
- Primary Food Group Building Blocks
 - Stage 2 and, 57–59
 - Stage 3, sensory sensitivity module,
 - and, 78–87
 - Stage 4 and, 103–112
- psychoeducation, 9, 27–29, 40–43
 - anxiety disorders and, 12
 - avoidance and, 88
 - case examples of, 117, 128,
 - 132–133
 - exposure and, 71–72, 88
 - fear of aversive consequences and,
 - 34
 - maintaining mechanisms and,
 - 34–39
 - nutrition deficiencies and, 55–57
 - sensory sensitivity and, 34
 - Stage 1 and, 33–46
 - Stage 2 and, 55–59
 - Stage 3, fear of aversive
 - consequences module, and,
 - 88
 - Stage 3, lack of interest in eating or
 - food module, and, 94–95
 - Stage 3, sensory sensitivity module,
 - and, 71–72
 - treatment engagement and, 34
 - weight gain and, 34
- psychosocial impairment
 - assessment and, 16
 - case examples of, 116, 118–119,
 - 123–124, 128, 130–131, 133
- randomized controlled clinical trials,
 - for CBT-AR, 136
- rapid response, CBT-AR and, 31
- regurgitation, 3
- relapse prevention plans
 - case examples of, 118, 122, 126–127,
 - 135
 - parents and, 115
 - Stage 4 and, 113–115
- reward schedules
 - case examples of, 120, 122, 129
 - weight gain and, 68–69
- rule-outs, for ARFID diagnosis, 16
- schedule, regular eating
 - lack of interest in eating or food and,
 - 48
 - parents and, 48
 - Stage 1 and, 47–48
- self-help, CBT-AR as, 137
- self-monitoring, 9
 - case examples of, 124–127, 133
 - discontinuing, 112
 - hunger and fullness ratings and,
 - 97–98, 100
 - patient not doing, 53
 - Stage 1 and, 53
 - session 1 in, 43–46
 - session 2 in, 46
 - session 3 in, 52
 - session 4 in, 52–53
 - Stage 2 and, 55–58
 - Stage 3, lack of interest in eating or
 - food module, and, 97–98, 100
 - Stage 4 and, 112
- sensory sensitivity, 47. *See also* Stage 3,
 - of CBT-AR, sensory sensitivity
 - module
 - assessment and, 15
 - biological contributors to, 21
 - case examples of, 116–127, 132–135
 - cognitive-behavioral model of
 - ARFID and, 21
 - formulation, individualized, and, 47
 - psychoeducation and, 34
 - smell and, 21
 - Stage 2 and, 68
 - supertasters and, 21
 - texture and, 21
 - Weber's law and, 22
 - weight and, 24
- sensory-specific satiety, 21–22, 40
- Sequential Oral Sensory approach
 - (SOS), 7–8, 71
- smell, sensory sensitivity and, 21
- social avoidance, 46–47
- social expectations, lack of interest in
 - eating or food and, 25
- SOS. *See* Sequential Oral Sensory
 - approach
- specific psychopathology, of ARFID,
 - assessment of, 14–16
- Stage 1, of CBT-AR, 27–29, 33–54
 - case examples of, 117, 120–121,
 - 124–125, 128–129, 132–133
 - checklist for moving on from, 54
 - overweight patients and, 33
 - session 1 in, 33–46
 - adherence ratings and, 140–141
 - agenda setting in, 33
 - at-home practice in, 43–46
 - food variety, focusing on, 43–46
 - food volume, focusing on, 43
 - handouts for, 34–35, 39–43

Index

- Stage 1, of CBT-AR (cont.)
 introductions in, 33
 parent monitoring in, 43
 patient, weighing of, in, 33
 psychoeducation in, 33–46
 self-monitoring in, 43–46
 session 2 in, 33
 adherence ratings and, 142–143
 at-home practice in, 51
 food variety and, 46
 food volume and, 46, 48–50
 formulation, individualized, in, 46–47
 preferred foods, increase in, and, 50
 schedule, regular eating, in, 47–48
 self-monitoring in, 46
 therapeutic meals planned in, 51
 weight gain, discussing
 importance of, in, 48–51
 session 3 in, 33, 51–53
 adherence ratings and, 144–145
 at-home practice in, 52–53
 food variety and, 52
 food volume and, 51
 self-monitoring in, 52
 therapeutic meals in, 51–52
 weight gain during, 51–52
 session 4 in, 33, 53
 adherence ratings and, 146–147
 at-home practice in, 52–53
 food variety and, 52–53
 food volume and, 52–53
 self-monitoring and, 52–53
 weight gain during, 53
 troubleshooting for, 53–54
 comorbidity and, 54
 dietitians and, 54
 general therapy, concurrent, and, 54
 medication and, 54
 self-monitoring and, 53
 weight gain and, 54
 underweight patients and, 29, 33, 40, 43, 47–53
 Stage 2, of CBT-AR, 28–29, 55–69
 case examples of, 117, 121, 125, 129, 133
 checklist for moving on from, 69
 session 1 in, 55–58
 adherence ratings and, 148–149
 at-home practice in, 58
 food variety and, 55–58
 handouts in, 55–57
 nutrition deficiencies and, 55–57
 psychoeducation in, 55–57
 self-monitoring in, 55–58
 session 2 in, 58–68
 adherence ratings and, 150–151
 at-home practice in, 59–68
 fear of aversive consequences and, 68
 handouts in, 59
 healthy diet and, 57–59
 lack of interest in eating or food and, 68
 parents and, 59
 Primary Food Group Building
 Blocks and, 57–59
 psychoeducation in, 58–59
 sensory sensitivity and, 68
 weight gain during, 68–69
 troubleshooting for, 68–69
 concluding treatment and, 69
 food variety and, 69
 parents and, 68–69
 underweight patients and, 29, 55, 58
 Stage 3, of CBT-AR, 28–29
 case examples of, 121–123
 checklist for moving on from, 102
 food volume and, 70
 maintaining mechanisms, focusing on, 70
 parents and, 71
 session 2 in, 74–75
 sessions, later, in, 92–94
 underweight patients and, 70–71
 weight gain and, 70–71
 Stage 3, of CBT-AR, fear of aversive consequences module, 88–94
 case examples of, 129–131
 preparing for next module in, 94
 session 1 in, 88–92
 adherence ratings and, 158–159
 at-home practice in, 92
 exposure hierarchy and, 88–93
 handouts in, 88
 psychoeducation in, 88
 sessions, later, in, 92–94
 adherence ratings and, 159–160
 at-home practice in, 94
 exposure and, 92–94
 troubleshooting for, 101
 weight gain and, 70
 Stage 3, of CBT-AR, lack of interest in eating or food module, 94–101
 case examples of, 121–122
 preparing for next module in, 101
 session 1 in, 94–100
 adherence ratings and, 161–162
 at-home practice in, 100
 Five Steps and, 97–100
 handouts in, 95
 interoceptive exposure and, 94–97
 parent monitoring and, 97–98
 psychoeducation in, 94–95
 self-monitoring in, 97–98
 sessions, later, in, 100–101
 adherence ratings and, 164–165
 at-home practice in, 101
 Five Steps and, 101
 interoceptive exposure in, 100–101
 self-monitoring in, 100
 troubleshooting for, 101–102
 Stage 3, of CBT-AR, sensory sensitivity module, 71–88
 case examples of, 117–118, 122, 125–127, 133–134
 operant conditioning and, 74
 preparing for next module in, 87–88
 session 1 in, 71–74
 adherence ratings and, 150–153
 at-home practice in, 73–74
 exposure in, 73
 food variety and, 71–74
 handouts in, 72–73
 parents and, 73
 psychoeducation in, 71–72
 session 2 in, 74–75
 adherence ratings and, 154–155
 at-home practice in, 75
 exposure and, 74–75
 food variety and, 74–75
 handouts in, 75–78
 incorporating *versus* tasting and, 75–78
 sessions, later, in, 75–88
 adherence ratings and, 163–166
 at-home practice in, 78–87
 exposure and, 78
 food variety and, 78–87
 incorporating and, 78–87
 Primary Food Group Building
 Blocks and, 78–87
 troubleshooting for, 101
 weight gain and, 70
 Stage 4, of CBT-AR, 28–29
 case examples of, 118, 122, 126–127, 131, 134–135
 session 1 in, 103–112
 adherence ratings and, 166–167
 CBT-AR, completing, and, 103–112
 formulation, individualized, and, 103
 parent monitoring and, 112
 Primary Food Group Building
 Blocks and, 103–112
 self-monitoring in, 112
 session 2 in, 112–115
 adherence ratings and, 168–169
 relapse prevention plans in, 113–115
 troubleshooting for, 115
 starvation syndrome, 24
 Subjective Units of Distress scale
 (SUDS scale), 88–94, 96–97, 100–101

| | | |
|---|---|---|
| SUDS scale. <i>See</i> Subjective Units of Distress scale | treatment interruptions, case examples of, 134 | Weber’s law, sensory sensitivity and, 22 |
| supertasters, 21, 34 | troubleshooting. <i>See</i> Stage 1, troubleshooting for; Stage 2, troubleshooting for; Stage 3, of CBT-AR, fear of aversive consequences module; Stage 3, of CBT-AR, lack of interest in eating or food module; Stage 3, of CBT-AR, sensory sensitivity module; Stage 3, troubleshooting for; Stage 4, troubleshooting for | weight, 3, 10–11, 21, 29–30 |
| supplement drinks, 17, 50–51 | | assessment and, 17 |
| assessment and, 17 | | classical eating disorders and, 3 |
| case examples of, 122 | | lack of interest in eating or food and, 24–25, 40 |
| systematic desensitization | | sensory sensitivity and, 24 |
| exposure and, 11 | | weight gain, 1–2 |
| pediatric feeding disorders and, 7–8 | | case examples of, 119–122, 128, 130–132 |
| play-based, 7–8 | | FBT and, 10–11 |
| SOS and, 7–8 | | maintaining mechanisms and, 70–71 |
| tasting, incorporating versus, 75–78 | tube feeding, 17, 137 | psychoeducation and, 34 |
| texture, sensory sensitivity and, 21 | | reward schedules and, 68–69 |
| therapeutic meals | underweight patients, 29 | Stage 1 and, 54 |
| active coaching in, 52 | case examples of, 119–123, 127–132 | session 2 in, 48–51 |
| case examples of, 121, 129 | FBT and, 10–11 | session 3 in, 51–52 |
| observation of, 51–52 | Stage 1 and, 29, 33, 40, 43, 47–53 | session 4 in, 53 |
| one more bite intervention in, 52 | Stage 2 and, 29, 55, 58 | Stage 2 and, 68–69 |
| Stage 1 and, 51–52 | Stage 3 and, 70–71 | Stage 3, fear of aversive |
| therapist stance, in CBT-AR, 31–32 | | consequences module, and, 70 |
| trauma, food-related, 22–23, 40 | visits, frequency of, CBT-AR and, 18–19 | Stage 3, sensory sensitivity module, and, 70 |
| treatment. <i>See also specific treatments</i> | vitamin deficiencies, 17 | Stage 3 and, 70–71 |
| for anxiety disorders, 11–13 | vitamin supplements, 17 | weight loss, 1–2, 4 |
| assessment and, 18–19 | vomiting, fear of, 88–93 | anorexia nervosa causing, 4 |
| for classical eating disorders, 9–11 | | case examples of, 130–131 |
| for panic disorder, 12 | | |
| for pediatric feeding disorders, 7–9 | | |
| treatment engagement, | | |
| psychoeducation and, 34 | | |