

Index

- abdominal and flank trauma.
See also renal injury
 anatomical assessment of, 213–14
 assessment approach to, 226
 causes of, 213
 diaphragmatic injury, 218–19
 differential diagnosis for, 216
 duodenal injury, 217–21
 evaluation of, 220–21
 with DPL, 225–26
 hepatic injury, 216–17
 imaging of, 217
 liver grading criteria for, 217–19
 management of, 218–28
 imaging of, 223–25
 with CT, 224–25
 of hepatic injury, 217
 with radiography, 223–24
 of renal injury, 221
 of splenic injury, 219
 with ultrasound, 223–24
 incidence rates for, 213
 laboratory testing for, 222–23
 management of, 226
 medical history of, 214–15
 pancreatic injury, 218, 221
 pathophysiology of, 214
 physical examination for, 215
 splenic injury, 217
 grading criteria for, 219–20
 from gunshot wounds, 220
 imaging for, 219
 from stab wounds, 216
 from vehicle accidents, 216
 abrasions. *See* corneal abrasions
 ACEP. *See* American College of Emergency Physicians
 Advanced Trauma Life Support (ATLS), 42
 AirTraQ, 35
 airway management, with trauma, 17–18
 for burns, 22, 308
 inhalation of toxic fumes, 22
 of cervical spine, 149
 for cervical spine injury, 21
 with closed head injury, 18–19
 complicating factors for, 17
 with difficult intubation, 34
 for direct airway trauma, 19–21
 with endotracheal tube introducer, 34–35
 with failed airway, 37–38
 cricothyrotomy for, 37–38
 with FFB, 35
 AirTraQ, 35
 with head trauma, 90–95
 induction in, 93
 post-intubation, 93
 pretreatment in, 92
 intervention indications, 18
 for intubation, 18
 through intubation
 intervention indications for, 18
 in primary survey, 5
 with maxillofacial trauma, 19–20
 with mechanical distortion of airway, 18
 with neck trauma, 171
 for pediatric trauma, 68
 post-intubation
 management, 32–34
 with paralytic agents, 33
 problem solving in, 34
 of sedation and analgesia, 32–34
 ventilator settings, 32
 pre-existing difficulties, 17
 during pregnancy, 57
 in primary survey, 5–6, 9
 intubation assessment, 5
 with pulmonary trauma, 175–76
 in-line stabilization procedures, 176
 supraglottic airways, 36–37
 with i-Gel, 37
 ILMA, 36–37
 LMA, 36–37
 for thoracic trauma, 21
 trauma immobilization with, 17–18
 through video laryngoscopy, 30–32
 with C-MAC, 31
 with GlideScope, 30–31
 tracheal placement, 31–32
 alveolar fractures, 107–8
 imaging for, 110
 American College of Emergency Physicians (ACEP), 322–23
 American College of Surgeons, 1
 American Society of Anesthesiology (ASA), 322–24
 ankle injuries, 299–300
 dislocation types, 299–300
 imaging for, 300
 anterior cord syndrome, 162
 anticoagulants, 50
 Coumadin, 50
 DOAC, 50–51
 with geriatric trauma, 83
 ASA. *See* American Society of Anesthesiology
 ATLS. *See* Advanced Trauma Life Support
 autologous transfusion, for pulmonary trauma, 180–82
 balanced resuscitation, 46
 blood products in, 46
 battle sign, 111–13
 Bennett's fracture, 275

Index

- bladder injury, 239–40
 - imaging of, 240
 - management of, 240
 - ruptures of, classification of, 239–40
- blast injuries, 87
- bleeding and blood loss, 7
- blood pressure control, with
 - head trauma, 95–98
 - hypertension, 97–98
 - hypotension, 89–97
 - neurogenic shock, 97
 - vasopressors, 97–98
- blood products
 - in transfusions, 42–43, 48
 - FFP, 46
 - PCC, 48
 - pRBCs, 44–46
 - recombinant factor VII, 48
 - whole blood, 52
 - FWB, 52
 - SWB, 52
- blunt cardiac trauma, 194, 197–99
 - cardiac dysfunction from, 199
 - cardiac rupture from, 197
 - commotio cordis injury, 199
 - coronary vessel injuries, 198
 - management of, 208
 - pericardial injury, 198
 - valvular damage from, 198
- blunt chest trauma, 175
- blunt injuries, neck trauma for, 166–70
 - Canadian Cervical Spine Rule criteria for, 167
 - causes of, 166–67
 - imaging for, 168
 - NEXUS Criteria for, 167
 - screening considerations, 168–69
- bones, upper extremity trauma for, 273–76
 - distal phalanges, 273–74
 - metacarpal, 274–75
 - proximal and middle phalanges, 274
- bradycardia, 42
- brainstem herniation, 88–89
- breathing, assessment of
 - with pediatric trauma, 70
 - during pregnancy, 57–59
 - in primary survey, 6, 9
- hemothorax issues, 6
- pneumothorax issues, 6
- Brown-Sequard syndrome, 161
- burns
 - airway management with, 22, 308
 - inhalation of toxic fumes, 22
 - carbon monoxide poisoning and, 313
 - cardiopulmonary injury from, 317
 - classification of, 309–10
 - clinical features of, 317–19
 - cutaneous injury from, 318
 - cyanide poisoning and, 313–14
 - depth estimation of, 308–9
 - electrical, 315–17
 - contact points for, 316–17
 - current effects on, 308–16
 - voltage characteristics, 308–16
 - emergency department management of, 308–17
 - escharotomies for, 312–13
 - considerations for, 312–16
 - evaluation of, 308
 - fluid resuscitation for, 309–12
 - gastrointestinal injury from, 317
 - incidence rates for, 307
 - from lightning, 319
 - lip, in children, 318–19
 - Lund-Browder diagram for, 309–11
 - management of, 308
 - muscle injury from, 317
 - nervous system injury from, 317
 - ocular cataracts and, 318
 - pathophysiology of, 307–8
 - Rule of Nines for, 310
 - size estimation for, 308–9
 - transfer to burn centers, 314–16
 - tympenic membrane rupture, 310–18
 - vascular injury from, 317
 - wound care with, 314–15
- burst fractures, 159
- calcaneal fractures, 301
- calcium, in transfusions, 48
- call-and-response principle, 1–2
- Canadian Cervical Spine Rule criteria, 146, 167
- carbon monoxide poisoning, 313
- cardiac missile, 197
- cardiac tamponade, 197
- cardiac trauma. *See also* blunt cardiac trauma
 - allied structures in, 193
 - anatomical assessment for, 194
 - causes of, 193–94
 - precordial penetrating wounds, 193
 - epidemiology of, 193
 - evaluation of, 200–1
 - with ECG, 200
 - through laboratory assessment of cardiac biomarkers, 200–1
 - great vessel injury in, 199
 - chest X-rays for, 202–3
 - CT imaging of, 203
 - management of, 208
 - imaging of, 201–3
 - with chest X-rays, 202–3
 - with CT, 203–4
 - with TEE, 201–2
 - with ultrasound, 201
 - incidence rates for, 193
 - management of, 203–8
 - of blunt trauma, 208
 - great vessel injury, 208
 - of penetrating trauma, 203–8
 - medical history of, 196
 - pathophysiology for, 195
 - physical examination for, 196
 - Trauma Association guidelines, 203–5
- cardiopulmonary resuscitation, 59–60
- carotid-cavernous fistula (CCF), 130–32
 - clinical presentation, 131
 - imaging of, 131
 - signs and symptoms of, 131
- cauliflower ear, 111–12
- CCF. *See* carotid-cavernous fistula
- Centers for Disease Control (CDCs), 3
- central cord syndrome, 162
- cerebral contusion, 87–91

Index

- cervical spine injuries, 21, 82
 tibial spine fractures, 294
 cervical spine trauma
 anatomy of, 144
 clinical clearance of, 150
 complications of, 149–50
 consultation indications, 150
 hypotension with, 150
 imaging of, 145–47
 assessment prior to, 145–46
 Canadian Cervical Spine Rule criteria in, 146
 NEXUS Criteria for, 145
 immobilization after, 145
 management of
 of airway, 149
 in emergency department, 147–51
 with steroids, 149
 medical history for, 144–45
 partial spinal cord syndromes, 149–50
 physical examination for, 144–45
 SCIWORA, 149
 spinal shock, 150
 types of fractures, 147
 Hangman's fracture, 148
 Jefferson burst fracture, 147
 Type 2 dens fracture, 148
 vascular injuries, 150
 chance fractures, 159–60
 chemical burns. *See* ocular chemical burns
 chest trauma. *See also* pulmonary trauma
 blunt, 175
 incidence rates for, 175
 chest tube removal, 180
 chest wall injuries, 81–82
 circulation
 during pregnancy, 59
 primary survey of, 6–7, 9
 Shock Index, 6
 closed head injury, 18–19
 C-MAC, for airway management, 31
 coagulopathy reversal, 99
 Colle's fracture, 277
 comminuted clavicular fractures, 285
 commotio cordis injury, 199
 compartment syndrome, 298–99
 compartments of the thigh, 290
 compression fractures, 158–59
 computed tomography (CT)
 scans, 11–12
 of abdominal and flank trauma, 224–25
 of cardiac trauma, 203–4
 for facial trauma, 109
 of genitourinary trauma, 234–36
 for great vessel injury, 203
 MDCTA, 165–66
 of renal injury, 237
 WBCT, 11–12
 conjunctival injury, 138–39
 from abrasions, 139
 conjunctival lacerations, 138–39
 from foreign bodies, 139
 subconjunctival hemorrhages, 138
 corneal abrasions, 136
 antibiotic treatment for, 138
 corneal foreign bodies, eye trauma from, 136–37
 antibiotic treatment for, 138
 physical examination of, 137–38
 coronary vessel injuries, 198
 corticospinal lesions, 155
 Coumadin, 50
 cranial nerves, 108
 cricothyrotomy, 37–38
 crystalloid resuscitation, in transfusions, 42–44
 CT scans. *See* computed tomography scans
 cuboid fractures, 302
 cuneiform fractures, 302
 cyanide poisoning, 313–14
 damage control resuscitation (DCR), 45
 components of, 45
 for PVI, 250–52
 delayed diagnosis
 definition of, 334
 universal scope of, 334–35
 dental injuries, 117
 Ellis fracture, 118
 depressed skull fracture, 95
 diagnostic peritoneal lavage (DPL), 225–26
 diaphragmatic injury, 218–19
 diffuse axonal injury, 87–90, 94
 direct airway trauma, 19–21
 direct oral anticoagulant considerations (DOAC), 50–51
 disability, primary survey of, with trauma, 8–9
 neurological assessment in, 9
 DOAC. *See* direct oral anticoagulant considerations
 dorsal column lesions, 155
 DPL. *See* diagnostic peritoneal lavage
 duodenal injury, 217–21
 ears, in facial trauma, 111–13
 battle sign, 111–13
 cauliflower ear, 111–12
 draining techniques, 111–12
 otorrhea, 112–13
 tympanic membrane perforation, 111
 ECG. *See* electrocardiogram
 eFAST. *See* Extended Focused Assessment with Sonography in Trauma
 elbow and forearm injuries. *See* upper extremity trauma
 elder abuse, 82–83
 elderly. *See* geriatric trauma
 electrical burns, 315–17
 contact points for, 316–17
 current effects on, 308–16
 voltage characteristics, 308–16
 electrocardiogram (ECG), for cardiac trauma, 200
 Ellis fracture, 118
 ELM. *See* external laryngeal manipulation
 emergency departments
 ACEP clinical policies, 322–23
 burns in, 308–17
 cervical spine trauma in, 147–51
 ED thoracotomy, for cardiac trauma, 203, 207–8
 genitourinary trauma in, 245–46
 head trauma evaluation in, 89–100
 goals of, 91
 management strategies for, 89–90
 pitfalls in, 100

Index

- emergency medical services (EMS), 2–4
 CDC guidelines, 3
 EMTs, 3–4
 information during
 transition of care
 from, 4
 emergency medical technicians (EMTs), 3–4
 EMS. *See* emergency medical services
 EMT. *See* emergency medical technicians
 endotracheal tube introducer, 34–35
 epidural hematoma, 87–90, 92
 escharotomies, 312–13
 considerations for, 312–16
 esophageal disruption, 190–91
 etomidate, 5, 325
 Extended Focused Assessment with Sonography in Trauma (eFAST), 10–11
 external laryngeal manipulation (ELM), 30
 extremity hemorrhage, with PVI, 249–50
 eye trauma
 acute visual threat
 identification with, 123
 from CCF, 130–32
 clinical presentation, 131
 imaging of, 131
 signs and symptoms of, 131
 conjunctival injury, 138–39
 from abrasions, 139
 conjunctival lacerations, 138–39
 from foreign bodies, 139
 subconjunctival hemorrhages, 138
 from corneal abrasions, 136
 antibiotic treatment for, 138
 from corneal foreign bodies, 136–37
 antibiotic treatment for, 138
 physical examination of, 137–38
 epidemiology of, 122
 examination for, 124–28
 with diagnostic imaging, 128
 of ocular region, 125–28
 of periocular area, 124–25
 of visual acuity, 124–28
 eyelid lacerations, 136
 types of, 136
 focused history with, 123
 mechanisms of, 122–23
 negative acuity outcomes, 123
 from ocular chemical burns, 128–29
 causes of, 129
 treatment of, 129
 open globe injury, 132–33
 physical examination of, 133
 optic nerve injury, 135
 from orbital compartment syndrome, 129–30
 imaging for, 130
 inferior cantholysis for, 130–31
 lateral canthotomy for, 130–31
 physical examination of, 130
 from orbital fractures, 111, 139–42
 management of, 140–41
 white-eyed fracture, 141
 patient assessment of, 123
 positive acuity outcomes, 123
 retinal trauma, 134–35
 traumatic hyphema, 133–34
 evaluation of, 133–34
 traumatic uveitis, 135
 vitreous hemorrhage, 134
 eyelid lacerations, 136
 types of, 136
 facial trauma
 affected vessels in, 107
 alveolar fractures, 107–8
 imaging for, 110
 anatomy of, 107–8
 assessment of, 105–7
 cranial nerves in, 108
 dental injuries, 117
 Ellis fracture, 118
 for ears, 111–13
 battle sign, 111–13
 cauliflower ear, 111–12
 draining techniques, 111–12
 otorrhea, 112–13
 tympanic membrane perforation, 111
 epidemiology of, 105
 examination of, 108–9
 history of injury for, 108
 imaging for, 105, 109
 of alveolar fracture, 110
 with CT, 109
 of dissection of carotid artery, 110
 management of, 109–11
 mandible fractures, 116–17
 maxillary fractures, 114
 Lefort fractures, 114–16
 nasal bone injuries, 113–14
 nasoethmoidal fractures, 107–8
 orbital fractures, 111. *See also* eye trauma
 with parotid duct, 117
 pitfalls in, 118
 TMJ injuries, 115–16
 reduction of, 116–17
 tripod fractures, 114–15
 zygomatic bone fractures, 114
 failed airways, 37–38
 cricothyrotomy for, 37–38
 falls, 81–82
 FAST. *See* focused assessment with sonography for trauma
 females, genitourinary trauma in, 245
 femoral injuries, 289–91
 compartments of the thigh, 290
 distal fractures, 290–91
 splinting of, 291
 traction for, 291–92
 mechanisms of, 289
 pediatric fractures, 291
 physical examination for, 289
 shaft fractures, 289–90
 fentanyl, 325
 fetal assessment, 61–62
 through FAST, 62
 through ultrasound examination, 62–63
 FFB. *See* flexible fiberoptic bronchoscope
 FFP. *See* fresh frozen plasma
 flail chest, 182–84
 flank trauma. *See* abdominal and flank trauma

- flexible fiberoptic
 bronchoscope (FFB), 35
 AirTraQ, 35
 focused assessment with
 sonography for trauma
 (FAST), 62
 for pediatric trauma, 74
 foot injuries, 300–1
 anatomical assessment of,
 301
 forefoot fractures, 302–3
 of hindfoot, 300–2
 calcaneal fractures, 301
 Hawkins classifications of,
 301
 imaging of, 301–2
 talus fractures, 300–1
 imaging for, 300
 of midfoot, 302
 cuboid fractures, 302
 cuneiform fractures, 302
 Lisfranc fracture, 302
 navicular fractures, 302
 phalanx fractures, 303
 forefoot fractures, 302–3
 fracture-dislocation injuries,
 160
 fractures. *See specific fractures*
 fresh frozen plasma (FFP), 8
 fresh whole blood (FWB), 52
- GCS. *See* Glasgow Coma Scale
 genitourinary trauma. *See also*
 renal injury
 assessment of, 232
 bladder injury and, 239–40
 imaging of, 240
 management of, 240
 ruptures of, classification
 of, 239–40
 diagnostic studies of, 233–36
 baseline renal function in,
 233
 with urinalysis, 233–34
 emergency department
 management of, 245–46
 in females, 245
 hematuria and, 233–34
 management of, 233–34
 imaging of
 with CT, 234–36
 with IVP, 236
 with MRI, 236
 with RUG, 236
 with ultrasound, 234–35
 incidence rates for, 232
- of penis, 244
 of scrotum and testicles,
 243–44
 signs and symptoms of,
 232–33
 surgical indications for, 246
 of ureter, 238–39
 urethral injury, 241–43
 classification of, 241–42
 management of, 242–43
- geriatric trauma
 anticoagulant use with, 83
 from elder abuse, 82–83
 epidemiology of, 77
 incidence rates for, 77–78
 injury mechanisms for,
 80–82
 cervical spine injuries, 82
 chest wall injuries, 81–82
 clinical implications of, 81
 falls, 81–82
 head injuries, 81
 from vehicle accidents, 82
 pathophysiology issues with,
 79–80
 comorbidities, 79
 shock, 79–80
 supporting evidence for,
 80
 research on, limits of, 78
 survey principles for, 78–83
 triage for, 77–79
 pre-hospital assessment,
 77–79
 after trauma center arrival,
 78–79
 Glasgow Coma Scale (GCS),
 8–9
 GlideScope, 30–31
 great vessel injury, 199
 chest X-rays for, 202–3
 CT imaging of, 203
 management of, 208
 gunshot wounds, 96
 renal injury from, 222
 splenic injury from, 220
- hand and finger injuries. *See*
 upper extremity trauma
 hanging, 170–71
 Hangman's fracture, 148
 hard signs, of PVI, 252
 head trauma
 airway management with,
 90–95
 induction in, 93
- post-intubation, 93
 pretreatment in, 92
 from blast injuries, 87
 blood pressure control with,
 95–98
 hypertension, 97–98
 hypotension, 89–97
 neurogenic shock, 97
 vasopressors, 97–98
 cerebral contusion, 87–91
 closed head injury, 18–19
 coagulopathy reversal after,
 99
 from depressed skull
 fracture, 95
 diffuse axonal injury, 87–90,
 94
 in elderly, 81
 in emergency department
 evaluation of, 89–100
 goals of, 91
 management strategies for,
 89–90
 pitfalls in, 100
 epidural hematoma, 87–90,
 92
 from gunshot wounds, 96
 herniation syndromes, 88–90
 brainstem herniation,
 88–89
 hyperosmolar therapy for,
 98
 hypothermia and, 100
 ICP after, 98–99
 hyperosmolar therapy for,
 98
 monitoring of, 98
 ultrasound for, 99
 incidence rates for, 87
 induced coma and, 100
 intracerebral hemorrhage,
 99
 from knife wounds, 96
 management strategies
 in emergency
 departments, 89–90
 pitfalls in, 100
 tiers of, 90–97
 from neurological injuries,
 87–90
 pediatric trauma and, 68–72
 physiologic goals, 87–89, 91
 with secondary injuries, 89
 seizure management, 99
 subarachnoid hemorrhage,
 87–90, 94

Index

- head trauma (cont.)
 subdural hematoma, 87–90, 93
 surgical interventions for, 99
 for specific injuries, 99
 TXA in, 100
 hematoma. *See* epidural hematoma; subdural hematoma
 hematuria, 233–34
 management of, 233–34
 hemorrhage
 blood transfusions and, 42
 ATLS and, 42
 bradycardia and, 42
 management of, 42
 extremity, with PVI, 249–50
 intracerebral, 99
 junctional, with PVI, 249, 251
 subarachnoid, 87–90, 94
 subconjunctival, 138
 vitreous, 134
 hemothorax, 6
 hepatic injury, 216–17
 imaging of, 217
 liver grading criteria for, 217–19
 management of, 218–28
 herniation syndromes, 88–90
 brainstem herniation, 88–89
 hindfoot injuries. *See* foot injuries
 hip dislocations, 268–72
 diagnosis of, 269–70
 reduction techniques for, 270–71
 treatment of, 270
 types of, 268
 hip fractures, 266–68
 anatomical assessment of, 266–67
 diagnosis of, 267
 management of, 268
 treatment for, 268
 hyperosmolar therapy, 98
 hypertension, 97–98
 hypocalcemia, 48
 hypotension, 7–8, 89–97
 with cervical spine trauma, 150
 differential for, 7
 permissive, 8
 REBOA and, 8
 hypothermia, head trauma and, 100
 iatrogenic injury, 197
 ICP. *See* intracranial pressure
 i-Gel, 37
 ILMA. *See* intubating laryngeal mask airway
 imaging. *See specific imaging modalities*
 incomplete spinal cord syndromes, 161–62
 anterior cord syndrome, 162
 Brown-Sequard syndrome, 161
 central cord syndrome, 162
 induced coma, 100
 inferior cantholysis, 130–31
 in-line stabilization procedures, 176
 innervation of the hand, 276
 intracerebral hemorrhage, 99
 intracranial pressure (ICP), 98–99
 hyperosmolar therapy for, 98
 monitoring of, 98
 ultrasound for, 99
 intravenous pyelogram (IVP), 236
 intubating laryngeal mask airway (ILMA), 36–37
 intubation, assessment for.
 See also rapid-sequence intubation
 in airway management
 intervention indications for, 18
 in primary survey, 5
 isolated fibular fractures, 300
 splints for, 300
 IVP. *See* intravenous pyelogram
 Jefferson burst fracture, 147
 junctional hemorrhage, with PVI, 249, 251
 ketamine, 5, 325
 ketofol, 325
 kidney injury. *See* renal injury
 knee dislocation, 291–92, 294–96
 associated injuries with, 295
 imaging of, 295
 management of, 295–96
 mechanisms of injury, 294–95
 physical examination for, 294
 knife wounds, 96
 lacerations. *See* eyelid lacerations
 laryngeal mask airway (LMA), 36–37
 laryngotracheal injuries, 169–70
 lateral canthotomy, 130–31
 Lefort fractures, 114–16
 lethal triad, 44–45
 pRBCs in, 44–45
 treatment for components of, 44
 lightning burns, 319
 lip burns, in children, 318–19
 Lisfranc fracture, 302
 liver injury. *See* hepatic injury
 LMA. *See* laryngeal mask airway
 low risk extremity injuries, 256
 lower extremity trauma. *See also* femoral injuries; foot injuries; tibial plateau fractures
 for ankle, 299–300
 dislocation types, 299–300
 imaging for, 300
 compartment syndrome, 298–99
 isolated fibular fractures, 300
 splints for, 300
 knee dislocation, 291–92, 294–96
 associated injuries with, 295
 imaging of, 295
 management of, 295–96
 mechanisms of injury, 294–95
 physical examination for, 294
 open knee joint, 297
 CT imaging for, 297
 saline load test for, 297
 patella dislocation, 296
 patella fractures, 296–97
 imaging of, 296–97
 management of, 297
 proximal fibula fractures, 299
 proximal third tibial fractures, 298
 tibial shaft fractures, 298
 tibial spine fractures, 294
 magnetic resonance imaging (MRI), of genitourinary trauma, 236

Index

- Mallampati classification, for RSI, 23–25
 mallet finger, 274
 mandible fractures, 116–17
 massive hemothorax, 180–82
 autologous transfusion for, 180–82
 massive transfusion protocols (MTPs), 46–48
 activation of, 47–48
 definition of, 46
 goal parameters, 47
 primary literature for, 47
 maxillary fractures, 114
 Lefort fractures, 114–16
 maxillofacial trauma, 19–20
 MDCTA. *See* Multi-Detector Computed Tomographic Angiography
 mechanical distortion of airway, 18
 midazolam, 325
 midfoot injuries. *See* foot injuries
 minimal volume normotension, 46
 missed traumatic injuries
 of abdomen, 339–40. *See also* abdominal and flank trauma
 imaging of, 340–44
 physical examination, 341
 definition of, 334
 of extremities, 344–45
 imaging of, 343–44
 physical examination, 342–43
 of head, 335. *See also* head trauma
 imaging of, 335–37
 physical examination for, 336–37
 of neck, 335–36. *See also* neck trauma
 imaging for, 336–39
 physical examination for, 338
 nerve compromise from, 344–45
 of pelvis, 340–44. *See also* pelvic trauma
 physical examination of, 341–44
 of spine, 336–37. *See also* cervical spine injuries;
 cervical spine trauma
 physical examination for, 337–38
 of thoracic region, 338–39.
 See also thoracic trauma
 imaging of, 339–42
 physical examination of, 339–40
 universal scope of, 334–35
 vascular compromise from, 344–45
 Monteggia fracture-dislocation, 282
 MRI. *See* magnetic resonance imaging
 MTPs. *See* massive transfusion protocols
 Multi-Detector Computed Tomographic Angiography (MDCTA), 165–66
 musculoskeletal system, in pediatric trauma, 69
 nasal bone injuries, 113–14
 nasoethmoidal fractures, 107–8
 navicular fractures, 302
 NCTH. *See* non-compressible truncal/torso hemorrhage
 neck fractures, 275
 neck trauma
 airway management with, 171
 anatomical assessment with, 164–65
 from blunt injuries, 166–70
 Canadian Cervical Spine Rule criteria for, 167
 causes of, 166–67
 imaging for, 168
 NEXUS Criteria for, 167
 screening considerations, 168–69
 from hanging, 170–71
 immobilization for, 168
 incidence rates for, 164
 laryngotracheal injuries, 169–70
 from penetrating injuries, 164–66
 MDCTA for, 165–66
 selective approach to, 165–66
 in Zone I, 165
 in Zone II, 165
 in Zone III, 165–66
 seatbelt signs, 169
 from strangulation, 170–71
 needle compression procedure, 176, 178–79
 neurogenic shock, 97
 NEXUS Criteria, 145, 167
 non-compressible truncal/torso hemorrhage (NCTH), 249, 251
 non-displaced bicondylar tibial plateau fractures, 293
 non-emergent transfusions, 42
 ocular cataracts, 318
 ocular chemical burns, 128–29
 causes of, 129
 treatment of, 129
 ocular trauma. *See* eye trauma
 open globe injury, 132–33
 physical examination of, 133
 open knee joint, 297
 CT imaging for, 297
 saline load test for, 297
 open pneumothorax, 180
 optic nerve injury, 135
 orbital compartment syndrome, 129–30
 imaging for, 130
 inferior cantholysis for, 130–31
 lateral canthotomy for, 130–31
 physical examination of, 130
 orbital fractures, 111, 139–42
 management of, 140–41
 white-eyed fracture, 141
 otorrhea, 112–13
 packed red blood cells (pRBCs), 44–46
 pancreatic injury, 218, 221
 paralytic agents, 29–30
 in post-intubation management, 33
 rocuronium, 29–30
 succinylcholine, 29–30
 vecuronium, 29–30
 parotid duct, with facial trauma, 117
 pars interarticularis fractures, 160
 partial spinal cord syndromes, 149–50

Index

- patella dislocation, 296
 patella fractures, 296–97
 imaging of, 296–97
 management of, 297
 PCC. *See* Prothrombin
 Complex Concentrate
 pediatric fractures, 291
 pediatric patients, PSA for,
 328–29
 discharge recommendations,
 329
 duration recommendations
 for, 329
 sedation scales, 329
 pediatric trauma
 anatomy in, 68–70
 airway assessment, 68
 musculoskeletal system, 69
 epidemiology of, 68
 FAST for, 74
 incidence rates for, 69
 management of, 75
 physiology of, 68–70
 breathing, 70
 of thermoregulation, 70
 regional evaluation for, 71–75
 of abdomen, 74
 of extremities, 74–75
 of head, 68–72
 of neck, 72–73
 pitfalls in, 75
 of thorax, 73
 pelvic fractures
 classification of, 262–64
 patterns of, 263
 pelvic trauma
 anatomical assessment of,
 260–61
 diagnosis of, 261–62
 hip dislocations, 268–72
 diagnosis of, 269–70
 reduction techniques for,
 270–71
 treatment of, 270
 types of, 268
 hip fractures, 266–68
 anatomical assessment of,
 266–67
 diagnosis of, 267
 management of, 268
 treatment for, 268
 hip joints and, 261
 imaging for, 262, 266
 pelvic fractures
 classification of, 262–64
 patterns of, 263
 treatment of, 262–66
 for hip dislocations, 270
 for hip fractures, 268
 penetrating cardiac trauma,
 196–97
 cardiac missile, 197
 cardiac tamponade, 197
 ED thoracotomy for, 203,
 207–8
 iatrogenic injury, 197
 management of, 203–8
 pericardiocentesis for, 203
 precordial wounds, 193
 penetrating injuries, neck
 trauma from, 164–66
 MDCTA for, 165–66
 selective approach to,
 165–66
 in Zone I, 165
 in Zone II, 165
 in Zone III, 165–66
 penile injuries, 244
 pericardial injury, 198
 pericardiocentesis, 203
 peri-mortem cesarean section
 procedure, 60–61
 peripheral vascular injury
 (PVI)
 evaluation of, 253–55
 extremity hemorrhage,
 249–50
 hard signs of, 252
 imaging for, 254–55
 incidence rates for, 249
 junctional hemorrhage, 249,
 251
 low risk extremity injuries,
 256
 management of, 253–55
 algorithms in, 255
 goals of, 253
 guidelines of, 253
 with tourniquets, 254
 NCTH, 249, 251
 resuscitation goals with, 249
 through DCR, 250–52
 physiological, 250–51
 secondary injuries with,
 251–53
 soft signs of, 252
 surgical indications for, 256
 types of, 249–50
 permissive hypotension, 8,
 45–46
 phalanx fractures, 303
 placental abruption, 63–64
 platelets and red blood cells
 (PRBCs), 8
 pleur-evac procedure, 176–80
 pneumothorax, 6
 Popeye sign of biceps, 284
 pRBCs. *See* packed red blood
 cells
 PRBCs. *See* platelets and red
 blood cells
 precordial penetrating wounds,
 193
 precordial wounds, 193
 pre-eclampsia, 63–64
 pregnancy, trauma during
 anatomical changes in, 56
 cardiopulmonary
 resuscitation, 59–60
 diagnostic evaluation, 63
 for pre-eclampsia, 63–64
 epidemiology of, 56
 evaluation of, 57–59
 of airway, 57
 of breathing, 57–59
 of circulation, 59
 pitfalls of, 64–65
 through secondary survey,
 61
 fetal assessment, 61–62
 through FAST, 62
 through ultrasound
 examination, 62–63
 injury mechanisms for,
 incidence rates for, 56
 management of, 57–59
 physiologic changes and,
 57–58
 pitfalls of, 64–65
 peri-mortem cesarean section
 procedure, 60–61
 physiologic changes in,
 57–58
 PSA for, 329–30
 therapeutic considerations,
 63
 for placental abruption,
 63–64
 uterine anatomy, 57
 primary survey, for trauma, 4–9
 or airway, 5–6, 9. *See also*
 airway management
 intubation assessment, 5
 or breathing, 6, 9
 hemothorax issues, 6
 pneumothorax issues, 6
 or circulation, 6–7, 9
 Shock Index, 6

- for disability, 8–9
 - neurological assessment in, 9
- for exposure of hidden injuries, 9
- GCS in, 8–9
- or hypotension, 7–8
 - differential for, 7
 - permissive, 8
 - REBOA and, 8
- of life-threatening conditions, 4–5
- overview of, 10
- for pulmonary trauma, 175
- for shock, 7–8
 - bleeding and blood loss, 7
 - blood products and, 8
- procedural sedation and analgesia (PSA)
 - ACEP clinical policy for, 322–23
 - American Society of Anesthesiology classifications, 322–24
 - discharge considerations, 325–28
 - criteria in, 328
 - for pediatric patients, 329
 - for elderly, 330
 - equipment for, 322–24
 - etomidate, 325
 - fentanyl, 325
 - indications for trauma patients, 322–23
 - ketamine, 325
 - ketofol, 325
 - medication agents, 325–27.
 - See also specific medications*
 - adjunctive treatments, 325–27
 - characteristics of, 327
 - midazolam, 325
 - in pediatric patients, 328–29
 - discharge recommendations, 329
 - duration
 - recommendations for, 329
 - sedation scales, 329
 - during pregnancy, 329–30
 - preparation for, 322
 - propofol, 325
- prothrombin complex concentrate (PCC), 8, 48
- proximal fibula fractures, 299
- proximal third tibial fractures, 298
- PSA. *See* procedural sedation and analgesia
- pulmonary contusion, 184–86
- pulmonary trauma
 - airway obstruction and, 175–76
 - in-line stabilization procedures, 176
 - esophageal disruption, 190–91
 - flail chest, 182–84
 - massive hemothorax, 180–82
 - autologous transfusion for, 180–82
 - open pneumothorax, 180
 - primary survey for, 175
 - pulmonary contusion, 184–86
 - secondary survey for, 175
 - tension pneumothorax, 176–80
 - chest tube removal, 180
 - needle compression procedure for, 176, 178–79
 - pleur-evac procedure for, 176–80
 - tracheobronchial disruption, 186–90
 - traumatic diaphragmatic injury, 184–89
- PVI. *See* peripheral vascular injury
- radiography, for abdominal and flank trauma, 223–24
- rapid-sequence intubation (RSI), 22–30
 - contraindications, 23
 - absolute, 23
 - relative, 23
 - with ELM, 30
 - emergency airway supplies, 24
 - paralyzing agents, 29–30
 - rocuronium, 29–30
 - succinylcholine, 29–30
 - vecuronium, 29–30
 - patient positioning with, 26
- predictions of difficult intubation, 23–25
 - LEMON law for, 23–24
 - Mallampati classification, 23–25
- pre-oxygenation in, 25–26
- preparation for, 25
- sedation agents, 26–29
- technique for, 22–23
- REBOA. *See* Resuscitative Endovascular Balloon Occlusion of the Aorta
- recombinant factor VII, 48
- red blood cells. *See* packed red blood cells; platelets and red blood cells
- renal injury, 219–20, 236–38
 - anatomical assessment of, 236
 - complications from, 238
 - evaluation of, 237
 - with CT imaging, 237
 - severity grading in, 237
 - grading criteria for, 221–22
 - from gunshot wound, 222
 - imaging of, 221
 - management of, 237–38
- resuscitation
 - balanced, 46
 - blood products in, 46
 - for burns, 309–12
 - cardiopulmonary, 59–60
 - crystalloid, 42–44
 - goals of, 50
 - for PVI, goals of, 249
 - through DCR, 250–52
 - physiological, 250–51
- Resuscitative Endovascular Balloon Occlusion of the Aorta (REBOA), 8
- retinal trauma, 134–35
- retrograde urethrogram (RUG), 236
- reverse Segond fracture, 293
- rocuronium, 29–30
- RSI. *See* rapid-sequence intubation
- RUG. *See* retrograde urethrogram
- Rule of Nines, for burns, 310
- scaphoid fracture, 278
- Schatzker Classification, of tibial plateau fractures, 293

Index

- SCIWORA. *See* spinal cord injury without radiographic abnormality
- scrotum and testicles, injuries to, 243–44
- seatbelt signs, neck trauma and, 169
- secondary survey, for trauma, 9–10
 overview of, 10
 during pregnancy, 61
 for pulmonary trauma, 175
- sedation agents, for RSI, 26–29
- Second fracture, 292–93
 reverse, 293
- seizure management, 99
- shaft fractures, of femur, 289–90
- shock, 7–8
 bleeding and blood loss, 7
 blood products and, 8
 geriatric trauma and, 79–80
- Shock Index, 6
- shoulder and upper arm region.
 See upper extremity trauma
- soft signs, of PVI, 252
- spinal cord injury without radiographic abnormality (SCIWORA), 149
- spinal shock, 150
- spine injuries. *See* cervical spine injuries; cervical spine trauma; lower extremity trauma; thoracolumbar trauma
- spinothalamic lesions, 155
- spinous process fractures, 160–61
- splenic injury, 217
 grading criteria for, 219–20
 from gunshot wounds, 220
 imaging for, 219
- splinting
 of femoral fractures, 291
 for isolated fibular fractures, 300
- stab wounds, abdominal and flank trauma from, 216
- steroids, for cervical spine trauma, 149
- stored whole blood (SWB), 52
- strangulation, 170–71
- subarachnoid hemorrhage, 87–90, 94
- subconjunctival hemorrhages, 138
- subdural hematoma, 87–90, 93
- succinylcholine, 29–30
- supraglottic airways, 36–37
 with i-Gel, 37
 ILMA, 36–37
 LMA, 36–37
- SWB. *See* stored whole blood
- talus fractures, 300–1
- TEE. *See* transesophageal echocardiography
- TEG. *See* thromboelastography
- temporomandibular joint (TMJ) injuries, 115–16
 reduction of, 116–17
- tension pneumothorax, 176–80
 chest tube removal, 180
 needle compression procedure for, 176, 178–79
 pleur-evac procedure for, 176–80
- testicles. *See* scrotum and testicles
- thermoregulation, with pediatric trauma, 70
- thoracic trauma, 21
- thoracolumbar trauma
 anatomic considerations for, 154–55
 burst fractures, 159
 chance fractures, 159–60
 classification of, 157
 of fractures, 158–62
 three column system, 158
 compression fractures, 158–59
 corticospinal lesions, 155
 dorsal column lesions, 155
 fracture-dislocation injuries, 160
 imaging of, 156–57
 incidence rates of, 154
 incomplete spinal cord syndromes, 161–62
 anterior cord syndrome, 162
 Brown-Sequard syndrome, 161
 central cord syndrome, 162
 pars interarticularis fractures, 160
- physical examination for, 156
 of motor groups, 156
 muscle grading in, 157
 reflex grading in, 156–57
- pre-hospital management, 155
 symptom assessment in, 155
- spinothalamic lesions, 155
- spinous process fractures, 160–61
- translational injuries, 160
- transverse process fractures, 160
- thromboelastography (TEG), 48–49
 properties, 49–50
 representation, 49–50
- tibial plateau fractures, 292–94
 associated injuries with, 293–94
 complications from, 293–94
 imaging considerations for, 293
 management of, 294
 mechanism of injury for, 292
 non-displaced bicondylar, 293
- Schatzker Classification of, 293
- Second fracture, 292–93
 reverse, 293
- tibial shaft fractures, 298
- tibial spine fractures, 294
- TMJ injuries. *See* temporomandibular joint injuries
- tourniquets, for PVI, 254
- tracheobronchial disruption, 186–90
- traction, for femoral fractures, 291–92
- tranexamic acid (TXA), 8
 in head trauma, 100
- transesophageal echocardiography (TEE), 201–2
- transfusions, in trauma
 anticoagulants, 50
 Coumadin, 50
 DOAC, 50–51
 autologous, 180–82
 balanced resuscitation, 46
 blood products in, 46
 blood products, 42–43, 48
 FFP, 46
 PCC, 48

Index

- pRBCs, 44–46
 recombinant factor VII, 48
 TXA and, 48
 whole blood, 52
 calcium and, 48
 complications from, 51–52
 classifications of, 51–52
 crystalloid resuscitation in,
 negatives of, 42–44
 DCR and, 45
 components of, 45
 hemorrhage and, 42
 ATLS and, 42
 bradycardia and, 42
 management of, 42
 hypocalcemia and, 48
 lethal triad in, 44–45
 pRBCs in, 44–45
 treatment for components
 of, 44
 massive hemothorax,
 180–82
 minimal volume
 normotension, 46
 MTPs, 46–48
 activation of, 47–48
 definition of, 46
 goal parameters, 47
 primary literature for, 47
 non-emergent, 42
 permissive hypotension,
 45–46
 pRBCs, 44–46
 resuscitation goals, 50
 TEG, 48–49
 properties, 49–50
 representation, 49–50
 trauma coagulopathy, 44–45
 whole blood in, 52
 FWB, 52
 SWB, 52
 transverse process fractures,
 160
 trauma. *See also* primary
 survey; *specific topics*
 final disposition, 12
 imaging for, 10–12
 CT scans, 11–12
 eFAST and, 10–11
 in trauma bays, 10–11
 maxillofacial, 19–20
 mortality rates, 1
 ongoing care for, 12
 secondary survey for, 9–10
 overview of, 10
 thoracic, 21
 Trauma Association, 203–5
 trauma coagulopathy, 44–45
 trauma teams, 1–2
 American College of
 Surgeons guidelines, 1
 call-and-response principle,
 1–2
 debriefing of, 2
 effective dynamics, 2
 EMS, 2–4
 CDC guidelines, 3
 EMTs, 3–4
 information during
 transition of care from,
 4
 human resource response
 suggestions, 1
 leaders, 1–2
 traumatic diaphragmatic
 injury, 184–89
 traumatic hyphema, 133–34
 evaluation of, 133–34
 traumatic uveitis, 135
 triage, for geriatric trauma,
 77–79
 pre-hospital assessment,
 77–79
 after trauma center arrival,
 78–79
 tripod fractures, 114–15
 Tufts fracture, 273
 TXA. *See* tranexamic acid
 tympanic membrane
 perforation, 111
 tympanic membrane rupture,
 310–18
 Type 2 dens fracture, 148
 ultrasound
 for abdominal and flank
 trauma, 223–24
 for cardiac trauma, 201
 fetal assessment with,
 62–63
 for genitourinary trauma,
 234–35
 for ICP, 99
 upper extremity trauma.
 See also wrist and distal
 forearm injuries
 Bennett's fracture, 275
 for bones, 273–76
 distal phalanges, 273–74
 metacarpal, 274–75
 proximal and middle
 phalanges, 274
 of elbow and forearm,
 280–84
 bone fractures and
 dislocations, 280–84
 Popeye sign of biceps,
 284
 of soft tissues, 284
 hand and finger injuries
 mallet finger, 274
 surgical consultation for,
 273
 incidence rates, 273
 innervation of the hand,
 276
 joint dislocations, 276
 neck fractures, 275
 of shoulder and upper arm
 region, 284–87
 anterior shoulder
 dislocation, 285–86
 bone fractures and
 dislocations, 284–87
 comminuted clavicular
 fractures, 285
 vessel injury, 287
 of soft tissues, 276–77
 in elbow and forearm,
 284
 in wrist and distal forearm
 injuries, 280
 Tufts fracture, 273
 ureter injuries, 238–39
 urethral injury, 241–43
 classification of, 241–42
 management of, 242–43
 urinalysis, 233–34
 uterine anatomy, during
 pregnancy, 57
 vascular injuries, 150
 vasopressors, 97–98
 vecuronium, 29–30
 vehicle accidents
 abdominal and flank trauma
 from, 216
 geriatric trauma from, 82
 vessel injury, 287
 vitreous hemorrhage, 134
 warfarin. *See* Coumadin
 WBCT scans. *See* whole body
 CT scans
 white-eyed fracture, 141
 whole blood, 52
 FWB, 52
 SWB, 52

Index

whole body CT (WBCT) scans, 11–12	of abnormal posterior fat pads, 280–81	scaphoid fracture, 278
wound care, with burns, 314–15	bone fractures and dislocations, 277–80	of soft tissue, 280
wrist and distal forearm injuries, 277–80	Colle’s fracture, 277	X-ray imaging
of abnormal anterior fat pads, 280–81	lunate dislocation, 280	of cardiac trauma, 202–3
	Monteggia fracture- dislocation, 282	for great vessel injury, 202–3
		zygomatic bone fractures, 114