



Contents

<i>List of Figures</i>	<i>xi</i>
<i>List of Tables</i>	<i>xv</i>
<i>Preface</i>	<i>xvii</i>
<i>Acknowledgments</i>	<i>xix</i>
1 Introduction	1
1.1 Characteristics of Remotely Sensed Imagery	3
1.1.1 Multi-spectral images	8
1.1.2 Panchromatic image	10
1.1.3 Hyper-spectral images	12
1.2 Low Spatial Resolution Imaging	14
1.3 Image Fusion in Remotely Sensed Images	17
1.4 Multi-resolution Image Fusion: An Ill-posed Inverse Problem	18
1.5 Indian Remote Sensing Satellites	20
1.6 Applications of Image Fusion	22
1.7 Motivation	25
1.8 Organization of the Book	25

2 Literature Review	27
2.1 Projection Substitution Based Techniques	29
2.2 Multi-resolution Based Techniques	34
2.3 Model Based Fusion Approaches	41
2.4 Hyper-spectral Sharpening Methods	48
2.5 Conclusion	50
3 Image Fusion Using Different Edge-preserving Filters	52
3.1 Related Work	53
3.2 Fusion Using Multistage Guided Filter (MGF)	54
3.2.1 Multistage guided filter (MGF)	55
3.2.2 Proposed approach using guided filter	57
3.3 Fusion Approach Using Difference of Gaussians (DoGs)	59
3.3.1 Difference of Gaussians (DoGs)	60
3.3.2 Proposed approach using DoGs	61
3.4 Experimental Illustrations	63
3.4.1 Experimentations: Ikonos-2 dataset	67
3.4.2 Experimentations: Quickbird dataset	71
3.4.3 Experimentations: Worldview-2 dataset	74
3.4.4 Computational complexity	78
3.5 Conclusion	78
4 Image Fusion: Model Based Approach with Degradation Estimation	80
4.1 Previous Works	81
4.2 Description of the Proposed Approach Using Block Schematic	83
4.3 Background: Contourlet Transform (CT)	85
4.4 Contourlet Transform Based Initial Approximation	85
4.5 Forward Model and Degradation Estimation	88
4.6 MRF Prior Model	91
4.7 MAP Estimation and Optimization Process	94
4.7.1 MAP estimation	94
4.7.2 Optimization process	95

Contents	ix
4.8 Experimentations	96
4.8.1 Effect of decimation matrix coefficients on fusion	100
4.8.2 Effect of MRF parameter γ_m on fusion	102
4.8.3 Fusion results for degraded dataset: Ikonos-2	103
4.8.4 Fusion results for degraded dataset: Quickbird	112
4.8.5 Fusion results for degraded dataset: Worldview-2	120
4.8.6 Fusion results for un-degraded (original) datasets: Ikonos-2, Quickbird and Worldview-2	126
4.8.7 Spectral distortion at edge pixels	133
4.8.8 Computational time	137
4.9 Conclusion	139
5 Use of Self-similarity and Gabor Prior	140
5.1 Related Work	141
5.2 Block Schematic of the Proposed Method	143
5.3 Initial HR Approximation	144
5.4 LR MS Image Formation Model and Degradation Matrix Estimation	150
5.5 Regularization Using Gabor and MRF Priors	152
5.5.1 Optimization process	155
5.6 Experimental Results	156
5.6.1 Experimental setup	158
5.6.2 Experimental results on degraded and un-degraded Ikonos-2 datasets	159
5.6.3 Experimental results on degraded and un-degraded Quickbird datasets	164
5.6.4 Experimental results on degraded and un-degraded Worldview-2 datasets	169
5.6.5 Comparison of fusion results with CS and TV based approaches	174
5.6.6 Computation complexity	178
5.7 Conclusion	179

6 Image Fusion: Application to Super-resolution of Natural Images	180
6.1 Related Work	181
6.2 Estimation of Close Approximation of the SR Image	184
6.3 Refining SR Using MAP–MRF Framework	188
6.4 MRF Prior and SR Regularization	190
6.4.1 Optimization process	191
6.5 Experimental Demonstrations	192
6.5.1 SR results on gray scale images	193
6.5.2 SR results on color images	197
6.6 Conclusion	202
7 Conclusion and Directions for Future Research	203
7.1 Conclusion	203
7.2 Future Research Work	207
<i>Bibliography</i>	<i>211</i>