Index

1,2,3 kg – 7,8,9 cm rule, 44
12 cm + ½ cm per year, 44
2–4–6-hour rule, 14

abdominal wall perforation, 152–153
acetylcholine, 196
activated charcoal, 206
acute lymphoblastic leukaemia, 120
ADARPEF study, 138, 155
adenotomy, 205
adequate anaesthetic plan, 8–10
admixture, spurious readings caused by, 24–25
adrenaline, 42, 44, 115–116, 138, 156
  intramuscular, 226
age-based formulas, insertion depth of ETT, 44–45
air embolism, 115–116, 122
  flushing, 123
air leak syndrome, 50
airway-related problems, 29–31
  Down syndrome, 212
airway trauma, prevention, 48–49
allergic non-haemolytic reactions, 250
altered metabolism, 191, 193
altered pharmacodynamics, 193
amnesia, 172
amniocentesis, 47
anaemia, 23, 250
anaesthesia
  depth, 60–61
  failed induction, 29–31
  optimal system for induction, 29
  outpatient, 84
  regional. *See* regional anaesthesia
  total intravenous, 165, 170–172
  trigger-free, 207
  wrong system, 29
anaesthetic plan, adequate, 8–10
anaesthetist
  experience, 60
  faulty manipulations, 30
  inattentive, 38
  open mind, 232
  supervising trainees, 157
  support from parents, 222
  unprofessional behaviour, 221–222
anaphylaxis, 225–227
  diagnosis, 226
  treatment, 226
angulation, left vena cava, 112
anisocoria, 135, 178
anthracycline, 215
antibiotics, 70, 91, 152, 155
anticholinergics, 196
anticipation, surgical complications, 17, 19
antidiuretic hormones, 180–181
antimicrobials, 229
anxiety disorder, 221
apnoea, 161
  obstructive sleep. *See* obstructive sleep apnoea
appendectomy, 109, 153
appendicitis, 180
Arndt blocker, 184
arteria lusoria, 11–12
arterial catheter, insertion, 121
arterial line, previously placed, 100
asparaginase, 120
aspiration
  foreign body, 72
  pulmonary, 68–70
asthma, 44–45, 211
atracurium, 100, 103–104, 161
atropine, 39, 87, 103
attention deficit hyperactivity disorder (ADHD), 223
autism spectrum disorder (ASD), 223
axillary artery, 121

bacteraemia, 91, 229
bacterial contamination, 249
bat ear surgery, 134, 195
Becker muscular dystrophy, 132
behavioural disorders, 221–223
benzocaine, 141
Berger, Professor Thomas M., ix, 1
beta-blockers, 9, 177
beta-2-agonists, 45
beta-stimulating agents, 87
Bier's block, 138
BIPAP ventilation, 130
BIS (bispectral index), 165, 195, 221
bladder augmentation, 86
Blalock–Taussig procedure, 121
blood transfusion
  hazards, 249
  reaction, 249–250
body temperature
  maintaining, 20–22
  raised, 237
brachial plexus block, 149–150
brachial plexus palsy, 240, 242
bradycardia, 212
brain oedema, 180, 219
bronchial blockers, 184
bronchial foreign body, 71–73
bronchial rupture, 50–51
  treatment modalities, 51
bronchoscopy, 50–51
  bronchial foreign body, 71–73
  tracheal stenosis, 81
bronchospasm, 225, 227
Broviac catheter, 115, 172
bupivacaine, 144
burns, iatrogenic, 243–244
button batteries, 78
butyrylcholinesterase, 191–193

calcium solutions, 98
can intubate, cannot ventilate, 35–37, 45
  differential diagnosis, 36
cannot intubate, cannot ventilate, 32–35
capital femoral epiphysis, slipped, 118

© in this web service Cambridge University Press www.cambridge.org

CAMBRIDGE

Cambridge University Press
978-1-316-62910-9 — Managing Complications in Paediatric Anaesthesia
Martin Jöhr
Index
More Information

capnography, 25
carbon monoxide, 168
cardiac arrest, 17, 85, 144, 187–188
  laryngospasm, 38
cardiac toxicity, 138–139, 174
cardiomyopathy, 132
caries, 53–54
cataracts, congenital, 177
catecholamine, 214
catheter, depth of insertion, 109
catheter tip, malposition, 109–111
caudal block, 131, 135, 143, 147
  complications, 155–157
  diagnosis, 155
  infections from, 156
  therapy, 155
Cavilon, 247
central anticholinergic syndrome (CAS), 196–197
central venous access
  femoral approach, 63
  peripheral insertion, 94, 110–111
  thrombosis, 118–119
cephalhaematoma, 106
cerebellar tumour, 180
cerebral palsy, 20, 22
cerebral salt-wasting syndrome, 181
cerebral toxicity, 138–139
cervical myelopathy, 212
cervical spine, extension, 44
Charcot–Marie–Tooth disease, 208
chemotherapy, 215
chloroprocaine, 138, 193
circle system, 29–30
circulatory overload, 250
circumcision, 56, 199
cleft palate, 11, 32–33, 65, 71–72, 74
  Down syndrome, 212
  impression, 72
  plate, 72
closed-loop communication, 162, 164
cold light, 62, 243–244
colonic puncture, 152–153
colour coding
  labelling, 161
  three-way stopcocks, 100
communication
  between team members, 185
  closed-loop, 162, 164
  failure, 164
  inadvertent drug administration, 168
  team performance, 164, 242–243
  with parents, 193
compartment syndrome, 91, 95, 146–149
  prevention, 147
  signs, 147
  situations at risk, 147
congenital cataracts, 177
congenital diaphragmatic hernia (CDH), 68–69, 79–85
congenital heart disease, 83, 112
  Down syndrome, 211
congenital naevus, resection, 41
contraceptives, oral, 232
convulsions, 137–138
Cook cannula, 50, 91
Cornelia de Lange syndrome, 11–12
corticosteroids, 42, 45, 226
cortisol concentration, 188
cranial nerve injury, 56
craniocervical instability, 212
craniosynostosis, 116, 217
creatine kinase, 146, 205, 237–238
cricothyroid membrane, perforation, 48–49
cuff pressure, 57
cuffed ETTs, 42–43
cyanosis, 180
  congenital heart disease, 83
  methaemoglobinaemia, 140, 142

dantrolene, 205–207
deciduous tooth, loose, 54
delayed awakening, 195–197
dental caries, 53–54
dental treatment, 211
depth of anaesthesia, 60–61
depth of insertion, catheter, 109
desflurane, 168, 199
desiccated absorbent, 168
desmopressin, 181
dexamethasone, 42, 215
dexmedetomidine, 165, 200, 212
diabetes, 217–219
  major and minor surgery and, 218
  management, 217
  treatment, 218
  type 1, 217–219
  type 2, 217
diaphragmatic hernia, 94
  congenital, 68–69, 79–85
diazepam, 188
differential diagnosis, 23
  can intubate, cannot ventilate, 36
  delayed awakening, 196
  oesophageal foreign body, 78
  pharyngeal foreign body, 74
  pulmonary aspiration, 69
  sepsis, 229
difficult venous access, 91–92, 100, 106
dilatation procedure, 106
Dinamap, 8
dosing errors, 162, 165, 184
  overdose. *See* drug overdose
double inferior vena cava, 113
Down syndrome, 211–212
  perioperative complications, 211
drug administration
  dosing errors, 162, 165, 184
  failed, 170
  inadvertent, 167–169
  inadvertent, of residual, 103–104
  total intravenous anaesthesia, 170–172
drug overdose, 164–166, 197
  prolonged paralysis, 193
  propofol, 164, 195
Duchenne muscular dystrophy, 130–131, 206, 209
duodenal atresia, 211
dyspnoea, 44
dystrophin, 132

echocardiography, neonates, 63
elbow fracture, 149–150
electric current, burns from, 243
electroencephalogram (EEG)
  emergence delirium, 199
  isoelectric, 164

© in this web service Cambridge University Press www.cambridge.org

Index

electroencephalogram (EEG) (cont.)
monitoring, 165, 170
supraglottic airway use, 60–61
electronic prescribing, 162
emergence delirium, 199–201
diagnosis, 199
pathophysiology, 199
prevention of, 200
treatment, 200
emphysema, subcutaneous, 50–51
endotracheal tube (ETT)
bronchial rupture, 50–51
cuffed, 42–43
inappropriate size, 41–43
insertion depth, 44–45
lost, 77
nasotracheal, 48
ENT surgery, 50, 178
body temperature, 22
enuresis, secondary, 217
Epican Paed, 155
epidural blockade, 135, 143
extension, 232
epiglottitis, 75
epispadias, 23–24
Escherichia coli, 228–229
ethyl ether, 247
etiquette, 222–223
EXIT (ex utero intrapartum treatment), 47–48
exothermic reaction, 168
external jugular vein, 95
extravasation, 94
prevention, 95
treatment, 95
eye drops, 177–178
EZ-IO system, 91–92

facial nerve palsy, 134
factor V Leiden mutation, 119
failed anaesthesia induction, 29–31
failed drug administration, 170
failure in communication, 164
fascia iliaca block, 137
fasciotomy, 95
fasting guidelines, 14, 16
fasting period, 14–16, 218
faulty manipulations, anaesthetist, 30
febrile non-haemolytic reactions, 250
femoral approach
malposition, 110
venous access, 63
femoral block, 137–138, 146–147
femoral fracture, 97, 137–138, 241
fentanyl, 138, 170–171
fetus, laryngeal perforation, 47
fibreoptic intubation, LMA, 32–33
fibrinogen, 19
fire risk, 244
fluid administration, 219
flushing
air embolism, 123
three-way stopcock, 103–104
Fogarty catheter, 183, 185
forced-air warming devices, 20–22, 243
foreign body
bronchial, 71–73
oesophageal, 12, 35, 77–79
pharyngeal, 74–76
pulmonary oedema, 65, 67
unobserved ingestion, 78
fractures, from positioning, 241
Frei endoscopy mask, 71

gastroenteritis, 91
gastro-oesophageal reflux, 77
gastroschisis, 62–63
giant neck mass, 47
Glasgow Coma Scale (GCS), 50
glucose, 188
concentration, 217, 219
requirements, 218
glucose-6-phosphate dehydrogenase deficiency, 142
Guillain–Barré disease, 209

haemangioma, 8–9
haemodynamic stability, 144
haemolytic transfusion reaction, 249
halothane, 177
heparin, 101, 118, 120, 122
high-frequency oscillatory ventilation (HFOV), 42
Hirschsprung disease, 211
histamine, 226
Horner syndrome, 134–135
hydatic disease, 184
hydration, sufficient, 15
hydrogen peroxide, 116
hyperchloraemia, 219
hyperchloraemic acidosis, 181, 183–185
hyperkalaemia, 187–189, 209
causes of, 187
diagnosis, 188
prevention, 187
treatment, 188
hypertension
oncologic disease, 214, 216
pulmonary, 84, 212
hyperthermia, malignant, 132, 205–207, 209, 237
hypertonic saline, 183
hypnosis, 172
hypnotic–saline–relaxant–saline sequence, 104
hypoganglionosis, 115
hypoglossal nerve, 56
hypoglycaemia, unrecognized, 218
hyponatraemia, 180–182, 206
hypospadias, 59, 199, 237–238
hypotension, 144
hypothermia, intraoperative, 20–21
hypotonic fluids, 180–181, 184
hypovolaemia, 17–18
hypoxia, opioid sensitivity, 82

iatrogenic burns, 243–244
iatrogenic hyponatraemia, 181–182
ilioinguinal nerve block, 152–153, 155
in vitro contracture, 206
inadequate technique, 130–132
inadvertent drug administration, 103, 105, 167–169
inattentive anaesthetist, 38
induction of anaesthesia
failed, 29–31
optimal system, 29
inferior vena cava anomalies, 113
infiltration of venous line, 94
infusion therapy, 115–116
inguinal hernia, 29, 53, 83–85, 100, 144, 153
inherited thrombophilia, 119
insertion depth, ETT, 44–45
insulin, 188, 217–218
Insyte, 106, 121

© in this web service Cambridge University Press www.cambridge.org

CAMBRIDGE

Cambridge University Press
978-1-316-62910-9 — Managing Complications in Paediatric Anaesthesia
Martin Jöhr
Index
More Information

internal jugular vein approach, 62
internal thoracic vein, malposition in, 114
intra-arterial injection, 98, 100–101
  cannula placement, 100
intraoperative hypothermia, 20–21
intraosseous cannulation, 91
  complications, 91
  indication for, 92
  semi-elective use of, 92
intravascular haemolysis, 249
invasive monitoring, 9
ischaemic extremity, 149–150
ischaemic fingers, 121–122
isotonic fluids, 181

Jackson Rees modification of the T-piece, 30

Kasabach–Merrit syndrome, 9
Kataria model, 165
ketamine, 98, 161, 200
  laryngospasm, 39
ketoacidosis, 217, 219

laboratory results, unexpected, 23–25
lactate concentration, 174–176
Lanoxin, 8
laparoscopy, 109, 116
laparotomy, 121, 170, 243
laryngeal mask airway (LMA)
  cuff pressure, 57
  fibreoptic intubation, 32–33
  lost, 77
  morbidities, 56
  nerve injuries, 57
  size, 57
laryngeal nerve damage, 57
laryngeal perforation, 47–49
laryngeal stenosis, 80
laryngoscopy, 33, 47
  impossible, 205
  pharyngeal foreign body, 74
laryngospasm, 38–39
  pathophysiology, 38
latex allergy, 225–226
LED (light-emitting diode) devices, 243–244
left vena cava, angulation, 112
leg ischaemia, 106
leucodepletion, 250
levobupivacaine, 137, 162
lidocaine, 39, 141
lingual nerve neurapraxia, 56–57
lip laceration, 38
lithotomy position, 238
local anaesthetic toxicity, 137–139, 156
local treatment, systemic effect, 177–178
lung hypoplasia, 84
lung separation, 184

malignancy, and thrombosis, 120
malignant hyperthermia, 132, 205–207, 209, 237
  blood chemistry, 205
  diagnosis, 206
  management, 206
  neuromuscular disease, 206
  testing, 206
malposition
  catheter tip, 109–111
  internal thoracic vein, 114
manual jet ventilation, 51, 80
massive bleeding, anticipating, 17–19
meatoplasty, 199
mediastinal mass syndrome, 215
medical charts, 232
medication errors, 161–164
  resilience against, 161
medication-related problems, 161–201. *See also specific drugs*
medulloblastoma, 174
metabolic acidosis, 174–175, 183
metabolism, altered, 191, 193
methaemoglobin, 137, 140
methaemoglobinaemia, 140–142
  drugs, 141
  treatment, 142
methotrexate, 162
methylene blue, 140, 142
methylphenidate, 223
metoclopramide, 197
MicroCuff, 42
midazolam, 100
milk, regurgitation, 14–16
mitochondrial disorders, 176
mitochondrial toxicity, 174
mivacurium, 104, 191–193
mock block initiative, 128
molecular genetics, 206–207
morphine, 171
MRI examination
  body temperature, 21
  burns from, 243
muscle biopsy, 206
muscle relaxants, 161–162, 209
muscular dystrophy, 130–131, 206, 209
myasthenia gravis, 208
mydriasis, 178
myeloproliferative disorder, 212
myotonia congenita, 206, 209
myotonic dystrophy, 208–209

nalbuphine, 162
naloxone, 188
Narcotrend, 165
nasotracheal intubation, 48
needle damage, 152–154
needles, types, 153
neonate
  air embolism, 116
  arterial catheter insertion, 121
  bronchial foreign body, 71
  cleft palate impression, 72
  Down syndrome, 212
  drug injections, 100
  echocardiography, 63
  extravasation, 94
  failed drug administration, 170
  ischaemic fingers, 121–122
  outpatient anaesthesia, 84
  pneumothorax, 62
  pulse oximetry screening, 83
  unknown pre-existing pathology, 83
  venous thrombosis, 118, 120
neostigmine, 103, 192
nerve injuries
  caudal block, 156
  cranial, 56
  lingual, 56–57
  LMA devices, 57
  peroneal, 241
  pre-existing deficit, 149
neuroblastoma, 214–216
neurodegenerative disorders, 20, 22
neuromuscular blockade, 60, 103, 193
  time course, 104

© in this web service Cambridge University Press www.cambridge.org

Index

neuromuscular disease, 208–209
classification, 208
malignant hyperthermia, 206
neutrophilia, 212
Niltac, 247
nitrous oxide, 167–168
non-depolarizing relaxants, 209
non-Luer connectors, 162
NSAIDs, 157
Nuss procedure, 240–241

obstructive sleep apnoea, 80–81, 84, 212
occipital craniotomy, 174
oesophageal dilatation, 35
oesophageal foreign body, 12, 35, 77–79
differential diagnosis, 78
omphalocoele, 64
oncologic disease, 214–216
open leg fracture, 249
open mind, 78, 232
open questions, 232
open system, 29–30
open vaporizer, 168
ophthalmic surgery, 178
opioids, 195, 200
high-dose, 170
hypoxia, 82
nasal, 221
oral contraceptives, 232
orchidopexy, 143, 155, 221
osteogenesis imperfecta, 54
osteomyelitis, 91, 161
osteotomy, 127
bilateral, 146
outpatient anaesthesia, 84
overdose. *See* drug overdose

Paedfusor model, 165, 195
pain
self-limited back pain, 156
severe, 187, 208, 221
pancuronium, 170, 193
paralysis, prolonged, 191–193
paraplegia, 209
paravenous drug injection, 97–99
correct positioning, 97
tissue damage from, 98
patient blood management, 250
patient identification, 5
percutaneous gastrostomy, 36
perioperative complications, 211
diabetes, 218
perioperative medicine, 242–243
peripherally inserted central catheters, 94, 111
peroneal nerve injuries, 241
persistent left superior vena cava (PLSVC), 112
petroleum ether, 246–248
phaeochromocytoma, 214
pharmacodynamic feedback, 170, 172
pharmacodynamics, altered, 193
pharyngeal foreign body, 74–76
differential diagnosis, 74
examples, 74–75
phenylephrine, 178
phrenic nerve, 134
physiology, understanding, 11–12
physostigmine, 196
Pierre Robin sequence, 32–33
planned spinal anaesthesia, 144
plastic cannula, 106, 121
platelet concentrates, 250
Plexufix, 153
pneumomediastinum, 50
pneumoperitoneum, 44
pneumothorax, 62–64, 71, 81, 243
diagnosis, 62
tension, 62, 71
treatment, 63
pocket lamps, 243
polyhydramnios, 47
polysomnography, 81
popliteal sciatic nerve block, 140
positioning
damage caused by, 238, 240–242
malposition in internal thoracic vein, 112
malposition of catheter tip, 109–111
post-extubation stridor, 42
post-obstructive pulmonary oedema, 65–67
prevention, 66
treatment, 66
potassium chloride, 162
potassium shift, 188
potassium, concentrated, 187
pre-existing neurologic deficit, 149
pre-existing pathology, unknown, 83–85
pregnancy, 231–232
potential existence, 231
unrecognized, 231
premature infants, anaesthesia, 84
pressure caused rhabdomyolysis, 237
pressure sores, 238, 241
pressure, damage caused by, 237–239, 241
prilocaine, 137, 140–141
primary dentition, 53
prolonged paralysis, 191–193
diagnostics, 192
overdose, 193
treatment, 192
propofol, 98, 100, 161
clinical use, 175
emergence delirium, 200
failure in administration, 170
laryngospasm, 38
overdose, 164, 195
propofol infusion syndrome, 174–176
pathophysiology, 174
re-exposure, 175
propranolol, 9
ProSeal, 59–60
psoas compartment block, 136, 144
psychogenic hyperventilation, 187
psychologist, 221–222
pulmonary aspiration, 68–70
differential diagnosis, 69
risk for, 69
steps to be taken, 70
pulmonary cystic echinococcosis, 183
pulmonary hypertension, 84, 212
pulmonary oedema, 250
post-obstructive, 65
pulse oximetry
methaemoglobinaemia, 140, 142
neonatal screening, 83
pressure injuries from, 238
pyloromyotomy, 103

© in this web service Cambridge University Press www.cambridge.org

CAMBRIDGE

Cambridge University Press
978-1-316-62910-9 — Managing Complications in Paediatric Anaesthesia
Martin Jöhr
Index
More Information

radial artery, 121
reassessments, importance, 18
rectal puncture, 155, 157
rectus sheath block, 152
red blood cells, rapid transfusion, 187
re-exposure propofol infusion syndrome, 175
regional anaesthesia, 127–157
  choice of technique, 153
  inadequate technique, 130–132
  masking compartment syndrome, 147
  needle damage, 152–154
  neuromuscular disease, 208
  renouncing, 149
  undesirable block extension, 134–136, 156
  wrong-site block, 127–128
regurgitation
  milk, 14–16
  supraglottic airway use, 59–61
remifentanil, 103, 171, 174, 196
respiratory syncytium virus (RSV), 39, 42
restraint, physical, 222
rhabdomyolysis, 132, 206, 209
  pressure-caused, 237
rhinovirus, 86–87
risk factors
  compartment syndrome, 147
  fire, 244
  pulmonary aspiration, 69
  wrong-site block, 127
rocuronium, 104, 209
ropivacaine, 137, 162
Rotem, 19
ryanodin-1 receptor (RYR1) gene, 206

safety rules, 5–6, 99
saline
  balanced solution, 181
  hypertonic, 183
  hypnotic–saline–relaxant–saline, 104
salmonella, 91
salt solution, balanced, 181
sampling technique, inadequate, 24
saphenous nerve, 130
Scandinavian fasting guidelines, 15
Schnider model, 165, 195
sciatic nerve block, 127–128, 146–147
  popliteal, 140
scintigraphy, full body, 161
scoliosis surgery, 130–131
Seldinger technique, 121
self-limited back pain, 156
sensitization, 226
sepsis, 228–230
  diagnosis, 228
  differential diagnosis, 229
sevoflurane, 130, 132, 167, 199, 212
SIADH (syndrome of inadequate secretion of antidiuretic hormone), 181
Sikich and Lerman scoring system, 199
skin blanching, 100, 106
skin disinfection, 247
skin flash, 161
skin laceration, 191
skin necrosis, 238, 241
  toxic, 246–248
slipped capital femoral epiphysis, 118
slow recovery, unexpected, 195–197
spermatic vein ligation, 152–153
spina bifida, 226
spinal anaesthesia, 130–131
  planned, 144
  total. *See* total spinal anaesthesia
spinal cord injury, 212
spurious laboratory results, 23–25
standard of care, 116
Staphylococcus aureus, 152–153
stop before you block initiative, 128
stress reaction, 180
stridor, post-extubation, 42
subclavian vein catheterization, 62
subclavian vein thrombosis, 109, 111
subglottic stenosis, 41–42, 211–212
succinylcholine, 98, 132, 191–193, 205, 209
sugammadex, 209, 232
supraglottic airway
  damage, 56–57
  regurgitation, 59–61
surgery
  anticipation of complications, 17, 19
  thrombosis, 120
  understanding of procedure, 11–12, 19
  wrong site, 5–6
surgical checklist, 5–6, 127
  before induction of anaesthesia, 6
  WHO, 128
surgical positioning, 238, 240–242
sympatholytic effects, 108
systemic effect, local treatment, 177–178

target-controlled infusion (TCI), 165
Taylor's approach, 131
team performance, 164, 243
teeth, damage to, 53–55
temperature
  maintaining, 20–22
  measuring, 21
  raised, 237
temporal artery catheters, 122
tension pneumothorax, 62, 71
tetralogy of Fallot, 83
thiopental, 98–101, 138
three-way stopcock, 103–105, 170
  colour coding, 100
  importance of flushing, 103–104
thrombectomy, 118
thrombophilia, inherited, 119
time-out, 5–6
timolol, 177
tissue damage, 98
TIVA-Trainer, 195
tonsillectomy, 80–81
total intravenous anaesthesia (TIVA), 165, 170–172
total spinal anaesthesia, 143–145, 156
  clinical signs, 144
  haemodynamic stability, 144
  incidence, 143–144
  prevention, 145
  speed of onset, 144
toxicity
  cardiac, 138–139, 174

© in this web service Cambridge University Press www.cambridge.org

Index

toxicity (cont.)
  cerebral, 138–139
  local anaesthetic, 137–139, 156
  mitochondrial, 174
  skin necrosis, 246–248
tracheal rupture, 50–51
tracheal stenosis, unanticipated, 80–82
tracheotomy, 80, 211
train-of-four stimulation, 103–104, 191
tranexamic acid, 106, 250
transfusion reaction, 249–250
transillumination, 62, 121, 243–244
transverse abdominal plane (TAP) block, 152
trigger-free anaesthesia, 207
trigonocephaly, 14
tropicamide, 178
tryptase, 226
tubing, wrong connection, 31
tumour lysis syndrome, 215
type 1 diabetes, 217–219
type 2 diabetes, 217
type and screen test, 249

ulnar artery, 121
ulnar nerve injuries, 240
ultrasound
  arterial catheter insertion, 121
  paravenous injection, 98
  pregnancy, 231
  vascular damage prevention, 108
  venous thrombosis, 119
umbilical artery catheter, 101
unanticipated tracheal stenosis, 80–82
uncooperative child, 222–223
understanding, surgical procedure, 18–19
undesirable block extension, 134–136, 156
unexpected slow recovery, 195–197
  anaesthesia chart, 196
  differential diagnosis, 196
  drug overdose, 195, 197
unknown pre-existing pathology, 83–85
unnecessary suffering, 221
upper respiratory tract infection, 86–88
  laryngospasm, 39
urological surgery, 246

vaccination reaction, 228, 230
vagal nerve, 134
vascular access, 91–122
vascular damage, 106–108
vascular insufficiency, 150
Vasofix, 121
venous access
  central. *See* central venous access
  difficult, 91–92, 100, 106
  infiltration of a line, 94
  malposition of catheter tip, 109–111
  paravenous drug injection, 97–99
  securing of, 66
  skin necrosis, 238
venous system, anomalies, 112–113
venous thrombosis, 118–120
  prophylaxis, 118, 120
Ventrain, 51, 81
vincristine, 162, 215
viral diseases, 249

WHO surgical checklist, 128
wrong anaesthetic system, 29
wrong blood in tube, 249–250
wrong-site block, 127–128
  incidence, 127–128
  prevention, 128
  risk factors, 127
wrong-site surgery, 5–6

© in this web service Cambridge University Press www.cambridge.org