# Index

Abbreviated Burn Severity Index (ABSI), 261-264 ABCDE protocol of ATLS, 18, 21 - 22Abdominal and Cardiac Evaluation with Sonography in Shock (ACES), 140 abdominal compartment syndrome, 240-241 abdominal trauma anatomic considerations, 232 angiographic embolization, 242-243 blunt trauma, 232 damage control laparotomy, 239-240 definitive repair surgery, 241 diagnostic studies, 234-235 early involvement of the anesthesiologist, 233 exploratory laparoscopy, 237 exploratory laparotomy, 237-239 great vessel injuries, 242 gunshot wounds, 233 hemodynamic instability in the preoperative period, 235 intraoperative management, 177 - 179intraperitoneal injuries, 241 key points, 244-245 management challenges, 232 mechanisms of injury, 232 - 233patient history, 233-234 penetrating trauma, 232-233 physical examination of the abdomen, 234 preoperative evaluation and management, 233-235 re-exploration surgery, 240-241 release of abdominal compartment syndrome, 240-241

resuscitative endovascular balloon occlusion of the aorta (REBOA), 243-244 retroperitoneal injuries, 242 stab wounds, 232 acid-base balance postoperative considerations, 169-170 acute respiratory distress syndrome (ARDS), 167 acute traumatic coagulopathy (ATC), 70-71 adductor canal block, 116-117 Advanced Cardiac Life Support (ACLS) pregnant patient, 304-305, 312 Advanced Trauma Life Support (ATLS), 18-22 ABCDE protocol, 18, 21-22 golden hour concept, 18 guidelines for initial airway and ventilatory management, 29-31 scene safety, 11 use of supraglottic devices, 37 Agency for Healthcare Research and Quality, 57 Aintree intubation catheter, 37-39 air bags injuries caused by, 8 prevention of trauma deaths, 7 airway compromise general anesthesia, 90 airway exchange catheter (AEC), 37 airway management abdominal trauma, 235 ASA difficult airway algorithm modified for trauma patients, 35-36 ATLS guidelines for initial management, 29-31 ATLS protocol, 18 awake intubation, 34 burn patients, 270-271

causes of obstruction or inadequate ventilation, 20 cervical spine injury, 34 concerns about the use of cricoid pressure, 33-34 considerations for trauma patients, 29 cricothyroidotomy, 40 difficult airway, 37 emergency airway management in maxillofacial trauma, 206-207 equipment required to secure the airway, 31 failed intubation, 40 flexible bronchoscopy, 40 general anesthesia, 88-90 geriatric patients, 296-297 indications for tracheal intubation, 30-31 induction agents, 32-33 key points, 42 laryngeal mask airway (LMA), 38-39 laryngeal tube airway (LTA), 39 maxillofacial trauma surgery, 209 - 210medications used in securing the airway, 31 neuromuscular blocking drugs (NMBDs), 32-33 oxygenation for RSI, 33-34 pediatric patients, 278-279 penetrating neck injury, 41 pregnant trauma patient, 309-310 prehospital care, 11 primary survey, 21 rapid sequence induction (RSI), 32-33 securing the airway in a trauma setting, 29-31 spinal cord injury (SCI), 189-193 supraglottic devices, 37-39 surgical airway, 40

317

# CAMBRIDGE

Cambridge University Press & Assessment 978-1-316-63671-8 — Essentials of Trauma Anesthesia Albert J. Varon, Charles E. Smith Index More Information

318

### Index

airway management (cont.) surgical airway in maxillofacial trauma, 208 tracheal tube introducers, 37 tracheostomy, 40 transfer from prehospital to hospital care, 29-31 use of manual in-line stabilization (MILS), 34 videolaryngoscopy, 35-37 albumin, 52 aminocaproic acid, 78 amnestic drugs, 84 amputation traumatic, 256 anemia traumatic brain injury (TBI) and, 181-182 anesthesiologists challenges of monitoring trauma patients, 124 challenges presented by trauma patients, 82 contribution to massive transfusion protocols, 76 role in initial evaluation and management, 18-20 role in trauma care, 16-17 role in traumatic brain injury (TBI), 173 anesthetic agents/management abdominal compartment syndrome, 240-241 abdominal trauma, 236-237 angiographic embolization, 242-243 burn patients, 96 burns, 269-273 cardiac injury, 95-96 damage control laparotomy, 239-240 exploratory laparotomy, 238-239 general anesthesia, 91-94 geriatric patients, 290-295, 299-302 great vessel injuries, 242 head injury, 94-95 hypovolemia, 91-94 intraperitoneal injuries, 241 maxillofacial reconstructive surgery, 210 musculoskeletal trauma, 247, 250 - 251ocular trauma, 202-204 pediatric patients, 212-213

pregnant trauma patient, 306-313 retroperitoneal injuries, 242 traumatic brain injury (TBI), 179 - 180angiographic embolization, 242-243 angiography during initial evaluation and management, 24-25 antecubital vein cannulation, 58 anterior cord syndrome, 192 antibiotics, 84 anticoagulated trauma patient, 78 - 80antifibrinolytic agents, 77-78, 195 aortic dissection, 95 diagnostic echocardiography, 144-145 aortic trauma, 228-229 arterial catheterization, 66-67 axillary artery, 67 brachial artery, 67 dorsalis pedis artery, 67 femoral artery, 67 indications for, 66 infection rates, 66 monitoring trauma patients, 126-129 radial artery, 66-67 sites for, 66-67 AS1 solution (adenine, glucose, mannitol, and sodium chloride), 71 ASA difficult airway algorithm modified for trauma patients, 35-36 supraglottic devices, 37 aspiration prophylaxis, 88-90 atropine, 84 awake intubation, 34 awareness monitoring, 128 axillary artery catheterization, 67 axillary block, 112-113 bag-valve mask (BVM), 30, 33 base deficit serum markers for, 135 - 136Beck's triad, 226

mechanisms of injury, 9 bicycle helmet use prevention of trauma deaths and injuries, 6-7 BIS (bispectral index) monitor, 91, 128 blast injuries mechanisms of, 10 Blaylock, Alfred, 44 bleeding prehospital care, 11–12 blood loss prevention spinal cord injury (SCI), 195-196 blood pressure (BP) non-invasive monitoring, 125 shock and, 45 blood products, 71 blood typing, 72-73 compatibility, 72-73 cross-matching, 72-73 cryoprecipitate, 71-72 fresh frozen plasma (FFP), 71 plasma products, 71 platelets, 72 red blood cells, 71 use of universal donor products, 72-73 whole blood, 72 blood typing, 72-73 blunt aortic injury diagnostic echocardiography, 144-145 brachial artery catheterization, 67 Brain, Archie, 38 breathing abdominal trauma, 235 ATLS protocol, 18 geriatric patients, 297 pediatric patients, 278-279 prehospital care, 11 primary survey, 21 Brown-Sequard syndrome, 192 burns Abbreviated Burn Severity Index (ABSI), 261-264 airway management, 270-271 anesthetic agents, 96 270 - 272anesthetic management, 269-273 chemical burns, 266

bicycle collisions

beta-blockers, 96

Index

319

classification of burn injuries, 261 electrical burns, 266 epidemiology, 261-262 estimation of extent and depth, 261 estimation of probability of survival, 261-264 initial management, 267-269 intrahospital transport considerations, 269 intraoperative fluid management, 272 key points, 273 monitoring burn patients, 269 - 270pathophysiology, 265 pharmacology for severe burns, 267 postoperative period, 272-273 regional anesthesia, 272 smoke inhalation, 266-267 specialist burn center referral, 261-265 tracheal extubation, 272 calcium chloride, 84 'cannot intubate, cannot ventilate' situation, 11 37-38 capnography, 125-126 cardiac arrest echocardiography monitoring of the precardiac arrest state, 151 cardiac injuries anesthetic agents and, 95-96 caused by chest trauma, 222-227 emergency department thoracotomy, 226-227 gunshot wounds, 226 myocardial contusion, 222 penetrating injuries, 222 tamponade, 225-227 cardiac output dynamic monitors of fluid bolus responsiveness, 132-134 echocardiography monitoring, 148-149 monitoring, 124-131 predicting 'recruitable' cardiac output, 124

cardiac tamponade, 49, 52, 225-227 diagnostic echocardiography, 142 - 144cardiogenic shock, 49 cardiopulmonary arrest guidelines for withholding or termination of resuscitation, 11-12 trauma patients found in, 11-12 cardiovascular management spinal cord injury (SCI), 193-194 catheters factors influencing flow, 56 cauda equina syndrome, 192 central cord syndrome, 192 central venous access, 57 59 - 60introducer catheters, 64-65 large-bore multi-lumen catheters, 65 large-bore single-lumen catheters, 64-65 sites of central line placement, 60-64 small-bore multi-lumen catheters, 64 types of central venous catheters, 64-65 central venous catheter monitoring trauma patients, 129 cerebral perfusion pressure (CPP) monitoring, 134 cervical spine precautions in general anesthesia, 90 cervical spine injury, 11 airway management, 34 use of manual in-line stabilization (MILS), 34 Cethrin, 198 chemical burns, 266 chest trauma aortic injury, 228-229 approach to management, 212 - 213blunt trauma, 212-214 cardiac injuries, 222-227 emergency department thoracotomy, 226-227 flail chest, 216-218 geriatric patients, 301 gunshot wounds, 212-213

hemothorax, 219-220 iatrogenic injuries, 213 indications for operative intervention, 212-213 key points, 229-230 mechanisms of injury, 212-214 mortality caused by, 212 one-lung ventilation, 220-224 pathophysiology, 213-214 penetrating injuries, 212-213 pneumothorax, 218-219 pulmonary contusion, 214-216 rib fractures, 216-218 stab wounds, 212-213 tracheobronchial injury, 218 vascular air embolism after, 220 child restraints injuries caused by, 8 prevention of trauma deaths, 7 children use of booster seats, 8 See also pediatric trauma circulation abdominal trauma, 235-236 ATLS protocol, 18 geriatric patients, 297-298 pediatric patients, 279-280 prehospital care, 11–12 primary survey, 21 citrate-phosphate-dextrose with adenine (CPDA-1), 71 coagulation effects of regional anesthesia, 103 coagulation assessment, 156-162 bedside point-of-care (POC) coagulation tests, 157-162 clinical presentation, 156 medical history, 156 rotational thromboelastometry (ROTEM), 157-162 Sonoclot, 157-162 standard lab coagulation tests, 157 thromboelastography (TEG), 157 - 162

320

Index

coagulation assessment (cont.) viscoelastic coagulation testing, 157-162 coagulation balance, 154 coagulation system activation by tissue factor (TF), 155 cell-based model of coagulation, 156 current concepts, 155-156 hemostatic mechanisms, 155 limitations of the cascade model, 155 role of thrombin, 155 coagulopathy acute traumatic coagulopathy (ATC), 70-71 early transfusion approaches in trauma, 73-75 goal-directed treatment, 74-75 key points, 80, 163 pathophysiology in trauma, 70 - 71pragmatic approach, 80 recombinant clotting factors, 78 trauma-induced coagulopathy, 70-71 traumatic brain injury (TBI) and, 182 treatment, 162 colloids, 52 Committee on Fluid Resuscitation for Combat Casualties, 51 compartment syndrome abdominal, 240-241 musculoskeletal trauma, 256-257 regional anesthesia and, 104 concentrated clotting factors, 78 - 79costs of trauma to society, 3-6 Cowley, R Adams, 45 craniotomy, 25 cricoid pressure, 90 concerns about the use of, 33 - 34cricothyroidotomy, 11, 40 crisis resource management (CRM) anesthesiologist familiarity with, 20 application to trauma team training, 27

crush injuries musculoskeletal trauma, 258-259 cryoprecipitate, 71-72 CT imaging during initial evaluation and management, 24 damage control concept in surgery, 25-27 damage control laparotomy abdominal trauma, 239-240 damage control hemostatic resuscitation, 54-55 damage control surgery general anesthesia, 97-99 definitive repair surgery abdominal trauma, 241 desflurane, 93-94 dexmedetomidine, 185 dextran, 52 difficult airway ASA algorithm modified for trauma patients, 35-36 supraglottic devices, 37 disability/exposure assessment ATLS protocol, 18-19 prehospital, 12 primary survey, 21 disseminated intravascular coagulation (DIC), 71, 182 pregnant trauma patient, 311 dorsalis pedis artery catheterization, 67 drinking age laws prevention of trauma deaths, 7 dual-lumen airways, 11 echocardiography in trauma air and fat emboli monitoring, 149 aortic dissection diagnosis, 144 - 145blunt aortic injury diagnosis, 144-145 cardiac output (CO) assessment, 148-149

assessment, 148–149 cardiac tamponade diagnosis, 142–144 clinical applications, 139–149 diagnostic applications, 139–146 FAST examination, 139–140 hemodynamic monitoring, 148–149

hemothorax diagnosis, 142 hypovolemia diagnosis, 145-146 indications in non-cardiac surgery patients, 138-140 intracardiac shunting, 149 intrapulmonary shunting, 149 key points, 151 monitoring applications, 146-149 myocardial contusion diagnosis, 145-146 myocardial ischemia monitoring, 148 performance of the perioperative echocardiography examination, 149-151 pericardial effusion diagnosis, 142-144 pneumothorax diagnosis, 140 - 141pre-cardiac arrest state, 151 pulmonary emboli monitoring, 149 range of applications, 138 regional wall abnormality (RWMA) monitoring, 148 shunting, 149 technological developments, 138 transesophageal echocardiography (TEE), 138 - 140transthoracic echocardiography (TTE), 138 - 140valvular lesion monitoring, 148 valvular regurgitation diagnosis, 145-146 ventricular function monitoring, 147 volume status monitoring, 146 - 147elbow block, 113 electrical burns, 266 electrocardiography (ECG), 126, 169 electromyography spinal cord injury (SCI), 197 emboli air embolism after chest trauma, 220

Index

echocardiography monitoring, 149 fat embolism syndrome, 256 emergency medical services (EMS) system prehospital care of trauma victims, 10-11 empiric transfusion ratios, 74 epidemiology of trauma, 1 - 6epidural block, 121-122 epinephrine, 84 epsilon-aminocaproic acid (EACA), 195 equipment for tracheal intubation, 31 trauma area setup, 17-18 esmolol, 185 etomidate, 32, 84, 88, 91, 94 burn patients, 270 physiologic effects, 92 exploratory laparoscopy abdominal trauma, 237 exploratory laparotomy abdominal trauma, 237-239 explosions mechanisms of injury, 10 exposure ATLS protocol, 18 primary survey, 22 extended FAST (eFAST) examination, 140 external jugular vein cannulation, 58 Factor III, 70, 156 Factor V, 155 Factor VII, 70, 78, 155 Factor VIII, 71 Factor IX, 156 Factor X, 70, 78 Factor Xa, 156 Factor XI, 155-156 Factor XIII, 71, 155-156 falls, 7-8 geriatric patients, 290 risk with regional anesthesia, 104-105 FAST examination, 139-140 during initial evaluation and management, 24 fat embolism syndrome, 256 femoral artery catheterization, 67 femoral nerve block, 116

femoral vein cannulation, 64 femur fracture management, 249-251 fentanyl, 84, 94, 185 fibrin, 77 fibrinogen, 71 fibrinogen concentrate, 78, 84 fibronectin, 71 flail chest, 216-218 flexible bronchoscopy, 40 frailty index, 296 fresh frozen plasma (FFP), 71 frontal air bags prevention of trauma deaths, 7 funding for trauma research, 6 gastrointestinal support postoperative care, 170-171 general anesthesia airway compromise, 90 airway management, 88-90 aspiration prophylaxis, 88 - 90burns and anesthetic agents, 96 cardiac injury and anesthetic agents, 95-96 cervical spine precautions, 90 consequences of hypothermia, 96-98 damage control surgery, 97-99 goals of general anesthesia for trauma, 87 head injury and, 94-95 hypovolemia and anesthetic agents, 91-94 key points, 100 maintaining normothermia, 96-98 monitoring, 84 musculoskeletal trauma, 247 operating room setup, 83-84 preoperative preparation, 82-86 selection of anesthetic and adjunct drugs, 86-87 geriatric trauma age-related changes in physiology, 290-295 aging population, 290 anesthesia for common injuries, 299-302

cardiovascular decline with age, 291-292 central nervous system changes with age, 293 chest trauma, 301 closed head trauma, 301-302 etiologies, 290 falls, 7-8, 290 hip fracture, 300-301 intraoperative management, 296-299 intraoperative monitoring, 298 key points, 303 medications taken by the patient, 295-296 orthopedic trauma, 299-301 outcomes, 302 pharmacological changes related to aging, 293-295 preexisting conditions, 295 preoperative evaluation, 295-296 renal function decline with age, 293 respiratory function decline with age, 292 rib fractures, 301 risk of postoperative delirium, 293 splenic trauma, 301 suicide with firearms, 290 temperature control, 299 traffic accidents, 290 traumatic brain injury (TBI), 301-302 Glasgow Coma Scale (GCS), 22 modified for children, 278 Glasgow Coma Scale (GCS) score, 12, 19, 29, 134, 168, 233 Glidescope, 36, 220 glucose control traumatic brain injury (TBI), 182 - 183golden hour concept, 18 shock, 45 great vessel injuries, 242 gunshot wounds abdominal trauma, 233 cardiac injuries, 226 chest, 212-213 death rates, 9 mechanisms of injury, 9 - 10

322

### Index

gunshot wounds (cont.) pregnant trauma patients, 304 suicide with firearms, 290 Hagen-Poiseuille equation, 56 head injury general anesthesia and, 94-95 hemodynamic instability regional anesthesia and, 102-103 hemodynamic monitoring echocardiography, 148-149 hemorrhage as a preventable cause of death, 69 key points, 80 mortality rates, 44-45 pragmatic approach, 80 resuscitative endovascular balloon occlusion of the aorta (REBOA), 243-244 hemorrhagic shock, 44-45, 47, 49, 69-70 early resuscitation, 53-54 later resuscitation, 54 mechanisms, 70 recognition of, 69 resuscitation, 52-55, 70 stages of, 70 treatment, 70 hemostasis abnormal, 154 key points, 163 mechanisms, 155 hemothorax, 219-220 diagnostic echocardiography, 142 hip fractures, 255, 300-301 hospital trauma area setup, 17-18 hydromorphone, 84 hydroxyethyl starch, 52 hyperfibrinolysis, 71, 77 hypofibrinogenemia, 72 hypothermia, 22 consequences in trauma patients, 96-98 maintaining normothermia, 96-98 temperature monitoring, 126-127 therapeutic, 183-184, 194-195 hypovolemia and anesthetic agents, 91-94 diagnostic echocardiography, 145 - 146

ICU delirium, 166 immobilization prehospital care, 12 induction agents, 32-33 infection postoperative, 171 risk with regional anesthesia, 104 informed consent, 102 infraclavicular block, 111-112 initial evaluation and management ABCDE protocol of ATLS, 18, 21–22 additional considerations, 24 - 25Advanced Trauma Life Support (ATLS), 18-22 angiography, 24-25 before the patient arrives, 17 - 18CT imaging, 24 damage control concept in surgery, 25-27 FAST examination, 24 first contact, 20 golden hour concept in ATLS, 18 imaging, 24-25 key points, 27 prehospital coordination with the receiving hospital, 17 primary survey, 20-22 prioritization of surgical management, 25-27 prioritization of trauma care, 18 - 25resuscitation phase, 22-23 role of the anesthesiologist, 16 - 20secondary survey, 23 teamwork of the trauma team, 27 trauma area setup, 17-18 injury three phases of, 1 internal jugular vein cannulation, 62-64 international normalized ratio (INR), 70 interscalene block, 109-110 intracardiac shunting echocardiography monitoring, 149 intracranial pressure (ICP) monitoring, 134-135

intraoperative awareness monitoring, 128 intraosseous (IO) access, 65 intraperitoneal injuries, 241 intrapulmonary shunting echocardiography monitoring, 149 introducer catheters, 64-65 isoflurane, 93-94 ketamine, 32, 84, 88, 91, 94-95 burn patients, 270 physiologic effects, 92 King LTS-D airway, 39 Kleihauer–Betke test, 312 labetalol, 185 lactate monitoring, 135 lap belts injuries caused by, 8 laparotomy, 25 laryngeal mask airway (LMA), 11, 30, 38-39 laryngeal tube airway (LTA), 30, 39 lateral femoral cutaneous nerve block, 117 Le Fort classification of midface fractures, 205 lethal triad (hypothermia, coagulopathy, and acidosis), 46, 70 lidocaine, 84, 119 local anesthetics selection for regional anesthesia, 119 toxicity, 120 lungs one-lung ventilation, 220-224 manual in-line stabilization (MILS) use in suspected cervical injury, 34 Marshall classification of traumatic brain injury (TBI), 174 mass casualty events, 6 massive transfusion activation of massive transfusion protocols, 76 complications, 77 definition, 75 role of massive transfusion protocols, 75-76

Index

323

scope of massive transfusion protocols, 76 maxillofacial trauma, 204 airway considerations for surgery, 209-210 anesthetic considerations for reconstructive surgery, 210 bleeding from injuries, 209 classification, 204-205 elective maxillofacial repair, 209-210 emergency airway management, 206-207 key points, 210-211 Le Fort classification of midface fractures, 205 mechanisms, 204 preoperative assessment, 205-207 preoxygenation, 208 rapid sequence induction (RSI) and tracheal intubation, 208 surgery considerations, 209 surgical airway, 208 McConnell's sign, 149 mechanisms of injury blasts, 10 explosions, 10 factors influencing severity of injury, 7 falls, 7-8 penetrating trauma, 9-10 transportation-related injuries, 8-9 mental status prehospital assessment, 11 - 12mepivacaine, 119 methylprednisolone, 198 midazolam, 84, 91, 94, 185 physiologic effects, 92 milrinone, 95 monitoring trauma patients abdominal trauma, 236 arterial catheter, 128-129 ASA Standards for basic monitoring, 124 awareness monitoring, 128 base deficit, 135-136 BIS (bispectral index) monitor, 128 burn patients, 269-270 capnography, 125-126 cardiac output, 130-131 central venous catheter, 129

cerebral perfusion pressure (CPP), 134 challenges for anesthesiologists, 124 dynamic monitors of fluid bolus responsiveness, 132-134 electrocardiography (ECG), 126 general anesthesia, 84 Glasgow Coma Scale (GCS) score, 134 intracranial pressure (ICP), 134-135 intraoperative awareness monitoring, 128 invasive monitors, 128-135 key points, 137 lactate, 135 neuromuscular monitoring, 128 non-invasive blood pressure monitoring, 125 non-invasive monitors, 125-128 predicting 'recruitable' cardiac output, 124 pulmonary artery catheter, 129-130 pulse oximetry, 125 selection of monitors to use, 124 serum markers of tissue perfusion and oxygen debt, 135-136 temperature monitoring, 126 - 127tissue acidosis, 135–136 tissue perfusion, 135 transesophageal echocardiography (TEE), 127 - 128trend monitoring, 124 venous saturation, 131-132 when monitors fail or are not effective, 136-137 morphine, 84 motor evoked potentials (MEP), 197 motor vehicle collisions bicycle collisions, 9 death and injury rates, 8 mechanisms of injury, 8-9 motorcycle collisions, 9 pedestrian collisions, 9

motor vehicle trauma preventive measures, 1 motorcycle collisions mechanisms of injury, 9 motorcycle helmet use prevention of trauma deaths and injuries, 6-7 musculoskeletal trauma choice of anesthetic technique, 247 compartment syndrome, 256 - 257crush injuries, 258-259 fat embolism syndrome, 256 femur fracture management, 251-252 general anesthesia, 247 hip fractures, 255 indications for urgent surgical treatment, 246 key points, 259-260 life-threatening injuries, 246 limb-threatening injuries, 246 multiple fractures, 255 open fractures, 255 pain management, 250 pelvic fracture management, 253-254 preoperative preparations, 247-249 regional anesthesia, 247, 250-251 traumatic amputation, 256 vascular injury, 256 myocardial contusion, 222 diagnostic echocardiography, 145-146 myocardial ischemia echocardiography monitoring, 148 myoclonus, 94 National Highway Traffic Safety Administration (NHTSA), 7 neuraxial blocks, 120-122 neurogenic shock, 45 neurologic disability ATLS protocol, 18-19 primary survey, 21 neurologic function pediatric patients, 280-282 neuromuscular blocking drugs (NMBDs), 11, 32-33, 84 abdominal trauma, 236 burn patients, 271

## CAMBRIDGE

Cambridge University Press & Assessment 978-1-316-63671-8 — Essentials of Trauma Anesthesia Albert J. Varon, Charles E. Smith Index More Information

#### 324 Index

neuromuscular blocking drugs (NMBDs) (cont.) ocular trauma, 203 pediatric patients, 285-286 neuromuscular monitoring, 128 nicardipine, 185 nitroglycerin, 96 nitrous oxide, 93-94 non-fatal trauma injuries falls, 7-8 rates of, 3 norepinephrine, 84 nutritional support postoperative care, 170-171 obstructive shock, 49 occult hypoperfusion syndrome, 45-46 ocular trauma, 200 anesthetic management, 202-204 aspiration prophylaxis, 202 emergence from anesthesia, 203-204 induction agents, 203 intraocular pressure (IOP), 201key points, 210-211 maintenance of anesthesia, 203 mechanisms, 200 neuromuscular blocking drugs (NMBDs), 203 oculocardiac reflex, 201-202 preoperative evaluation, 200 - 201rapid sequence induction (RSI) and tracheal intubation, 202 oculocardiac reflex, 201-202 open fractures, 255 open pneumothorax, 11 opioids, 94, 96 burn patients, 271 pain management, 84 musculoskeletal trauma, 250 postoperative, 165-167 paravertebral block, 122

patient medications potential complications related to, 176–177 pedestrians mechanisms of injury in motor vehicle collisions, 9 pediatric trauma airway and breathing, 278-279 anesthetic management, 283 - 287circulation, 279-280 equipment sizes for children, 275-277 induction agents, 285 induction technique, 285 initial assessment and resuscitation, 275-283 injury patterns in children, 275 intraoperative fluid management, 287-288 key points, 288 length-based weight conversion devices, 275 - 277maintenance of anesthesia, 286-287 medication dosage for children, 275-277 mortality among children, 275 neurologic function, 280-282 neuromuscular blocking drugs (NMBDs), 285-286 postoperative care, 288 preoperative evaluation and preparation, 283-285 primary survey, 275-282 role of the anesthesiologist, 275 secondary survey, 282-283 spinal cord injury without radiographic abnormality (SCIWORA), 282 types of trauma in children, 275 pelvic fracture management, 253-254 penetrating neck injury airway management, 41 penetrating trauma mechanisms of injury, 9-10 pericardial effusion diagnostic echocardiography, 142-144 peripheral venous access, 57-59 phenylephrine, 84 plasma products, 71 plasmin, 77 platelet dysfunction, 71 platelets (blood products), 72

pneumothorax, 218-219 diagnostic echocardiography, 140 - 141prehospital treatment, 11 posterior cord syndrome, 192 postoperative care, 164 acid/base considerations, 169 - 170cardiovascular concerns, 168-169 gastrointestinal support, 170 - 171ICU delirium, 166 infection and sepsis, 171 key points, 171-172 neurologic considerations, 168 nutritional support, 170-171 pain management, 165-167 patient disposition, 164-165 renal considerations, 169-170 secondary resuscitation, 165 sedation, 165-167 transport from the operating room, 164-165 ventilator management, 167 - 168postoperative delirium risk in geriatric patients, 293 pregnant trauma patient Advanced Cardiac Life Support (ACLS), 312 airway management, 309-310 anesthesia technique, 306 anesthetic management, 306-313 antifibrinolytics, 311 blunt trauma, 304 burns, 304-305 causes of trauma, 304 complications, 305-306 disseminated intravascular coagulation (DIC), 311 emergency cesarean delivery during cardiac arrest, 312 fetal monitoring, 313 fetal outcome, 305-306 imaging, 306 incidence of trauma in pregnancy, 304 induction of anesthesia, 309 key points, 316 laboratory evaluation, 311-312

Index

325

mechanisms of injury, 304-305 mortality related to trauma, 304 multidisciplinary management, 313-315 normal physiologic changes in pregnancy, 306-308 obstetric issues, 305-306 patient history, 306 penetrating trauma, 304 physical examination, 306 placental abruption, 305 positioning prior to induction, 308 premedication, 306-308 primary survey, 306-308 resuscitation, 309-311 Rh sensitization, 312 transfusion, 309-311 uterine rupture, 305 prehospital care airway management, 11 assessment of mental status, 11 - 12bleeding, 11-12 breathing, 11 circulation, 11-12 coordination with the receiving hospital, 17 disability/exposure assessment, 12 guidelines for withholding or termination of resuscitation, 11-12 key points, 14-15 patient immobilization, 12 pneumothorax, 11 role of the emergency medical services (EMS) system, 10-11 trauma patients found in cardiopulmonary arrest, 11 - 12triage, 12-14 ventilation management, 11 prevention of trauma deaths challenges of, 6-7 primary survey of the trauma patient, 20-22 procoagulation adjuncts, 77-78 propofol, 32, 84, 88, 91, 94-95, 185 burn patients, 270 physiologic effects, 92

protein C, 71, 155 prothrombin complex concentrates, 78-79, 84 prothrombin time (PT), 70 prothrombinase, 70 public health initiatives prevention of trauma deaths and injuries, 6-7 pulmonary artery catheter monitoring trauma patients, 129-130 pulmonary contusion, 214-216 pulmonary emboli echocardiography monitoring, 149 pulse oximetry, 125 pupillary response, 12 radial artery catheterization, 66-67 rapid sequence induction (RSI), 32-33 failed intubation, 40 oxygenation and cricoid pressure, 33-34 Rapid Ultrasound for Shock and Hypotension (RUSH), 140 recombinant clotting factors, 78 recombinant Factor VIIa, 78, 84, 195 red blood cells antigens, 72-73 red blood cells (blood products), 71 re-exploration surgery abdominal trauma, 240-241 regional anesthesia adductor canal block. 116-117 ASRA guidelines, 103 axillary block, 112-113 burns, 272 catheter placement, 119 coagulation status and, 103 compartment syndrome, 104 concerns in trauma patients, 102-105 elbow block, 113 epidural block, 121-122 fall risk, 104-105 femoral nerve block, 116 general considerations, 102 hemodynamic instability, 102 - 103

infraclavicular block, 111-112 interscalene block, 109-110 key points, 122-123 lateral femoral cutaneous nerve block, 117 local anesthetic selection, 119 local anesthetic toxicity, 120 lower extremity blocks, 114-119 musculoskeletal trauma, 247, 250-251 neuraxial blocks, 120-122 obtaining informed consent, 102 paravertebral block, 122 risk of infection, 104 sciatic nerve blocks, 117-119 subarachnoid block, 120-121 supraclavicular block, 110 - 111traumatic nerve injury, 103 upper extremity blocks, 107 - 113use of ultrasound, 105-107 regional wall abnormalities (RWMAs) echocardiography monitoring, 148 remifentanil, 185 renal function acute kidney injury (AKI), 169-170 postoperative care, 169-170 research funding for trauma research, 6 resuscitation antifibrinolytic agents, 77-78 assessing the need for fluid resuscitation, 49-50 damage control hemostatic resuscitation, 54-55 dynamic monitors of fluid bolus responsiveness, 132-134 from hemorrhage, 52-55 goals of, 52 guidelines for prehospital withholding or termination, 11-12 hemorrhagic shock, 70 in initial evaluation and management, 22 key points, 55 predicting 'recruitable' cardiac output, 124

326

#### Index

resuscitation (cont.) risks associated with, 47 secondary resuscitation, 165 small volume resuscitation, 51 - 52timing and rate of fluid administration, 50-51 use of colloids, 52 resuscitative endovascular balloon occlusion of the aorta (REBOA), 243-244 retroperitoneal injuries, 242 Rh sensitization pregnant trauma patient, 312 rib fractures, 216-218 geriatric patients, 301 riluzole, 198 rocuronium, 32-33, 84, 89 ropivacaine, 119 rotational thromboelastometry (ROTEM), 54, 74-75, 157-162 Ryder Trauma Center emergency airway algorithm, 29 - 31massive transfusion protocol, 75 - 76tracheal intubation methods, 41 scene safety, 11 sciatic nerve blocks, 117 gluteal approach, 118 lateral popliteal approach, 118-119 scopolamine, 84, 91 seat belts injuries caused by, 8 prevention of trauma deaths and injuries, 6-7 secondary survey, 23 ATLS protocol, 18 sedation postoperative, 165-167 Seldinger method of placing catheters, 59 sepsis postoperative, 171 septic shock, 47 serum markers of tissue perfusion and oxygen debt, 135-136 sevoflurane, 93-94 shock assessing the need for fluid resuscitation, 49-50

blood pressure (BP) and, 45 cardiogenic shock, 49 compensated shock, 45 current understanding of, 44 - 45damage control hemostatic resuscitation, 54-55 diagnosis, 47-49 estimating blood loss, 49-50 estimating the likelihood of ongoing hemorrhage, 49 - 50etiologies, 44-45 fluid therapy, 49-52 goals of resuscitation, 52 golden hour concept, 45 hemorrhagic shock, 44-45, 47, 49, 69-70 hemorrhagic shock resuscitation, 52-55 history of thought on, 44 irreversible shock, 46 key points, 55 neurogenic shock, 45 obstructive shock, 49 occult hypoperfusion syndrome, 45-46 organ failure, 46 oxygen debt, 44 recognition of, 47-49 risks associated with resuscitation, 47 septic shock, 47 small volume resuscitation, 51 - 52spinal shock, 47 use of colloids, 52 shoulder restraints injuries caused by, 8 shunting echocardiography monitoring, 149 smoke inhalation, 266-267 sodium bicarbonate, 84 somatosensory evoked potentials (SSEP), 196 Sonoclot, 157-162 spinal cord injury (SCI) airway management, 189-193 blood loss prevention, 195-196 cardiovascular management, 193 - 194classification, 189-192 clinical presentation, 189-192

early surgical decompression, 194 electromyography, 197 electrophysiological monitoring, 196-198 epidemiology, 187-188 etiologies, 188 initial assessment, 189 key points, 198-199 motor evoked potentials (MEP), 197 non-traumatic etiologies, 188 pathophysiology, 188 pharmacological therapeutic agents, 198 somatosensory evoked potentials (SSEP), 196 spinal cord syndromes, 189-192 Stagnara wake-up test, 196 therapeutic hypothermia, 194-195 traumatic etiologies, 188 spinal cord injury without radiographic abnormality (SCIWORA), 282 spinal shock, 47, 192 splenic trauma, 301 stab wounds abdominal trauma, 232 chest, 212-213 death and non-fatal injury rates, 10 mechanisms of injury, 10 pregnant trauma patients, 304 Stagnara wake-up test, 196 START triage system, 14 subarachnoid block, 120-121 subclavian vein cannulation, 60-62 succinylcholine, 32-33, 84, 88-89, 91, 94, 96, 203, 271 sugammadex, 84, 90, 203 supraclavicular block, 110–111 supraglottic devices, 37-39 surgical airway, 40 maxillofacial trauma, 208 surgical management damage control concept, 25 - 27emergent cases, 25 non-urgent cases, 25 prioritization in the trauma patient, 25-27 urgent cases, 25

Index

327

systemic air embolism after chest trauma, 220 systemic inflammatory response, 46 Task Force for Advanced Bleeding Care in Trauma, 162 teamwork of the trauma team, 27 temperature monitoring, 126 - 127tension pneumothorax, 11, 49, 52 terrorism mass casualty events, 6 thiopental, 32, 84, 88, 91, 94-95 physiologic effects, 92 thoracic endovascular aortic repair (TEVAR), 228-229 thoracotomy, 25, 226-227 thrombin, 70 role in the coagulation system, 155 thromboelastography (TEG), 54, 74-75, 157-162 thrombomodulin, 71 tissue acidosis serum markers for, 135-136 tissue factor (TF), 155 tissue perfusion monitoring, 135 tracheal intubation Aintree intubation catheter, 38-39 awake intubation, 34 basic equipment, 31 failed intubation, 40 indications for, 30-31 penetrating neck injury, 41 tracheal tube introducers, 37 videolaryngoscopy, 35-37 tracheal tube introducers, 37 Aintree intubation catheter, 37 tracheobronchial injury, 218 tracheostomy, 40 tranexamic acid, 77-78, 84, 195 transesophageal echocardiography (TEE), 127-128, 138-140 See also echocardiography in trauma. transfusion early approaches in trauma, 73-75

goal-directed treatment, 74-75 key points, 80 massive transfusion, 75-77 pragmatic approach, 80 risks associated with, 73-74 transient paralysis, 192 transportation-related injuries, 8-9 transthoracic echocardiography (TTE), 138-140 See also echocardiography in trauma. trauma as a leading cause of death, 1 - 4costs to society, 3-6 definition, 1 epidemiology, 1-6 known risk factors, 1 mass casualty events, 6 modification of risk factors, 1 mortality rates, 82 non-fatal injury rates, 3 prevention of trauma deaths, 1 three phases of injury, 1 viewed as a disease, 1 trauma area setup, 17-18 trauma care prioritization in initial management, 18-25 role of the anesthesiologist, 16 - 17trauma-induced coagulopathy, 70 - 71trauma team teamwork, 27 training in crisis resource management (CRM), 27 traumatic brain injury (TBI) anemia, 181-182 anesthetic technique, 179-180 coagulopathy, 182 decompressive craniectomy, 184 definition, 173 emergence from anesthesia, 184-185 epidemiology, 173 fluid management, 181 geriatric patients, 301-302 glucose control, 182-183 hemodynamic management, 181

empiric transfusion ratios, 74

intracranial pressure (ICP) monitoring, 134-135 intraoperative management, 177-179 intraoperative monitoring, 180-181 key points, 185 laboratory investigations, 177 Marshall classification, 174 non-neurological surgery in TBI patients, 184 oxygenation, 179 pathophysiology, 173-174 postoperative airway protection, 167 pre-existing conditions, 176-177 preoperative considerations, 175-177 primary injury, 173-174 role of the anesthesiologist, 173 secondary injury, 174 therapeutic hypothermia, 183 - 184ventilation, 179 traumatic nerve injury regional anesthesia and, 103 triage prehospital care, 12-14 START triage system, 14 ultrasound needle approaches to nerves, 106-107 physics of, 105-106 use in regional anesthesia, 105 - 107See also echocardiography in trauma. Ultrasound guided procedures subclavian vein cannulation, 61 vascular cannulation, 56 - 58universal donor blood products, 72-73 valvular lesions echocardiography monitoring, 148 valvular regurgitation diagnostic echocardiography, 145 - 146vascular air embolism after chest trauma, 220

328 Index

vascular cannulation arterial access, 66-67 factors influencing flow, 56 importance of vascular access in trauma management, 56 intraosseous (IO) access, 65 key points, 68 ultrasound-guided procedures, 56-58 venous access, 57-65 vascular injury musculoskeletal trauma, 256 vasopressin, 84 vecuronium, 84 venous access, 57-65 antecubital vein, 58

central access, 57, 59-60 external jugular vein, 58 femoral vein, 64 internal jugular vein, 62-64 peripheral access, 57-59 Seldinger method, 59 sites for central line placement, 60-64 sites for peripheral catheter placement, 58-59 subclavian vein, 60-62 types of central venous catheters, 64-65 venous saturation monitoring, 129-132 ventilation management prehospital care, 11 ventilator management postoperative, 167-168

ventricular function echocardiography monitoring, 147 vicious cycle (hypothermia, coagulopathy, and acidosis), 70 videolaryngoscopy, 35-37 Vietnam War, 49 viscoelastic coagulation testing, 157-162 viscoelastic hemostatic assays (VHAs), 74–75 volume status echocardiography monitoring, 146-147 von Willebrand factor (VWF), 71 VX-210, 198

wake-up test, 196