

## Index

- Aharonov–Bohm effect, 111, 121  
 with conduction electrons, 111–14  
 anomalous dispersion  
 in electron wavefunctions, 90–4, 97  
 in optics, 87–90  
 anti-localisation, 128
- Boltzmann equation, 26, 33  
 Boltzmann theory of electrical conduction,  
 25–32  
 Born approximation, 36, 39, 51
- Ca–Al metallic glasses  
 band structure, 212–13  
 characteristic parameters, 218–19  
 conductivity of, 211–23  
 dephasing times, 222  
 electronic heat capacity and Hall  
 coefficient, 213  
 magnetic impurities in, 222  
 magnetoresistance, 213–23  
 resistivity, temperature dependence,  
 222–3  
 effect of spin–orbit scattering on, 220–2  
 value of  $F$ , 218–19, 221  
 closed paths, probability of, 115–16, 157–9  
 coherence of electron wavefunction, 111–12,  
 115–20  
 collective electron modes, 70  
 comparison of experiment and theory,  
 200–24  
 conductivity, electrical (*see also* resistivity),  
 calculation by Boltzmann theory, 25–32  
 of Ca–Al metallic glasses, 211–23  
 of Cu–Ti metallic glasses, 201–5  
 and diffusion, 32–3  
 effect of Coulomb interaction on, 167–75  
 Cooper channel, 174  
 particle–hole channel, 167–70  
 Einstein relation, 32  
 and weak localisation, 105–39  
 temperature dependence due to weak  
 localisation, 117–18  
 conduction electrons, 21–3  
 Cooper channel interactions, 162–4, 174  
 and magnetoresistance, 181–2  
 Coulomb interaction between conduction  
 electrons, 143–62  
 Coulomb anomaly (*see also* enhanced  
 interaction effect), 140–75  
 Coupling constant in exchange interaction,  
 154–6  
 for Hartree terms, 159–60, 230–1  
 in Cooper channel, 162–4, 174,  
 229–31  
 Cu–Ti metallic glasses, conductivity of,  
 201–5
- Debye–Waller factor, 58–60  
 density of states (for free electrons),  
 24–5  
 changes due to enhanced electron  
 interaction, 141 *et seq.*  
 changes with temperature, 165  
 dephasing processes, 117–18, 120–3, 207,  
 211  
 characteristic time in Ca–Al glasses, 222  
 diffusion, effect on electron wavefunctions,  
 149–55  
 importance of, in enhanced interaction  
 effect, 161–2  
 diffusion coefficient of electrons, 32–3, 115  
*et seq.*, 200–1  
 values in Ca–Al glasses, 216, 218–20  
 values in Cu–Ti glasses, 205  
 values in a range of metallic glasses, 207  
 diffusive screening, 154–5, 171
- Einstein relation, 32, 165, 200

- electrical conductivity (*see* conductivity, electrical)
- electron, independent, picture, 63–4
- electron transport in metals, conventional theory, 20–33
- electron–electron interaction, collective electron modes, 70  
 in crystalline metals, 64, 231–2  
 in disordered metals, 183, 231–2  
 mediated by phonons, 67–9  
 via screened Coulomb potential, 38
- electrons, conduction, 21–3
- electrons, collective modes, 70
- enhanced interaction effect, 140–75  
 and conductivity, 167–75  
 and density of states, 140–66  
 and Hall effect, 184–7  
 and magnetoresistance, 176–82  
 effect of spin–orbit scattering, 171–4
- exchange terms in particle–hole channel, 147–53  
 Hartree terms in particle–hole channel, 156–9
- exchange contribution to interaction energy, 145–56
- $F$  coupling constant, 159–60, 228–30
- Fermi electrons, 24  
 energy, 22–4  
 surface, 23, 48, 80, 201
- Fourier transforms, 37–8, 227–8
- glass, insulating, 6  
 nature of, 6  
 properties in general, 7  
 transition temperature, 9
- glasses, metallic (*see* metallic glasses)
- Hall coefficient, Hall effect, 81–99, 184–7  
 hybridisation theory of, 87–94, 97–9, 234  
 influence of enhanced interaction effect, 184–7  
 in metallic glasses, 81–2, 86, 97–9  
 in terms of Fermi surface, 85  
 simple theory, 81–3  
 with two types of carrier, 84–5
- Hartree terms in interaction energy, 145–6  
 contribution to density of states, 157–9
- hybridisation, of electron wavefunctions in crystalline metals, 73–5  
 of s- and d-character in transition metal glasses, 87–94  
 theory of Hall coefficient, 87–94, 97–9, 234
- independent electron picture, 63–4
- inelastic scattering, 116–18, 120, 124, 201, 207, 222
- interaction effect (*see also* enhanced interaction effect), 140–87
- interaction mechanism in Hartree terms of particle–hole channel, 156–7, 168–9
- interactions and quasi-particles, 63–70
- Kohler’s rule in magnetoresistance, 102–3  
 $k$ -space, 23  
 $k$ -states in disordered metals, 44, 48–9
- Lindhard screening, 43–4, 53
- liquid metals, Ziman theory, 48–54
- localisation, weak (*see* weak localisation)
- magnetoresistance, 100–4  
 due to weak localisation, 121–5  
 at high fields, 122–3  
 at low fields, 123–5  
 due to enhanced interaction effect, 176–84  
 at high fields, 176–7  
 at low fields, 178  
 in the Cooper channigel, 181–2  
 spin–orbit effects, 180–1  
 temperature dependence, 178–80
- Kohler’s rule, 102–3
- of metallic glasses, 204–23
- simple theory, 100–4
- two-band model, 100–2
- Maki–Thompson correction, 183, 209–12, 229–31
- melt spinning, 10–11
- metallic glasses, atomic deposition techniques, 11  
 Hall coefficient, 81–2, 86, 97–9, 184–7  
 how produced, 9–11  
 magnetoresistance, 204–23  
 range of stability, in composition, 11–12  
 in temperature, 9  
 resistivity (*see also* conductivity), 105–9, 201–4  
 structure of, 13–19  
 thermopower, 194–9  
 what are they? 6
- metals, liquid, Ziman theory of, 48–54
- Mooij correlation, 107–9, 118
- Mott model of transition metal, 74–6
- normal modes in glasses, 55
- particle–hole pairs, 66–7
- Peltier heat, 188–93
- phonon drag, 58, 194

phonons in disordered systems, 55–62  
 phonon ineffectiveness, 60–1  
 pseudopotential, 44–7  
   applied to liquid metals, 49, 51–2  
 Principle of Least Action, 109–10

quantum interference, 109–11, 121  
   and Aharonov–Bohm effect, 111, 121  
   with conduction electrons, 111–4  
   and Principle of Least Action, 109–10

quasi-particles, 64–5

radial distribution function, 17–18

resistance, resistivity, electrical (*see also* conductivity),  
 Mooij correlation, 107–9, 118  
 magnetic field dependence,  
   due to enhanced electron interaction, 176–83  
   due to weak localisation, 121–5  
 temperature dependence due to enhanced electron interaction, 167–75  
   with spin–orbit scattering, 171–4  
 temperature dependence due to weak localisation, 117–18  
   with spin–orbit scattering, 127–31  
 sources of, 20–1

s- and d-electrons, 71–6, 78, 86–7, 90–4, 97–9, 103, 124, 201, 226–7

scaling theory and weak localisation, 136–9

scattering, 34–47  
   angle, influence on resistivity, 39–40, 51  
   of electrons by phonons, 56  
   at low temperatures, 57–8  
   at high temperatures, 61–2  
   inelastic, 116–18, 120, 124, 201, 207, 222  
   influence of Pauli principle, 40–1  
   by phonons in disorder systems, 140

screening by electrons in metals, 41–4  
   influence of diffusion, 154–5, 171  
   Thomas–Fermi, 41–3, 45, 144, 201

self-energy of electrons, 65, 142–3

short-range order, 15, 18–19

singularity in density of states, 153, 161–2

skew-scattering of electrons, 94–6

spectral function of electrons, 49

spin–orbit interaction, 94–6

spin–orbit scattering,  
   in the enhanced interaction effect, 171–4  
   in weak localisation, 126–35

spin–spin or spin–flip scattering, 135–6, 222

structure factor, 49–53

superconducting fluctuations, 163 174, 183

superconductivity mechanism via phonons, 67–9

temperature coefficient of resistance,  
   negative, 106–9

thermopower, 188–99  
   absolute, 189–90  
   calculation of, in metals, 190–4  
   electron phonon enhancement, 195–9  
   general and definitions, 188–94  
   in metallic glasses, 194–9  
   Mott formula, 193

Thomas–Fermi screening, 41–3, 45, 144, 201

transition metals and alloys, 71–80

crystalline, 71–4  
   Fermi surface in glasses, 79–80  
   glasses, 77–80  
   Mott model, 74–6

virtual processes, 68

weak localisation, 105–39  
   and diffusion of electrons, 115–17  
   and conductivity, some formulae for temperature and magnetic field dependence, 225–6  
   and electron spin, 125–36  
   and quantum interference, 115  
   and scaling theory, 136–9  
   and spin–orbit scattering, 126–35  
   and Zeeman splitting, 182–3  
   with s- and d-bands, 226–7

Zeeman splitting in magnetoresistance,  
   due to enhanced electron interaction, 177–8  
   due to weak localisation, 182–3

Ziman model of liquid metals, 48–54  
   applied to metallic glasses, 105