

Index

- absolute distance between points, 5–6, 34
- acridine orange, 190–96
- age distribution theory, 130–32, 150–51
- amplifiers, 94–95
 - linear, 94
 - log, 94–95
- analogue-to-digital conversion, 95–97, 114
- anthracyclines, 231–36
- association, χ^2 , 49–53
- autofluorescence, 99
- average, representation of, 101–3
 - see also* central tendency
- Bessel's correction, 35, 49, 55, 80
- binding-site modulation, 85–86
- binomial distribution, 16–18, 28–30
- biological sources of variation, 99–100
 - autofluorescence, 99
 - inherent variation, 100
 - light scatter, 90°, 99
 - non-specific binding, 100
- bleaching, 82–83, 89
- bromodeoxyuridine, 173–87
 - continuous labelling, 185–87
 - pulse labelling, 176–85
- calcium, 228–31
- cell cycle, 126–27
- cell-cycle kinetics, 145–96
 - acridine orange, 190–96
 - bromodeoxyuridine, 173–87
 - FPI analysis, 171–72
 - human tumour kinetics, 187–90
 - mitotic selection, 146–49
 - modelling population kinetics, 149–55
 - multiparameter optimisation, 155–71
 - stathmokinetic analysis, 145–46
 - synchrony progression, 146–55
- central tendency, 4–5, 101–3
 - mean, 4, 6, 102–3
 - median, 5–6, 102–3
 - mode, 5–6, 102–3
- change of variable, 79
- chi-squared (χ^2), 49–53, 56–57, 76, 113, 122
- chloroethylnitrosoureas, 238–40
- chromosome sorting, 28–30
- coefficient of variation (CV), 96–97, 102, 105–11, 152–53, 155, 163–71
- coefficients of polynomial, 77–78
- coincidence correction, 91–94
- compensation, 97–99
- constant CV analysis, 105–14
 - histogram deconvolution, 112–14
 - skewed-normal distribution, 105–12
- constant-variance analysis, 114–25
 - labelled-fraction mean calculation, 116–20
 - ratio analysis of means, 114–16
 - statistical verification, 120–22
- correlated bivariate density function, 242
- correlation coefficient, 63–64, 66, 70
 - significance of, 64
- correlation ratio, 65–66, 268
- cumulative frequency, 20–23, 34, 104–5, 136
 - tables, 21, 250–51
- cytoplasmic enzymes, 205–12, 217–19
 - esterases, 206–8, 217–19
 - glutathione S-transferase(s), 208–12
- data transforms, 68–74, 217
 - inverse exponential, 68
 - inverted exponential, 68–71, 217
 - logistic equation, 73–74
 - multitarget cell-survival curve, 71–73
- D_{crit} , 45, 262–64
- degrees of freedom, 35, 38
- de Moivre, 14
- direct linear plot, 204
- dispersion, 7–9
 - mean deviation, 8
 - standard deviation, 8
 - variance, 8

286 *Index*

- distributions, 12–30, 130–32, 150–51
 age distribution, 130–32, 150–51
 binomial, 16–18, 28–30
 Gaussian, 14–16, 20
 mathematical transforms, 12
 Poisson, 18–19, 26–28
- DNA degradation, 86
- DNA histogram analysis, 126–44, 150–51
 age distribution theory, 130–32, 150–51
 cell cycle, 126–27
 multiple Gaussian, 133–34
 polynomial, 135
 rectilinear integration, 132–33
 single Gaussian, 135–43
 TCW analysis, 143–44
- double reciprocal plot, 202
- drug resistance, 208, 233–38
- dynamic cellular events, 197–240
 anthracyclines, 231–36
 calcium, 228–31
 chloroethylnitrosoureas, 238–40
 enzyme kinetics, 198–204
 flow cytoenzymology, 204–28
 incorporation of time, 197–98
 membrane transport, 231–40
 methotrexate, 236–38
- Eadie–Hofstee transform, 202–3
- Eisenthal–Cornish–Bowden transform, 204
- electronic sources of variation, 90–99
 analogue-to-digital conversion, 95–97
 coincidence correction, 91–94
 compensation, 97–99
 linear amplifiers, 94
 log amplifiers, 94–95
 triggering and thresholds, 90–91
- enzyme kinetics, 198–204
- enzyme progress curves, 202, 207, 211, 213
- error function (erf), 20, 138, 154, 249
- esterases, 206–8, 217–19
- ethidium bromide, 173–74
- Euclidean hyperspace, 7, 9, 161–71
- F*-ratio, 37–38, 65, 72
 tables, 39, 252–55
- filter combinations, 90
- Fisher, R. A., 34
- fit criteria, 53–57
 Student's *t*, 55–56
 $\Sigma \chi^2$, 56–57
- flow cytoenzymology, 204–28
 cytoplasmic enzymes, 205–12
 inhibition kinetics, 217–19
 membrane enzymes, 212–17
 short time-scale kinetics, 220–28
- fluorescein diacetate, 206–8
- fluorescein phosphate, 3 O-methyl, 212, 216–17
- fluorescence excitation, as source of variation, 86–89
 bleaching, 89
 hydrodynamic instability, 88
 illumination constancy, 87–88
 light-source stability, 86–87
 fluorescence of filters, 89
 fluorescent methotrexate, 236–38
 FPI analysis, 171–72
- Gaussian distribution, 14–16, 20–23, 106, 116, 133–35, 150, 153
 cumulative frequency, 20–23
 population sampling, 23–24
 standard error of the mean, 24–25, 33
 standard error of the standard deviation, 25
 skew and kurtosis, 26
- glutathione S-transferase(s), 208–12
- Gosset, W. S., 34
- gp¹⁷⁰, 233
- growth fraction, 126
- histogram deconvolution, 112–14
- Hoechst dyes (33342, 33258), 173–75, 185–87
- human tumour kinetics, 187–90
- hydrodynamic instability, 88
- illumination constancy, 87–88
- immunofluorescence data, 32, 101–25
 constant CV analysis, 105–14
 constant-variance analysis, 114–25
 conventional analysis, 103
 errors with conventional methods, 122–24
 Kolmogorov–Smirnov assessment, 103–5
 representation of the average, 101–3
- inhibition kinetics, 217–19
- inverse exponential, 68
 inverted exponential, 68–71, 217
- K*_m (Michaelis constant), 200–4, 213–16
- Kolmogorov–Smirnov (K–S), 44–46, 103–5, 121, 262–64
 tables, 263–64
- kurtosis, 26
- Law of Leibniz, 272–73
- Laplace, P. S., 14
- light-collection efficiency, 89
- light scatter, 90°, 99, 191–93
- light-source stability, 86–87
- linear amplifiers, 94
- linear regression, 58–67, 266–68
 correlation coefficient, 63–64, 66
 minimising squared deviations, 61–63
 testing linearity, 65–67, 266–68

- x* on *y*, 64–65
- y* on *x*, 61–63
- Lineweaver–Burke plot, 202
- log amplifiers, 94–95
- log transform, 13
- logistic equation, 73–74, 226–27
- Mann–Whitney
 - test, 39–43
 - U* tables, 42, 256–61
- mathematical transforms, 12–13, 15–16, 68–74, 79, 217, 273
 - inverse exponential, 68
 - inverted exponential, 68–71, 217
 - logistic equation, 73–74, 226–27
 - multitarget cell-survival curve, 71–73
 - u*-transform, 79
 - z*-transform, 15–16
- MDR-1 (multidrug resistance gene), 233–35
- mean, 4, 6, 102–3
- mean deviation, 8
- median, 5–6, 102–3
- membrane enzymes, 212–17
- membrane transport, 231–40
- methotrexate, 236–38
- mode, 5–6, 102–3
- modelling population kinetics, 149–55
- monochlorobimane, 208–12
- Michaelis constant (K_m), 200–4, 213–16
- Michaelis–Menten rectangular hyperbola, 201
 - minimising squared deviations, 61–63
 - mitotic selection, 146–49
 - multidrug resistance, 233–35
 - multiparameter optimisation, 155–71
 - conventional space, 156–61
 - Euclidean hyperspace, 161–71
 - multitarget cell-survival curve, 71–73
 - multivariate analysis, 241–45
 - correlated bivariate density function, 242
 - density function, 241–42
 - two-dimensional surface fitting, 242–44
 - non-linear regression, 67–81
 - data transforms, linear, 68–74
 - polynomial regression, 77–81
 - successive approximation, 74–76
 - non-parametric statistics, 38–49, 256–64
 - Kolmogorov–Smirnov, 44–46, 262–64
 - Mann–Whitney, 39–43, 256–61
 - rank correlation, 46–49
 - non-specific binding, 100
 - normal distribution probabilities, 21, 250–51
 - see also* Gaussian distribution
 - normal equations of polynomial, 77–79
 - null hypothesis, 34, 45
 - optical design, as source of variation, 89–90
 - collection efficiency, 89
 - filter combinations, 90
 - fluorescence of filters, 89
 - p62^{c-myc}, 42–43, 180–85
 - p-glycoprotein, 233
 - parametric statistics, 32–38
 - standard error of difference, 32–34
 - Student's *t*, 34–37
 - variance assessment, *F*-ratio, 37–38
 - percent labelled mitosis (PLM) curve, 126
 - pH, 12–13, 83
 - photolysis, 83
 - Poisson distribution, 18–19, 26–28
 - polynomial, 77–81, 135
 - coefficients, 77–78
 - in DNA histogram analysis, 135
 - non-linear regression, 77–81
 - normal equations, 77–79
 - population sampling, 23–24
 - potential doubling time (T_{pot}), 156–57, 188–90
 - preparation and staining, as source of
 - variation, 82–86
 - binding-site modulation, 85–86
 - quenching, 82–83, 89
 - stoichiometry, 83–85
 - probability, 9–12
 - probability functions, 20–30
 - binomial distribution, 28–30
 - Gaussian distribution, 20
 - Poisson distribution, 18–19, 26–28
 - propidium iodide, 176–78, 211
 - Pythagoras, 7, 8, 33, 162
 - quenching, 82–83, 89
 - rank correlation, 40, 46–49
 - rate equations, 198, 200, 210, 213–14, 234–35, 272–74
 - rates of change, 198–99
 - ratio analysis of means, 114–16
 - reaction velocity, 201–4
 - rectangular hyperbola, 201
 - rectilinear integration, 132–33
 - regression analysis, 58–81, 265–68
 - computer routine, 265
 - linear, 60–67
 - non-linear, 67–81
 - resonant energy transfer, 85
 - S-phase probability function, 138
 - short time-scale kinetics, 220–28
 - significance tests, 31–53
 - association, χ^2 , 49–53
 - parametric, 32–38
 - non-parametric, 38–49

288 *Index*

- skew, 26
skewed-normal distribution, 105–12
sources of variation, 82–100
 biological, 99–100
 electronic, 90–99
 fluorescence excitation, 86–89
 optical design, 89–90
 preparation and staining, 82–86
Spearman’s rank correlation coefficient, 46
standard deviation, 8, 39
standard error
 of difference, 32–34, 37–38, 113
 of estimate, 63
 of mean, 24–25, 33, 37
 of standard deviation, 25
stathmokinetic analysis, 145–46, 165–67
statistical verification, 120–22
 Kolmogorov–Smirnov, 121
 simulation with $\Sigma \chi^2$, 122
 Student’s *t*, 121
stoichiometry, 83–85
Student’s *t*, 34–37, 48–49, 55–56, 80, 113, 121
successive approximation, 74–76
synchrony progression, 146–55
TCW analysis, 143–44
time of flight, 93
time, incorporation of, 197–98, 220–28
 T_{pot} (potential doubling time), 188–90
transforms, *see* mathematical transforms
transverse emission mode (TEM), 87
triggering and thresholds, 90–91
t-table, 36
two-dimensional surface fitting, 242–44
U-test, 40–43
variance, 8
variance assessment, *see* *F*-ratio
 V_{\max} , 201–4
Yates’s correction, 52