

ANALYTICAL TABLE OF CONTENTS

PREFACE	page vii
EDITORIAL NOTE	ix
GENERAL INTRODUCTION	xi
LIST OF PLATES	xxxix

PART 1

RESEARCHES INTO FLUXIONS AND INFINITE SERIES (1670–1671)

INTRODUCTION	3
--------------	---

Barrow reports on a new Newtonian treatise of infinite series (December 1670), 3. Newton tells Collins he has begun to ‘new methodiz’ the *De Analysi* but that other ‘buisnesse’ has delayed him (July 1671), 5. The revise now (Christmas 1671) to be appended to the edition of his ‘20 Dioptrick Lectures’, 7. Prompted by ‘wrangling disputes’ Newton abruptly decides (May 1672) to publish neither, 8. Sherburne still advertises the treatise on series as ‘expected’ (in 1673), 10. The manuscript copied by Craige (1685?) and Jones (1710?): Halley and Raphson still plan to print it (1691), 11. Pemberton’s intended publication thwarted by Newton’s death (1727): Colson’s English (1736) and Horsley’s Latin (1779) *princeps* editions, 13. Contemporary quotation by Newton of the 1671 tract, 14. His conception (inspired by Barrow?) of time and fluxional increase, 17. Quadrature of curves defined by trinomial equations (1676?), 18.

APPENDIX (ULC. Add. 3968. 39: 538^r). Newton’s citation of the 1671 tract in a draft preface to his *Commercium Epistolicum* (late 1712): quotations from Collins’ contemporary correspondence with James Gregory, Bertet, Borelli, Vernon and Newton (1670–2), and from Newton’s *epistola prior et posterior* for Leibniz (1676), 20.

1. PRELIMINARY SCHEME FOR A TREATISE ON FLUXIONS (In private possession)	28
-----------------------------------------------------------------------------	----

Two general problems on the direct and inverse methods of fluxions, followed by applications (21 problems) to the determination of extreme values, tangents, quadratures, curvature and rectifications and to the comparison of curves.

2. THE TRACT ‘[DE METHODIS SERIERUM ET FLUXIONUM]’	32
----------------------------------------------------	----

§1 (restored from contemporary transcripts by Jones and David Gregory). The missing first leaf of Newton’s autograph manuscript. Introduction: algebra as universal arithmetic, with infinite series the analogue of untermiated decimal sequences, 32. An example of reducing an algebraic fraction by infinite division, 36.

§2 (ULC. Add. 3960. 14: 3–132). The main autograph text. Further examples of reduction to infinite series by division, 38. Examples of reduction by extracting roots as infinite series (binomial expansion for index $\frac{1}{2}$), 40. The roots of equations reduced to infinite series: the numerical instance of $y^3 - 2y - 5 = 0$, 42. Extension of the method to general algebraic equations: preliminary observations, 46. The parallelogram rule for isolating ‘fictitious’ equations at any stage, 50. The method exemplified at length (x assumed small), 54. Used to invert the series expansions of $\log(1+y)$ and $\sin^{-1}y$, 58. Rounding off to desired accuracy: exemplified as before, but the fictitious cubic now supposed irreducible, 60. Adjustments to be made when x is assumed large, 64. Tentative (inaccurate) generalities on convergence: Newton generalizes Gregory’s concept of a (double) *series convergens*, 68. Axiomatic assumption of ‘time’ as an intuitively known, uniformly fluent base variable to measure the instantaneous ‘fluxion’ (speed of motion) at a point: Barrow’s evident influence, 70. Introduction of literal fluxions of fluent variables (typically l, m, n, r of v, x, y, z), 72.

Problem 1. Given an equation to find the corresponding fluxional derivative. Use of modified Huddonian multipliers in simple cases, 74. Geometrical justification and Fermatian ‘proof’ of the technique (by considering limit-increments om, on, or of x, y, z , where o is the time-increment), 78.

Problem 2. The inverse problem: given a fluxional equation, to determine its fluent form. Use of inverse Huddonian multipliers to determine (finite) solutions by inspection in simple cases of first-order fluxional equations involving two variables (x, y), 82. The solution of more general first-order cases: distinction of ‘related’ and ‘correlate’ quantities (to avoid confusion), 86. Case 1 (two fluent variables but the related quantity itself not present): straightforward reduction to integrable form by extracting n/m as the root of an algebraic equation, 90. Case 2 (two fluent variables, both present): preliminary reduction of fractions to equivalent infinite series is advised, 94. Derivation of a ‘*solutio particularis*’ of the reduced equation as an infinite series (x assumed small): for the most part examples are general de Beaune equations (of form $n/m = A + By$), 98. Elimination of fractional powers of the related quantity by substitution of a new one, 102. The ‘*solutio generalis*’ exemplified in the case of a de Beaune equation ($A = 1 - 3x + x^2, B = 1 + x$), 104. A ‘*solutio particularis*’ of the ‘Riccati’ equation $n/m = 3 + 2y - x^{-1}y^2$, 106. Exact ‘*solutiones particulares*’ of two de Beaune equations, 108. Examples where series solution is valid for large x : illustrated by a de Beaune equation ($B = x^{-2}$), 110. Fractional powers of the related variable eliminated by substitution of a new one: ‘this is very rarely of use’, 110. Case 3 (more than two fluent variables are involved): reduced to the preceding by appropriate elimination of all but two of the fluents, 112. ‘Demonstration’ of the validity of these solutions by taking their fluxions: the original fluxional equation must be obtained (equivalently as an infinite series), 112. The problem of the arbitrary constant of integration clarified on the geometrical model of points moving in line-segments, 114.

Problem 3. Determination of maxima and minima by the Fermatian technique of equating the fluxion to zero: simple examples, 116. List of nine problems to which the method may be applied, 118.

Problem 4. Determination of tangents to curves defined in nine ‘modes’ of coordinate system by considering the ratios of limit-increments of the co-ordinate variables (which are those of their fluxions). Mode 1, standard (oblique) Cartesian co-ordinates: examples include the pair of a line and an ellipse, the Cartesian trident and the conchoid, 120. The inflexion points in the conchoid determined by maximizing the subtangent, 126. Examples in more complex ‘mechanical’ curves (notably the quadratrix and Archimedean spiral)

Analytical Table of Contents

xxxix

and composite curves (influence of James Gregory and Barrow), 128. Mode 2, ‘focus-directrix’ co-ordinates: examples include the pair of a conic and quartic and the conchoid, 132. Mode 3, standard bipolars: instanced in the Cartesian oval, 136. Modes 4/5/6, variants on a simple monopolar system (the last a generalization of Cartesian co-ordinates): no examples, 138. Mode 7, polar co-ordinates: examples include the Archimedean and Fermatian spirals, 140. Mode 8, ‘quadratrix’ co-ordinates: the quadratrix itself is Example 2 (variant proof of Barrow’s construction of the subtangent), 144. Mode 9, ‘trochoidal’ co-ordinates (taken from Gregory): Example 2 is the general cycloid, 146. Enunciations of seven allied problems involving consideration of tangents (instantaneous direction of a curve at a point), 148.

Problem 5. Determination of the curvature of a curve at any point. Preliminary considerations: the curvature is that of the osculating circle, 150. Possible defining ‘symptoms’ of the centre of curvature: limit-meet of normals in the vicinity of a point, instantaneous centre of motion, centre of the circle having 3-point contact, 152. Analytical measure of the radius of curvature in Cartesian co-ordinates (centre defined as limit-meet of normals): examples include the general conic, cissoid, conchoid, 156. Curvature in the cycloid: properties of its evolute (a congruent cycloid), general observations on angles of contact stemming from the (infinite) curvature at its cusp, 160. The quadratrix, 168. Analytical measure of curvature in polar co-ordinates: instances in the Archimedean and general Fermatian spirals, 168. The (cumbrous) equivalent method of measuring curvature by constructing the circle having 3-point contact (worked through on the simple Apollonian parabola), 174. ‘Certain related questions’: to determine the points on a curve where there is a given curvature, 178. ‘Straight’ points (general inflexions, of zero curvature) and cusps (of infinite curvature), 178. Location of points of extreme curvature, 180. Determination of the evolute: the cycloid and parabola are cited as examples, 182: The diacaustic of refraction at an interface, 184. Radius of curvature at a vertex (the limit value of the subnormal): examples in the ellipse and conchoid, 184.

Problem 6. The ‘quality’ of curvature at a point. Justification of the (intrinsic) measure of ‘inequability’ of curvature by the ratio of the fluxion of the curvature radius to that of the arc-length (zero in the circle, constant in the logarithmic spiral), 186. Evaluation of this measure ($d\rho/ds$) in the general conic (varies as the ordinate), 188. The inequability of curvature in the logarithmic spiral and cycloid compared with that in a given parabola ($y^2 = 6x$), 192. Related problems: to determine the points of a curve where the inequability of curvature has a given value, and to identify members of a family of curves (especially conics) which have given curvature and variation of curvature at a point, 194.

Problem 7. Determination of the area of curves capable of exact algebraic quadrature. Statement of the fundamental theorem: the fluxion of the area of a curve is proportional to its ordinate, 194. Examples: the Apollonian and other simple parabolas, 196.

Problem 8. General transformations which preserve area. Transform of

$$s = \int v \cdot dx \rightarrow t = \int y \cdot dz$$

by $f(s, t, v, x) = 0$ and $\phi(s, x, z) = 0$: general observations, 198. Examples in geometrical curves: the circle $v^2 = ax - x^2$ and hyperbola $v^2 = x^2 + c^2$, 198. The cissoid $v^2 = x^3/(a-x)$, 202. Use of integration by parts to evaluate the area under mechanical curves (of algebraic slope): exemplified in the cycloid and sinusoid, 204. Cavalierian area-preserving convolution of a Cartesian curve into a corresponding polar spiral: the Apollonian parabola and its convolute (an Archimedean spiral), 206. The main use of Problems 7 and 8 is to draw up catalogues of the areas of curves, 208.

Problem 9. Methods of determining the area of any curve proposed. Further examples of simple exact quadrature in the case of general parabolas and hyperbolas, 210. By reducing the ordinates to convergent infinite series all curves may be reduced to integrable form: examples are mostly taken over from the introduction, 212. Integral bounds: in all cases the area is the difference of the values of the integral corresponding to the upper and lower limits of the base, 216. Interpretation of negative areas, 218. ‘Square’ notation for hyperbola-area (to be evaluated by an infinite series) introduced and exemplified, 218. Numerical calculation of areas expressed by infinite series: parallel computation of the hyperbola/circle segments $\int_0^{\frac{1}{2}} \sqrt{[x \pm x^2]} \cdot dx$, 222. An evaluation of π (correct to 15 D) is a corollary, 226. Computation of the area $\log(1+x)$ under the hyperbola $y = 1/(1+x)$, x small: a simplified version of his 1665 calculations, 226. Construction on this basis of a canon of natural logarithms (of low primes): their conversion to common form by dividing through by $\log 10$, 230. A method of subtabulating means in this table, 232. A hint that it may be generalized to ‘interpolating’ arbitrary means, and that similar methods will produce tables of logarithmic sines and tangents without prior construction of the natural sines and tangents, 234. A ‘prior’ catalogue of curves of form

$$y = dz^{\lambda\eta-1}(e+fz^\eta)^\mu$$

and their exact areas ‘constructed by the help of Problem 7’, 236. And of the ordinates of curves whose area is $z^\theta(e+fz^\eta+gz^{2\eta})^\lambda(h+iz^\eta)^\mu$, ‘by means of which the way is opened up to higher things’, 238. Preliminary explanation of a diagram depicting instances of the general central conics $d = (e+fx)v$ and $v^2 = e+fx+gx^2$, 240. A ‘posterior’ catalogue, ‘constructed with the help of Problem 8’, evaluating the areas of curves of form

$$y = dz^{\lambda\eta-1}(e+fz^\eta+gz^{2\eta})^\mu$$

and $y = dz^{\lambda\eta-1}(e+fz^\eta)^\mu(h+iz^\eta)^\nu$ in terms of the areas s (and σ) of these, 244. Observations on the two catalogues: unknowns may be positive or negative, 258. Ways of indefinitely extending either table, 260. Duplicate entries are retained because of their usefulness, 262. Negative areas: correctly choosing bounds, 264. Similar tables might be constructed involving comparisons with curves of higher order, 264. The latter catalogue used to square special curves: the *versiera* (witch of Agnesi), 266. The kappa (Gutschoven) quartic, 268. The cissoid and conchoid, 270. The quartic $y^2(c^2-z^2) = (z^2+bz-c)^2$, 272. The quadrature of mechanical curves is (where possible) to be reduced to that of allied geometrical ones by means of Problem 8 (integration by parts when slope is algebraic, or reverse convolution from polar to Cartesian form), 274. An integration by parts yields the area of the ‘conic-arc’ curve, 276. Justification of the quadratures of the four latter curves by *ad hoc* limit-increment methods, 278. Equivalent fluxional arguments in the case of the kappa quartic and cissoid, 282. Scholium. The area and ordinate of a curve is but a canonical instance of the relationship of a fluent quantity to its fluxion: by choosing appropriate units the two preceding catalogues may be used in all problems of integration (rectifying arcs, measuring solid content, and so on), 284. Related topics: the ‘mechanical’ approximation of areas, 288. Given the area z of a curve $r = f(x)$ as an infinite series in terms of its base x , to find the ordinate r (or \dot{z}) as a series in z , 290.

Problem 10. To find arbitrarily many curves of finite (algebraic) arc-length. Five preliminary observations on evolute properties: the length of a convex evolute’s arc is the difference of the curvature radii at the points on any involute of it corresponding to its end-points, 292. Measure of the curvature radius in a Cartesian co-ordinate system: rectification of Neil’s semicubical parabola by determining the curvature of its involute

Analytical Table of Contents

xxxiii

Apollonian parabola, 296. The same performed in semi-intrinsic 'Newtonian' co-ordinates: an equivalent rectification of Neil's cubic by means of the involute parabola, 298. An insufficient criterion for a cusp, 300.

Problem 11. To find arbitrarily many curves whose arc-lengths may be expressed algebraically in terms of that of a given curve. First method: the arc-length v of the given curve $f(s, t) = 0$ is related to that of the curve $F(x, y) = 0$ determined in semi-intrinsic 'Newtonian' co-ordinates by some transform $x = \alpha(s)$, $y = \beta(s, v)$, 304. Exemplified for a circle ($f \equiv s^2 + t^2 - as$) and hyperbolas ($f \equiv st - a^2$ and $(a/c)s^2 - t^2 - a^2$), 306. Second method: the Gregorian length-preserving 'involution' of a Cartesian curve into an equivalent polar 'spiral', 308. The primary example is Roberval's theorem that the involute of an Apollonian parabola is an Archimedean spiral, 314.

Problem 12. The rectification of curves defined in Cartesian co-ordinates. The arc-length t of the curve $f(y, z) = 0$ is found by 'integrating' its fluxion q (or t) = $\sqrt{[y^2 + z^2]}$, 314. Examples (finite) of the rectification of simple cubics (including Neil's parabola) and quartics, 316. 'Logarithmic' rectification of the cissoid by relating its arc-length to the area under a constructed hyperbola, 320. That of an 'elliptical' sinusoid, 322. Rectification of a curve defined in terms of the (logarithmic) complementary area of a hyperbola is reduced to finding the area under a conic, 322. Arc-lengths determined as infinite series in the base variable: examples of the hyperbola, ellipse and quadratrix, 236.

§3 (ULC. Add. 3960. 4: 33–46). An intended addendum on geometrical fluxions (to Problem 9 preceding). Proofs by fluxions appear more 'natural' than those by limit-increments, 328. Four axioms on the fluxions of uniformly fluent magnitudes, 330. The fluxion of the product of two magnitudes is equal to the sum of the products of each multiplied into the other's fluxion, 330. Nine corollaries extend this fundamental theorem to finding the fluxion of the n -th power of a magnitude, 334. The fluxional form of Pythagoras' theorem on right triangles: (cancelled) remarks on the fluxion of an area, 338. The fluxions of the 'legs' of a triangle of given base, whose vertex is on a given straight line. 340. Those of the intercepts cut off on a given line by a rotating radius vector, 342. The fluxion of the area of a rectangle and triangle as one side 'flows' parallel to itself, 344. That of the area generated by a rotating radius vector, 346. That of an arc intercepted by the radius vector, 348. The contact point of a tangent 'rolling' round a curve is the instantaneous centre of motion, 350. Equivalent fluxional proofs of Examples 2, 3 and 4 of Problem 9 of the 1671 tract, 350.

APPENDIX (ULE. David Gregory MS, A56). Gregory's 'Tractatus de Seriebus infinitis et Convergentibus' (c. 1690, copied by Craige?). The opening pages duplicate those in the 1671 tract, 354. Summary of Newton's 'reduction of affected equations', 358. Problems 1 and 2 drastically summarize those of Newton, 364. Problems 3, 4 and 5 are short extracts from Newton's Problems 9, 7 and 8, 366. Problem 6 repeats his method of constructing the circle of curvature (Problem 5) as that having 3-point contact with a curve, 369. Loose (English) jottings from Newton's Kinckhuysen 'Observations' and his 1676 *epistola prior* to Leibniz, 370.

3. THE QUADRATURE OF CURVES DEFINED BY POLYNOMIALS 373

§1 (ULC. Add. 3962. 6: 77^v). First calculations, 373.

§2 (ULC. Add. 3962. 6: 74^r–76^r). First extended draft. Transformation of the area

$s = \int v \cdot dx$ under the curve $f(x, v) = 0$ into that, s/τ , under $f(z^\tau, yz^{1-\tau}) = 0$: determination of the ordinate of a curve from its area, and the converse (simple polynomial examples), 374. ‘The Quadrature of all Curves whose \mathcal{A} equations consist of but three termes’: threefold determination of the area $s = \int v \cdot dx$ under $bv^\alpha + cv^\beta x^\epsilon + dx^\zeta = 0$ from that, $s/\tau = \int y \cdot dz$, under its transform $(Ay^\alpha + By^\beta)^\gamma = z$: a simple example is given, 375. Reducing ‘y^e index of y^e dignity’ to ‘y^e least fraction’ and so determining ‘y^e simplest curve wth w^{ch} y^e pro-pounded curve may be compared’, 376.

§3 (ULC. Add. 3962. 6: 70^r–72^r). The improved, augmented version. Reduction in the general trinomial case, 380. ‘The Quadrature of many Curves whose \mathcal{a} equations consist of more then three terms’, 384. Another rule for binomials, 385.

PART 2

MISCELLANEOUS RESEARCHES

(Early 1670’s)

INTRODUCTION

389

Newton’s restyling of Book 2 of Euclid’s *Elements* and ‘A fragment relating to the comparison of curved surfaces’ (intended for professorial lectures?), 389. Simple harmonic motion induced by gravity in a vertical cycloid: Newton’s possible debt to Huygens, 390.

APPENDIX. Huygens, Brouncker and Pardies on cycloidal motion. Galileo and Mersenne on the (near) isochronism of the circular pendulum: Huygens’ prior analytical investigation (December 1659) and James Gregory’s published series solution (1672), 392. Huygens’ 1659 discovery that (for constant gravity to a distant point) the cycloid is the true tautochrone, 394. Brouncker’s first attempts to confirm Huygens’ result (January/February 1662), 395. Hooke’s criterion (1666) for isochronous motion, 398. Brouncker’s third proof that the cycloid is the tautochrone (May 1673): Huygens’ reservations, 398. Pardies’ independent proof (pre-1673), 400.

1. THE SECOND BOOK OF EUCLID’S ‘ELEMENTS’ REWORKED

402

(ULC. Add. 3959. 2: 22^r; Add. 3970. 11: 635^v/637^v). Condensed geometrical restatement (using squares and rectangles) of Propositions 1–10, 402. ‘Improved’ version of Propositions 12 and 13: diagrams for Propositions 11 and 14 (repeated from Barrow’s 1655 edition), 406.

2. RESEARCH INTO THE ELEMENTARY GEOMETRY OF CURVED SURFACES

408

(ULC. Add. 3963. 16: 184^r–185^v). Axiomatic bounds to surface area and solid content, 408. The area of a circle sector and the volume of a prism standing upon it, 408. The curved surface and content of a cylindrical wedge: introductory lemmas and proofs, 412. The curved surface of a conic frustum, 416. The area of the zone of a sphere (regarded as the limit of a ‘pumpkin’) and of a ‘spherical triangle’ (one side of which is a small circle parallel to the tangent plane at the opposite vertex), 418.

Analytical Table of Contents

xxxv

3. HARMONIC MOTION IN A CYCLOIDAL ARC 420

(ULC. Add. 3958, 5: 89^r–91^v). The force ('power') induced by simple gravity in a vertical cycloid is proportional to the distance of fall along it to its lowest point, 420. Hence all bodies, wherever let fall from rest in it, reach that point in the same time, 422. Cycloidal 'cheeks' of the (congruent) evolute deflect a pendulum bob into a cycloidal path, 424. Newton uses a (Huygenian) limit-increment argument to obtain an explicit measure of the total time of fall, 428. Harmonic motion in the cycloid corresponds to uniform motion in the generating circle, 430.

PART 3**RESEARCHES IN GEOMETRICAL OPTICS***(c. 1670–1672)***INTRODUCTION 435**

Newton's '20 Dioptrick Lectures': the autograph version retained and the (variant) copy deposited in 1674, 435. They intentionally pursue the theme of Barrow's preceding lectures, 437. In amplification, Newton expounds his own hypothesis that 'white' light is a congeries of elemental coloured rays, each separately refracted, 438. The preamble to his first lecture (January 1670), 438. Mathematical aspects of the two versions of the 'Lectiones': construction of refraction points in a plane interface, models for the variation of dispersion with that of the density of the refractive medium, 440. Refraction at a spherical interface: the theory of thin lenses and explanation of the radius and colour-sequence of the n -ary bow (especially $n = 1, 2$), 441. Newton's failure to construct an achromatic refractor (but he is not convinced that a colour-free compound lens is impossible), 442.

APPENDIX 1. Newton's exordium to his optical lectures (ULC. Add. 4002: 1–2): variants in the deposited copy (ULC. Dd. 9. 67), 444.

APPENDIX 2. Pemberton's (or Jones'?) editorial description of the 'Opticæ Pars prima'. Résumé of Newton's optical discoveries, 445. Breakdown of this first part of his lectures (as deposited), 446. Appreciation of Barrow's influence, 449.

1. EXTRACTS FROM NEWTON'S LECTURES ON OPTICS (1670–1671) 450

§1 (ULC. Add. 4002: 99–100, 92–5/101–2, 115–20). Refraction at a plane interface. Construction of the refraction point: the resulting quartic is solved in Cartesian style by the meets of a circle and parabola, 450. Barrow's variant construction in lecture 5 of his *Lectiones* (used by Newton in his revise), 452. Barrovian centres of radiation: their construction, 456. The locus of these centres (a Dioclean cissoid) as the refractive index varies, 462. Extremes of dispersion in refractive media of varying density: the basic geometrical model, 464. The problem of locating maximum dispersion is 'solid' (reducing to a cubic equation), 468. The 'plane' construction of this maximum when the dispersion is small, 472.

§2 (ULC. Dd. 9. 67: 64–77). Refraction at a curved interface. Construction of the principal focus of rays refracted at a spherical interface, 476. Extension to several interfaces: application to the construction of image points in telescopes and microscopes suggested, 480. Lemma: use of the subnormal to construct the centre of curvature at a conic's vertex, 480. Finding the principal focus for rays incident on a general interface of known vertex

curvature from a point in its axis, 480. The error by which parallel rays refracted at a spherical interface digress from the principal focus: expression of this spherical distortion as an infinite series, 482. Corollaries: measures of the depth of focus of a spherical interface and of its 'lateral' error, 484. The image of a point when rays incident from it are refracted over a small region of a spherical interface: the Barrovian condition for the spherical diacaustic, 488. The particular case where the object point is at infinity: the condition $di/dr = \tan i/\tan r$, 492. Generalization to any curved surface whose curvature is known, 494. Descartes' refractive 'ellipses', 496. Lemma: determination of $\tan i/\tan r$, given $\sin i/\sin r$ and $\cos i/\cos r$, 498. Determination of the angular radius of the primary bow (Harriot's priority), 500. And of the secondary bow, 504. Generalizations to bows of higher orders, 506. Estimate of the chromatic distortion of a spherical interface, both laterally and in depth, 508. Comparison of spherical and chromatic distortion for a telescope lens (object at infinity): the latter is 'far greater', 510. Possibility of eliminating chromatic distortion in a compound lens by 'contrary refractions', 512.

2. MISCELLANEOUS RESEARCHES INTO REFRACTION AT A CURVED INTERFACE

514

§1. Theory of spherical lenses. [1] (ULC. Add. 4004: 71^r-72^v). 'Optical theorems'. The basic refraction condition $1/r - 1/u = (I/R)(1/r - 1/v)$: the case of reflection ($I = -R$), 514. Fluxional derivation of this theorem, 516. Problems: given a spherical lens (plano-convex or doubly convex), to find the refractive index and the curvature of the surfaces, 520. The method exemplified for two telescopes (one Babington's), 524. [2] (ULC. Add. 4004: 73^r). 'Outline of a new telescope': a coated concave lens is used (in place of a simple mirror) in a Newtonian reflector to minimize chromatic distortion, 526.

§2 (ULC. Add. 4004: 90^r). Descartes' problem of twofold refraction resolved. Given one refracting interface, the refractive index, radiating point and focus, to define the second refracting interface through any point: Newton's construction implicitly invokes the common diacaustic of the two points, 528. Extension to three or more refracting interfaces, 530.

APPENDIX 1. Miscellaneous optical calculations. [1] (ULC. Add. 4004: 72^v/73^r [insert]). Construction of the general diacaustic point, 532. [2] (ULC. Add. 3958. 2: 32^r). Fluxional evaluation (nullified by a slip) of the 'error' between diacaustic point and principal focus of a spherical interface: Cartesian notation for refractive index used, 532. [3] (ULC. Add. 4004: 72^v/73^r [insert]/3970. 11: 634^r). Straightforward computation (again nullified by a small slip) of this refractive error, 534. [4] (ULC. Add. 3970. 11: 638^r). A successful attempt to evaluate the error (Barrow's notation is used) by means of a diacaustic involute, 536. [5] (ULC. Add. 3970. 11: 638^r/638^v). The revised version with diacaustic involute eliminated (substantially the one incorporated in the 'Optical problems'), 537. [6] (ULC. Add. 3970. 11: 633^r/634^v). A preliminary worksheet for the 'new' catadioptrical telescope correcting for chromatic distortion: a small error vitiates the construction, 539. [7] (ULC. Add. 4004: 72^v/73^r [insert]). Revised computations for the 'new' telescope (Cartesian notation used): a numerical example is given, 541.

APPENDIX 2. Newton's explanation of the rainbow. [1] (ULC. Add. 3975: 14). Observations, based on chapter 8 of Descartes' *Meteores* (Latin version), of the colour of the refractions of a 'large glasse Globe filled wth water', 543. [2] (ULC. Dd. 9. 67: 96-100). Newton's deduction of the colour sequence, bow width and angular radii of the primary

Analytical Table of Contents

xxxvii

and secondary rainbows: the light refracted after three internal reflections is 'scarce strong enough to cause a sensible Bow', 544.

APPENDIX 3. (ULC. Add. 3970. 3: 428 *bis*^r/3970. 9: 615^r-617^r). The general problem of twofold refraction, point to point, as solved in the *Principia* (Book 1, Section 14, Propositions 97-8). The problem resolved for a single interface (by Descartes' ovals), 549. Construction of the second interface in twofold refraction, 551. But chromatic aberration (rather than spherical distortion) remains the main 'impediment' to the perfection of dipotrics, 552

APPENDIX 4. (ULC. Add. 3970. 3: 471^r). Correcting (1692?) for spherical distortion by means of a compound (glass-water) lens: but 'y^e different refrangibility of several rays' still prevents the perfection of telescopes, 553.

APPENDIX

NEWTON'S MATHEMATICAL CORRESPONDENCE

(1670-1673)

Topics in Newton's contemporary mathematical correspondence, not dealt with in his extant papers, are solved by applying general techniques developed earlier, 558. The two problems proposed in late 1669 by Collins, 559. Newton's use of 'Gunters line' (July 1672) to locate the roots of numerical equations (approximately), 560. Summing a general 'musical' series: exact computation (January 1670) when the harmonic progression has only a few terms, 561. Logarithmic approximation (February 1670) by comparison with the area under a rectangular hyperbola: justification of Newton's harmonic and geometric bounds (July 1671), 563. Solution of Collins' annuity problem (February 1670): 'I could give exacter solutions but . . . have noe leisure', 566. Series solution (February 1670) to Dary's problem of finding the area of a zone of a circle: extension (late 1670) to computing the 'second segment' of a spheroid, 567. Newton's approximation (August 1672) of that of an ellipsoid as the second segment of a 'parabolick spindle': the rule is 'not exact but approaches . . . exactly enough for practice', 568. His communication to Collins (August 1672, without proof) of his organic description of conics, 569. Dary's problem (September 1673) of constructing an equilateral 24-gon in an ellipse: Newton calculates that its construction would require solution of an equation of 324-th degree, 570. His contemporary optical correspondence and replies to Collins' request for algebraic enlightenment, 571.

INDEX OF NAMES

573