

Index

- acetylaminofluorenes, structures, 231
- acidic soft waters
- aluminium, mechanism of toxicity, 116–24
 - chronic sublethal exposure, acclimation by trout, 143–67
- acids, uptake rate vs pH of water, 5
- adenovirus E1A and CREB, 191
- aflatoxins
- carcinogenicity, 238–9
 - structures, 231
- Agent Orange, 269
- Ah *see* aryl hydrocarbon (Ah)
- alkylphenol polyethoxylate surfactants (APEs)
- nonylphenol, 205
 - oestrogenic activity, 210–11
 - oestrogenic products, 211
- aluminium
- in acidic soft water, mechanism of toxicity, 116–24
 - adrenocortical response, 124–5
 - dietary exposure, 33–6
 - excretion, 35–6
- ammonia
- accumulation, toxic action of copper, 99–102
 - oxidation to nitrite, 170
 - and swimming performance, 102–6
- Anguilla japonica see* eel, Japanese
- antioxidant enzymes, 233–5
- AP1 genes, 191
- Arnt protein (Ah receptor nuclear translocator), 270
- arochlor-1221
- oestrogenic activity, 208
 - structure, 206
- aryl hydrocarbon (Ah), 269
- and P4501A1, 271–6
- aryl hydrocarbon receptor, 269
- Astacus astacus see* crayfish
- β -HCH (β -hexachlorocyclohexane), 206, 208
- bacterial mutagenicity assay, 231
- bacterial nitrification *see* nitrite
- bass
- activation of benzopyrenes, 231
 - oxygen consumption and TCB uptake rate, 8–14
 - recovery from nitrite intoxication, 182
- benzo(a)pyrenes (BaP), 234, 259
- structure, 231
- BHC, 208
- biotransformation caused by pollutants, 226–35
- biotransformation enzymes, 229
- birds, mixed-function oxygenase (MFO) systems, 228, 229
- blood:water distribution coefficient K_b , 6
- cadmium
- bioaccumulation, 49
 - dietary exposure, 36–42
 - environmental monitoring, 196–8
 - metallothionein synthesis, 192
- calcium, fish, uptake, shared pathway with zinc, 72–3
- Carassius auratus see* goldfish
- carcinogens, 226, 228
- pro-carcinogens, structures, 231
- Carcinus maenus see* crab
- carp
- copper toxicity, 100
 - nitrite: acid–base balance, 178; entry into red cells, 172–4; uptake and accumulation, 170–2
- catecholamines
- interactions with cortisol, 125
 - monitoring in AI-stressed fish, 127
 - release, 125
 - in stress, 16
- catfish
- antioxidant enzymes, 233–5
 - nitrite intoxication, 182
 - oxygen consumption and TCB uptake rate, 8–14

280 *Index*

- catfish (*cont.*)
 reactive oxygen species (ROS), 236
 sublethal Cu toxicity, 46
 caudal neurosecretory system (CNS),
 urotensins, 132
 copper, 84–113
 in aquatic environments, 84–6;
 mobilization in acid episodes, 84;
 surface water, 84
 aquo ions, 86
 Cu²⁺ toxicity, 86
 dietary exposure, 42–6
 intracellular, regulation, 263
 metallothionein synthesis, 192
 toxic action, 87–91: ammonia,
 swimming performance, 103–6;
 ammonia accumulation, 99–102;
 sublethal effects, 94–9; trout:
 freshwater, 91–3, 94–7; seawater,
 93–4
Coregonus clupeaformis see lake
 whitefish
 cortisol
 induction of metallothionein synthesis,
 193
 interactions, 125
 coumestrol, structure, 206
 crab, shore
 activation of pro-carcinogens, 232
 nitrite effects, 170–86
 crayfish
 nitrite uptake and accumulation, 170–2
 recovery from nitrite intoxication, 182
 CYP450s
 cytochrome P4501A1 (CYP1A1), 267–
 78; environmental monitoring, 271–
 6; mRNA, induction kinetics, 261,
 273; variables affecting, in fish,
 274–6
 metabolism, 259–64
 see also mixed-function oxygenase
 (MFO) systems
Cyprinus carpio see carp
 dab, antioxidant enzymes, 233–5
 DDT
 induction of vitellogenin synthesis,
 219
 oestrogenic activity, 208–9
 structure, 206
 2,6-dibromo-4-nitrophenol (DBNP),
 uptake rate vs pH of water, 5
Dicentrarchus labrax see bass
 3,4-dichlorobenzoic acid (DCBeA),
 uptake rate vs pH of water, 5
 2,4-dichlorophenol (DCP), uptake rate
 vs pH of water, 5
 dietary exposure to toxic metals, 29–60
 diffusion coefficient, defined, 7
 7,12-dimethylbenzanthracene, structure,
 206
 dioxin
 PAH product, effects, 251
 structure, 206
 see also polycyclic aromatic
 hydrocarbons
 DNA damage
 indicators, 232–5
 mechanisms, 235–9
 eel, Japanese, aluminium in diet, 36
 environmental monitoring
 cytochrome P4501A1, 271–6
 cytogenetic aberrations, 241
 cytotoxicity testing, 254–9
 metallothioneins, 187–203:
 detoxification of metals, 196–8
 epithelium: water distribution coefficient
 K_m, 6
 ethinyloestradiol, structure, 206
 flounder
 cadmium exposure, 197
 copper exposure, 94
 cytochrome p450 reductase, 235
 reactive oxygen species, 236
 starry flounder: carcinogens, 230;
 DNA adducts, 237–8
 free radical scavengers, 232–3
Gambusia affinis see mosquito fish
 gas chromatography, ECD–GC, 19
 genetic toxicants, 225–51
 biomarkers of exposure, 239–43
 DNA damage, 232–9
 gills, teleost
 acidic soft water plus Al, chronic
 sublethal exposure, 144–51
 Al exposure, dimensions, 151
 calcium and zinc uptake, model, 74
 damage by copper, 88
 microenvironment, 3–5
 water chemistry, 1–16
 zinc and calcium uptake, 74
 glucocorticoid responsive elements
 (GREs), 191–2
 goldfish, oxygen consumption and TCB
 uptake rate, 8–14
 haemocytometry, 254
 HCB, 208

Index

281

- heavy metals, binding *see* metallothioneins
- β -hexachlorocyclohexane
oestrogenic activity, 208
structure, 206
- Ictalurus punctatus see* catfish
- immune system, nitrite immune system, 182
- kepone
activity, 208
structure, 206
- 2-keto-4-methylbutyric acid (KMBA), 234
- lake whitefish, cadmium
bioaccumulation, 49, 50
- lead bioaccumulation, 49
- Limanda see* dab
- lindane, oestrogenic activity, 208–9
- lipids, analysis, 19–28
- liver, nitrite damage, 182
- log uptake rate, bioaccumulation of xenobiotics, 3–4
- marine mammals
bioaccumulation of oestrogenic substances, 212
DNA adducts, 241
mixed-function oxygenase (MFO) systems, 228, 229
- mercury
bioaccumulation, 49
dietary exposure, 29–33
metallothionein synthesis, 192
- metal responsive elements (MREs), 191
- metallothioneins
detection, 190
endogenous regulation, 193–6
gene promoters in fish, 191–2
gene transcription, induced, 192–3
induction in trout fry, 262
metal detoxification, 196–8
metal responsive elements (MREs), 191
properties, 188–90
structure, 187–8
use in environmental monitoring, 187–203
zinc ligand, 67–70: model of synthesis, elimination and functions, 68
- metals
detoxification, 196–8
dietary exposure, 29–60; relative contributions of diet and waterborne sources, 48–50
- methoxychlor, oestrogenic activity, 208–9
- Micropterus salmoides see* bass
- minnow
DNA damage, 239
metabolism of BaP, 261
oxygen consumption and TCB uptake rate, 8–14
reactive oxygen species, 236
TCB accumulation, 23–8
- Mirex, oestrogenic activity, 208
- mixed-function oxygenase (MFO) systems, 228, 229
activation by toxic compounds, 259
see also CYP450s
- molluscs
mixed-function oxygenase (MFO) systems, 228, 229
reactive oxygen species, 236
- mosquito fish
DDT, induction of vitellogenin synthesis, 219
zinc bioaccumulation, 49
- mussel
activation of xenobiotics, 231, 232
DNA adducts, 241
DNA damage, 239
micronuclei, 242
reactive oxygen species, 236
- nitrite, 170–86
and acid–base balance, 178
in aquatic environments, 170–2
electrolyte balance, 178–81
immune system, damage, 182
liver damage, 182
N-nitroso compounds, 181–2
recovery from intoxication, 182
toxic action, 170
uptake and accumulation, 170–2
- nonylphenol
alkylphenol polyethoxylate surfactant degradation, oestrogenicity, 205, 213
structure, 206
- noradrenaline, induction of metallothionein synthesis, 193
- octanol:water partition coefficient K_{ow} , 3
- oestradiol
feminization of fish, 215, 217
induction of metallothionein synthesis, 193
role in fish, 214–15
structure, 206, 205

282 *Index*

- oestrogenic substances
 aquatic environment, 205–24
 bioconcentration and bioaccumulation, 212–13
 defined, 205
 effects on aquatic organisms, 213–19
 natural and synthetic, 211–12
 nonylphenol, alkylphenol
 polyethoxylate surfactants, 205
 relative potencies, 216
 in sewage, 212
 structures, 206
- oestrogens, of action, 215–16
- Onchorhynchus tshawytscha* *see* salmon (Chinook)
- Oncorhynchus mykiss* *see* trout (rainbow)
- organochlorine pesticides, oestrogenic activity, 208–9
- osmoregulation, prolactin, 127–9
- oxygen data bank (OXYREF), 7–8, 12–13, 24–5
- oxygen transport, effects of nitrite:
 crustaceans, 177–8; fishes, 174–7
- P4501A1 *see* cytochrome P4501A1
- Parophrys vetulus* *see* sole
- perch
 micronuclei, 242
 reactive oxygen species, 236
- phenol, uptake rate vs pH of water, 5
- phosphorus, P-32 post-labelling, 238
- Pimephales promelas* *see* minnow
- plaice
 benzo(a)pyrenes metabolism, 260
 P4501A1 cycle, 275
 zinc bioaccumulation, 49
- Platichthys stellatus* *see* flounder (starry)
- Platyichthys flesus* *see* flounder (common)
- Pleuronectes platessa* *see* plaice
- pollutants
 biotransformation, 226–35
 cytotoxicity testing, 254–9
see also environmental monitoring:
 xenobiotics
- polychaete worm, biomarker of DNA damage, 232
- polychlorinated biphenyls (PCBs)
 bioaccumulation, 229
 effects, 251
 oestrogenic activity, 209, 212–13
- polychlorinated dibenzodioxins (PCDDs), oestrogenic activity, 209–10, 212–13
- polycyclic aromatic hydrocarbons, halogenated (HPAHs), 267, 268
- polycyclic aromatic hydrocarbons (PAHs)
 activation of receptor, 269–71
 bioaccumulation, 229
 DNA adducts, 237–8
 oestrogenic activity, 209–10, 212–13
 P4501A1 levels, 272–6
 reactive oxygen species production, 235
 structures, 268
- progesterone, induction of
 metallothionein synthesis, 193
- prolactin, monitoring in Al-stressed fish, 127–9
- quantitative structure–activity relationships (QSARs), 254
- reactive oxygen species (ROS), 233, 239
 NADPH-dependent, 234
- Salmo gairdneri* *see* trout (rainbow)
- Salmo trutta* *see* trout (brown)
- salmon
 aluminium in diet, 36
 Chinook, copper toxicity, 93
 oxygen transport, and swimming performance, 150
 thyroid hormones, 129–30
- Salmonella* mutagenicity assay, 231
- Salmonella typhimurium* *umu*-test, 232
- Scophthalmus maximus* *see* turbot
- sole
 carcinogens, 230
 DNA adducts, 237–8
 DNA damage, 239
 spectrophotometry, 254
 spot, zinc bioaccumulation, 49
- starfish, activation of pro-carcinogens, 232
- sunfish, PAH pollution, strand breaks in DNA, 243
- superoxide radical, 233
- surfactants, oestrogenic products, nonylphenol, 205
- swimming performance, and oxygen transport, 150–4
- 2,3,7,8-TCDD *see* dioxin
- teleost fish, nitrite effects, 170–86
- 1,2,4,5-tetrachlorobenzene (TCB)
 analysis, 8, 18
 bioaccumulation, 18–28
 uptake rate in fish, 8–12: uptake due to skin binding, 13

Index

283

- tetrachloroguaiacol (TCG), uptake rate
in fish, 12–13
- thyroid hormones, teleosts, 129–32
- tilapia
PAHs, 240
prolactin monitoring, 128
- toxic equivalency factor (TEF), 269
- 2,4,6-trichlorophenol (TCP), uptake rate
in fish, vs pH of water, 5
- trout
*note: references are undifferentiated
since they may apply to more than
one species of trout*
acidic soft water: aluminium, 115–42:
chronic sublethal exposure,
acclimation, 143–67
aluminium in acidic soft water, 115–
42: mechanism of toxicity, 116–24;
physiology and metabolic costs,
143–67
aluminium in diet, 36, 37
antioxidant enzymes, 233–5
cadmium in diet, 36–42, 49
cadmium exposure, 197
caudal neurosecretory system,
urotensins, 132
copper: in diet, 38, 42–6; toxic action
of copper, 91–4
feeding, effects of Al/acid, 154–5
lead bioaccumulation, 49
mercury in diet, 38, 49
metallothionein levels, 194–6
nitrite uptake and accumulation, 170–2
oestrogenic substances, response to
sewage effluent, 217, 218
oxygen consumption: and TCB uptake
rate, 8–14, 18–28; and TCG uptake
rate, 12–13
oxygen transport, and swimming
performance, 150–4
prolactin responses, 128–9
protein turnover and growth, effects
of Al/acid, 155–8
reactive oxygen species, 236
recovery from nitrite intoxication, 182
RTG (rainbow trout gonad) cell line,
261
- STE (steelhead trout embryo) cell
line, 261
- TCB accumulation, 8–14, 18–28
- thyroid hormones, 129–132
- zinc: bioaccumulation, 46–8, 49;
dietary requirements, 48
- turbot, dietary zinc, 48
- UBC respirometer, 18
- uptake rate, log uptake rate and log
 K_{ow} , 4
- urotensins, caudal neurosecretory
system, 132
- vitellogenin
as measurement of exposure to
oestrogenic substances, 216–19
production by liver, 214
- vitellogenesis, excessive, 219
- water chemistry, gill surfaces, 1–16
- xenobiotics
bioaccumulation, log uptake rate, 3–4,
18–28
biotransformation, 226–35
examples, 227–8
metabolic fate and molecular toxicity,
227
uptake at gill surfaces, 1–16:
modelling, 5–14
- zinc
in aquatic environments, 61, 70–1
fish: adaptation to waterborne zinc,
73–5; bioaccumulation, 49; dietary
exposure, 46–8, 71; dietary
requirements, 48, 61–2; internal
regulation, 65–70; physiology and
toxicology, 61–84; toxicity,
mechanisms, 70–2; uptake: routes of
uptake, 62–5; shared pathway with
calcium, 72–3; various tissues, 63
intracellular, regulation, 263
ligand, metallothionein, 67–70
mammals, 62
metallothionein synthesis, 192