

Index

- Acaena*, description of, 37
 on South Georgia, 37
- Acaena magellanica*, effects of reindeer
 grazing, 218, 224
 exclusion experiment 228–229, 233–240,
 242
 nitrogen content, 113
 nutrient dynamics, 96–97
 on Barff Peninsula, 96–97
 on South Georgia, 38, 93–97, 102, 259
 propagation, 238
 seasonal use as forage, 102, 107–111,
 113, 198–200, 257
- Acaena tenera*, on Barff Peninsula, 95–96
 on South Georgia, 231
 propagation, 238
- Accidental deaths, 157, 160–162, 187
- Adak Island, *see* Aleutians
- African buffalo, *Syncerus caffer*, disease,
 177
- African elephant, *Loxodonta africana*,
 dental disease, 170
- Age determination, 47–48
- Age of first breeding, in reindeer and
 caribou, 67, 194
 in ungulates, 193
- Age structure, 47–48, 152, 157–159
 in Barff herd, 48, 158
 in Busen herd, 48
 in Royal Bay herd, 48
- Alaska, caribou, 60, 61, 68
 introduction of reindeer, 24–27
 numbers, 15, 25
 range occupied, 25
- Alaskan caribou, calving season, 60
 dental disease, 170
 diet, 121
 distribution, 5, 15
 irruptions, 16
 migration, 120
- mortality, 179–180
 numbers, 15
 taxonomic division, 4
- Alces alces*, *see* moose
- Aleutians, introduction of reindeer, 24, 26,
 185–186, 188
- Amsterdam Island, climate, 246
 effect of introduced mammals, 264–265
 endemic species, 246
 human impact, 246
 introduced mammals, 246, 251–252, 254,
 260
 natural history, 246–247
 sovereignty, 246
- Anticosti Island, introduction of reindeer,
 24, 26
- Antipodes Islands, climate, 246
 endemic species, 246
 human impact, 246
 introduced mammals, 30, 246, 251–252
 natural history, 246
 sovereignty, 246
- Antler cycle, 83–88, 154–155, 195
- Antler growth, 83, 86, 88, 124–130
- Antler pedicle, in red deer, 83–85
 in reindeer, 83–87
- Antler shape, 3, 4, 7, 90, 123–127
 compressicornis group, 4
 cylindricornis group, 4
- Antler size, 128–130, 149, 195–197
 relationship with shoulder height in male
 cervids, 6
- Artiodactyl, 2
- Atka Island, *see* Aleutians
- Auckland Islands, climate, 246
 effect of introduced mammals, 261,
 264–265
 endemic species, 246
 human impact, 246
 introduced mammals, 30, 246, 251–252

308 *Index*

- Auckland Islands (*cont.*)
 natural history, 246–247
 sovereignty, 246
- Baffin Island, introduction of reindeer, 24, 26, 185–187
- Barff herd, age structure, 48, 158
 antler size, 123–130, 195–197
 areas occupied, 33, 49–50
 body size, 123–149, 195
 culling, 47–48
 density, 50–51, 189, 202, 206
 dental disease, 167–178, 206
 diet, 103–119, 198
 exclusion experiment, 226, 240
 historical summary, 43
 introduction, 42
 mortality, 150–162, 206–207
 overgrazing, 219–221, 223–224, 258
 pregnancy rates, 66, 194
 rates of decline, 191, 203, 258
 rates of increase, 190
 rumen contents, 103–115, 198–200
- Barff Peninsula, area occupied by reindeer, 33, 49–50
 flora, 95–96, 199
 overgrazing, 219–220
 snow depths, 97
 soil erosion, 216–217
 tussock grassland, 222–223
- Barren-ground caribou, age of first breeding, 73
 calving season, 60
 dental disease, 167, 170
 diet, 121, 203
 distribution, 5, 15
 ecology, 11, 179–180
 fat reserves, 144, 204
 gestation period, 61
 luteal scars, 74
 migration, 9, 120
 mortality, 179–189, 204
 numbers, 15
 ovaries, 71–72
 parasites, 165
 pregnancy rates, 68, 73–74, 214
 range, 9, 213
 rates of increase, 213
 reproductive failure, 75
 snow effects, 98
 taxonomic division, 4
- Black rat, *Rattus rattus*, introduced to southern islands, 250–251
- Black-tailed deer, 7
 breeding group size, 8
 calving season, 88
 feeding ecology, 119
 growth patterns, 147
 habitat, 8
 origin, 8
 pregnancy rates, 89, 193
 seminiferous tubules, 88
 sexual dimorphism, 181
 testis weights, 89
- Body condition, 203–205
 effect of disease, 177–179, 205
 effect of photoperiod, 255–256
 effect of fertility, 145–147, 194
 fat reserves, 140–147, 157, 159–160, 177, 203–205, 211
 serum protein, 141–143, 145
- Body growth, 194–198
 in relation to pregnancy rates, 194
 seasonal patterns, 123, 124, 134–140, 147–149, 203–204
- Body size, 129, 133, 151, 195–197, 203–205
 changes, 134–140, 146–149
 effect of disease, 177–179
 effect on reproductive success, 149
 relationship with antler size, 127, 129–130
 sexual dimorphism, 123–149, 181
- Bouvetøya, climate, 246
 endemic species, 246
 human impact, 246
 introduced mammals, 246
 natural history, 246
 sovereignty, 246
- Breeding group size, 6–7, 8, 89–90
- British Antarctic Survey, 40, 45
- Brown hare, *Lepus europaeus*, introduced to southern islands, 30, 251–252
- Brown rat, *Rattus norvegicus*, introduced to South Georgia, 40
 introduced to southern islands, 250–252
- Bryophytes, exclusion experiment, 231–233, 236
 in plant communities, 231–233
 nitrogen content, 113
 nutrient dynamics, 97
 on Barff Peninsula, 96
 on South Georgia, 33, 93, 198, 227, 231–233, 236
 seasonal use as forage, 107–108, 110–113, 121, 198
- Busen herd, age structure, 48
 areas occupied, 33, 49–51
 body size, 195
 counts, 43, 45–47, 51, 188, 206
 culling, 47–48
 density, 50–51, 188–189, 202, 206
 dental disease, 170, 178, 206
 diet, 198

Index

309

- exclusion experiment, 226, 240
 historical summary, 43
 mortality, 156, 206
 overgrazing, 219–221, 223–224, 258
 pregnancy rates, 66
 rate of increase, 190
 rate of decline, 258
 rumen contents, 199–200
 tussock grassland, 222–223
- Calving season, of mountain sheep, 62
 of reindeer and caribou, 57, 60, 52–63, 88–89
 of ungulates, 88
- Campbell Island, climate, 246
 effect of introduced mammals, 261, 264–265, 267–269
 endemic species, 246
 human impact, 246
 introduced mammals, 30, 246, 251–253
 natural history, 246–247
 sovereignty, 246
- Canada, barren-ground caribou, 60, 68, 203–204
 introduced reindeer, 24, 25–27, 60, 61
 numbers, 15
 parasitic diseases in reindeer and caribou, 165–166
- Peary caribou, 68
 predators, 179
 woodland caribou, 60, 68, 165
- Capreolus capreolus*, *see* roe deer
- Carrying capacity, 20, 21, 188–190, 193, 201, 225
- Cat, *Felis catus*, as predators, 262–263, 265–266
 births, 256
 effect on ecosystems, 265–266
 food preferences, 259–261
 introduced to South Georgia, 40
 introduction to southern islands, 30–31, 249, 251–253, 255–261
- Cattle, *Bos taurus*, effect on ecosystem, 264
 food preference, 259–260
 introduced to South Georgia, 40
 introduced to southern islands, 30, 251–253, 259–260
 parasitic disease, 164
- Cave art, 12–13
- Cephenomyia trompe*, *see* warble flies
- Cervus elaphus*, *see* red deer
- C. e. canadensis*, *see* wapiti
- China, herded reindeer, 15, 60
- Chorisodontium aciphyllum*, on South Georgia, 93
- CL, *see* corpus luteum
- Cladonia*, 10, 92
 on South Georgia, 95, 227
- Climate, effect on introduced reindeer, 207–209
 of Arctic regions, 7, 9
 of South Georgia, 32–36, 97–101, 208, 210, 246
 of southern islands, 245–247
- Compressicornis group, 4
- Cook Glacier, 45, 46
- Corpus luteum, description, 63, 66, 71–74
 diameter, 70
- Crash die-off, 217–218
 Iceland, 191–192, 203
 Nunivak Island, 191
 St Matthew Island, 29, 192, 207–209
 St Paul Island, 29, 191
- Culling, of reindeer, 47
- Cylindricornis group, 4
- Dama dama*, *see* fallow deer
- Density, affecting mortality, 150–151, 162–163, 205–206
 effect on reindeer body size, 197–198
 of reindeer, 51, 188–191, 194, 224–225
 on South Georgia, 49–51, 162–163, 188–191, 200–203
- Dental disease, effects, 177–179
 eruption of teeth, 167–168, 173–175
 in Barff herd, 167
 in Busen herd, 170
 in cervidae, 167
 in reindeer, 167–179
 in Royal Bay herd, 170
 mandibular swellings, 167–181, 205–207, 257
 molar damage, 167–169
 osteoporosis, 171–176
- Deschampsia antarctica*, effects of reindeer grazing, 218, 224, 229–230
 exclusion experiment, 229–230, 232–239
 nitrogen content, 113
 nutrient dynamics, 96–97
 on Barff Peninsula, 96
 on South Georgia, 93–95, 198–200
 propagation, 238
 seasonal use as forage, 107–113, 198
- Diet, 99–121
 influence on dental disease, 170–171
 nitrogen content, 112–113
 seasonal changes, 103–115, 119, 198–200
- Disease, as cause of mortality, 150–151, 162–181, 205
 as a limiting factor, 150, 177–181, 205

310 *Index*

- Distribution, sub-species of *Rangifer tarandus*, 5, 15
- Dog, *Canis familiaris*, introduced to South Georgia, 40
introduced to southern islands, 30, 251–253
- Domestication, of caribou, 14
of reindeer, 13
- Donkey, *Equus* spp., introduced to southern islands, 30, 251–253
- Dry meadow, exclusion experiment, 226, 228–235
on Barff Peninsula, 96
on South Georgia, 93–94, 98–99, 101–102, 104, 224–239
overgrazing, 218, 224
seasonal use as forage, 101–102, 104
snow depth and hardness on, 98–99
- Dwarf-shrub sward, exclusion experiment, 226, 231–235
on Barff Peninsula, 96
on South Georgia, 93, 226
overgrazing, 218–221, 224
propagation, 238
- East Greenland caribou, extinction, 4, 11, 179
taxonomic division, 4
- Ecology, differences in mortality, 179–181
impact from introduced mammals, 30–31, 261–270
- Elaphostrongylus rangiferi*, *see* nematode
- Elephant seal, *Mirounga leonina*, industry, 40
on South Georgia, 37, 227
- Elk, 7
- Emigration, influence on declines of reindeer, 202, 207–281
of Royal Bay herd, 189, 191
- Endemic species, 244
of southern islands, 246–247, 250, 264–265, 267
- Environment, of South Georgia, 32–37
- Epididymis, in reindeer and caribou, 77–82
- Eruptions in reindeer populations, *see* eruptions
- Establishment of introduction, 20–21
of mammals on southern islands, 255–261
of reindeer on Arctic islands, 185–193
- Eurasia, dental diseases in reindeer, 171
domestication of reindeer, 13
elk (moose), 7
fallow deer, 7
herded reindeer, 15, 60
herding practices, 13
parasites, 163
roe deer, 7
saiga, 6
- Eutrophic mire, exclusion experiment, 226, 231, 234–235
on Barff Peninsula, 96
on South Georgia, 93–94, 99, 101, 104
seasonal use as forage, 101, 104
snow depth and hardness on, 99
- Exclusion experiment, on Campbell Island, 267–268
on South Georgia, 225–243, 269
- Extinction, of East Greenland caribou, 4, 11, 179
of native avifaunas, 263–266, 270
of plant species, 241, 263–267, 270
of Queen Charlotte Island caribou, 4
- Failed introductions, of mammals, 19, 31, 245, 252–255
of reindeer, 23–24, 42, 186–187, 209
- Falkland Islands, climate, 246
effects of introduced mammals, 264–265
endemic species, 246
human impact, 246, 250, 263
introduced mammals, 30, 246, 251–253
natural history, 246
sovereignty, 246
- Falkland fox, *Dusicyon australis*, 247
- Fallow deer, 7
breeding group size, 8
calving season, 88
dental disease, 167, 171
feeding ecology, 119
growth patterns, 147
habitat, 8
origin, 9
reproductive changes in male, 82
seminiferous tubules, 88
testis weights, 89
- Fat reserves, 140–148, 157, 159–160, 177, 203–205, 211
- Feeding craters, 99, 128
- Fellfield, on Barff Peninsula, 96
on South Georgia, 38, 94, 99, 101, 103–104
on southern islands, 247
overgrazing, 218, 224
seasonal use as forage, 101, 103–104
snow depth and hardness on, 99
- Fennoscandinavia, herding practices, 13
- Festuca contracta*, exclusion experiment, 228–229, 232, 235–239
nitrogen content, 113
nutrient dynamics, 96–97
on Barff Peninsula, 96

Index

311

- on South Georgia, 39, 93–94, 102, 198–200
- propagation, 238
- seasonal use as forage, 102, 107–108, 110–113, 198
- Finland, forest reindeer, 60
- herded reindeer, 60
- numbers, 15
- Forage availability, 19, 21, 29, 147–148
 - as cause of mortality, 150, 160, 179, 184, 187, 193
 - for reindeer and caribou, 92–103, 119–121, 198–200, 208–209, 212–213, 215–243
- Forage selection, by introduced mammals, 258–261
 - by reindeer and caribou, 110–115, 120–122, 198–200, 240, 259–260
- Forbs, 92, 227
 - forage for caribou, 121
 - in plant communities, 231–233
 - on Barff Peninsula, 95–96
 - propagation, 238
 - reindeer exclusion experiment, 231–233, 237–238
- Forest reindeer, calving season, 60
 - distribution, 5, 13, 15
 - mortality, 179–180
 - numbers, 15
 - taxonomic division, 4
- Founder population, of mice, 256
 - of ungulates, 256–257
- Fur seal, *Arctocephalus* spp., on South Georgia, 37
 - on southern islands, 250
- Galapagos Islands, introduction of mammals, 244
- Geology, of South Georgia, 32, 33
- George River herd, irruption, 16
- Gestation period, 54, 61–62
- Goat, *Capra hircus*, dental disease, 170
 - effect on ecosystems, 264
 - food preferences, 259–260
 - introduced to South Georgia, 40
 - introduced to southern islands, 30, 251–253, 259–260
- Golden eagle, *Aquila chrysaetos*, 11
- Gough Island, climate, 246
 - effect of introduced mammals, 265
 - endemic species, 246
 - human impact, 246
 - introduction of mammals, 30, 246, 251–254
 - natural history, 246–247
 - sovereignty, 246
- Grasses, forage for reindeer and caribou, 92, 121
 - on Barff Peninsula, 96
 - propagation, 238
- Great Cloche Island, introduction of reindeer, 24, 26, 187
- Greenland, antler casting, 87
 - barren-ground caribou, 11, 60
 - extirpation of caribou in NW, 17
 - introduction of reindeer, 24, 26, 185–187
 - muskox, 6
 - numbers, 15
 - predators, 179
 - reindeer density, 51
 - warble flies, 163
- Growth, antler, 124–130
 - seasonal patterns in body, 123–124, 134–140
 - skeletal, 130–134
- Grytviken, 40
 - climate, 210
 - plant communities, 93–94
 - snow depths, 98
 - whaling station, 37
- Guanaco, *Lama guanicoe*, introduced to southern islands, 30, 251–253
- Hare, *see* brown hare
- Heard and McDonald Islands, climate, 246–247
 - endemic species, 246
 - human impact, 246
 - introduced mammals, 246, 252–253
 - natural history, 246
 - sovereignty, 246
- Herbivores, damage to areas of vegetation, 31, 215, 219
 - introduction to southern islands, 30, 244–255
- Herded reindeer, calving season, 60
 - gestation periods, 61
 - in Alaska, 27
 - in Canada, 27
 - in Eurasia, 13, 60, 61, 165
 - in Fennoscandinavia, 13
 - in Norway, 14, 68
 - in Sweden, 68
 - mortality, 156
 - numbers, 15
 - parasitic disease, 165
 - pregnancy rates, 68
 - reintroduction, 185
- Himalayan tahr, *see* Himalayan thar
- Himalayan thar, *Hemitragus jemlahicus*
 - introduction to New Zealand, 22
 - irruption of, 22–23, 193, 216

312 *Index*

- House mouse, *Mus musculus*, births, 256
 effect on ecosystems, 266
 food preferences, 259–260
 impact on environment, 263
 introduced to South Georgia, 40
 introduced to southern islands, 30, 250–251
- Horse, *Equus caballus*, introduced to South Georgia, 40
 introduced to southern islands, 30, 251–253
- Hound Bay, area occupied by reindeer, 49
 snow depths, 99, 101
 snow hardness, 99, 101
- Hunting, 180, 200, 211
 of caribou, 14
 of predators, 184
 of reindeer on South Georgia, 45, 150, 152
- Husvik Harbour, 42
- Iceland, calving season, 63
 decline, 209
 diet, 202
 distribution, 191, 207–208
 introduction, 24, 26, 27, 185–187
- Iles Amsterdam, introduced mammals, 30
- Iles Crozet, climate, 246
 effect of introduced mammals, 264–265
 endemic species, 246
 human impact, 246
 introduced mammals, 30, 246, 248, 251–254, 256, 258–260
 natural history, 246
 sovereignty, 246
- Iles Kerguelen, climate, 246
 effects of introduced mammals, 264–265
 endemic species, 246
 human impact, 246
 introduced mammals, 30, 246, 251–254, 256–260
 introduced reindeer, 26, 30–31, 186, 225
 natural history, 246
 sovereignty, 246
- Iles St Paul, climate, 246
 endemic species, 246
 human impact, 246
 introduced mammals, 30, 246, 251–254
 natural history, 246
 sovereignty, 246
- Inbreeding, in ungulates, 150, 170
- Industry, on South Georgia, 37, 40
- Infertility, 67
- Introduced mammals, accidental, 250–251, 253
 deliberate, 250–251
 establishment of, 255–261
 failure of, 19, 31, 245, 252–255
 irruptions of, 19–24
 on South Georgia, 40, 183
 on southern islands, 30–31, 183, 224–225, 244–270
 success of, 186–187, 209, 245, 252–255
- Introduced reindeer, body growth, 194–198, 203–205
 calving seasons, 60, 62
 densities, 200–201
 diet, 198–200, 202–203
 emigration, 207–209
 failure of, 23–24, 42, 186–187, 209
 gestation periods, 61
 in continental areas, 24–27, 51, 60
 irruptions of, 24–31, 183–214
 mortality, 205–209
 on islands, 12, 24, 51, 60, 185–209
 on St Paul, 23–24, 51
 on St Matthew, 23–24, 51
 on South Georgia, 12, 40–52, 41, 183
 parasites, 165
 pregnancy rates, 68, 194
 rates of decline, 201–202
 success of, 186–187, 209
- Invertebrates, 36–37, 215, 247, 259–260, 266–267
- Irruptions, definition of, 17–18
 in barren-ground caribou, 16
 in established populations, 16, 17, 183
 in Iceland, 191, 202
 in natural reindeer, 184–185, 210–214
 in tundra reindeer, 16
 in woodland caribou, 16
 of introduced mammals, 19–24
 of introduced reindeer, 24–31, 50–51, 183–214
 stages of, 19–23, 29, 50–51, 194–198, 216–219, 223, 240–241, 257–258
- Isla Navarino, introduced mammals, 30–31
 introduced reindeer, 187
- Island classification, 27, 185–187, 245–247
- Island ecosystems, effects from introduced mammals, 245, 261–270
 of St Matthew Island, 199
 of South Georgia, 33, 36, 37, 92–104, 162–163
 of southern islands, 215, 244–247, 261–270
- Japan, introduction of reindeer, 26
- Juncus scheuchzerioides*, exclusion experiment, 231–232, 238
 in plant communities, 231–232
 nitrogen content, 113

Index

313

- on Barff Peninsula, 96
- on South Georgia, 93–94, 198–200, 231–232, 238
- propagation, 238
- seasonal use as forage, 107–111, 113, 198–200
- Kerguelen Island, climate, 246
- effect of introduced mammals on ecosystems, 264–266
- endemic species, 246
- human impact, 246
- introduction of mammals, 30, 246, 251–254, 256–260
- introduction of reindeer, 26, 30–31, 186, 225
- natural history, 246
- sovereignty, 246
- Kuriles, introduction of reindeer, 24, 26
- Lactation, 145–148
- Leith Harbour, 42
- Lichen, 9, 10, 11, 92, 196
- effect of introduced mammals, 264
- exclusion experiment, 234, 237
- influence on reindeer ecology, 12, 119, 187, 215
- on Barff Peninsula, 96
- on South Georgia, 33, 39, 93–95, 102–103, 109
- overgrazing, 218, 224, 240–241
- seasonal use as forage, 102–103, 107–110, 120–122, 202–203, 210, 259
- Lungworm, 164
- Luteal scars, 65–66, 73–74, 146–147, 194
- Lynx, *Felis lynx*, 11, 180
- Mackenzie Delta, comparison of introduced and native caribou, 62
- introduction of reindeer, 24–27
- Macquarie Island, climate, 246
- effect of introduced mammals on ecosystems, 262–265
- endemic species, 246
- human impact, 246
- introduced mammals, 30, 246, 249, 251–254, 256, 259–260
- natural history, 246
- soil erosion, 219
- sovereignty, 246
- Man, impact on southern islands, 246, 253
- influence on ecosystems, 244–245
- influence on reindeer ecology, 12–16, 180, 210–211, 258
- Marion and Prince Edward Islands, climate, 246
- effect of introduced mammals on ecosystems, 263–266, 269
- endemic species, 246
- human impact, 246
- introduced mammals, 30, 246, 251–254
- natural history, 246
- sovereignty, 246
- Mediterranean area, fallow deer, 7
- Mesic meadow, exclusion experiment, 226–235
- on Barff Peninsula, 96
- on South Georgia, 93–95, 99, 101, 103–104
- overgrazing, 218
- seasonal use as forage, 101, 103–104, 110
- snow depths and hardness on, 99
- Michigan, introduction of reindeer, 25
- Migration, 9, 11, 13, 89, 92, 120, 210
- Mink, *Mustela vison*, introduced to southern islands, 252–253
- Moose, 7
- breeding group size, 8
- calving season, 88
- diseases, 170, 177
- feeding ecology, 119
- growth patterns, 147
- habitat, 8
- mortality, 180
- origin, 8
- pregnancy rates, 89, 193
- seminiferous tubules, 88
- testis weights, 89
- Mortality, causes, 150–151, 157–181, 204–205
- differential, 151–153, 159, 180–181, 205–207
- patterns, 151–156, 162, 205
- perinatal, 153–154, 156–157, 205–207
- rates, 150–151, 156–162, 201–202, 205, 207, 211
- seasonal, 154–155
- Mossbank, exclusion experiment, 226, 229–237, 242
- on Barff Peninsula, 96
- on South Georgia, 93–95, 98–99, 101, 103–104
- overgrazing, 218
- seasonal use as forage, 101, 103–104
- snow depth and hardness on, 98–99
- Mouflon, *Ovis ammon*, births, 256
- food preferences, 259
- introduced to southern islands, 30, 251–253, 256, 259
- Mountain or tundra reindeer, calving season, 60
- diet, 105

314 *Index*

- Mountain or tundra reindeer (*cont.*)
 distribution, 5, 13, 15
 ecology, 11
 irruptions, 16
 migration, 120
 mortality, 179
 numbers, 15
 predation by man, 14
 pregnancy rates, 68
 snow effects, 98
 taxonomic division, 4
- Mountain sheep, *Ovis canadensis*, 6
 feeding ecology, 119
 lambing season, 62
- Mouse, *see* house mouse
- Mule deer, *see* black-tailed deer
- Mushrooms, forage for reindeer and caribou, 121
- Muskox, *Ovibos moschatus*, 6
 feeding ecology, 119
- Navarino Island, introduction of reindeer, 26
- Nelchina herd, caribou density, 51
 irruption, 16
- Nematode, 163–167
 protostrongylid, 163, 180, 187
 strongylate, 164–167, 180
- New Fortuna Bay, introduction of reindeer, 42
- Newfoundland, introduction of reindeer, 24, 26, 185
 rutting activities, 57
 woodland caribou, 60
- New Zealand, effect of introduced mammals on ecosystems, 263
 introduction of Himalayan thar, 22–23, 216
 irruption of introduced mammals, 19, 21, 30, 240–241
 red deer, 55
 reversal of breeding season in introduced ungulates, 55, 62
 soil erosion, 219
- Nitrogen balance, 115, 137
 in captive deer, 147
- Nitrogen content, of forage, 112–114, 120, 199
 of rumen, 116–119, 148–149, 198, 200
- North America, dental diseases in caribou, 171
 habitat of caribou, 11
 hunting of caribou, 14
 irruptions of introduced mammals, 240
 moose, 7
 mountain sheep, 6
 mule deer, 7
 muskox, 6
 parasites, 163
 white-tailed deer, 7
- Norway, climate, 36, 98
 dental disease, 174
 herded reindeer, 14, 68, 165
 numbers, 15
 parasitic diseases, 165
 reindeer density, 51
 wild reindeer, 60, 68
- Norway rat, *Rattus norvegicus*, introduced to southern islands, 250–251
- Novosibirsky Islands, tundra reindeer, 11, 179
- Numbers, herded reindeer, 15
 wild reindeer and caribou, 15
- Nunivak Island, introduction of reindeer, 24–28, 185, 191
 overgrazing, 207, 209
 reindeer density, 51
- Nutritional limitations, in reindeer and caribou, 115–119
- Ocean Harbour, area occupied by reindeer, 49
 introduction of reindeer, 42
 snow depths, 99, 101
 snow hardness, 99, 101
- Odocoileus hemionus*, *see* black-tailed deer
- Odocoileus virginianus*, *see* white-tailed deer
- Oedemagena tarandi*, *see* warble flies
- Oestrous cycles, 54, 57, 59
 in red deer, 55
- Oligotrophic mire, exclusion experiment, 226, 231–235
 on Barff Peninsula, 96
 on South Georgia, 38, 93–94, 99, 101, 103–104, 226
 seasonal use as forage, 101, 103–104
 snow depths and hardness on, 99
- Origin, of cervidae, 5–6, 8
 of *Rangifer*, 5–6
- Ostertagiasis, 164–166
- Organ weights, 134–140, 148–149, 195
- Otter, *Lutra* spp., introduced to southern islands, 252
- Overgrazing, 201, 203
 definition, 215
 in Iceland, 208
 of lichens, 29, 203
 of tussock grassland, 202
 on St Matthew Island, 29, 209
 on South Georgia, 45, 94, 96, 162, 183, 193, 218–225

Index

315

- Ovary, changes in yearling and adult females, 71
description, 63–65
of barren-ground caribou, 71–72
productivity, 71–74
removal, 87
weights, 69–70, 72
- Parasite burden, as cause of mortality, 180–181, 187
effects of, 177, 180–181
lungworms, 164
nematodes, 163–167
on South Georgia, 162–167
ostertagiasis, 164–166
seasonal patterns, 166
warble flies, 163
- Peary caribou, calving season, 60
diet, 121
distribution, 5, 15
ecology, 11
migration, 120
mortality, 179
numbers, 15
pregnancy rates, 67, 68
range, 9
relationship with Spitzbergen reindeer, 4, 67
taxonomic division, 4
- Perissodactyl, 2
- Phleum alpinum*, exclusion experiment, 233, 235–236, 238
nitrogen content, 113
on Barff Peninsula, 96
on South Georgia, 39, 94, 198–200, 233, 235–236, 239
propagation, 238
seasonal use as forage, 107–108, 110–113, 198
- Photoperiod, effects on breeding season, 43–45, 54–55, 255
effects on moult, 44
effects on seasonal inappetance, 147–148
- Pig, *Sus scrofa*, effect on ecosystems, 264–266
food preferences, 259–260
introduced to South Georgia, 40
introduced to southern islands, 251–253, 258–260
- Plant communities, on South Georgia, 33, 93–104, 201, 215–243, 246–247
on southern islands, 261–270
- Pleistocene fauna, 5
- Pneumostrogylus tenuis*, see nematode
- Poa annua*, effects of introduced mammals, 264
effects of overgrazing, 218, 224, 240
exclusion experiment, 229–230, 232, 235–240
nitrogen content, 113
on Barff Peninsula, 96, 240
on South Georgia, 94–97, 198–200, 219–221
propagation, 238
seasonal use as forage, 107–113, 198
use as forage on southern islands, 259, 261–263
- Poa flabellata*, effects of introduced mammals, 261, 264–265
effects of grazing, 218, 261
exclusion experiment, 228, 232, 235–240
exposure above snow, 99
nitrogen content, 112–113
nutrient dynamics, 96–97, 112–115
propagation, 238
on Barff Peninsula, 96, 240
on South Georgia, 33, 36, 38, 93–97, 99, 198–200, 202, 259
seasonal use as forage, 107–115, 118, 120
- Polygyny, 7, 90, 149
- Polytrichum alpestre*, on Barff Peninsula, 96
lack of on South Georgia, 37, 150
- Predation, 89, 128, 130, 184, 200–201, 211–212
as cause of mortality, 150–151, 179–180, 211–212
as limiting factor for population size, 11–12, 152, 156, 192, 193, 210–213, 263
in South Georgia, 37
of wildebeest calves, 89
- Predators, bear, 11, 180
cat, 262–263, 265–266
golden eagle, 11
lynx, 11, 180, 211
man, 14, 180, 210–211
rodents, 263, 265–266
weka, 265
wolf, 11, 12, 180, 211–212
wolverine, 11, 180
- Pregnancy rates, 65–69, 73–75, 145–147, 150, 193–194, 210–211, 214
- Pribilof Islands, introduction of reindeer, 24, 26, 27, 185–186
see also St George Island, St Paul Island
- Prolactin, 87
in cervidae, 147
- Propagation, of South Georgian vascular plants, 238
- Protostrongylid nematode, 163
cause of mortality, 180, 187

316 *Index*

- Pseudocyphellaria*, on South Georgia, 39, 95
- Pteridophytes, on Barff Peninsula, 95–96
on South Georgia, 33
- Queen Charlotte Island caribou, extinction of, 4
taxonomic division, 4
- Rabbit, *Oryctolagus cuniculus*, as prey, 262
births, 256
effect on ecosystems, 264–267
food preferences, 259–260
introduced to South Georgia, 40
introduced to southern islands, 30, 248, 250–254, 256–260, 267
myxomatosis, 267
- Raised beaches, on South Georgia, 95, 219, 222–223
plant communities, 95, 219, 222–223
tussock grassland, 222–223
- Rangifer tarandus caribou*, see woodland caribou
- R. t. dawsoni*, see Queen Charlotte Island caribou
- R. t. fennicus*, see forest reindeer
- R. t. granti*, see Alaskan caribou
- R. t. groenlandicus*, see barren-ground caribou
- R. t. oegroenlandicus*, see East Greenland caribou
- R. t. pearyi*, see Peary caribou
- R. t. platyrhynchus*, see Spitzbergen reindeer
- R. t. tarandus*, see mountain reindeer
- Rat, as a predator, 263, 265–266
effect on ecosystems, 265–266
introduced to South Georgia, 40
introduced to southern islands, 30, 250–251
- Rate of decline, of introduced reindeer, 190–193
- Rate of increase, of introduced reindeer, 189–194, 197, 199
- Red deer, 7
antler cycle, 88
antler pedicle formation, 83–85
breeding group size, 8
calving season, 88
dental disease, 167
feeding ecology, 119
growth patterns, 147
habitat, 8
mortality, 153
nutrition, 117
origin, 8
pregnancy rates, 89
reproductive changes in males, 82–83
reversal in breeding season, 55, 62
rutting behaviour, 81
seminiferous tubules, 88
sexual dimorphism, 181
testis weights, 89
- Reproduction, 54–91
age of first breeding, 67, 193–194, 197
calving season, 57, 59–60, 62–63, 88–89
changes in females, 63–75
changes in males, 76–82
corpus luteum, 63–66, 70–74
epididymis, 77–82
failure, 75
gestation period, 54, 61–62
infertility, 67, 69
luteal scars, 65–66, 73–74, 194
oestrous cycles, 54, 57, 59
ovary, 63–65, 69–74, 87, 194
pregnancy rates, 65–69, 73–75, 193–194, 210–211, 214
prolactin, 87
puberty, 77, 81
reversal of breeding season, 55–63, 255
rutting behaviour, 55–58
seminiferous tubules, 76–79, 81, 88
sexual characters, 81, 83
sexual maturity, 69–71, 77–81
testis, 76–83, 87–90
testosterone, 77–79, 81, 88
- Reproductive changes, in females, 63–75
in males, 76–82
- Reproductive failure, 75
- Reproductive potential, 89–90
- Reversal of breeding season, in reindeer, 43–44, 55–63
in red deer, 55
- Ridgetops, plant communities, 95, 219, 222–223
on South Georgia, 95, 219, 222–223
tussock grassland, 219, 222–223
- Rodents, as predators, 263, 265–266
breeding, 255–256
effect on ecosystems, 265–266
food preferences, 259–260
introduced to South Georgia, 40
introduced to southern islands, 250–251, 255–256, 259–260
- Roe deer, 7
body size, 197–198
breeding group size, 8
calving season, 88
feeding ecology, 119

Index

317

- growth pattern, 147
- habitat, 8
- mortality, 180
- origin, 8
- pregnancy rates, 89, 193
- semiferous tubules, 88
- sexual dimorphism, 181
- testis weights, 89
- Rostkovia magellanica*, exclusion
 - experiment, 233
 - in plant communities, 233
 - nitrogen content, 113
 - nutrient dynamics, 97
 - on Barff Peninsula, 96
 - on South Georgia, 38, 93–97, 198–200, 233
 - propagation, 238
 - seasonal use as forage, 107–113, 198
- Royal Bay herd, age structure, 48
 - antler size, 195–197
 - areas occupied, 33, 49–50
 - body size, 195–197
 - counts, 45–47, 50, 188, 206
 - culling, 47–48
 - density, 50–51, 189, 201, 206
 - dental disease, 170, 178, 206
 - diet, 198
 - emigration, 189, 191
 - exclusion experiment, 226, 240
 - mortality, 156–157, 206
 - organ weights, 195
 - overgrazing, 50, 216–217, 219–221, 223–224
 - pregnancy rates, 66, 194
 - rumen content, 198–199
 - tussock grassland, 222–223
- Rumen contents, 103, 105–110, 115–119, 176, 198–200
- Ruminantia, 2
- Rushes, forage for caribou, 121
 - on Barff Peninsula, 95–96
 - propagation, 238
 - reindeer exclusion experiment, 237
- Rutting behaviour, 55–58, 90, 148–149
- Saiga, *Saiga tatarica*, 6
- St George Island, introduction of reindeer, 26–28, 188–189, 191
- St Lawrence Island, introduction of reindeer, 24, 26–28, 188–191
 - overgrazing, 207, 209
- St Matthew Island, crash die-off, 192, 207–209
 - ecosystem, 199, 202–203
 - grazing by reindeer, 240–241
 - introduction of reindeer, 23–24, 26–29, 185
 - reindeer density, 51, 188–189
 - reindeer numbers, 28, 188–190, 192, 194–195, 207
- St Paul Island, ecosystem, 202–203
 - grazing, 207, 240
 - introduction of reindeer, 23–24, 26–28
 - reindeer density, 28, 51, 188–189
 - reindeer numbers, 28, 188–191
- Scotland, introduction of reindeer, 24, 26, 60, 61, 185
- Seabirds, effects of introduced mammals, 265–266, 268
 - on South Georgia, 37, 265
- Seals, on South Georgia, 37, 227
- Sedges, 92
 - exclusion experiment, 237
 - forage for caribou, 121
 - in plant communities, 231
 - on Barff Peninsula, 95–96
 - on South Georgia, 199, 231
- Semiferous tubules, 76–79, 82, 88
- Serum protein, 141–143, 145
- Sex differences and dimorphism, 53, 89–90, 92–93, 151, 179–181
 - antler cycle, 85–87, 88, 123–130, 195–197
 - body growth, 135–140, 149, 195–197
 - body weights, 135–136, 149, 181
 - fat reserves, 141–145, 149
 - mortality, 151–153, 159, 205
 - organ weights, 135–140
 - reproductive phenology, 148
 - rumen contents, 117–119
 - skeletal growth, 130–134
- Secondary sexual characters, in males, 81, 83
- Sexual maturity, in females, 69–71, 194
 - in males, 77–81
- Seychelle Islands, introduction of mammals, 244
- Sheep, *Ovis aries*, births, 256
 - dental disease, 174, 178
 - effect on ecosystems, 264–265, 268
 - exclusion experiment, 268
 - food preferences, 259
 - introduced to South Georgia, 40, 242
 - introduced to southern islands, 30, 251–253, 256–259
- Shinshiru Island, *see* Kuriles
- Shrubs, 92
- Silver fox, *Vulpes vulpes*, introduced to South Georgia, 40
 - introduced to southern islands, 30, 251–252
- Skeletal growth, 130–134, 195, 197

318 *Index*

- Skunk, *Canepatus* spp., introduced to southern islands, 252
- Slave River, introduction of reindeer, 24–25
- Snow, depths, 98–101, 119–120
hardness, 98–101
influence on reindeer ecology, 7, 9, 12
on South Georgia, 33, 93, 97–101, 105, 210, 220–221, 225, 246
on southern islands, 246
restricting forage availability, 92–93, 97–103, 105, 109, 179, 202, 208, 224
- Soil erosion, 241
Barff Peninsula, 216–217
on South Georgia, 218, 220–221
on southern islands, 218–221, 263
- Sörling Valley, area occupied by reindeer, 49
snow depths, 98, 99, 101
snow hardness, 99, 101
- South Georgia, climate, 32–36, 97–101, 208
endemic species, 246
ecosystems, 33–37, 92–104, 162–163, 238
environment, 32–37
geology, 32–33
history, 37, 40
human impact, 246
industry, 37, 40
introduction of reindeer, 12, 26, 30, 40–45, 183
introduction of mammals, 30, 40, 183, 246, 251–253
plant communities, 93–104, 201–203, 224–225
reindeer herds, 33, 43–52, 156
sovereignty, 40, 246
topography, 32, 34, 219–223
see also Grytviken
- Southampton Island, introduction of reindeer, 24, 26, 185
- Southern islands, characteristics, 245–250
climate, 245–247
effects of introduced mammals on ecosystems, 261–270
endemic species, 246–247, 250
floras, 215, 224–225, 246–247
human impact, 246
introduction of mammals, 30–31, 219, 244–270
overgrazing of, 216, 224
soil erosion, 219
- Sovereignty, of southern islands, 246
of South Georgia, 40, 246
- Spitzbergen reindeer, body condition, 140, 144
calving season, 60
distribution, 5, 15
irruption, 17
migration, 120
mortality, 152, 179–180
numbers, 15
parasites, 165–166, 180
predation by man, 14
pregnancy rates, 67–68
range, 9
relationship with Peary caribou, 4, 67
taxonomic division, 4
- Status, 15
- Stromness, Bay, 42, 49
whaling station, 35
- Strongylate nematode, 164–167, 180
- Starvation, 160, 180
- Successful introductions, of mammals, 245, 252–255
of reindeer, 25–26, 186–187, 209
- Svalbard, density, 51
mortality, 151–154, 160, 162
parasitic diseases, 164–166
predators, 179
- Rangifer*, 11
- Spitzbergen reindeer, 9, 17, 60, 68, 142, 167
- Sweden, herded reindeer, 60, 68, 165
numbers, 15
parasitic diseases, 165
- Taxonomy, 2–4
antler shape, 3, 4
- Tasmania, introduced mammals, 30
- Testis, 76–83, 87–90
- Testosterone, 77–79, 81–83, 87–88
- Tooth annulations, for age determination, 47–48
- Topography, influence on forage availability, 119–121
of South Georgia, 32, 34, 219–223
- Tortula robusta*, on Barff Peninsula, 96
on South Georgia, 93
- Tristan da Cunha Islands, climate, 246
effect of introduced mammals on ecosystems, 262–265
endemic species, 246
human impact, 246, 250
introduction of mammals, 30, 246, 251–252
natural history, 246–247
sovereignty, 246
- Tundra, Arctic, 4, 37, 92, 120, 199–201
subantarctic, 37, 92–96, 199–201, 247
- Tundra reindeer, *see* mountain reindeer

Index

319

- Tussock grassland, effect of introduced mammals, 264
 exclusion experiment, 226–235
 in Royal Bay, 216–217
 on Barff Peninsula, 96, 216–217
 on South Georgia, 33, 38, 45, 93–94, 98–101, 103–104, 162, 201–202
 on southern islands, 247, 261
 overgrazing, 218–223
 seasonal use as forage, 101, 103–104, 109–110, 112, 148, 162, 216, 225, 240–241, 260
 snow depths and hardness on, 98–100
- Ungulate, 2, 77, 92, 119, 128, 150, 157, 162–163, 170–171, 177, 183–184, 193, 201, 205, 240–241, 245, 252, 256, 258
 artiodactyl, 2
 perissodactyl, 2
- Unmah Island, *see* Aleutians
- Upland goose, *Chloephaga picta*, introduced to South Georgia, 40
- Usnea*, on South Georgia, 95
- USSR, introduction of reindeer, 24, 26, 62
 herded reindeer, 61, 65
 herding practices, 13
 numbers, 15
 parasitic diseases, 165
 tundra reindeer, 68
- Vegetation, influence of reindeer, 215–243
- Wapiti, 7
 breeding group size, 8
 calving season, 88
 dental diseases, 167
 feeding ecology, 119
 growth patterns, 147
 habitat, 8
 origin, 8
 pregnancy rates, 80
 seminiferous tubules, 88
 sexual dimorphism, 181
 testis weights, 89
- Warble flies, 163
- Warrah, *see* Falkland fox
- Waterbuck, *Kobus ellipsiprymnus*, dental disease, 170
- Weka, *Gallirallus australis*, effect on ecosystems, 265
- West Kola herd, density, 51
 irruption, 16
- Whaling stations, on South Georgia, 35–37, 41–42, 95
- White-tailed deer, 7
 breeding group size, 8
 calving season, 88
 disease, 171, 177, 181
 feeding ecology, 119
 growth patterns, 147
 habitat, 8
 origin, 8
 pregnancy rates, 89, 193
 seminiferous tubules, 88
 sexual dimorphism, 181
 testis weights, 89
- Wildebeest, *Connochaetes taurinus*, calf predation, 89
- Wolf, *Canis lupus*, 11, 12, 180
- Wolverine, *Gulo gulo*, 11, 180
- Woodland caribou, calving season, 60
 diet, 121
 distribution, 5, 15
 fat reserves, 144
 feeding ecology, 119–120
 gestation period, 61
 irruptions, 16
 mortality, 179–180
 numbers, 15
 parasites, 165
 pregnancy rates, 68
 taxonomic division, 4