

Index

Abbreviations: HAN = high-altitude native(s),
LAN = low-altitude native(s)

- abortion and miscarriage rates, in HAN: Andes, 90, 91, 324, 326, 327–8; Ethiopia, 91–2, 104, 327; Nepal, 327; Tajikistan, 91; Tien Shan and Pamirs, 311
- acclimatization, to altitude, 179–83
- ACTH: change of altitude and production of, 76; responses of HAN and LAN to injection of, 76
- aerobic capacity (ability to deliver oxygen to working muscles), 174–9; age and activity effects on, 10; as integrated measure of oxygen intake, transport, and utilization, 183; physiological factors related to, 183–5; upward migration and, 10, 179–83
- age: and aerobic capacity, 10, 175, 177; increased blood pressure with, less marked in HAN, 7, 304; at migration to high altitude, and aerobic capacity, 180, 181; vital capacity and residual volume of lung decrease with, in HAN, 302
- agriculture: in Andes, 117, 118, 219; in Ethiopia, 26–7, 138; in Himalaya, 35–6, 39, 147; vertical zonation of (Andes), 33–4
- air: insulating properties of, increase with decrease of density, 262
- alcohol: consumption of, and skin temperature of foot exposed to cold, 285, 291
- alkalosis, respiratory, at high altitude or in hypoxia, 86; in foetus, 94, 95; and sex ratio in mice, 103; on transfer to high altitude, 208, 209
- 'Altiplanides', 190, 199–200; irreversible specialization of? 212
- altiplano*, Andes, 20, 31, 34; ice-free by time of arrival of man, 49, 67
- alveolar ventilation, altitude and, 300, 301
- Amhara people, NW Ethiopia, 4, 7, 22, 26, 138, 195
- cAMP: lower level of, in HAN? 76
- anaemia: haematological findings may be affected by presence of, 196, 200; raised blood methaemoglobin in, 201; sickle-cell, 55, 58, 350
- Andean region, 18; agriculture and stock-raising in, 33–4, 117, 118, 219; date of arrival of man in, 49, 67, 96; genetic distances between HAN of, and neighbouring LAN, 53–4; genetic drift in, 57, 60; genetic polymorphisms in, 49–50; geology of, 20; history of, 28–31; inbreeding in, 61; migration in, 58, 60; population of, 31–2, (above 2,500 m), 318
- 'Andides', 190, 200–2
- anti-erythropoietin, inhibits polycythaemic response to hypoxia in mice, 193
- Argentina: estimated population of, above 2,500 m, 318
- arm tissues, Andean HAN; area of, 121–2; development of bone, marrow, muscle, and fat in, 122
- ascorbic acid: intake of, by Andean HAN, 239, 241
- athletics, HAN in, 179
- Aymara people, NE Chile and Bolivia, 4, 7, 57; birth rate of, 324; exogamy among, 61; genetic distances between other tribes and, 54; maturation and body size in two groups of boys of, 135, 136; possible evidence for selection pressure in white-cell antigens of, 57; proteinogram of, HAN *in situ* and after downward migration, 207
- babies (first year of life): body measurements, bone development, and psychomotor development of, in HAN and LAN, rural and urban, Mestizo and Indian, Andes, 104–5
- Baltis people, Karakoram, 60
- basal metabolic rate: altitude and, 229–30, 279, 280–1, 291; in LAN upward migrants, 279–80
- Bhutan, estimated population of, 319
- bicarbonate of blood, altitude and, 302
- birth rate: for HAN, Chile, 324; for HAN and LAN, Peru, 320
- birth spacing, HAN and LAN, Peru, 326
- birth weight: altitude and, 83, 92–4, 95–6, 106; altitude, and ratio of placenta weight to, 83–4, 87; of babies born at high and low altitudes to same mothers, 93, 95; in HAN and LAN, Andes, 88, 89; maternal stature 95; and neonatal mortality, 101–2; sex and, in HAN and LAN, 100; and survival, in HAN and LAN, 342, 346
- blood circulation time, altitude and, 303
- blood pressure, 334; altitude and, 284, 313; in HAN downward migrants, 337; increase of, with age, less marked in HAN, 7, 304, 335–6; in LAN upward migrants, 10
- body measurements: of Andean HAN and LAN, rural and urban, Mestizo and Indian, (babies) 104–5, (newborn) 99–100, (2 years old upwards) 119–27; of Ethiopian HAN and LAN, 140–4; of Tien Shan HAN and Kirov LAN, 156–9
- Bohr effect, of pH on affinity of haemoglobin for oxygen, 208; in Andean HAN and acclimatized Europeans, 10, 61, 185, 209–10; 2,3-diphosphoglycerate of red cells in, 201, 208–9; in HAN, 183; in LAN upward migrants, 208–9; in Sherpas and Europeans, 210–12
- Bolivia, 61; estimated population of, above 2,500 m, 318
- bone development: in arm, Andean HAN, 122; in metacarpal, Sherpa HAN and Tibetan children at low altitude, 152, 153
- bone marrow: altitude or hypoxia, and erythropoiesis in, 123, 191–5; of arm, development of, in Andean HAN, 122–3; hypoxia and size of, 123
- breast feeding: of children up to 2 or 3 years old, Andean HAN, 241; of infants in HAN and LAN, Andes, 105, 106
- calcium: absorption of, 242; intake of, by Andean HAN, 239, 241–2; ratio of, to phosphate in diet, 242

Index

- capillary blood vessels: in muscle of HAN, 183, 185, 304–5, 335; in placentae of HAN, 85
- carbohydrates: intake of, by Andean HAN, 236; percentage of calorie intake as, 237
- carbonic anhydrase, altitude and activity of, 302
- cardiac cycle: during exercise and recovery, in groups from different altitudes, 308–10; in HAN children, 312–13
- cardiac defects, in HAN newborn, 103
- cardiac index, altitude and, 303
- cardiovascular disease, altitude and, 334–5, 344
- Caucasoid ancestry: in Andes, very small, 55; in Ethiopia, 55; in Himalaya, 51, 54
- cell division, hypoxia and rate of, 70, 165, 208
- cereals, staple food up to 3,500 m in Andes (rice to 2,000 m, maize 2,000–3,500 m), 221, 222, 223
- cestode infestation, at different altitudes in Peru, 331
- chemoreceptor sensitivity to oxygen breathing, altitude and, 300, 302, 310
- chenopodiums (*quinoa* and *cañihua*), in diet of Andean HAN, 221, 222, 224
- chest: circumference of, Ethiopian HAN and LAN, 141, 147; circumference/height ratio, in Andean HAN at 4,000 m and above 4,500 m, 125, in Sherpa HAN and Tibetan downward migrant children, 150, 151, and in Tien Shan HAN and Kirov LAN, 158; enlargement of, genetically determined in Andean HAN, 8, 345; enlargement of, not general characteristic of Himalayan HAN, 166; width and depth of, Andean HAN, 119, and Tien Shan HAN and Kirov LAN, 157, 162–3
- Chile, 61; estimated population of, above 2,500 m, 318
- cholesterol, blood content of, 334; in Andean HAN, and HAN downward migrants, 337
- chorionic gonadotrophin, human, response to injection of: in HAN downward migrants, before and after return to high altitude, 75; in HAN and LAN, 76, 77
- chromosome abnormalities, none found in Andean HAN, 208
- climates, in high-altitude zones, 251; Andes, 33, 253–6; Ethiopia, 258; Himalaya, 34–5, 256–7
- clothes at high altitudes, 262–3, 290; Andes, 263–4; Himalaya, 264–5
- coca leaf chewing, by Andean HAN, 284; and response of body temperature to exposure to cold, 286–8, 291
- cold stress, in Andean HAN, 232, 266–73; influence of drugs on tolerance to, 284–8; interaction of, with hypoxic stress, 281–2; interaction of, with low-humidity stress, 288–9; laboratory studies of 274–9
- Colorado, population in high altitude areas of, 4
- colour-blindness, rare or absent in Andean HAN, 50
- congenital defects: altitude and, 103–4, 313; in Andean HAN, 342
- cor pulmonale: chronic, in some HAN (Tien Shan and Pamirs), 313
- cortisol: normal production of, in HAN, 78
- cosmic radiation, at high altitudes, 56, 69, 101
- cultures, altitudinal similarities in? 42
- cytochrome, microsomal, in oxygen transport, 89–90
- demographic structure of HAN: age and sex distribution, 319–22; total numbers, 317–19
- dental development: in Ethiopian HAN and LAN, 145–6; in Sherpa HAN and Tibetan children at low altitude, 154–6; in Tien Shan HAN and Kirov LAN, 160–1
- Diego allele: distribution of gene frequency for, in Andean HAN and neighbouring LAN, 58, 59, 60
- diet: composition of, in different groups of Andean HAN, 236–44
- 2,3-diphosphoglycerate content of red cells: in Bohr effect, 201, 208–9; did not increase on transfer to high altitude when alkalosis was prevented, 209; in polycythaemia, 203; in Sherpas and Europeans, 211, 212
- diseases, infectious: in babies, Andean HAN and LAN, 105; in Ethiopian HAN and LAN, 138, 139; and food requirements, 233, 245; mosquito-borne, HAN free of, 330, 334; often less common in HAN, 69
- divorce rate, in Ethiopia, 27
- dogs, hypoxia and foetal haemoglobin in, 94
- ductus arteriosus, patent: in HAN, 313, 329–30
- Echinococcus* infestation, in Andean HAN, 332
- Ecuador, 61; estimated population of, above 2,500 m, 318
- emphysema, and polycythaemia, 204
- energy balance, in Andean HAN, 235
- enzymes of oxidative chain, in HAN, 183, 185
- eosinophilia: in Ethiopian LAN, indicating nematode infestation, 197
- erythropoiesis: in bone marrow, altitude and, 123, 305; in mice, hypoxia and, 191–5
- erythropoietic islands: hypoxia and numbers of, in bone marrow and spleen of mice, 191, 193
- erythropoietin: increased secretion of, at high altitude, 191–4, 201, 305; in LAN upward migrants, 202
- Ethiopia, 18; economic and political structure of, 26–7; estimated population of, 319; ethnic and linguistic diversity of population of, 25–6; exogamy in, 61; genetic distances between HAN and neighbouring LAN in, 55; genetic polymorphisms in, 51–2; geology of, 20–1; migrants between altitudes in, 11, 58; occupied by man longer than other highlands, 51
- exogamy, 61
- fat: of animals, in diet of Andean HAN, 221, 222, 226; of arm, development of, in Andean HAN, 122; of body, rural–urban differences in, 135; intake of, by Andean HAN, 236, 237–8, (apparent absorption of) 235, (percentage of calorie intake as) 237
- fertility, 5; altitude and, 322–8; differential, apparently less important than differential mortality in selection pressure estimate, 56–7; possible

Index

- fertility (*cont.*)
 effects on, at high altitude, of increased loss of early embryos by resorption or abortion, 90, 92, 103–4; of Spaniards first coming to high altitudes in Andes, 67, 322; *see also* offspring, number of fibrinolytic activity, altitude and, 205, 305
 foeto-placental unit, 87–90
 food(s): altitude and requirements for, 226–33; factors other than altitude affecting requirement for, 233–4; habits in consumption of, Andes, 223–6; varieties and amounts of, utilized in three neighbouring towns at same altitude, Andes, 220, 222; and in four towns at different altitudes, Andes, 220–2
 footwear: scarcely used in Andes, 263, or by Sherpas, 264
 gametogenesis, change of altitude and, 70–3
 gene flow, 56; *see also* migration
 genetic distances, between HAN and neighbouring LAN, 53–4
 genetic drift, random: estimates of, for two populations in Andes, 57, 60
 genetic polymorphisms, 47–9; in Andes, 49–50, and nearby LAN, 52–4; in Ethiopian highlands, 51–2, and nearby LAN, 55; in Himalaya, 51, and nearby LAN, 54; possibility of estimating selection pressure from distribution of, 57–8
 genetics, 6; limitations of studies of, 47
 γ -globulin, in blood of HAN and HAN downward migrants, 207–8
 glucose: altitude, and content of, in blood of newborn, 90, 312; blood content of, in HAN and HAN downward migrants, 206
 glucose 6-phosphate dehydrogenase: no abnormal types of, in HAN, 55
 glycogen content, in liver of newborn animals: hypoxia of mother and, 102
 goats: anaerobic metabolism of foetus during acute hypoxia of mother, 95
 growth: altitude and rate of, 5, 8, 117, 312; and development at high altitudes, 117, 163–6, (Andes) 117–34, (Ethiopia) 138–47, (Himalaya) 147–55, (Tien Shan) 156–63; dietary factors in, 245; in first year of life, 104–6; of foetus, 92, 94, 95, 106, (sheep) 81, 92; genetic origin of rate of, in HAN downward migrants, 10; hypoxia and, in animals, 165
 guinea pigs: hypoxia, and length of gestation and birth weight in, 92; hypoxia, and vascular changes in placenta in, 82
 haematocrit measurements (red cell volumes): at and after birth, 94, 97–8; change in, with age, in HAN, 98–9, 200; in HAN, 184, 191, 305, 306, 312; in HAN, and HAN downward migrants, Andes, 206; in HAN and LAN, Ethiopia, 195, 196; in HAN and LAN, and LAN upward migrants, Himalaya, 198–9; in HAN and LAN, US, 203; in hypoxic mice, 192
 haematology: adaptive and inherited factors in, 345; factors other than altitude affecting, 204–12; of HAN, less affected by change of altitude than that of LAN, 8, 10; studies of, in HAN, Andes, 199–202, Ethiopia, 195–7, Himalaya, 197–9, and US, 203–4
 haematopoiesis, *see* erythropoiesis
 haematopoietic stem cells: hypoxia and numbers of, in bone marrow of mice, 191, 192, 194
 haemoglobin
 Bohr effect in, *see* Bohr effect
 concentration of, in blood: of HAN, and HAN downward migrants, Andes, 206; of HAN and LAN, 184, 243, (Ethiopia) 195, 196, (US) 203; of HAN and LAN, and LAN upward migrants, Himalaya, 198; of Sherpas and Britons, 197, 198
 dissociation curve of: in HAN, and LAN upward migrants, Andes, 202, 210; in polycythaemic HAN, US, 204; temperature and, 291
 foetal, increased in hypoxia in some species, 94
 sickle-cell: high altitude effect on, 330; lacking in HAN, 55; natural selection and, 58
 total circulating amount of, altitude and, 306
 hamster: anoestrus caused by cold in, 73; hypoxia, and congenital defects in, 103
 health: of LAN upward migrants, 328; and work performance, of LAN, and LAN and HAN migrants, 332–3
 heart: altitude, and rate of growth of, 138; increased size of right ventricle of, in HAN, 183, 184, 303–4, (children) 313; increased myocardial contractility of, in HAN, 304
 heart rate: altitude and, 284, 303, (children) 312; exercise and, in groups from different altitudes, 306
 heat: atmospheric humidity and loss of, 288; estimated production of, by sedentary and active LAN and active HAN, 229; extra loss of, in respiration at high altitude, 227–9
 height: of Andean HAN and LAN, (Indian) 119, 123, 126, (Mestizo) 126–7; of Ethiopian HAN and LAN, 140–3; parent–offspring correlations in, Quechua, 136–8; rate of growth in, Andean HAN and US standard, 120; of Sherpa HAN and Tibetan children at low altitude, 148–50, 151, 155; of Tien Shan HAN and Kirov LAN, 156, 157
 high altitude: defined as above 2,500 m, 17–18, 317; definitions of natives of, 189–90; estimated population living at, 319; rough terrain of, and calorie requirements of HAN, 232–3; simpler, less stable ecosystems at, 69; temperatures at, 68–9; variations in conditions at, 189
 Himalaya region, 34–5; agriculture of, 35–6; as genetic cross-roads, 51; genetic distances between HAN of, and neighbouring LAN, 54; genetic polymorphisms in, 51; geology of, 21–2; migration in, 39, 60; population of, 35; *see also* Nepal
 houses: Andes, 259–60; Ethiopia, 261; Himalaya, 261
 humidity of air, and heat loss, 288
 hydroxysteroid dehydrogenase, of cultured placenta: effect of hypoxia on activity of, 88
 hyperventilation: in newcomers to high altitude, 174, 183; in pregnant women at high altitude, 89
 hypoxia: interaction of cold stress and, in HAN, 281–2; of normal foetus, 88

Index

- inbreeding, in HAN, 60–1
 Incas, in Andes, 28, 30–1, 60, 67; inbreeding in dynasty of, 61
 Indian Government: Border Roads Organization of, in Himalaya, 36
 insulation, unit of (for clothing), 262
 iodine, in diet of HAN, 244
 iron: blood content of, in HAN and HAN downward migrants, Andes, 206; intake of, by Andean HAN, 239, 243; intestinal absorption of, promoted by increase in erythropoiesis, 201
 irrigation, in Himalaya, 35, 38, 40

 Karakorum ranges, Pakistan, 51, 54, 60
 Kashmir province, estimated population of, 319
 Kirghizia, 77–8

 lactic acid, produced in foetus in acute hypoxia of mother, 95
 Ladakh province, estimated population of, 319
 leprosy, in Nepal, 333
 leucocytes, in HAN children, 312
 livestock: exposure of HAN to climatic stress in herding of, 265–6; introduced by Spaniards, at first failed to reproduce at high altitudes, 68; lodged on ground floor of Himalayan houses, 261; raising of, in Andes, 34, 117–18, 219, in Ethiopia, 26–7, 138, and in Himalaya, 36, 38–9, 40–1
 llama: altitude, and size of placenta in, 83
 lung capacity, 163–4, 165–6; effect on, of age at acclimatization to high altitude, 133, 134; residual volume as percentage of, higher in HAN, 302
 lung forced vital capacity, in Andean HAN and LAN: chest circumference and, 132–3; compared with US standard, 130–1; and forced expiratory volume, 131–2
 lung vital capacity: not affected by altitude, 302; and tidal volume and respiratory volume, in Tien Shan HAN and Kirov LAN, 161, 162, 163
 lung oedema, lung ventilation, *see* pulmonary oedema, pulmonary ventilation
 luteinizing hormone: altitude, and production of, 76, 130
 lymphocytosis, in HAN, 197, 305

 magnesium, in red cells of HAN, 207
 malaria: agent of natural selection on sickle-cell haemoglobin, 58; at different altitudes in Peru, 330; eradication of, in Nepal, 39; in Ethiopian HAN and LAN, 138
 measles: child mortality from, Bolivian HAN, 343
 meat and other animal products, in diet of Andean HAN, 221, 222, 223, 224, 225–6, 244
 menarche, altitude and age of, 77–8, 311; in Andes, 128–9; in Himalaya, 154; in Tien Shan HAN and Kirov LAN, 159–60
 menstrual cycle, change of altitude and, 73
 Mestizos, Peru, 32, 99–100, 118; growth of children of, HAN and LAN, 126–7
 methaemoglobin: in normal, anaemic, and polycythaemic HAN, 201
 mice: alkalosis in, and sex ratio, 103; effects of hypoxia in, on congenital defects, 103, on erythropoiesis, 191–4, on foetal haemoglobin, 94, on gametogenesis, 71, 72, on mating behaviour, 74, and on survival and abnormalities of ova, 79–80, 90, 92
 migration (gene flow): in Andes, 58, 60, 321, 345–6; in Ethiopia, 11, 58; in Himalaya, 39, 60
 mining, in Andes, 220
 mitochondria: numbers of, in skeletal muscle of HAN, 183, 185
 Mongoloid ancestry, in Himalaya, 51, 54
 monogamy, serial, in Andes, 91
 morbidity at high altitudes: infant and neonatal, 340–2; childhood, 342–3; adult, 343–5
 mortality: death rates, Andes, (by age) 338, (age- and sex-specific) 339, (children) 342–3; differential, apparently more important than differential fertility in selection pressure estimate, 56–7
 mortality, infant: birth weight and, HAN and LAN, 341, 342; of first generation Spaniards at Potosi (4,000 m), 67; in HAN and LAN, Andes, 56, 106, 340–1; hypoxia and, in animals, 102; increased in HAN, 101, 106, 244, 399–40; sex and, HAN and LAN, 100–1, 321–2
 mountain climbing: oxygen utilization during, by HAN and LAN, 178
 mountain sickness, altitude-specific disease, 329
 muscle: of arm, development of, in Andean HAN, 122, 124; capillarity of, in HAN, 183, 185, 304–5, 335; number of mitochondria in skeletal, of HAN, 183, 185
 mutations: cosmic radiation and, 56, 69; not detected in HAN populations, 56

 natural selection, possibility of estimating intensity of: from differential survival and fertility rates, 56–7; from distribution of genetic polymorphisms, 57–8
 nematode infestation: at different altitudes, Peru, 331; in LAN, Ethiopia, 197
 Nepal, 19, 37; agriculture and stock-raising in, 38–9, 40–1; estimated population of, 319; population distribution in, 37; vegetational zones of, 37, 38
 neutrophils: excess segmentation of, at high altitudes, 305; numbers of, in HAN children, 312
 newborn: measurements of, in HAN downward migrants, 10, and in HAN and LAN, Andes, 99–100; sex differences in measurements of, 100; weight of, *see* birth weight
 niacin: intake of, by Andean HAN, 239, 240
 night vision, altitude and, 18
 nitrogen balance, in Andean HAN, 235
 Nuñoa district, S. Peru, 31–2, 118; population structure in, 120
 nutrition, 5, 163; and growth, Andes, 134–5; of infants, HAN and LAN, 104–5; of Sherpa HAN, and Tibetan children at low altitude, 147, 155

 oestrogens: altitude, and concentrations of, in blood of mother and of newborn, 87–8, 102; produced by placenta, 86–7
 offspring, numbers of; in Andean HAN and LAN, 90–1, 323–6, 345; in Ethiopian HAN, 327; in

Index

offspring (*cont.*)

- Nepal, 327; in Tien Shan, 311; percentage surviving to adult age, Andes, 345
- oogenesis, change of altitude and, 71
- ova: hypoxia, and survival and abnormalities of, in mice, 79–80, 90, 92
- oxygen: altitude and consumption of, 300; cost in, per unit of work, in groups from different altitudes, 308; exercise ventilation and uptake of, in HAN, and in LAN upward migrants, 182–3; in foetal blood, 94; hyperventilatory response to breathing of, altitude and, 300, 302, 310; partial pressure of, at different altitudes, 225–7; transport of, across placenta, 88–90; utilization of, in mountain climbing, by HAN and LAN, 178
- oxygen-carrying capacity of blood: effect of hypoxia on, in HAN, and in LAN upward migrants, 184
- Pamir Mountains: meteorological data for different altitudes in, 299; peoples of, 4, 51
- Pan-American Health Organization, and high-altitude project, 3
- parasites, intestinal: in Andes, in HAN, 234, and in populations at different altitudes, 331, 332; in Ethiopian LAN, 146, 163
- Peru, estimated percentage of population of, above 2,500 m, 317–19
- phosphogluconate dehydrogenase, in HAN and LAN, Ethiopia, 55
- phosphorus: blood content of, HAN and HAN downward migrants, Andes, 206; intake of, by Andean HAN, 239, 242–3; ratio of calcium to, in diet of Andean HAN, 242
- physical activity, habitual: and aerobic capacity, 175, 177; and hypoxic ventilatory drive, 182; and oxygen utilization in mountain climbing, 178; and ventilatory exchange ratio, 182
- placenta: altitude, and infarcts in, (guinea pigs) 82, (humans) 85–6; altitude, and ratio of weight of, to birth weight, humans, 10, 83–4, 87; altitude, and size and function of, (animals) 80–3, (humans) 83, 84, 85, 87; oxygen transport across, 88–90; perfusion of, with various concentrations of oxygen, 84–5; production of hormones by, 86–8
- placenta praevia: altitude, and frequency of, 86, 97
- plasma proteins, polymorphisms of, 48
- plasma volume, in LAN upward migrants, 202
- platelets, in HAN, 205, 305
- polyandry, in Nepal, 327
- polycythaemia: in HAN, 184–5, 190–1, 194–5, 203–4; in hypoxic mice, 191–4; only apparent in Andean HAN in pathological conditions, 199
- population, of high-altitude areas of world, 317–19
- population genetics, 47
- population structure, HAN of Peru compared with total population, 126
- potassium: blood content of, in HAN downward migrants and LAN upward migrants, 206–7
- potatoes (fresh or dehydrated): in diet of Andean HAN, 221, 223–4; as source of ascorbic acid, 241
- Potosi city, Bolivia (4,000 m): first generation Spanish children in, all died at birth or soon after, 67
- precipitation: Andes, monthly averages, 253; Ethiopia, of monsoon pattern, 258; Himalaya, 35, 40, (monthly averages) 257
- pregnancy, altitude and duration of: in guinea pig, 92; in human, 93
- pregnenediol: excretion of, by HAN and LAN women, 77
- progesterone, 77; produced by placenta, 86
- protein: altitude and requirement for, 234; hypoxia or altitude, and specific dynamic action of, 230; intake of, by Andean HAN, 236, (as percentage of calorie intake), 237
- proteinograms: for blood of two ethnic groups of HAN, before and after downward migration, 207–8
- protozoan infestations, at different altitudes in Peru, 331
- psychomotor development of babies, Andean HAN and LAN, 105, 106
- puberty, altitude and age of, 77–8, 312; in Andean HAN and LAN, 129–30; in Ethiopian HAN and LAN, 146; in Tien Shan HAN and Kirov LAN, 159–60; genetic origin of late, in HAN downward migrants, 10
- pulmonary oedema, altitude-specific disease, 328–9
- pulmonary ventilation: altitude and, 300, 301; exercise and, in groups from different altitudes, 307
- Quechua people, Bolivia and S. Peru, 4, 10, 57, 182–3, 207; genetic distances between other tribes and, 54; large chests of, genetically determined, 8; migration of, in 15th century, 59–60
- rabbits: hypoxia and foetal haemoglobin in, 94
- rats, hypoxia in: and embryonic deaths, 79; and gametogenesis, 71; and litter size and birth weight, 92; and lung development, 133–4; and placenta, 82
- red cell antigens, polymorphisms of: in Andes, 48, 49–50; in Ethiopia, 48, 52; in Himalaya, 48, 51; *see also* Diego allele, Rhesus system
- red cell proteins, polymorphisms of: in Andes, 48, 49; in Ethiopia, 48, 52; in Himalaya, 48; *see also* glucose 6-phosphate dehydrogenase, haemoglobin
- red cells: altitude, and numbers of, 303, 305, 306, 312; numbers of, HAN and LAN, Ethiopia, 105, HAN, LAN, and LAN upward migrants, Himalaya, 198, and LAN upward migrants, Andes, 202; size of, in HAN, 305, 312; volumes of, *see* haematocrit measurements
- respiration: altitude, and loss of heat in, 227–9; percentage loss in, of heat produced by LAN (sedentary and active), and HAN (active), 229
- respiratory centre, sensitivity threshold to hypoxia of: in HAN, and in LAN acclimatizing to high altitude, 300, 302; in HAN returned to high altitude after period at low altitude, 310
- respiratory rate, altitude and, 300, 301
- respiratory disease: in HAN, 332, 333, 334; in HAN downward migrants, 344
- reticulocytes: altitude, and numbers of, 305, 306, 312; altitude, and maturation rate of, 305, 306

Index

- Rhesus system: distribution of frequency of segments of, from north to south along Andes, 49–50, 58, 59, 60; in Himalaya, 54; as indicator of genetic drift in Andes, 60; possible action of natural selection through variation in resistance of haplotypes of, to salmonella infections, 57–8
- riboflavin: intake of, by Andean HAN, 239, 240
- rural–urban differences, Andes: in body fat, 135; in skinfold thickness and muscle development, 124; in stature and weight, 123
- salmonella infections, variation in resistance of Rhesus haplotypes to, 58
- Sardinia, sickle-cell haemoglobin in HAN and LAN in, 58
- serum proteins: in Andes, 48, 50; in Ethiopia, 48, 52; in Himalaya, 48
- sex: and birth weight in HAN and LAN, Andes, 100; and body measurements of babies, HAN and LAN, 106; and development of arm tissues, Andean HAN and US, 122; and neonatal mortality, HAN and LAN, 100–1; and response to cold stress, 278–9
- sex hormones, altitude and, 73–8
- sex ratio of births: in HAN and LAN, 103, 321; in mice, alkalosis and, 103
- sheep, at high altitude: foetal growth rate in, 92; foetal haemoglobin in, 94; metabolism of foetus in, 95; placenta in, 81, 82, 85
- Sherpa people, Nepal, 4, 7; aerobic capacity of, 178, 179; growth and development of, compared with Tibetans at low altitude, 147–55; population structure of, 321, 322
- Sikkim, estimated population of, 319
- silicosis, and polycythaemia, 204
- skeletal age and chronological age: in Andean HAN, 127–8; in Ethiopian HAN and LAN, 144–5; in Sherpa HAN and Tibetan children at low altitude, 153–4, 155
- skin colour, of HAN and LAN in same area, 289–90
- skinfold thickness: in Andean HAN, 120–1, (rural and urban) 124; in Ethiopian HAN and LAN, 143–4; in Sherpa HAN and Tibetan children at low altitude, 151, 153, 155, 164; in Tien Shan HAN and Kirov LAN, 157, 159, 162
- smoking, haematological effects of, 204
- socio-economic factors, 66; may obscure effects of altitude, 164–5; may be related to altitude, 69–70
- sodium: blood content of, in HAN, HAN downward migrants, and LAN upward migrants, 207
- soil erosion, in Nepal, 40
- solar radiation, altitude and, 252–3, 289
- spermatogenesis, change of altitude and, 70–1, 72–3
- survivorship curves, for US, Peru, and Peruvian HAN, 338
- Tajikistan, 78, 91
- temperature of body: in cold stress, remains higher in extremities of HAN than of LAN, but rectal temperature decreases more in HAN, 274–5, 291; on different parts of skin of HAN, 271–2; of HAN during sleep in different conditions, 266–9, and during normal activities, 269–70; of hands and feet exposed to cold, in HAN and LAN, 272–3, 277–8; relation of body weight to, 272; sex differences in, 278–9
- temperature of surroundings: altitude and, 252; diurnal variation in, increases with altitude; seasonal variation diminishes, 251–2; maximum, at three altitudes, Ethiopia, 258; mean annual, over 40 years, Andes, 256; mean monthly, in Andes at three altitudes, 254, 255–6; in Ethiopia at three altitudes, 258, and in Himalaya, 256–7; within houses, Andes, 259–60
- testosterone: altitude, and production and excretion of, 74–6, 130
- thalassaemia, lacking in HAN, 55
- thiamin: intake of, by Andean HAN, 239, 240
- thrombo-cythaemia, in LAN upward migrants, 205
- thrombo-embolic accidents: rare in HAN, frequent in LAN upward migrants, 205, 329
- thyroid, in increased basal metabolic rate of HAN?, 280
- Tibet, plateau of, 18–19, 22; estimated population of, 319
- Tibetan migrants, Nepal, 4, 10; growth and development of children of, compared with Sherpa HAN, 147–55
- Tien Shan Mountains, 19; meteorological data for different altitudes in, 299; peoples of, 4, 51
- Tigrean (Tigrinyan) people, Ethiopia, 22, 26, 195
- trade, in Himalayan region, 36, 39
- trematode infestation, at different altitudes in Peru, 331
- tuberculosis, in HAN downward migrants, 333, 344
- tubers, staple food in Andes above 3,500 m, 221, 222, 223–4
- typhoid fever, variation in resistance of Rhesus haplotypes to, 58
- typhus, at different altitudes in Peru, 330, 332
- ultraviolet radiation, at high altitudes, 69, 289; and vitamin D requirements of HAN, 233, 245
- United States: haematology of HAN and LAN in, 203–4
- uric acid content of blood: in HAN and HAN downward migrants, Andes, 206
- vascular resistance, altitude and, 304, 310
- vasodilation: alcohol and, 285; in extremities of HAN during cold stress? 278; in placenta in hypoxia, 89; in uterus in hypoxia, 86
- ventilatory drive, hypoxic; habitual physical activity and, 181, 182
- ventilatory exchange ratios, habitual physical activity and, 182
- vitamin A: intake of, by Andean HAN, 228, 229, 230, 245
- vitamin D: and calcium absorption, 242; ultraviolet radiation at high altitudes, and requirement of, 233, 245
- water: effect of lowered boiling point of, at high altitudes, on cooked foods, 226, 240, 245; extra loss of, in respiration at high altitudes, 227–8
- weight: of Andean HAN, rural and urban, 123; of

Cambridge University Press

978-0-521-11195-9 - The Biology of High-Altitude Peoples

Edited by P. T. Baker

Index

[More information](#)*Index*weight (*cont.*)

Ethiopian HAN and LAN, 140–3; of Sherpa HAN and Tibetan children at low altitude, 148–50, 151, 155; *see also* birth weight

white cell antigens (HLA system), polymorphisms of: in Andes, 48, 50; estimate of selection pressure from frequency of haplotypes of, 57; in Ethiopia, 48; in Himalaya, 48, 51

white cells: temporary increase of, in LAN upward migrants, 204–5

wool textiles: resistance to heat flow of, compared with that of cotton textiles, 262

work capacity, altitude and, 5, 166, 173–4, 307–10; in HAN after a period at low altitude, 310–11

World Health Organization, and high-altitude project, 3

yellow fever, at different altitudes in Peru, 330