

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)

## INDEX

**NB** References in bold type refer to large passages or chapters.

References in italic type refer to figures &amp; tables.

- Abies alba*, 4, 23, 29  
*Abies amabilis*, 56  
*Abies balsamea*, 3, 29, 69, 70  
*Abies firma*, 4, 18; mineral cycling in, 344, 390  
*Abies fraseri*, 14  
*Abies grandis*, 215  
*Abies mariesii*, 4, 18  
*Abies procera*, 56  
*Abies* spp., 26  
*Acacia catechu*, 183  
*Acacia* spp., 7  
*Acer campestre*, 3, 21, 574  
*Acer circinatum*, 247  
*Acer japonicum*, 21  
*Acer macrophyllum*, 5, 56  
*Acer pennsylvanicum*, 32  
*Acer platanoides*, 4, 5, 21, 22  
*Acer pseudoplatanus*, 147  
*Acer rubrum*, 5, 21, 23, 29, 32, 559; photosynthetic efficiency, 515  
*Acer saccharum*, 5, 21, 23, 32, 325; and environmental influences, 69, 70, 74, 75, 78  
*Acer sieboldiana*, 21  
*Acer spicatum*, 21, 23, 32  
*Acer* spp., 14, 21; mineral cycling in, 344, 398  
*Acer tataricum*, 5, 21, 23  
*Achromobacter* spp., 317  
*Actinodaphne lancifolia*, 18  
*Adenanthos terminalis*, 7  
adsorption, of chemicals, by vegetation, 298–9  
*Aegopodium podagraria*, 21  
aerodynamic exchange: and carbon dioxide fluxes, 104–5; and heat fluxes, 98–9  
*Aesculus* spp., 14  
age, *see* stand age  
air turbulence *see* turbulence, air  
albedo: of global radiation, 120–4; of snowpack, 220  
*Albizia* spp., 10  
algae, blue-green, 300  
*Allenblackia floribunda*, 3, 4, 10  
*Allanblackia* spp., 10  
*Allium ursinum*, 23  
*Alnus barbata*, 21  
*Alnus crispa*, 29  
*Alnus rubra*: mineral cycling in, 344, 363, 406  
*Alnus* spp., 300  
*Alstonia scholaris*, 182  
aluminium: atmospheric input to soil, 297; litter input to soil, 301  
aluminium buffer zone, in soil, 306–7  
*Amelanchier arborea*, 29  
Anacardiaceae, 13  
Andersby Angsbackar, Sweden, 4, 616; radiation balance, 175; temperate forest of, 15, 23  
Andrews Experimental Forest, Oregon, USA, 245, 246–7, 656; and leaching effects, 373; mineral cycling, 382; partial cutting simulation, 250–2; temperate forest of, 18–19  
*Anemone nemorosa*, 18, 22, 23  
*Anemone ranunculoides*, 21, 23  
angiosperms: biomass, 431–3  
Annonaceae, 575  
*Anogeissus latifolia*, 3, 13, 365  
Arabari Range, Bengal, 182  
*Aralia nudicaulis*, 32  
Ashu, Kyoto, Japan, 4, 21, 601  
*Asparagus acutifolius*, 8  
*Asperula odorata*, 22, 23  
autotrophs, 509; respiration of, 523–5; *see also* photosynthesis  
Azerbaijan, USSR, 5, 622–4  
*Azotobacter* spp., 300  
Bab, Czechoslovakia, 3, 580; radiation flux, 139–40; water balance, 189  
Babadag 1, Rumania, 4, 15, 23, 611  
Babadag 2, Rumania, 4, 23, 612  
Baku, 21  
Banco, Ivory Coast, 3, 598; tropical rain forest, 9–11; water balance, 182  
*Baphia* spp., 10, 95  
Barringtoniaceae, 11  
Beaver Creek Experimental Watershed, Arizona, USA, 245, 247–8; simulations, 249, 250, 251, 253  
*beech forest*: biomass dynamics, 480; carbon dioxide balance, 152–4, 155  
canopy drip, 269; carbon metabolism,

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- 504; mineral cycling, 373; nitrogen fixation, 372; radiation and heat flux, 149; stand age, 413, 419; water balance, 190–1, 270, 273, 280
- Belgorod, USSR, 5, 626–7
- Belize, 176
- Berberis nervosa*, 247
- Berlinia confusa*, 3, 10
- beta density function, 43
- Betula alleghaniensis*, 325
- Betula ermani*, 4, 18
- Betula glandulosa*, 21
- Betula lutea*, 5, 23, 69–70
- Betula papyrifera*, 29, 379
- Betula pendula*, 5, 23
- Betula pubescens*, 5, 23
- Betula* spp.: mineral cycling in forests of, 344, 398, 404
- Betula verrucosa*, 4, 18, 23, 27
- Bialowieza, Poland, 4, 608; radiation balance, 175; temperate forest, 22
- biomass: angiosperm v. gymnosperm, 431–3; annual net primary production, 417–18, 573–671; *see also* productivity; branch and bole comparisons, 479–85; COMEX *see* COMEX; distribution among components, 573–671, dynamics of *see* DCLM; forest floor, 350–7, 358–9, 571; FORET *see* FORET; heterotroph to autotroph ratio, 445–7; standing crop, 573–671; and mineral cycling, 364–8; of roots, 485–6; *see also* litter
- biosphere, and radiation flux *see* radiation flux
- birch forests, water balance of, 194
- boreal forests, 26–9, 31, 32; evergreen, 26–9, 31, 32; and linear model of biomass dynamics, 473–6; litter turnover, 441; mineral cycling in, 344; nitrogen accumulation, 353–8; organic matter, 444; radiant energy, 172–6, 177; and water balance, 188, 193–6; *see also* individual sites
- Boswellia* spp., 10
- Brachypodium pinnatum*, 18, 23
- Brachypodium ramosum*, 8
- Brookhaven, New York, USA, 5, 23, 652; and carbon metabolism, 506–7; photosynthetic processes, 510–11
- Buchanania lanzan*, 3, 13
- buffering systems, in soil, 305–7
- Bursera* spp., 10
- Burseraceae, 3, 4, 10, 11, 575
- calcium: atmospheric input, 296–7, 301, 302; cycling of, 346, 349, 350–3, 355–9, 361–2, 365–6; exchangeable cation, 305–7; site studies, 376–407
- Calluna vulgaris*, 18
- Calyptranthe petioli*, 16
- Canarium* spp., 10
- canopy drip, 267, 268, 275; and throughfall, 110, 243, 267, 268
- carbonate buffer zone, of soil, 304–6
- carbon dioxide flux, 152–4, 155; absorption into soil, 300; determination of, 104–9; and forest balance, 112, 113–16; and photosynthate allocation, 520; and respiration, 522–30; in spruce forest, 160, 164–8, 169; *see also* carbon metabolism
- carbon metabolism, 499–532; budgets of, 503–30; and CONIFER model, 552–8; of forests, 502–3; in mesophytic deciduous forest, 503–6, 508; in oakpine forest, 506–7, 508; and photosynthesis, 509–22; and respiration, 522–30; *see also* carbon dioxide
- Cardamine pratensis*, 574
- Cardamine* spp., 21
- Carex lanceolata*, 18
- Carex michelii*, 22
- Carex montana*, 22
- Carex pilosa*, 21
- Carex silvatica*, 22
- Carex trisperma*, 22
- Carissa opaca*, 13
- Carpinus betulus*, 3, 4, 21, 22, 574
- Carpinus caroliniana*, 21
- Carpinus caucasica*, 21
- Carpinus laxiflora*, 4, 21
- Carpinus* spp., 21, 215
- Carya orata*, 24
- Carya* spp., 344, 395
- Carya tomentosa*, 5, 24
- Castanopsis chrysophylla*, 5, 19, 247
- Cascade Head, USA, 344, 392
- Castanopsis cuspidata*, 4, 15, 18
- Caucasus Birch, Azerbaijan, USSR, 15, 622–3
- Cedar River, USA, 344, 370
- Centaurea stenolepis*, 23
- Central Forest Reserve, USSR, 5, 28, 176, 628
- Cercis canadensis*, 24, 505
- Chakia, Varanasi, India, 3, 587, 588; radiant energy in, 171, 172–3, 175; tropical deciduous forest, 9, 11, 13
- Chamaecyparis obtusa*, 4, 16
- chemical elements, dynamics of, in soil, 293–318, 325–6; adsorption of, by vegetation, 298–9; anion adsorption, 309; budgets for forests, 376–407;

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- buffering systems, 305–7; cation exchange equilibria, 307–8; gas adsorption, 300–1; internal processes, 303–15; leaching, 295–8; litterfall, 301–2; output routes, 316–18; pathways of chemical input, 293–303; phosphate equilibria, 309–11; redox equilibria, 309–12; transport in soil, 314–15; weathering, 312–14; *see also* mineral cycling
- chlorine, 297
- Chrysanthemum corymbosum*, 22
- Chrysophyllum* spp., 3, 10
- Circaea quadrisulcata*, 24
- Cladonia rangiferina*, 18, 32
- Cladonia* spp., 28, 29
- classification, of forest systems, 2–33, 573–671; boreal forest, 26–32; Mediterranean woodlands, 6–8; temperate forests, 14–25; tropical forests, 8–14
- Clethra barbinervis*, 16
- Cleyera japonica*, 16
- climatic factors, and forest diversity, 66–7
- Clostridium* spp., 300
- Cola* spp., 10
- Combretaceae, 13
- Combretodendron africanum*, 3, 10
- Combretodendron* spp., 10
- COMEX computer code, 454–5, 459–60, 461, 469, 489–92; numerical methods, 490–2
- Commiphora* spp., 10
- computer simulations *see* models
- CONIFER model, 522–8, 560
- coniferous forests: and mineral cycling, 360–4, 375; photosynthetic allocation, 519; radiation flux, 123; and stand age, 413, 414; succession modelling in, 52–65; water flux, 192, 194
- Coptis groenlandica*, 32
- Cornus florida*, 24, 505, 515
- Cornus mas*, 3
- Cornus sanguinea*, 18
- Corydalis halleri*, 21
- Corylus americana*, 24
- Corylus avellana*, 4, 5, 21, 23, 574
- Corylus cornuta*, 32
- Corylus* spp., 23
- Cotinus coggygria*, 23
- Coula edulis*, 3, 4, 10
- Coula* spp., 10
- Coweeta, USA, 5, 245–6, 653–4; clearcut simulation, 248, 249, 250; and conversion to conifers, 253; mineral cycling in, 344, 373, 389, 397; radiation balance, 175; temperate forest, 19–20, 23
- Crataegus currisepala*, 21
- Crataegus oxyacantha*, 3
- Crataegus* spp., 574
- Cyathea* spp., 15
- cycling, of elements *see* chemical elements
- Dacryodes klaineana*, 3, 4, 10
- Dactylis polygama*, 22
- Dakar: radiation balance, 176
- Darcy's law, 287–8
- deciduous forests: and biomass dynamics, 474, 480; carbon budget, 503–6, 507–8; and chemical input to soil, 296, 302, 328; and litter fall, 420–1; and mineral cycling, 360–4, 375; radiation flux, 123, 124, 132; stand age, 414, 415; and succession modelling, 52–65; temperate, 16, 20–5, 31; tropical, 11–13; *see also* individual sites
- defoliation, effects of: using CONIFER model, 555–9; using TEEM-Prosper model, 547–52
- Dehra Dun, India, 183
- denitrification, of soils, 317–18
- Dennstaedtia punctilobula*, 23
- Dentaria glandulosa*, 23
- Deschampsia flexuosa*, 23
- Dicranum* spp., 29
- differential-difference equation models, of stand development, 41–2
- Diospyros melanoxylon*, 13
- Diospyros* spp., 10
- Dipterocarpaceae, 4, 11
- Dipterocarpus* spp., 11
- Distylium racemosum*, 15
- diversity of woodlands, 67–70, ecosystem, 70–44
- donor-controlled linear compartment models (DCLM), and biomass dynamics, 452–94; COMEX *see* COMEX; general linear model, 455–69
- Dryanobalanus* spp., 11
- Dryopteris austriaca*, 28
- Dryopteris intermedia*, 32
- Dryopteris spinulosa*, 23
- Duke Forest, USA, 5, 19–20, 655
- Eastern Deciduous Forest Biome (EDFB), 65–6, 152–4
- Ebenaceae, 13
- Ebersberger, Germany: carbon dioxide sampling in, 106, 107–8, 160, 164–8, 169; heat flux, 142, 143, 147, 148–51, 152; radiation flux, 139, 148–51, 152; sensible and latent heat flux, 157–60
- ecosystem composition, woodland, 65–79; climatic and biogeographic elements, 66–7; local edaphic control, 67–70;

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- primary production, 70–4; and time-related influences, 74–9  
 ECOWAT, hydrology model, 243, 245–55  
 eddy correlation method: and carbon dioxide flux, 108; and heat flux, 103  
*Elaeagnus angustifolia*, 5, 21  
 elements, chemicals *see* chemical elements  
 El Verde forest, Puerto Rico, 180, 181–2  
 energy balance, model of, 112, 153  
 environmental influences: and ecosystem diversity of woodlands, 67–70; models of, 111–20  
 equatorial forest biome: and litter fall, 420; radiation energy, 170–1, 172, 173, 174, 175–6; water balance in, 178, 180, 182  
*Equisetum* spp., 29  
*Erianthus* spp., 21  
*Eucalyptus globulus*, 183  
 Eucalyptus forest: biomass dynamics, 483; water balance, 183  
*Eucalyptus obliqua*, 3, 6, 7, 573  
*Euonymus europaeus*, 574  
*Euonymus* spp., 21, 22  
 Euphorbiaceae, 4, 11  
*Eurya japonica*, 16  
 evaporation, 214, 249, 546; of intercepted water, 215–16, 271, 275; from litter, 225, 277; from soil, 277  
 Evaportron, and latent heat flux determination, 103  
 evapotranspiration, 272, 273; and productivity, 417; simulation of, 288–93, 546, 548–9, 551  
 evergreen forests, 15, 16–20, 30, 31; biomass dynamics, 475–5, 483; litter fall, 420, 421; temperate, 15, 16–20, 30, 31  
  
 Faeglharting, 28, 665  
 Fagaceae, 4, 11  
*Fagus crenata*, 4, 21  
*Fagus grandifolia*, 5, 21, 23, 32, 325  
*Fagus* spp., 14; mineral cycling in, 344, 398; water storage, 215  
*Fagus sylvatica*, 3, 4, 6, 18, 21, 22, 23, 24, 302, 574; mineral cycling in, 344, 363, 401, 402, 403, 405; and root organic matter turnover, 320  
 fertilization, and chemical input to soil, 303  
*Festuca heterophylla*, 22  
 Feterosso Site, Scotland, 128  
 Flacourtiaceae, 11  
 Fluxatron, and latent heat flux determination, 103  
 Fontainebleau, France, 3, 22, 583  
 FORET computer model, and biomass allocation, 481–2  
*Fragaria vesca*, 22  
  
*Frangula alnus*, 4, 22  
*Fraxinus excelsior*, 4, 5, 21, 23  
 fungivore respiration, 529  
  
*Gagea lutea*, 21  
*Galium dasypodium*, 23  
*Galium schultesii*, 22  
*Galium* spp., 24  
 Gapon equation, 307–8  
 gas adsorption, by soil and trees, 300–1  
*Gaultheria hispidula*, 32  
*Gaultheria procumbens*, 23  
*Gaultheria shallon*, 247  
*Gaylussacia baccata*, 506  
 general linear model, for biomass dynamics, 455–69  
*Geocaulon lividum*, 29  
 geological input, to soil, 302  
 Georgia, Piedmont, USA, deciduous-coniferous succession model, 53, 57–8, 61  
 global radiation (short wave radiation flux), 120–31, 148–51; and effect on hydrological cycle, 211–12; extinction through canopy, 124, 126–7; interception by forest biomass, 122–3, 124, 125; photosynthetically active radiation (PAR) flux, 127–31; reflection on upper surfaces, 120–4, 125; transmission to soil, 127  
 Gorakhpur, India, 3, 589–97  
 grasses, perennial *see* individual spp.  
 Great Smoky Mountains, Tennessee, USA, 14  
 growth, of trees, 39; evolution of forest stand modelling, 40–50; simulation models, 43–50; succession modelling *see* succession modelling  
 Guttiferae, 10, 11  
*gymnosperms*, 431–2  
  
*Hakea* spp., 7  
*Hannoa* spp., 10  
 heat balance of atmosphere-biosphere system, 95; and effect of air turbulence, 142, 159–60, 161; instrumentation, 98–103; sensible heat, 141–2, 143, 147–52; *see also* radiation flux, sensible and latent heat  
*Hedera helix*, 8, 21, 574  
 herbs *see* individual species  
 Hestehaven, Denmark, 3, 581; radiation balance, 175; temperate forest, 22  
 heterotrophs, 509; respiration of, 525–30, 531  
 H. J. Andrews site *see* Andrews site  
 Honshu, Japan, 16  
*Hopea* spp., 11  
 Hubbard Brook, New Hampshire, USA, 5, 651; mineral cycling in, 344, 373, 398

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)

## Index

- Hydrangea arborescens*, 505  
hydrologic cycles, 205–55, 265–6;  
  ECOWAT, 243, 245–55; hydrology  
  models, 242–53; PROSPER, 242, 245–  
  55, *see also* TEEM; structure and  
  function of a watershed, 207–42  
hydrology, of terrestrial strata, 213–41;  
  interception by foliage, 213–18; litter  
  and soil surface, 223–7; plant water  
  relations, 227–35; snowpack, 218–23;  
  soil water balance, 235–41; *see also*  
  water, dynamics of  
*Hylocomium proliferum*, 29  
*Hylocomium splendens*, 29
- Illex aquifolium*, 18  
*Illicium religiosum*, 16, 18  
impaction, and element contribution to soil,  
  299  
*Imperata* spp., 21  
infiltration, 223–7  
instrumentation: and carbon dioxide flux,  
  104–9; in forest studies, 98–103; of  
  heat flux, 98–103; of radiation, 100–1;  
  and water exchange processes, 109–11  
interception, of global radiation, 122–3,  
  124, 125  
interception, of precipitation, 213–18,  
  267–8, 271, 275–7; storage and evapora-  
  tion processes, 215–16  
invertebrates, respiration of, 529–30  
iron: atmospheric input to soil, 297; litter  
  input to soil, 301 isotopic techniques:  
  of mineralisation of soil, 303–4, 322;  
  of root dynamics, 325  
Ispina, Poland, 4, 609
- Juniperus communis*, 28  
*Juniperus deppeana*, 247
- Kalmia latifolia*, 23  
Kampinos National Park, Poland, 4, 18,  
  19, 610  
Karelia, USSR, 5, 629–45; mineral cycling,  
  383  
*Kennedia prostrata*, 7  
Koinas site, USSR, 4, 28, 176, 622  
Koiwai, Japan, 600  
Kongalund, Sweden, 4, 617–18; boreal  
  forests, 26, 28; mineral cycling in, 344,  
  384, 405; radiation balance, 175; tempe-  
  rate forest, 23; water balance, 187  
Kuala Lumpur, Malaysia, 12
- Lamium galeobdolon*, 21, 23, 574  
landscape dynamics model, 60–3  
Langarod, Sweden, 4, 23, 619
- Larix leptolepis*, 4, 16  
latent heat flux: long-term measurements  
  of, 147–52; in spruce forest, 157–60; in  
  snowpack, 221  
*Lathyrus niger*, 22  
*Lathyrus vernus*, 22  
Lauraceae, 3, 11, 575  
leaching, of chemical elements: and loss of  
  nitrogen, 372–4; from root zone, 318, 325;  
  site studies, 386–407; to soil, 295–8, 316  
leaf area index: and Stand Energy model,  
  541–2; and TEEM-Prosper model, 544  
Lecythidaceae, 3, 11, 575  
*Ledum groenlandicum*, 29  
Leguminosae, 3, 4, 9, 11, 300, 575  
*Lepidosperma canescens*, 7  
Le Rouquet, France: radiation flux, 130,  
  175; water balance, 184, 185, 186  
light, available: and photosynthesis, 510;  
  and productivity, 422–4  
*Ligustrum* spp., 22  
*Linnaea borealis*, 28  
Linnebjerg, Sweden, 4, 620; radiation  
  balance, 175; temperate forest, 23  
*Liriodendron* spp., 14; carbon budget,  
  503–4, 505; and general linear model for  
  biomass dynamics, 455–7, 478–9; and  
  microflora respiration, 526–8; mineral  
  cycling in, 344, 393; photosynthetic  
  processes, 511; root organic accumula-  
  tion, 323, 324, 328  
*Liriodendron tulipifera*, 5, 21, 23–4, 25, 26,  
  505; carbon balance, 154; mineral cycling  
  in, 393, 394; photosynthetic efficiency,  
  515, 519, 520; and root organic matter  
  turnover, 320, 321, 322, 326  
Lisbon, Portugal, 184, 185  
litter layer and litter fall: and chemical  
  element input to soil, 301–2, 304; and  
  CONIFER model, 553; and mineral  
  cycling, 349, 351, 354, 356; and produc-  
  tivity, 417–22, 429–31, 439–41; site  
  studies, 376–407; water balance in,  
  223–7, 243–4  
*Lomandra dura*, 7  
*Lonicera* spp., 24  
*Lonicera xylosterum*, 23  
Lotka-Volterra theory, 42  
Lubumbashi, Zaire, 6, 9, 11, 175, 672  
*Luzula pilosa*, 28  
*Lycopodium lucidulum*, 32  
lysimeter techniques, 303–5, 313–14, 315
- Madeleine, France, 3, 7–8, 584  
*Madhuca indica*, 13  
magnesium: atmospheric input to soil, 296,  
  297, 301; cycling of, 346, 353, 355,

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- 356–7, 258–9, 361–2, 365–6; site studies, 376–407
- Magnolia salicifolia*, 21
- Maianthemum bifolium*, 18
- Majanthemum* spp., 28
- Malus sylvestris*, 21
- Manaus, Brazil, 3, 575; radiation balance, 176; tropical rain forests of, 10, 11; water balance, 180, 181
- manganese: atmospheric input to soil, 297
- Marquesia macroura*, 6, 11
- Massa Me, Ivory Coast, 10
- Meathop Wood, England, 5, 647; mineral cycling, 404; radiation flux, 175; temperate forest of, 23; water balance, 190
- Mediterranean woodlands: classification of, 3–8, 30–1; and linear model of biomass dynamics, 473, 475–6; nitrogen accumulation, 353–8, 359; radiant energy, 172–3, 174–6; radiation flux, 130; water balance, 180, 184–5, 186
- Meerdink, Netherlands, 4, 22, 607
- Meliaceae, 11
- Melica uniflora*, 22
- Mercurialis perennis*, 23, 574
- Mercurialis* spp., 21
- Merlewood, Great Britain, 344
- meteorological instrumentation, 106
- Michigan, USA: succession modelling in deciduous-coniferous forest, 53–65 micro-organisms: and nitrogen adsorption into soil, 300, 304; respiration of, 526–8
- mineral cycling, in forest ecosystems, 341–407; additions and losses, 368–74; and annual production, 358–9; and biomass, 350–7; and differences between deciduous and coniferous spp., 360–4; and forest productivity, 364–8, 442; methods used, 345–7; sites and forest types studies, 344, 376–407; and stand maturity, 348–50; *see also* chemical elements
- mineralization studies, of soil, 303–5
- Miombo woodland, Africa, 11, 13
- models, and simulations: COMEX *see* COMEX; CONIFER *see* CONIFER; donorcontrolled linear compartment (DCLM), 452–4; of forest growth and water balance, 537–63; FORET *see* FORET; of forest stand dynamics, 38–52, 538–43; general linear model, 455–69; of physical environment, 111–20; reproduction models, 50–1; stand energy models, 538–43; succession *see* succession modelling; TEEM-Prosper model *see* TEEM; tree simulation models, 43–50; of watershed behavior *see* ECOWAT, PROSPER; of woodland ecosystem composition, 65–79
- Moraceae, 11, 575
- Morus bombycis*, 16
- Mount Disappointment, Australia, 3, 6–7, 573
- Mpanga Reserve forest, Uganda, 183
- Myrianthus* spp., 10
- Myrtaceae, 4, 11
- Myrtillosum* spp., 427
- Nakoma Urban Forest, Wisconsin, USA, 5, 664
- Napoleona* spp., 10
- Narcissus pseudonarcissus*, 574
- Narcissus* spp., 21
- Neolitsea sericea*, 15
- Nephroma arcticum*, 29
- net radiation balance, 94–6, 135–40; above canopy, 135–6; at forest floor, 137–8; long-term measurement of, 138–40; of snowpack, 222–3; vertical profile of, 136–7
- New Brunswick, Canada, environmental factors and woodland diversity, 68–70
- Newnham's model, of tree growth, 46–7
- nitrogen: atmospheric input to soil, 297, 300, 369–70; fixation of, 370–2; and loss by leaching, 372–4; mineralization of, 303–4; and recycling, 346, 348, 349–53, 355–63, 365–6; site studies, 376–407
- Noe Woods, Wisconsin, USA, 5, 24, 663; radiation balance, 176; woodland ecosystems, factors affecting, 65–6, 71–2, 73, 76
- Nothofagus truncata*, 302
- nutrient cycling *see* mineral cycling
- Nyssa sylvatica*, 24
- oak forest: annual energy flow, 154, 156; carbon metabolism, 504, 506–7; radiation and heat flux, 150–1, 153; water balance, 189–91; *see also* *Quercus* spp.
- Oak Ridge, Tennessee, USA, 5, 657–61; carbon dioxide balance, 154; carbon metabolism, 505–6; general linear model for biomass dynamics, 445–7; mineral cycling, 343, 344, 388, 393–5, 396; photosynthetic processes, 510–11, 514, 516–17; radiation balance, 176; root dynamics, 325, 326, 327; temperate forest of, 23, 24
- Okinawa, Japan, 4, 15, 601
- Okita, Japan, 4, 18, 602
- Oleaceae, 9, 10, 11
- Ontario, Canada, 3, 29, 32, 576–9

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- Oregon, USA, 5; succession modelling in coniferous forest, 52–65  
 organic matter: and biomass production, 443–7, 485–6. 487; and nitrogen accumulation, 353, 354–5  
*Ostrya virginiana*, 75  
 Oulu, Finland, 3, 26, 29, 176, 582  
 Oved, Sweden, 4, 23, 621  
*Oxalis acetosella*, 18, 23, 28, 29  
*Oxalis griffithii*, 21  
*Oxalis montana*, 32  
*Oxalis* spp., 28  
*Oxydendron arboreum*, 24  
 oxygen absorption, by soil, 300
- Paeonia peregrina*, 23  
 Palmae, 3, 9, 11, 575  
 Panama tropical forests, 182  
 Paris, France, 187  
*Parrotia persica*, 5, 21  
*Parthenocissus quinquefolia*, 24, 505  
 Pasoh, Malaysia, 4, 8, 10, 11, 175, 606  
*Paltigera aphosa*, 29  
*Paltigera scobrosa*, 29  
 percolation, simulation of, 288–93  
 phosphorus: atmospheric input to soil, 296–7, 301; and cycling of, 346, 355–9, 361–2, 365–6; and litter input to soil, 304; and phosphate equilibria in soils, 309–11; site studies, 376–407  
 photosynthesis: and carbon metabolism, 509–22; and moisture availability, 511–13; photosynthate allocation, 518–22; and Stand Energy model, 541, 542; and TEEM-Prosper model, 548–9  
 photosynthetically active radiation (PAR), flux of, 127–31  
*Phragmites* spp., 21  
*Phryma leptostycha*, 24  
 physical environment, modelling Of, 111–20; atmosphere and plant, 111–16; plant-water-soil relationships, 116–19  
 physiognomy and phytosociology of woodlands, 1–35  
 phytosociology, 1–35  
*Picea abies*, 4–6, 26, 27; biomass, 427; carbon dioxide flux, 160; mineral cycling in, 344, 348–51, 363, 373; radiation flux, 129; water storage by canopy, 215; *see also* spruce forests  
*Picea engelmannii*, 17  
*Picea excelsa*, 3, 4, 22, 29  
*Picea glauca*, 3, 29; and environmental factors, 69–70; radiation flux, 123  
*Picea mariana*, 3, 5, 29; and environmental factors, 69–70; mineral cycling, 344, 376–8  
*Picea pungens*, 129–30  
*Picea rubens*, 3, 14, 29  
*Picea sitchensis*, 17, 123, 229, 234  
*Picea* spp., 26  
 pine forests: carbon metabolism, 504, 507; and chemical input to soil, 302; radiation flux, 149; water balance, 183, 192; *see also* *Pinus* spp.  
*Pinus banksiana*, 142  
*Pinus contorta*, 142  
*Pinus densiflora*, 4, 16, 18  
*Pinus echinata*, 5, 24, 25, 26; mineral cycling in, 344, 388; photosynthate allocation, 519, 520  
*Pinus flexilis*, 17  
*Pinus lambertiana*, 5, 19  
*Pinus longifolia*, 183  
*Pinus ponderosa*, 235, 247  
*Pinus radiata*: biomass dynamics, 477; carbon dioxide flux, 154; heat flux, 142; net radiation balance, 135; Weibull density function of, 43  
*Pinus resinosa*: and global radiation transmission to soil, 127; heat flux, 142; water storage, 215, 228  
*Pinus rigida*, 5, 23, 506  
*Pinus strobus*, 3, 5, 20, 29; biomass dynamics, 477; mineral cycling in, 344, 389, 397; water storage, 215  
*Pinus sylvestris*, 4, 18, 27, 302; heat flux, 142, 147; and mineral cycling, 348; root organic matter turnover, 320; transpiration, 230; water balance, 192, 215  
*Pinus taeda*, 5, 19, 507; carbon dioxide balance, 154; global radiation transmission to soil, 127; heat flux, 147, 148–51; interception, 124; and mineral cycling, 348; photosynthetic processes, 511, 540–3; radiation flux, 122, 131, 132, 135, 138; and root organic matter turnover, 320–1; and water-soil relationships, 117–19  
*Piptadeniastrum africanum*, 4, 10  
*Piptadeniastrum* spp., 10  
 plant, water movement in, model of, 116–19  
*Pleurozium schreberi*, 29  
*Poa pratensis*, 24  
*Polystichum acrostichoides*, 505  
*Polystichum munitum*, 247  
*Polytrichum commune*, 28  
*Polytrichum perinum*, 29  
*Polytrichum* spp., 28  
*Populus deltoides*, 129, 348  
*Populus pruinosa*, 5, 21  
*Populus tremula*, 147  
 potassium: atmospheric input to soil, 296,

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- 297; litter and input to soil, 304; and mineral cycling, 346, 349–53, 355–9, 361–2, 365–6; site studies, 376–407
- precipitation, 249, 266–71, 546; and chemical input to soil, 294–302, 299–300, 315; and productivity, 416–17; site studies, 376–407; *see also* rain, snow
- primary production, and ecosystem diversity, 70–4; *see also* productivity
- production *see* productivity
- productivity, 411–46; and annual average temperature, 415–16; and available light, 422–4; and effects of defoliation using models, 547–52; and litter fall, 417–22; and mineral cycling, 364–8; and precipitation, 416–17; and stand age, 413–15; *see also* biomass
- profile methods: of carbon dioxide flux determination, 104; instrumentation for, 106–8
- PROSPER hydrology model, 242, 245–55, 545–7; *see also* TEEM-Prosper model
- Protium* spp., 10
- Prunus avium*, 3, 21, 574
- Prunus grayana*, 16, 21
- Prunus serotina*, 5, 24
- Pseudomonas* spp., 317
- Pseudoscleropodium purum*, 18
- Pseudotsuga menziesii*, 5, 17, 19, 56, 246; mineral cycling in, 344, 348–9, 350, 363, 380–1, 382; and root organic matter turnover, 322; water potential gradient, 233–4; water storage, 215
- Pteridium aquilinum*, 18, 32
- Ptilium crista-castrensis*, 29
- Pulmonaria rubra*, 23
- Quercus alba*, 5, 20, 23, 24, 505; photosynthetic efficiency, 515
- Quercus acuta*, 16
- Quercus castaneifolia*, 5, 21
- Quercus cerris*, 3, 21, 22
- Quercus coccifera*, 8
- Quercus coccinea*, 5, 21, 23
- Quercus gambelli*, 247
- Quercus ilex*, 3, 8, 130; mineral cycling in, 344, 407; water balance, 186, 195
- Quercus macrocarpa*, and environmental influences, 75, 75
- Quercus mongolica*, 4, 21
- Quercus pedunculiflora*, 4, 21, 23
- Quercus petraea*, 3, 4, 5, 21, 22, 23
- Quercus prinus*, 5, 21, 23, 25–6; mineral cycling in, 344, 396
- Quercus pubescens*, 4, 21, 23
- Quercus robur*, 3–5, 18, 21, 22, 23, 574
- Quercus rubra*, 21, 24, 505, 519, 520, 559
- Quercus salicina*, 16
- Quercus serrata*, 18
- Quercus* spp., 14, 20, 74; and mineral cycling in, 344, 394–5, 399–400, 404; *see also* oak forests
- Quercus velutina*, 5, 21, 23, 25, 75–6, 505
- radiant energy transfer in forests, 140–57, 168–76; air temperature gradients, 142–3; sensible heat, 141–2, 147–52; and Stand Energy model, 540, 541; in turbulent air, 142; vapor tension gradients, 145; *see also* radiation flux
- radiation: and the Earth, 89–91; and forest characteristics, 432–6
- radiation flux, 87–98, 569; from biosphere to space, 91, 93–4; energy transfer in forests, 140–57; entropy flow and entropy production, 97–8; instrumentation, 98, 100–1; long wave, 131–5; net radiation balance, 94–6, 135–40; short wave, 120–31, *see also* global radiation; from space to biosphere, 91, 92–3; and Stand Energy model, 541; 542; *see also* heat balance, radiant energy
- rain, 270–1; instrumentation, 109–10
- reproduction models, of tree growth, 50–1
- respiration: autotrophic, 523–5, *see also* photosynthesis; heterotrophic, 508–9, 525–30; of invertebrates, 529–30; of microflora, 526–8
- Rhizobium* spp., 300
- Rhododendron macrophyllum*, 247
- Rhododendron maximum*, 23
- Rhus ambigua*, 21
- Ribes alpinum*, 23
- Rocklin, California, USA; 184, 185, 186
- roots, dynamics of: biomass, 485–6; organic accumulation, 323–6; sloughing, 327–8; and turnover root organic matter, 320–3; uptake of chemical elements, 316–17
- root zone water, 227–8, 240
- Rosa acicularis*, 29
- Rosaceae, 9, 11, 575
- Rosa sempervivens*, 8
- Rosa* spp., 21
- Rouquet, France, 3, 7–8, 585, mineral cycling in, 344, 407
- Rubiaceae, 3, 11, 575
- Rubus idaeus*, 28
- Rubus saxatilis*, 28
- Rubus spidigerus*, 22
- Ruscus aculeatus*, 8, 22
- Ruscus* spp., 21
- Russia (USSR): boreal forests, 26, 28; temperate forests, 15

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- Saigon, Vietnam, 180, 181  
 Sal Plantation, India, 589–91  
*Salix* spp., 29  
 San Juan, Spain, 4, 18, 615  
*Santiria* spp., 10  
 Sapotaceae, 3, 11, 13, 575  
 saprovores respiration, 529  
*Sasa* spp., 21  
 Saxapahaw, USA, 5, 19, 20; and simulated results of forest stand, 119–20  
*Schefflera octophylla*, 15  
*Schizophragma hydrangeoides*, 21  
*Scilla bifolia*, 574  
*Scilla sibirica*, 21  
 Scots pine forest, 122–3  
*Scottelia chevalieri*, 4, 10  
*Scotellia* spp., 10  
 Scytopetalaceae, 11  
 Site, Seattle, USA, 175  
*Scytopetalum thieghemii*, 4, 10  
 sensible heat flux, 141–2, 145, 146; long-term measurements of, 147–52; in spruce forests, 157–60; in snowpack, 221–3  
 Shigayama, Japan, 4, 16, 18, 175, 603  
*Shorea curtisii*, 182  
*Shorea robusta*, 3, 13, 14, 182–3  
 shrubs *see* individual species  
 Sikfokut, Hungary, 3, 22, 586  
 silicon: and litter input to soil, 301  
 simulations *see* models  
 Sinaia 1, Rumania, 4, 23, 175, 613  
 Sinaia 2, Rumania, 26, 29, 175, 614  
 Siskiyou Mountains, Oregon, USA, and hydrologic cycle, 208, 209  
 sites, of forest ecosystems, 3–6, 573–671; and linear model of biomass dynamics, 473–6; *see also* individual sites  
 snow, 271; and CONIFER model, 553; measurement of, 110  
 snowpack dynamics, 218–23, 243; and CONIFER model, 553; energy and water balance of, 220–3; structure, 218–20  
 sodium: atmospheric input, 296, 297  
 soil, heat flux to, 115–16; global radiation received, 127; water balance in, 223–7, 235–41, 244  
 soil processes, 265–328; chemical elements, dynamics of, 293–318; dynamics of water, 265–93, *see also* water; root dynamics, 318–28  
 soil types *see* individual sites  
 soil water, 283–8; measurement of, 110–11; water balance in, 223–7, 235–41, 244, 280–3; water potentials, 283–4  
 Solling Project, Germany, 6, 666–71; boreal forests of, 26, 28; interception, 124, 125; heat flux, 143, 147, 148–51; mineral cycling in, 344, 385–7, 400–3; net radiation balance, 135–6, 137, radiation flux, 122, 148–51, 176; temperate forests of, 23, 24; water balance, 190, 192  
*Sorbus aucuparia*, 4, 22  
*Sorbus* spp., 23  
 SPAC (soil-plant-atmosphere) as a hydraulic continuum, 272  
*Sphagnum capillaceum*, 32  
*Sphagnum girgenschnii*, 29  
*Sphagnum palustre*, 32  
 spruce forests: canopy drip, 269; carbon dioxide balance, 160, 164–8; heat flux, 147; net radiation balance, 136, 137–8, 148; and nitrogen cycling, 348, 349, 373; sensible and latent heat, 157–60, 161–2; water balance, 192, 270, 292  
*Stachyurus praecox*, 16  
 stand age: effect of, on forests, 413–15; stand height, 425, 428–9  
 stand, dynamics of, 38–52; evolution of growth modelling, 40–51; and mineral cycling, 348–50; and Stand Energy model, 538–43; succession modeling *see* succession modelling; woodland ecosystem composition, 65–79  
 Stand Energy model, of forest growth, 538–43  
*Stellaria holostea*, 21, 28  
*Stellaria nemorum*, 23  
 stem flow, 266–71, 295; site studies, 376–407  
 Sterculiaceae, 11  
*Stipa pubescens*, 7  
 stochastic processes, of stand development, 42–3, 50–1  
 stocking density, 436–9  
 stomatal control, 231–5, 511, 512, 513  
*Strombosia glaucescens*, 3–4, 10  
*Strombosia* spp., 10  
 structural characteristics, 424–32  
*Styrax japonica*, 15  
 subsurface run-off, 271, 277, 278–80  
 succession modelling: in deciduous and coniferous forests, 52–65; landscape dynamics model, 60–3; land-use and forest management scenarios, 63–5; natural models, 54–9  
 sulphur: atmospheric input to soil, 297  
 Sangai Lui Catch, Malaysia, 182  
 Supramonte di Orgosolo, Sardinia, Italy, 125  
 surface run-off, 271, 277  
 Sverdrup method: for determination of carbon dioxide flux, 105, 106; for determination of heat flux, 99, 102

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)

## Index

*Symphytum cordatum*, 23*Syringa vulgaris*, 24

Tallish Oak, Baku, 15

*Tamarix ramosissima*, 21*Tarrietia utilis*, 4, 10*Tectona grandis*, 3, 13, 14, 183, 497

TEEM-Prosper model: of forest water balance, 543–52, 560; and effects of defoliation, 548–9, 550–2

Tellermanovsky, USSR, 190

temperate forests, 14–25, 30–1; carbon budget, 503; classification of, 15, 16–19, 20–5, 30–1; energy flow, 154, 156; and general linear model of biomass dynamics, 473–6, 483; and litterfall, 420–1, 441; mineral cycling in, 344; nitrogen accumulation, 354–4, 355–9; radiant energy, 172–7; stand age, 414; water balance, 185–8, 189–91; *see also* individual sites temperature: and function of a watershed, 210–11; and productivity, 415–16, 435–7; site studies, 376–407

tensiometers, 110–11

*Terminalia tomentosa*, 3, 13*Thelyerania* spp., 21*Thelypteris phegopteris*, 28*Themeda australis*, 7

Thompson Research Center, Seattle, USA, 5, 622; mineral cycling, 380–1, 406; radiation balance, 175; temperate forest of, 19, 20

throughfall, of water, 266–71, 295; measurement of, 110, 243

*Thuja occidentalis*, 3, 29, 69–70*Thuja plicata*, 5, 19

Tigrovaya Balka, USSR, 5, 646; radiation balance, 175–6; temperate forests of, 15, 21

*Tilia cordata*, 4–5, 21, 22*Tilia* spp., 14, 23, 147

time-related influences, and ecosystem composition, 74–9

transpiration, 230–1, 233, 244, 249, 271–2, 274–5, 280, 292, 551–2; *see also* evapotranspiration

Triangle, Carolina, USA, 655; carbon dioxide balance, 154; heat flux, 147, 148–51; photosynthetic processes, 511

*Trientalis europaea*, 28*Trigonobalanus* spp., 11tropical forests, 8–14, 30; classification, 9–14; linear model of biomass dynamics, 473, 475–6; and litter turnover, 441; organic matter, 444; radiant energy, 171, 172–6; water balance, 180, 183; *see also* individual sites*Tsuga canadensis*, 3, 14, 29*Tsuga diversifolia*, 4, 18*Tsuga heterophylla*, 5, 17, 19, 20, 56, 247; mineral cycling in, 344, 392*Tsuga mertensiana*, 56*Tsuga sieboldii*, 4, 16; mineral cycling in, 344, 391

turbulence of air, and heat flux, 142; and Stand Energy model, 538–43; in spruce forest, 159–60, 161–2

*Turreanthus africana*, 3, 10

Ulmaceae, 21

*Ulmus campestris*, 4*Ulmus carpinifolia*, 3*Ulmus scabra*, 5, 21*Vaccinium augustifolium*, 32*Vaccinium brittonii*, 32*Vaccinium myrtilloides*, 32*Vaccinium myrtillus*, 18, 28*Vaccinium uliginosum*, 28, 29*Vaccinium vacillans*, 23, 506*Vaccinium vitis-idaea*, 29

vapor pressure, and hydrologic cycle, 210–11, 212

Varanasi, India, 3, 13

variable density yield tables, and growth modelling, 41

*Viburnum alnifolium*, 23*Viburnum opulus*, 16*Vinca major*, 8

Violaceae, 11, 575

*Viola vaginata*, 21

Virelles, Belgium, 3, 574; air temperature above canopy, 143, 144–5; heat balance, 95; heat flux, 141–2, 147, 148–52; interception, 124, 125; mineral cycling, 399, 400; net radiation balance, 135, 138–9, 175; radiation flux, 121–2, 133–4, 135, 148–51, 152; temperate forest, 21; transmission of global radiation to soil, 127; water balance, 189, 191

von Karmen's constant, 115, 116

Vorskla River sites, USSR; 21

Washington, USA; 17

water, dynamics of, 177–96, 265–93; in boreal coniferous forest, 188, 192–4, 195–6; CONIFER model, 522–8; at continental surface, 178, 179; in equatorial forest, 178–80, 181–2; input variables, 266–71; instrumentation of, 190–11; internal foliage storage, 228–30; measurement in soil, 110–11; in Mediterranean forests, 180–1, 184–6; and movement in plants model, 116–19; output variables,

Cambridge University Press

978-0-521-11220-8 - Dynamic Properties of Forest Ecosystems

Edited by D. E. Reichle

Index

[More information](#)*Index*

- 271–80; simulations, 288–93; TEEM-Prosper model *see* TEEM; in temperate forest, 185, 187–91; in tropical forests, 180, 181, 183; water balance model, 112, 116–19; *see also* hydrological cycles
- watershed, structure and function, 206, 207–42; behavior of, *see* PROSPER, ECOWAT; climatic variation over, 208–13; hydrology of terrestrial strata, 213–41; and weathering of soil, 313
- weathering, of soil, 312–14
- Weibull density function, 43
- wind, and effect on heat flux, 142; and effect on watershed, 213; in spruce forest, 159–60, 161–2
- Woodlands Data Set (WDS), 411–12, 439–47
- Wyoming, USA, 17
- Xanthorrhoea australis*, 7
- Xoiwai, Japan, 175
- Yangambi, Zaire, 176
- Yapo, Ivory Coast, 4, 9–11, 182, 599
- yield tables, for growth modelling, 41
- Yusuhara Kubotaniyama, Japan, 16–18, 604
- Yusuhara Takatoriyama, Japan, 4, 18, 390–1, 605
- Zelkova carpinifolia*, 5, 21
- Zizphus xylophyrus*, 13