

# Index

- ABH (Se) secretor system**, 11  
 in Gujerat, 16  
 in Siberia, 60, 76; under selection pressure of differentiating type, 78
- ABO blood group system**, 9–10  
 in Africa: in calculation of genetic affinities, 198, 199; climate and, 192, 193; in Pygmy–Pygmoid and farmer pairs, 202  
 in Amerindians, B absent, 28  
 associations of, with goitre, 304, with leprosy, 16, and with smallpox, 16  
 in Indian tribes, 15–16  
 in Malaysia, 18  
 in Siberia, 60, 63, 76, 79; distribution of, (A) 80, 82, (B) 80, 83, (O) 80, 84; selection pressure of stabilising type acting on, 78, 79; stabilising selection, mechanism shown by frequency distribution, 88–92, 84  
 in Yanomama, only O present, 113
- acculturation to Western influences**, in Solomon Islands, 147–8, 149, 151–2  
 and blood pressure, 159, 160  
 and serum cholesterol and uric acid, 157
- acetylase system**, in Korea, 19
- acid phosphatase**  
 in calculating genetic affinities of African peoples, 198  
 in circumpolar populations, 29  
 and malarial infection, in Papua New Guinea, 304  
 in Yanomama, 114, 122
- adaptation to environment**, 4
- adenosine deaminase**: search for variants of, in South American tribes, 127–9
- adenylate kinase**  
 in calculating genetic affinities of African peoples, 198  
 search for variants of, in South American tribes, 127–9
- Africa**  
 genetic affinities of populations of centre of, for genes unresponsive to climate, 198–202  
 sets of populations evolved in, 192–3  
 surveys in, 19–21; of peoples of northern semi-arid zone, *see* Dogon, Fali, Fulani, and Kurumba; of rain forest, *see* rain forest
- age**, and blood pressure  
 in African northern semi-arid zone, 260–1  
 in Solomon Islands, 159–61
- Ainu people**, 19, 26
- Åland Islands**, 14
- aldosterone excretion**, by Yanomama, 133
- Aleuts**, 28, 29  
 gene frequency diagrams for two populations of, settled in Komandorski Islands in 1820s, 98–101
- Allen's 'rule'**, on length of protruding body parts at different latitudes, 244–5
- altitude**, and blood-factor genes in Ethiopia, 21
- Amarar people (East Sudan)**: genetic affinities of, show Arab element, 199, 200, 201
- Amerindians**  
 calculation of date of separation of, from Siberians, 51, 101, 103  
 gene frequency polygons for Siberians and, 98–101  
 genetic relations of different tribes of, 114–15  
 resemblances between Siberians and, 28, 29  
 of South America, 109–11; *see also* Yanomama tribe
- Amhara people (Ethiopia)**, 21  
 genetic affinities of, 198, 199, 200, 201
- anaemia**, in Papua New Guinea, 300–1
- angina**, in Tokelau non-migrants, 179
- Arabs, Yemenite**: and Yemenite Jews, 15
- arid zone, Africa**: stature and skull characters in, 190–1
- Arid Zone Research, Negev Institute for**, 224, 233
- arm-folding pattern**, in African northern semi-arid zone, 257–8
- arm length**  
 in African northern semi-arid zone, 250, 251  
 in Pygmy–Pygmoid and farmer pairs, 203, 204  
 related to mean annual temperature, 252
- asthma**: in Tokelau non-migrants, 180, and in children of migrants, 182, 183
- astigmatism**, not found in Solomon Islands, 156
- Australia**, surveys in, 24, 25
- Australia antigen**: variable occurrence of, in Yanomama, 135
- Australian Aborigines**, resemblances between inland peoples of New Guinea and, 25
- auto-immune diseases**, and histocompatibility antigens, 11
- Aymara tribe**, Peru, 23

*Index*

- Bantu-speaking peoples  
 in Angola: hand-clasping pattern in, 257;  
 arm-folding pattern in, 258  
 spread of, 21, 189
- Badik people (East Senegal), genetic affinities of, 198, 199, 200
- Bergmann's 'rule', on relation between size and latitude, 244
- Binga Pygmoids, 273  
 genetic affinities of, 200, 201, 202, 205
- Binga-Mbimu pair of hunter-farmer peoples, in African rain forest, 194  
 body measurements compared, 202-6
- Bira people, genetic affinities of, 205  
*see also* Mbuti-Bira pair
- birth rates, *see* fertility
- blood pressure  
 in African northern semi-arid zone, 259-61  
 in Maoris, and peoples of Pukapuka, Rarotonga, and Tokelau Islands, 166  
 in Papua New Guinea, 301  
 in Solomon Islands: relations with age and weight, 159-60, and with salt intake, 160-1  
 in Yanomama and Cayapo tribes, 133
- body measurements  
 in Africa: different climatic zones, 192, 193, northern semi-arid zone, 249-56; rain forest, Pygmy-Pygmoid and farmer pairs, 202-6  
 of Siberians, 56-9  
 in Solomon Islands, 149, 154-5  
*see also* height, weight, *etc.*
- Bolivia, surveys in, 22, 23
- Bougainville Island, Papua New Guinea, 145-6  
 survey in, *see* Solomon Islands
- Brazil  
 Negroes in: arm-folding pattern among, 258; hand-clasping pattern among, 257  
 surveys in, 22-3, 24
- breast-feeding of infants, in Tokelau migrants and non-migrants, 183
- Burma, surveys in, 17
- Bushmen (San), 20  
 similarities between Pygmies and, 283
- Bushong, Bantu farmers south of rain forest, 190  
 genetic affinities of, 205  
*see also* Cwa-Bushong pair
- cadmium, in blood of Yanomama and in US city samples, 134
- Canary Islands, surveys in, 14
- cardiovascular diseases, very rare in Solomon Islands, 156-7
- cargo cult philosophy, 306
- Caroline Islands (Micronesia), survey in, 26
- Caucasoid people, in North and East Africa, 21
- Cayapo tribe, Brazil, 23, 115, 120, 129
- ceruloplasmin: search for variants of, in South American tribes, 128-9
- chest radiography, in Solomon Islands, 153, 156
- children, health of: in Tokelau migrants to New Zealand, 182, 183, 185
- Chile, surveys in, 22
- Chinese, in Malaysia, 18
- chloride, in blood of Yanomama, 133
- cholesterol in serum  
 increases with acculturation in Solomon Islands, 156-7, 158  
 in Tokelau migrants, 181, 184-5, and non-migrants, 175, 176
- chromosome damage: extent of, in mitotic leucocytes of Yanomama, 132
- Chukchi people, 28
- circumpolar populations, 26-9
- coconuts, in diet on Tokelau Islands, 167, 180, 184
- colour blindness, not found in Siberians, 56
- congenital defects, in Yanomama, 133
- copper, in blood of Yanomama and in US city samples, 134
- cranial abnormalities, in Neolithic and present-day Siberian populations, 95-6  
 gene frequency polygon for, 96-8
- Cwa (Kuba) Pygmies, genetic affinities of, 205
- Cwa-Bushong pair of hunter-farmer peoples, in African rain forest, 190  
 ABO groups in, 202  
 body measurements compared, 202-6
- Czechoslovakia, surveys in, 14
- death rates  
 of children in African northern semi-arid zone, 248-9  
 of Yanomama, 117
- demography, 3  
 in African northern semi-arid zone, 247-9  
 in Papua New Guinea, 293-4, 297  
 of Pygmies, 276-9  
 in Solomon Islands, 148, 149  
 of Tokelau migrants and non-migrants, 172, 185  
 of Yanomama, 116-19
- dendrograms  
 linguistic, for Siberians, 52  
 for South American tribes, 114  
 for Yanomama villages, 122

*Index*

- dental morphology**  
 in Solomon Islands, 147  
 of Yanomama, 113
- dermatoglyphics**  
 in African northern semi-arid zone, 259  
 not correlated with climate in Africa, 192, 193  
 of Yanomama, 113
- diabetes**, in Tokelau non-migrants, 178, 179
- Diego blood group system**  
 in Burma, 17  
 in calculating genetic affinities of South American tribes, 114–16  
 in Eskimos and Aleuts, 28  
 in Siberians, 62, 76; under selection pressure of stabilising type, 78  
 in Yanomama, 114, 122
- diet**  
 changes of, occurring in Papua New Guinea, 304–5  
 energy content of, apparently lower than energy expenditure, on Kar Kar Island, 298  
 of Oto, Twa, and Sara-Majingay, 197, 210  
 in Solomon Islands, Western items in, 151, 152  
 of Tokelau non-migrants, 167, 180, 184  
 of Yanomama, 112
- Dogon people** (semi-arid zone, Mali), 241–2  
 body measurements of, 250; skinfold thicknesses, 255, 256  
 estimate of total number of, 247  
 nutrition of, 246
- Duffy (Fy) blood group system**  
 in Amhara, shows gene flow from Arabia, 198  
 in calculating genetic affinities of South American tribes, 114–16  
 in Siberians, 62, 76  
 in Pygmies, 279
- ecological approach**, 6, 7
- environment**  
 adaptation to, 4  
 population system less affected by, than single populations, 102
- enzymes of red cells**, 9, 10  
 search for variants of, in South American tribes, 127–9
- epidemiological approach**, 5
- ergodicity**: principle of, extended from physics to population genetics, 49, 95
- Escherichia coli*: untypable strains of, from Yanomama, 134
- Eskimo people**, 28–9
- ethical issues**, raised by surveys, 306
- Ethiopia**, mixture of Caucasoid and Negro peoples in, 21
- ethnonyms**, marking gene exchange between populations, 69
- Evenk people** (Tungus-speaking), population sizes and inbreeding coefficients of, 66–7
- exercise**  
 cardiovascular and respiratory response to, in Papua New Guinea, 303  
 gas exchange response to, in Kurdish and Yemenite Jews in Israel, 226  
 heart rate in response to, in Fali and Fulani, 261–3
- Fali people** (semi-arid zone, North Cameroon), 243  
 arm-folding pattern in, 258  
 blood pressure in, 260  
 body measurements of, 250, 251  
 demography of, 247  
 hand-clasping pattern in, 256–7  
 heart rate in response to work in, 262  
 lung function in, 263–6
- fertility rates**, 6–7  
 in African northern semi-arid zone, 248  
 in Papua New Guinea, 297  
 of Pygmies, 278–9  
 of Sara-Majingay: body measurements and, 212–13; compared with Oto and Twa, 197  
 of Yanomama, 117
- filariasis**  
 in Kar Kar Island, 302  
 in Pygmies, 280
- Finland**, Finns, 14, 27, 28
- fluid balance**: usually negative in Papua New Guinea (made up by metabolic water), 303
- Fulani people** (semi-arid zone, North Cameroon)  
 arm-folding pattern in, 258  
 body measurements of, 251  
 hand-clasping pattern in, 256–7  
 heart rate in response to work in, 262  
 lung function in, 263–6
- gene frequency**, large variations between Yanomama villages in, 119–20
- gene frequency distribution**, stationary mathematical theory of, 87–8  
 in Siberian population system, 88–9
- gene frequency polygons**  
 for cranial abnormalities, Neolithic and present-day Siberian populations, 96–8  
 for 28 genes, Siberian and Amerindian populations, and two Aleut populations

*Index*

- settled in Komandorski Islands in 1820s, 98–102
- gene frequency variance, in Siberian populations  
as measure of genetic differentiation, 74–9  
predictions of, 73–4
- gene migration coefficients, between Siberian populations  
island model, 71–2, 78  
stepping-stone model, 72–3, 78
- gene migrations in Siberia, 67–70  
equilibrium between random drift and, 79, 88
- genealogies, of Tokelau migrants to New Zealand, 170–1, 185
- genetic distances between populations, 12–13, 18, 20, 22, 100–1  
between Central African peoples, 198–202  
between villages of a South American tribe, 120–1  
between Yanomama and other tribes, 114–16  
calculated from different sets of characteristics, 123–5
- genetics of populations, the evolution and stability of a population in the light of, 48–50
- genogeography, 49  
of Siberian populations, 79–80, 85–6
- Gerbich-negative gene, in New Guinea, 25, 296
- Germany, surveys in: East, 14; West, 13–14
- globulins  
 $\gamma$ , in blood of Pygmies, 280  
unusual, in Yanomama, 131
- glucose-6-phosphate dehydrogenase of blood, 10  
in Greece, 14  
in indigenous Kurds and Kurdish Jews, 15  
in Korea, 19  
in Papua New Guinea, and malarial infection, 304
- goitre  
among Majingay (absent from Twa–Oto), 197  
associations between ABO system and, 304  
on Kar Kar Island, but not at Lufa (Papua New Guinea), 300
- gout, in Tokelau males, 175, 179, 184
- Greece, surveys in, 14
- Greenland, Scandinavian genes of Eskimos in south of, 29
- group-specific component  
incidence of, in cases of leprosy, 17  
in Siberians, 62, 76; under selection pressure of differentiating type, 78
- growth hormone: normal in blood of Pygmies, but some evidence for decreased responsiveness to, 281, 283
- Guaymí tribe (South America), and Yanomama, 114, 116
- Gujerat, tribes in, 15, 16
- Hadza people, Tanzania, 20–1
- haemoglobin  
blood content of, in Solomon Islands, 153, 154–5  
polymorphisms of, 9, 10
- haemoglobin A: search for variants of, in South American tribes, 128–9
- haemoglobin E, in Senoi of Malaysia, 18
- haemoglobin J, in Papua New Guinea, 296  
and malarial infection, 304
- haemoglobin S  
in Bantu-speaking peoples, 21  
in New Guinea, 25  
not correlated with climate in Africa, 192, 193  
in Pygmies, 280  
and susceptibility to malaria, 13  
in tribes of Tamilnadu and Gujerat, 16
- haemoglobin  $\Delta$ , in Pygmies, 280
- hair  
colour of, in Siberians, 57, 59  
in Solomon Islands, 146, 147
- hand-clasping pattern, in African northern semi-arid zone, 256–7
- haptoglobins  
of Ainu, 19  
in calculating genetic affinities: of Central African peoples, 198; of South American tribes, 114–16  
lack of, in many Pygmies, indicating haemolytic anaemia caused by parasites? 280  
in Micronesia (Caroline Islands), 26  
in Papua New Guinea, and malarial infection, 304  
in Siberia, 62, 76; compared with Amerindians, 28; under selection pressure of stabilising type, 78
- health  
in children of Tokelau migrants, 183, 185  
in Papua New Guinea, 293–5, 299–302; social status and, 304–5  
of Pygmies, 280–1  
in Solomon Islands, 152–3
- heart rate in response to work, in Fali and Fulani, 261–3
- heat stress  
response of Kurdish and Yemenite Jews in Israel to, 226

*Index*

- response of Papua New Guinea populations to, compared with Europeans, 302, 303, 308
- height (stature) of adults  
 in Africa, Central: increases with air temperature and dryness, 191, 192; in Pygmy–Pygmoid and farmer pairs, 203, 204, 281–3; relation of weight and, 206–10  
 in Africa, northern semi-arid zone, 250, 251; and lung function, 264–5  
 in Solomon Islands, 154–5  
 in Tokelau non-migrants, 176  
 trend towards increase of, with acculturation, in Solomon Islands, 149, 150, 152  
 of Yanomama, 113
- hip width  
 in Africa, Central: increases with air moisture, 191, 192; in Pygmy–Pygmoid and farmer pairs, 203, 204  
 in Africa, northern semi-arid zone, 250, 251
- histocompatibility antigens, 9, 11
- humidity: correlation between body measurements and, Central Africa, 190–2, 214
- Hungarians, 27, 28
- hunting territories, of Pygmies, 277
- Hutu people (Rwanda)  
 genetic affinities of, 199, 200, 201  
 weight–height relations in, 210
- Iceland, surveys in, 28
- immunoglobulin Gm groups of plasma, 10  
 in Ainu, 19  
 in Amhara, show gene flow from Arabia, 198  
 in Pygmies, 279  
 in Senoi of Malaysia, 18  
 in Yanomama, 113
- inbreeding  
 coefficients of, for Evenk population, Siberia, 66–7  
 in Yanomama, 122–3
- India, surveys among tribes in, 15–16
- infanticide, among Yanomama, 117
- interdisciplinary approaches, 6
- intestinal parasites  
 in Pygmies, 180  
 in Solomon Island peoples, 153, 154–5
- Ireland  
 Icelandic connections with, 28  
 surveys in, 14
- Irian Barat, surveys in, 25
- iron tools, obtained by Pygmies from Bantu farmers, 188, 275
- isocitrate dehydrogenase: search for variants of, in South American tribes, 127–9
- isolates (isolated populations)  
 in equilibrium with environment, 47–8  
 population genetics of, 48–50
- Israel: study of environmental and genetic factors in physiological responses of individuals in, (original form) 219–22, (modified form) 222–4, (conclusions) 238–9  
 duration of, 232–3  
 field work in, 224–5, 227–8  
 laboratory work in, 225–6  
 organization of, 228–32  
 results of first stage of (Negev), 233–5  
 second stage of (Judean Hills), 236–8
- Japan, survey in, 19
- Jews, Kurdish and Yemenite  
 hand-clasping patterns in, 257  
 in Israel, genetically dissimilar, and living in adjacent villages: Judean Hills, 236; Negev, 223  
 two physiological responses of, not affected by genetic differences, 235, 238
- Kar Kar Island, Papua New Guinea, 287–8, 290, 292  
 Kaul village in, 289  
 surveys in, *see* Papua New Guinea
- Kell blood group system  
 in Siberians, 62, 76; under selection pressure of stabilising type, 78  
 in Yanomama, 113
- Khoikhoi people (South Africa), genetic affinities of, 198, 199
- Khoisan-speaking people (South Africa), 20, 193, 198, 199, 200
- Kidd gene, in calculating genetic affinities of South American tribes, 114–16
- Konda people, *see* Twa-Oto pair
- Korea, surveys in, 19
- Kuba-Cwa people, *see* Cwa
- Kung people (South Africa), genetic affinities of, 198, 199
- Kurds of Iran, and Kurdish Jews, 15
- Kurumba people (semi-arid zone, Upper Volta), 242  
 arm-folding pattern in, 257–8  
 blood pressure in, 260  
 body measurements of, 250; skinfold thicknesses, 255, 256  
 demography of, 247–9  
 hand-clasping pattern in, 256–7  
 lung function in, 263–6  
 nutrition of, 245–6
- kwashiorkor, among Pygmy infants, 281

Cambridge University Press

978-0-521-11262-8 - Population Structure and Human Variation

Edited by G. A. Harrison

Index

[More information](#)*Index*

- lactate dehydrogenase: search for variants of, in South American tribes, 127–9
- languages
- in African rain forest, 189; of Pygmies, 277
  - in Kar Kar Island, Papua New Guinea, 291–2
  - in Solomon Islands, 147
  - tracing Siberian population genealogy from, 51, 52
  - see also* Yanomama language
- Lapps, 14, 28
- are something other than Caucasoid-Mongoloid hybrids, 27
- lead, in blood of Yanomama and US city samples, 134
- leg length
- in African northern semi-arid zone, 250, 251
  - related to mean annual temperature, 252
- leprosy, associations of genetic markers with, 16, 17
- Lewis blood group system, in Siberians, 60, 76
- life expectations, Yanomama, 117, 118–19
- Lufa, in New Guinea mountains, 287, 292
- village scene in, 291
- surveys in, *see* Papua New Guinea
- lung function
- in Fali, Fulani, and Kurumba, 263–6
  - in Papua New Guinea, 303
- Maca tribe, Paraguay, 23
- Macushi tribe, South America, 115, 129
- Madagascar, 21
- Maharashtra, tribes in, 15, 16
- Majingay group of Sara people (savanna of South Chad), 190, 195–7
- genetic affinities of, 198, 199, 200
  - goitre among, 197
  - nutritional status and work capacity of, 210–11
  - skin reflectance in, 213–14
  - stabilising and directional selection in, 211–13
- Makiritare tribe, Venezuela, 23, 115, 120, 129
- Malaita Island, British Solomon Islands Protectorate, 145, 146
- survey in, *see* Solomon Islands
- malaria
- and anaemia, 300
  - association of, with genetic markers, 304
  - haemoglobin S, and susceptibility to, 13
  - in Papua New Guinea, 300, 307
  - in Pygmies and farmers of same area, 280
  - in Sardinia, 14
- malate dehydrogenase
- fast-migrating form of, in New Guinea, 25
  - search for variants of, in South American tribes, 127–9
- Malaysia, surveys in, 17–18
- Maoris
- blood pressure, and serum cholesterol and uric acid in, 175
  - hypertension, coronary heart disease, obesity, diabetes, and gout among, compared with Europeans in New Zealand, 166, 178, 184
- Mari people (USSR), Finno-Ugric speaking, 27, 28
- mating patterns, and gene frequencies, 3–4, 13
- Mbimu people, genetic affinities of, 205
- see also* Binga-Mbimu pair
- Mbuti Pygmies, 273
- genetic affinities of, 200, 201, 202, 205
- Mbuti–Bira pair of hunter–farmer peoples, in African rain forest, 190, 194
- ABO groups in, 202
  - body measurements compared, 202–6
- measles: response of Yanomama to, and to vaccine for, 132–3
- Melanesia, compendium of genetic data for, 25–6
- mercury, in blood of Yanomama and in US city samples, 134
- MNSs blood group system
- in Burma, 17
  - in calculating genetic affinities of South American tribes, 114–16
- M: in Siberians, 79–80, 81, (compared with Amerindians) 28, (frequency distribution) 88–9; in South America, 24; in South-east Asia and New Guinea, 25
- MN: in calculation of genetic affinities of Central African peoples, 198, 199; in Siberians, 60–1, 79
- N: in Micronesia (Caroline Islands), 26; in New Guinea and Australia, 25
- Ss: in Siberians, 60–1
- morbidity patterns, 6
- mutations: attempts to study, among Yanomama, 125–6, 127–9
- Mycobacterium tuberculosis*: positive reactions to, and to atypical mycobacteria, in Solomon Islands, 153, 156
- myopia, not found in Solomon Islands, 156
- Naga people, Burma, 17
- natural selection, 2–3
- attempts to study, among Yanomama, 125–7
  - detection of effects of, 13
  - and gene marker systems, 7

Cambridge University Press

978-0-521-11262-8 - Population Structure and Human Variation

Edited by G. A. Harrison

Index

[More information](#)*Index*

- Negro peoples of Africa, 20–1
- Neolithic period, frequency distribution of genes in Siberian populations of present day and of, 96, 97, 103
- net hunting, by Pygmies, 277, 278
- Neu Guinea Kompagnie, 286, 288
- New Guinea, *see* Papua New Guinea
- nicotinic acid, in serum of rural and urban Pedi and Venda males, 322, 323
- Nilotes, of Sudan, 20
- nose breadth, correlated with climate in Africa, 192, 211
- nutrition  
in Africa: northern semi-arid zone, 245–7; rain forest (hunter-farmer pairs), 210–11; Sara–Majingay group, 210–11  
in Papua New Guinea, 297–9  
in Solomon Islands, 152–3
- ophthalmological surveys  
in Negev, 235  
in Solomon Islands, 156
- Orkney Islands, population of Westray in, 27
- Oto (Konda) people, genetic affinities of, 205  
*see also* Twa–Oto pair
- ovalocytosis  
in Burma, 17  
in Senoi of Malaysia, 18
- P blood group system, in Siberians, 62, 76, 80, 85  
integrating role of gene migration in frequency distribution of, 93  
under selection pressure of differentiating type, 78, 79
- Pacific Islands, surveys in, 26
- Palaeolithic period, separation of Siberians and Amerindians during, 98–101, 103
- Papua New Guinea, surveys of Kar Kar and Lufa populations in, 25, 286–7, 304–8  
body measurements, genetic markers, 296–7  
demography, 297  
health, 293–5, 299–302  
nutrition, 297–9; changes in, 304–5  
physiological tests, 302–4
- parotid glands, symmetrical enlargement of, in Lufa population (Papua New Guinea), 300
- Pedi, Bantu people, chosen for survey (males 20–50), 316  
demography: rural, 316–17, 317–18; urban, 317, 318–19  
genetic affinities of, 198, 199  
results of survey for rural and urban populations: age distribution, 320; biochemical data, 321; disease patterns, 325; nutritional status, 322, 323; perceptual and conceptual tests, 323–4; work capacity, 323
- peptidase  
A and B: search for variants of, in South American tribes, 127–9  
C: in Pygmies, 280
- permafrost: border between continuous and irregular, associated with change in concentration of P<sub>1</sub> gene, 80, 85, 86
- Peru, surveys in, 22, 23
- phenyl alanine, in plasma of Pygmies, 281
- phenyl thiocarbamide, ability to taste, 11, 56  
in circumpolar populations and Amerindians, 29  
in Indian tribes, 16  
in Siberians, 62, 76  
in South American tribes, 24
- phosphoglucomutase  
in calculating genetic affinities of Central African peoples, 198  
in circumpolar populations, 29  
in Pygmies, 280  
search for variants of, in South American tribes, 127–9
- 6-phosphogluconate dehydrogenase  
in calculating genetic affinities of African peoples, 198  
at different altitudes in Ethiopia, 21  
search for variants of, in South American tribes, 127–9
- phosphohexose isomerase: search for variants of, in South American tribes, 127–9
- phylogenetic techniques, computer-based, of Cavalli-Sforza and Edwards, 13, 22
- Piaroa tribe, South America, 115, 129
- plasma proteins, 9, 10  
search for electrophoretically detectable variants (mutations) of, in South American tribes, 127–9
- Poland, surveys in, 14
- political issues, raised by surveys, 306
- polygamy, and fertility  
in African northern semi-arid zone, 248  
in Papua New Guinea, 297
- populations  
concept of effective size of, in calculations of genetic structure and outcomes, 130–1  
genetic distances between, *see* genetic distances  
as immortal super-organisms, 48  
system of, descended from a common ancestral population, for study of genetic dynamics, 86–7, 94, 96, 102–3

Cambridge University Press

978-0-521-11262-8 - Population Structure and Human Variation

Edited by G. A. Harrison

Index

[More information](#)*Index*

- potassium  
 in blood of Yanomama, 133  
 urinary excretion of, by Tokelau non-migrants, 177, 178
- pottery, obtained by Pygmies from Bantu farmers, 188, 275
- protein, in diet in Papua New Guinea, 299
- Pygmies of African rain forest (hunter-gatherers), 20, 273–5  
 ABO groups in, 202  
 compared with their suzerain peoples (body measurements), 202–6  
 demography of, 276–9  
 as earliest inhabitants, now living as vassals to Bantu farmers, 188–90, 275  
 genetic studies on, 279–80  
 languages of, 276  
 nutrition of, 281  
 parasite infection of, 280  
 stature of, 281–3, and skull characters, 191  
*see also* Cwa, Mbuti, Twa
- Pygmoids of African rain forest (hunter-gatherers), defined in socio-economic terms, 189  
 ABO groups in, 202  
 compared with their suzerain peoples (body measurements), 202–6  
*see also* Binga
- Quechua tribe, Peru, 23
- rain forest, African  
 migration of populations into, 5, 187–8  
 Pygmy and Pygmoid hunter-gatherers as original inhabitants of, Bantu farmers as later arrivals, now living in vassal-suzerain pairs, 188–90  
 stature and skull characters in, 191  
 stature and weight in, 209  
 surface-weight ratio for populations of, 252–3
- renin, in plasma of Yanomama, 133
- respiratory disease, chronic non-specific; in Papua New Guinea, 301
- Rhesus blood group system  
 in Amhara, shows gene flow from Arabia, 198  
 in Burma, 17  
 in calculating genetic affinities: of African peoples, 198, 199; of South American tribes, 114–16  
 in Micronesia (Caroline Islands), 26  
 in New Guinea and Australia, 25  
 in Papua New Guinea, 296  
 in Pygmies, 279  
 in Siberians, 61, 76, 78; compared with Amerindians, 28  
 in South American tribes, 24  
 Romania, surveys in, 14
- salt intake  
 and blood pressure, 160–1  
 and serum uric acid, 157–8  
 in Solomon Islands, 158, 161  
 Sandawe people, Tanzania, 20–1
- Sara people  
 body measurements of, 207, 251  
 fertility data for, 248  
 genetic affinities of, 205, 206  
 stabilising and directional selection in, 211–13, 215  
*see also* Majingay
- Sardinia, surveys in, 14
- savanna, Africa  
 stature and skull characters in, 190–1  
 stature and weight in, 208  
 surface-weight ratio of populations of, 252–3
- Scotland, Icelandic connections with, 28
- Senoi people (Burma and Malaysia), 17, 18
- serum albumin  
 'private' polymorphism of, in Yanomama, 114, 122  
 search for variants of, in South American tribes, 128
- Shangan people (South Africa), genetic affinities of, 198, 199, 200, 201
- shoulder width  
 increases with air moisture, in Africa, 191, 192  
 in African northern semi-arid zone, 250, 251  
 in Pygmy–Pygmoid and farmer pairs, 203, 204
- Siberia: isolated indigenous populations of, chosen for study, 50–2; planning of study, 52–4; sampling of populations, 54–5; techniques used, 55–6; conclusions, 102–3  
 duration of stable equilibrium in, 95–102  
 gene frequencies in: predicting variance of, 73–4; stationary distribution of, 86–94; variance of, as measure of genetic differentiation, 74–9  
 gene migrations in, 67–73  
 genetic-anthropological characteristics of, 56–63  
 genogeography of, 79–80, 85–6  
 sizes of indigenous populations (1959), 63–7
- skin colour  
 in African northern semi-arid zone, 245  
 in Sara, Oto, and Twa, 213–14  
 in Solomon Islands, 146, 147



Cambridge University Press

978-0-521-11262-8 - Population Structure and Human Variation

Edited by G. A. Harrison

Index

[More information](#)*Index*

- skinfold thicknesses, for estimates of body fat, 253–5  
   in Africa: rain forest, 210; semi-arid zone, 255–6  
   in Solomon Islands, 154–5  
 skull characters (head, face, and nose breadths)  
   increase with air moisture and decrease with air temperature, 191, 192  
   in Pygmy–Pygmoid and farmer pairs, 203, 204  
 smallpox, relative incidence of ABO blood groups in cases of, 16  
 societies: demographic characteristics of ‘young’ and ‘primitive’, compared with those of USA, 148, 149  
 sodium  
   in blood of Yanomama, 133  
   urinary excretion of, by Tokelau migrants, 181, 184, and non-migrants, 177, 178, 184  
 Solomon Islands (British Solomon Islands Protectorate, and Bougainville Island, Papua New Guinea), study of eight populations in, 143–6; future of project, 161–2  
   biological variation in, related to history and geography, 146–7  
   blood pressure in, 158–61  
   body measurements in, 149, 154–5  
   cardiovascular disease in, 156–8  
   chest radiography in, 153, 156  
   demography in, 148, 149  
   health and nutrition in, 152–3, 154–5  
   salt intake in, 158  
 Sotho people (Lesotho), genetic affinities of, 200  
 South Africa, study of adaptation to urbanisation in, 313–15  
   choice of groups for study, 315–16  
   survey: of rural groups, 316–17; of urban groups, 317–19; results of survey, 320–5; evaluation of survey, 325–8  
 South America, surveys in, 22–4  
   *see also* Yanomama  
 Spain, surveys in, 14  
 spleen, enlargement of  
   in Negev, 235  
   in Papua New Guinea, associated with genetic markers, 304  
   in Solomon Islands, 153, 154–5  
 stress, social and psychic, experienced by migrants, 169  
*Strongylocentroides füllebornii*, new parasite found in Pygmies and other Africans, 280  
 superoxide dismutase gene, in populations of Scandinavian-Finnish descent, 27  
 surface area of body: ratio of, to weight, in different African climatic zones, 207, 213, 252–3  
 Surinam, survey of Negroid population in, 24  
 sweat production in response to heat  
   in populations of Papua New Guinea, compared with Europeans, 303  
   in tests on Kurdish and Yemenite Jews in Israel, 226  
 Tamilnadu, India: tribes in, 15–16  
 temperature, mean annual: correlations between body measurements and, in Africa  
   Central, 190–2, 214  
   northern semi-arid zone, 244, 252  
 Thailand, association of genetic markers with leprosy in, 16–17  
 Tokelau Islands, 167; settlement in New Zealand of migrants from, 5, 165–6, 168  
   basis for study of migrants and non-migrants, 68–70; development of study (1966–73), 170–1; methods of study, (anthropological) 172, (medical) 172–4; populations studied, 171–2  
   data from study: anthropological, 183; diet of non-migrants, 181–3; medical, for migrants, 181–3, and non-migrants, 174–80  
   discussion of results of study, 183–5  
 transferrins  
   in calculating genetic affinities of African peoples, 198, 199  
   in circumpolar populations, 28  
   in Siberians, 62, 76; under selection pressure of stabilising type, 78  
 triglycerides, in serum of Tokelau migrants, 181, 185, and non-migrants, 175, 176, 181  
 tropics, human adaptability in, 286–7  
 Tswana people (South Africa), genetic affinities of, 198, 199  
 tuberculosis-like disease, in Solomon Islands, 153, 156  
 Tutsi people (Rwanda), genetic affinities of, 199, 200, 201  
 Twa (Konda) Pygmies, 275  
   genetic affinities of, 205  
 Twa–Oto pair of hunter–farmer peoples, African rain forest, 190, 194–5  
   comparisons between: body measurements, 202–6; nutritional status and work capacity, 210–11; skin reflectance, 213–14

Cambridge University Press

978-0-521-11262-8 - Population Structure and Human Variation

Edited by G. A. Harrison

Index

[More information](#)*Index*

- urban environments, effects of, 8
- uric acid in serum  
 acculturation and, in Solomon Islands, 157, 158  
 salt intake and, 157–8  
 in Tokelau migrants, 181, and non-migrants, 175, 177, 184; correlated with weight, 184
- Veddas, Ceylon, 17
- Venda, Bantu people, chosen for survey (males 20–50), 316  
 results for rural and urban populations:  
 age distribution, 320; biochemical data, 321; demography, 316–17; disease patterns, 325; nutritional status, 322, 323; perceptual and conceptual tests, 323–4; work capacity, 323
- Venezuela, surveys in, 23
- vitamin C, in serum of rural and urban Pedi and Venda males, 322, 323
- Wapishana tribe, South America, 115, 129  
 weight of adults  
 in Africa, Central: relation of height and, in open country and rain forest, 206–10  
 in Africa, northern semi-arid zone, 250, 251; seasonal changes of, 246–7  
 declines with age: in Papua New Guinea, 297; in Solomon Islands, 159, 160  
 inversely related to mean annual temperature, 244  
 ratio of surface area to, in different climatic zones of Africa, 253–4  
 of rural and urban Pedi and Venda males, 322  
 in Solomon Islands, 154–5; related to blood pressure, 159  
 in Tokelau migrants, 181, and non-migrants, 175, 176; correlated with serum uric acid, 184
- work capacity  
 of hunter–farmer pairs, African rain forest, 210–11  
 of rural and urban Pedi and Venda males, 322, 323, 324
- Xavante people, Brazil, 22, 120, 121
- Yanomama language  
 calculation of date of separation of, from neighbouring languages, 113  
 evolution of local varieties of, 129–30
- Yanomama tribe, Brazil, 23, 111–13  
 attempts to study selection and mutation in, 125–9  
 comparison of genetic distances for, based on different sets of data, 123–5  
 concept of effective population size applied to, 130–1  
 demography of, 116–19  
 genetic markers in, showing lack of admixture with Caucasoids or Negroids, 113–16  
 microdifferentiation and population structure in, 119–23  
 plans for further observations on, 135–6
- yaws  
 endemic among Pygmies, 280  
 in Kar Kar Island, 302
- Yupa tribe (South America), and Yanomama, 114