

INDEX

- Aardwolf. *See* *Proteles cristata*
- Absolute configuration, 309
- Acarosides (nematode pheromones)
potential use, control, 266
- Accessory gland proteins (Acps), 88
- Accessory olfactory bulb (AOB)
combinatorial processing, 179
- Accessory olfactory system (AOS), 193
- (5*R*,6*S*)-6-Acetoxy-5-hexadecanolide, 106
- Acomys cahirinus* (spiny mouse)
kin recognition cues, 131
- Acromyrmex* (leaf-cutter ant)
genetic CHC differences between patrines, 130
- Acromyrmex insinuator* (ant), CHC insignificance, 257
- Actias luna* (silk moth)
antenna as sieve or paddle, 184
- Active space, 227
- Acyrtosiphon pisum* (pea aphid)
alarm pheromone
ecological selection on response, 168
induces winged offspring, 169
suicide hypothesis, 168
- Adoxophyes honmai* (smaller tea tortrix moth), resistance to
pheromone control, 272
- Aeolidia papillosa* (nudibranch sea slug)
predator marked by prey pheromone, 167
- Aethia cristatella* (crested auklet, bird)
odor concentration & male rank, 75
- African mole rats. *See* *Heterocephalus glaber* (naked mole rat)
- Aggregation pheromones
Allee effects, 105
aposematic insects, 107
bark beetles, 107
larval pheromones, 108
definition, 105
dilution of risk, 105
ecophysiological benefits, 107
in space
feeding advantages, 110
for defense, 105
oviposition pheromones, 105–106
settlement of marine invertebrates, 109
reproductive benefits, 109
in time (synchronization)
barnacle egg hatching pheromone, 110
co-ordinating external fertilization, 69, 110
larval release, 110
intra-specific eavesdropping
males only signal until females arrive, 109
use in pest monitoring, 263
- Agrotis ipsilon* (black cutworm moth)
male response activated by juvenile hormone, 207–208
males stop responding to female pheromone after
mating, 43
- Agrotis segetum* (turnip moth)
regional variation in pheromone blend and co-varying male
receptors, 92
- Ailuropoda melanoleuca* (giant panda)
captive breeding, 262
male handstand, honest signal, 35
- Alarm cues, 166–167
behavioral responses, 166
fish
“Schreckstoff”, 166
chondroitin, 166
hatchery trout learn predator odors, 262
neural circuits, 166
inter-specific responses, 166
public chemical information, 165, 166
survival benefits to responding, 166
use in pest management, 269
- Alarm pheromones
alert signals, 165
aphids, 167
brain processing, 183
common lack of species specificity, 168
costs to responding
aphid drop, 168
diffusion from source, ants, 229
evolution from pre-existing chemical cues, 165, 170
evolution in unrelated individuals. *See* Alarm cues
fish. *See* Alarm cues; fish
larval insects in family groups, 165
multicomponent, 229
ant, 229
predator labeling with host pheromone, 167
propaganda, 251–258
social insects, 169–172
evolved from defensive molecules, 20–21
social species, 169
suicide hypothesis (aphids), 168
use for pest management, 270
- Alcelaphus buselapus cokei* (Coke’s hartebeest)
self-marking and presentation, 120

- Aleochara curtula* (rove beetle)
 she-male anti-aphrodisiac, 87
- Alkaloids, 76
- Alkylmethoxypyrazine, 107
- Allee effects, 105
 defense against predation, 105
 definition, 105
 ecophysiological benefits, 107
 feeding, 110
 for control, invasive pests, 267
 for wildlife conservation, 262
 mating, 109
 external fertilization, 109
 internal fertilization, 109
 overcoming tree defenses
 bark beetles, 107
- Allelochemicals
 definition, 6
- Allocholic acid, 24
- Allohormone pheromones, 45, 88–89, 213
- Allomarking, 40, 132
- Alloparental care, 130, 146, 219
 egg dumping, 248–249
- Allostatins, 212
- Alpheus heterochaelis* (snapping shrimp) modulation of visual
 threat signals by female pheromones, 44
- Alternative mating strategies, 86–87
 eavesdropping parasitoids select for, 246
 satellite males, 246
 she males, 87
- Altruistic behavior toward kin, 126
- Alzheimer's disease
 sense of smell, loss, 283
- Ambystoma tigrinum* (tiger salamanders)
 male condition vs. MHC, mate choice, 86
- Amniotic fluid, 16, 130, 216, 282
- Amphibians
 alarm cues
 tadpoles, learn predator odors and dangerous times
 of day, 167
 pest management (cane toad), 269
- Amphid sensillae (nematodes), 210
- Amphiprion* spp. (anemonefish)
 symbiotic with sea anemones, 249
- Amygdala
 integration of main olfactory and VNO outputs, 199–201
- n*-Amyl acetate, 185
- Anastrepha ludens* (fruit fly)
 oviposition marking pheromone, 110
 pest management, 269
- Anastrepha oblique* (fruit fly)
 pest management, 269
- Anastrepha suspensa* (Caribbean fruit fly)
 exaggerated signaling limited by predation, 37
 lekking, 79
- low cost of pheromone signaling, 34
- Andrena nigroaenea* (solitary bee), duped by
 orchid, 252
- Androstadienone (androsta-4,16-dien-3-one), 71
 “putative human pheromone”, 296–299
 perception, 294–295, 297–298
- Androstenol (5 α -androst-16-en-3 α -ol), 290
 “putative human pheromone”, 296–297
- Androstenone (5 α -androst-16-en-3-one)
 “putative human pheromone”, 296–299
 perception, 294–295, 297–298
- Anemonefish and sea anemones, mutualism, 249–250
- Animal welfare, 262–263
 animal husbandry, 263
 lab mice, 262
- Anomala osakana* (Osaka beetle), 25
- Anonymous, pheromones, 7
- Anosmias. *See also* under Humans:olfaction
 specific, 293
- Antagonists (inhibitors), 179
- Ant-decapitating flies (phorids), 246
- Anteroventral periventricular (AvPv) nucleus, 201
- Anthonomus grandis* (boll weevil)
 low cost of pheromone signaling, 34
- Anthopleura elegantissima* (sea anemone)
 alarm pheromone, 167
- Anthopleurine, 167
- Anti-aphrodisiac pheromones, 87–88
 “chemical mate-guarding”, 87
 eavesdropped by parasitoids, 245
- Antifouling coatings, 270
- Ants
 colony recognition, 129–130
 enemy specification, 172
 processing alarm pheromone, 183
 queen pheromones
 primer effects, 212–213
- Aphaenogaster (Novomessor)* (ant)
 multimodal signals
 recruitment for foraging, 44
 recruitment, 157
- Aphaenogaster cockerelli* (ant)
 queen detects cheating workers, 37
- Aphids
 alarm pheromones, 167–169
 (*E*)- β -farnesene, 168
 ecological selection on response, 168
 induce winged offspring, 169
 not species specific, 24
 social species, 169
 suicide hypothesis, 168
 use for pest management, 270
 sex pheromones (multicomponent, species-specific), 24
- Apis cerana japonica* (Japanese honeybee), 170–171
 co-ordinated defense against hornet, 170–171

- Apis mellifera* (honeybee)
- alarm pheromone
 - immediate and longer term effects, 208
 - anarchistic colonies, 142, 214
 - bee-keeping
 - using pheromones to manage swarms etc, 260
 - blending of colony odor on comb wax, 128
 - brood pheromone, 136, 145, 214
 - effects on worker nurse–forager transition, 214–215
 - cell capping, pheromones, 136, 145
 - colony odor sources, 128
 - colony recognition, 127–128
 - complexity of pheromones in colony, 134
 - control of worker reproduction
 - maintaining, 214
 - differences between queens and workers, 213
 - gene expression, 214
 - larval pheromones
 - E- β -ocimene, 214
 - multiple paternity in nest, 128
 - olfactory receptors
 - AmOR11, male receptor for 9-ODA, 62
 - large number of different, for an insect, 189
 - pheromone responses change with age, 43
 - hormones underlying, 208
 - pheromones lead to changes in worker bee brain gene expression, 62, 172
 - pheromones summary, 142–145
 - queen mandibular pheromone (QMP), 142, 214
 - as sex pheromone, 135
 - effects on worker nurse–forager transition, 214–215
 - multiple signals, parsimony, 25
 - primer effects, 43
 - rapid colony response to loss, 141, 144
 - releaser effects, sex pheromone, 43
 - swarming, 145
 - transmission, 141, 144
 - queen pheromones
 - mechanisms, 213–215
 - queen retinue pheromone (QRP), 142
 - recruitment pheromones with waggle dance, 155
 - response to QMP depends on early exposure, 16
 - royal jelly, 45, 144, 213
 - royalactin, 45, 213
 - RNAi used in discovery, 60
 - worker nurse–forager transition, 214–215
 - changes in brain gene expression, 214
 - worker task allocation behavior related to *forager* and *malvolio* gene expression, 215
- Aplysia* (sea slug)
- oviposition pheromones, 105
 - peptide sex pheromones, 23
- Apocephalus paraponerae* (phorid fly), eavesdrops host alarm pheromone, 246
- Aposematic insects
- aggregation pheromones, 107
- Applications of semiochemicals
- animal husbandry
 - biostimulatory effects, 261
 - primer effects, 271
 - animal welfare, 262–263
 - beneficial insects
 - bumblebees, 260
 - honeybees, 260
 - captive breeding rare species, 262
 - commercialization, challenges to, 272–273
 - greenhouse IPM, virtuous spiral, 263
 - pest management
 - alarm cues, 269
 - alarm pheromones, 270
 - deterrent odors, 269
 - lure and kill or mass trap, 266–269
 - marine antifouling, 270
 - mating disruption, 264–266
 - combined with biological control, 266
 - mechanisms, 264
 - monitoring, 263–264
 - pest resistance to pheromones, 272
 - primer effects, 271
 - push–pull (stimulo–deterrent diversionary) strategies, 270
 - self-protecting plants, 270
 - slow–release formulations, 264
 - trail pheromones, 270
- Aquaculture, 262
- pollution risk, from, 262
- Arabidopsis thaliana* (plant)
- transgenic, producing aphid alarm pheromone, 270
- Arginine vasopressin, 218
- Argyropelecus hemigymnus* (deep-sea hatchet fish), mate location, 223
- Armpit effect (self-inspection or matching), 38
- Arms race, 258
- Arrestment, 107, 224, 270
- Arthropodin (settlement-inducing protein complex, SIPC), 109
- Artificial insemination (AI)
- use of Boarmate™, 261–262
- Ascarosides (nematode pheromones)
- multicomponent, identification, 58
- Assembly pheromone
- ticks, 107
- Assortative mating (like with like), 90
- Asymmetric tracking, 94
- Atta* leaf-cutter ants, trunk trails, 158
- Atta texana* (ant)
- sensitivity of workers to trail pheromone, 154
- Atta vollenweideri* (leaf-cutter ant)
- trail pheromones
 - macroglomerulus in worker brain, 183
 - sensitivity to, 154, 230
- Axillary malodor releasing enzyme (AMRE), 289

- Bacteria. *See* Microbes
- Bactrocera dorsalis* (oriental fruit fly)
adult feeding key to male success, 75
- Badgers. *See* *Meles meles* (European badger)
- Badges of status, *Nauphoeta cinerea* (cockroach), 72
- Balanus balanoides*. *See* *Semibalanus balanoides*
- Bark beetles. *See also* Aggregation pheromones; Allee effects;
Dendroctonus; Eavesdropping; *Ips*
aggregation pheromones
evolved from detoxification of tree defenses, 20
climate change leading to tree stress and beetle epidemics, 107
larval aggregation pheromones, 108
mass attack, 107
mass trapping, 266
- Barnacles
disrupting settlement, 270
egg hatching pheromone, 110
larval settlement, 52, 109
transcriptome studied, 62
- Beavers. *See* *Castor*
- Begging, 36
- Benzaldehyde, 255
- Benzyl cyanide, 88, 245
- Bet-hedging
marine larval settlement, 110
- Bicyclus anynana* (butterfly)
male pheromones and mate choice, 99
- Bioassays. *See also* Semiochemical identification
appropriate concentrations, 53–54, 179
behavioral analysis software, 53
brain imaging, 55
functional magnetic resonance imaging (fMRI), 55, 199, 297, 299
optical, 55
positron emission tomography (PET) scans, 55, 297, 299
- challenges for
demonstrating individual recognition, 132–133
identification and synthesis must include
stereochemistry, 25
learning, 277
multicomponent pheromones, 54–55
multimodal signals, 54
non-physiological concentrations, 298
randomization and blinding, lack of, 53
replication, lack of, 53
signal redundancy, 54
- chemical profiles and signature mixtures, 52–53
direct manipulation of chemical profiles, 53, 127
distinguishing individuals, 52
principal components analysis, 53
recognition, aggression bioassays, 53
- ecological relevance, 50
- electrophysiology, 55
does not always predict behavioral response, 55
electroantennogram (EAG), 55
electro-olfactogram (EOG), 55
electrovomeroogram (EVG), 55
single cell recording (SCR), 55
- humans
organoleptic tests, 289
T-shirt sniff tests, 278
- loss-of-function mutants as tool, 50, 58
- multicomponent pheromones
subtractive method, 54–55
- olfactometers, 54
- primer effects, 51
- standardization, necessary but problematic, 51
- test details can affect results, 51–52
aggression bioassay in ants, 53
discrimination vs. habituation task, 51
lab animal strain, 51
prior experience and learning, 51
- Biomarkers, 58
- Biostimulatory effects, livestock, 261
- Birds
chemical profile differences in related species, 91
cryptic female choice, 85, 89
eavesdropping vole scent marks by ultraviolet cues, 245
major histocompatibility complex (MHC)
mate choice, 84, 85
mate choice by odor, 75
olfactory receptor (OR) gene repertoire similar to mammals, 189
preen gland, 57
semiochemicals, 3
tracking plumes, 227, 242
uropygial gland secretions differ between the sexes, 66
- Blattella germanica* (cockroach)
uses CHC profiles to avoid sibs, 81
- Boarmate™, 261, 262, 263
- Body condition, 19, 36, 75, 76
- Bombus (Psithyrus) bohemicus* (cuckoo bumblebee), 258
- Bombus (Psithyrus)* spp. (cuckoo bumblebee), 257
- Bombus* spp. (bumblebees)
footprints on flowers, 111
- Bombus terrestris* (bumblebee)
foraging recruitment pheromone, 155, 260
- Bombykol, 2, 49, 55
- Bombyx mori* (silk moth), 2, 184
- Bos taurus* (cattle)
estrus detection, 261
- Bostrichthys sinensis* (black sleeper fish)
aquaculture, 262
- Boundary layer, 228, 229
effects on sense organ design, 184
flick and sniff to mitigate, 186
- Bourgeonal, 298
- Brain imaging, 55
functional magnetic resonance imaging (fMRI), 199, 297, 299

- Brain imaging (cont.)
 optical, 55
 positron emission tomography (PET) scans, 55, 297, 299
- Brainbow (visualizing neural circuits), 62
- Brevicomin, 24
- Bruce effect, 133, 146, 219–221
 does it occur in wild?, 220
- Bufadienolide toxins, 269
- Buteo lagopus* (rough-legged buzzard)
 visual eavesdropping, 245
- n*-Butyl-*n*-butyrate, 139, 212
- 2-*sec*-Butyl-4,5-dihydrothiazole, 11, 27, 193
- Caenorhabditis elegans* (nematode)
 aggregation pheromones, 105
 ascarosides, 58
 dauer pheromone, 25
 primer effects, 210–211
 metabolomics to identify multicomponent pheromones, 58
 orientation behavior, 226
 pheromonal parsimony, signal meaning depends on
 concentration, 25
 selection for change in dauer pheromone receptor at high
 density, 190
 sex pheromone
 multicomponent, 25, 28
- Callinectes sapidus* (blue crab)
 orientation in plume, 239
- Callithrix jacchus* (common marmoset)
 subordinate female ovulation suppressed, 146
- Camouflage. *See* Deception
- Camponotus floridanus* (ant)
 alarm pheromone, brain imaging, 183
 colony recognition, processing in brain, 216
 queen pheromone passed round colony on queen's eggs, 141
- Camponotus japonicus* (ant)
 sensilla involvement in colony recognition, 216
- Camponotus obscuripes* (ant)
 alarm pheromone, brain imaging, 183
- Camponotus socius* (ant)
 group recruitment, 153
- Cane toad. *See* *Rhinella marina*
- Canis lupus* (grey wolf)
 border marks in territory, 121
- Canis lupus familiaris* (domestic dog)
 “dog-appeasing pheromone”, 262
 detectors
 estrus, cows, 261
 human disease, 283
 distinguishing humans by smell, 278
 female advertisement pheromone, 66
 female sex pheromone, still unidentified, 5
 forensic use of smell, 282
 nasal flow design for olfactory sensitivity, 186
 olfactory sensory sensitivity, compared to human, 185
 track trails and detect direction, 233
- Canis sinensis* (Ethiopian wolf)
 border marks in territory, 121
- Capra aegagrus hircus* (domestic goat)
 male effect, 201, 221
 males mark females when mounting, 124
- Carassius auratus* (goldfish)
 chemical duets to co-ordinate external fertilization, 69
 combinatorial processing of multicomponent
 pheromones, 179
 endocrine-disrupting pollution affects response to
 pheromone, 46
 multicomponent sex pheromones, 27
 primer effects, bioassay, 51
 scramble competition, 70
- Carboxylic acids, 290
- Carcinus maenas* (crab)
 mate guarding of pre-molt female, 71
 priming effect of female pheromone,
 sensitizes, 207
- Cardiocondyla obscurior* (ant)
 alternative strategies, wingless males with mandibles and
 females mimic males, 87
- Castes. *See also* Ants; *Apis mellifera*; Social insects; Termites
 development, 207, 211–213
 differences
 brain structures, 183, 207
 pheromone secretions, 32, 135
 responses to pheromones, 169, 171
 self-organization models for proportions of different castes, 163
 specialist soldier-guards to resist robber bees, 255
- Castor canadensis* (beaver)
 anal glands, 120
 avoids marks at low population densities, 115
 castoreum gland, 120
 kin recognition, 120, 131
 over-marking, 120
 pest management, 264, 269
 scent-matching, 120
 territories
 scent matching hypothesis, 120
- Castor fiber* (Eurasian beaver)
 territories, scent-matching, 120
- Castoreum, 120
- Cataglyphis niger* (desert ant), territories “owner
 advantage”, 116
- Cats. *See* *Felis catus*
- Cattle. *See* *Bos taurus*
- Central-place foragers, 150
- Ceratitis capitata* (Mediterranean fruit fly)
 benefits of female choice unclear, 81
 eavesdropped by predator, 245
 cost to signaling, 37
 leks, 79
 used as pheromone decoys, 264

- Ceratovacuna lanigera* (social aphid)
 - soldiers, 169
- c-fos*, 218
- Chemical profiles
 - definition, 2
 - do social species have more complex?, 41–42
 - how diversity maintained, 42–43
 - identification. *See* Semiochemical identification
 - mate choice, MHC, 82
 - social insects and social mammals, 126
- Chemical senses
 - compared with other senses, 4
- Chemosensory receptors
 - contrasted with opsins, 4
 - evolution, 188–192
 - “birth-and-death” process, 189
 - independent in insects & vertebrates, 188, 195–197
 - opportunistic co-option, many receptor families, 188
 - selection on receptor repertoire and sensitivities, 189
 - insects
 - gustatory receptors (GRs), 197
 - ionotropic receptors (IRs), 197
 - odorant receptors (ORs), 196
 - unrelated to vertebrate receptors, 196
 - use ionotropic signaling pathways, 196
 - mammals
 - all receptor types are GPCRs, 192
 - formyl peptide receptors (FPRs), 195
 - trace amine-associated receptors (TAARs), 192, 193
 - vomerolateral receptors (V1Rs and V2Rs), 192, 193–195
- Chemosensory systems
 - invertebrates and vertebrates
 - taste (gustation) vs. smell (olfaction), 175
- Chiloglottis* spp. (orchids)
 - speciation, 253
- Chiloglottones (2,5-dialkylcyclohexan-1,3-diones), 253
- Chiral molecules, 25, 306
- Chocolate trails, humans follow, 233
- Cholesta-5,7-dien-3-ol (provitamin D₃), 21
- 5 β -Cholestan-3-one, 152
- Chondroitin, fish alarm cue, 166
- Cimex lectularius* (bed bug)
 - aggregation pheromone
 - monitoring, 263
 - alarm pheromone, pest management, 270
 - arrestment by pheromone, 224
- cis*-regulatory DNA, 29, 91
- cis-trans* isomers, 310
- Citral, 255
- Cleptobiosis, 255
- Cleptotrigona* (stingless bee), uses propaganda pheromones, 255
- Clethrionomys glareolus* (bank vole)
 - females prefer odors of dominant male, 75
- Climate change
 - ocean acidification, 46
 - tree stress and bark beetles, 107
- CNV. *See* Copy number variation
- Coccinellidae (ladybugs)
 - detect larvae, lay fewer eggs, 111
- Coccinella septempunctata* (ladybug)
 - aggregation pheromone, 107
- Cockroaches
 - aggregation
 - ecophysiological benefits, 107
 - sex pheromones
 - highly varied between genera, related within, 28
- Co-evolution
 - social parasites and host social insects, 255–258
- Colletes cunicularius* (solitary bee), duped by orchid, 252
- Colony level selection, 214
- Colophina monstifica* (social aphid)
 - soldiers, 169
- Communication, 18, 32 *See also* Pheromones; Signals
- Composite signals. *See* Signals; multimodal
- Concentration
 - pheromonal parsimony, 24–25
 - signal meaning depends on, 24–25, 54, 157, 210
 - too high, response specificity lost, 53, 54, 179
- Conservation
 - captive breeding, 262
 - using scent marking to manipulate mate choice, 262
 - vertebrate scent marks to census, 263–264
- Contact chemoreception, 175
- Coolidge effects, 86
 - mechanisms, 218
- Co-operative breeding, 146, 219
- Copulatory plug, 23, 88, 89
- Copulins, 299–300
- Copy number variation (CNV), 191, 292
- Coremata, 66, 76
- Corpora allata, 139, 207, 212, 214
- Cortisol, 89, 279
- Corynebacteria*, 288
- Corynebacterium striatum*, 289
- Cosmophasis bitaeniata* (spider)
 - gets colony-specific CHC from host ant, 257
- Cosmopolites sordidus* (banana weevil)
 - aggregation pheromone, for control, 266
- Costs (of signaling). *See* Signals, costs
- Countermarking. *See* Over-marking
- p*-cresol, female horse pheromone, 261
- Critical periods
 - behavioral development, 16
 - caste development, 213
- Crocota crocata* (spotted hyena)
 - scent marking patterns, variation by habitat, 121
- Cross-fostering, 85, 131
 - social insects
 - sources of colony odors, 128
 - species-specific mate recognition, 16

- Crotalaria* (plant), source of alkaloids, 76
- Crozier's paradox, 42
- Crustaceans
- aggregation for defense, 105
 - antennal flicks (sniffs)
 - scaling with increasing size in lifetime, 186
 - avoiding predator odors, 248
 - distributed chemosensory system on claws, 175
 - individual recognition, 72
 - mediated by aesthetascs pathways, 215
 - larval settlement from plankton, 109
 - lobster flicks antennae to smell, 186
 - male lobsters fight for ownership of shelters, 72
 - mate guarding, 71
 - orientation behavior
 - copepods following pheromone trails, 230
 - crabs orienting in odor plume, 239
 - integration of olfactory and hydrodynamic cue inputs, 44
 - pheromones
 - contact sex pheromones (copepods), 8
 - metabolomics may be fruitful approach to identification, 58
 - modulation of visual threat signals by female pheromones, 44
 - uridine diphosphate, shore crab female pheromone, 71
 - pollution
 - chemosensory disruption from ocean acidification, changes ligand interaction, 46
 - male responses to female odors are reduced, 46
 - symbiosis with sea anemones, 250
 - synchronized larval release, 110
 - transcriptome studied, 62
- Cryptic choice, 89–90
- Cryptotermes secundus* (termite)
- differences in gene expression, queens vs. kings and workers, 139
 - queens and nymphs, different CHC profiles, 138
- Cues
- definition, 18
- Culex* (mosquito)
- oviposition pheromone, 24, 106
- Cuticular hydrocarbons (CHCs), 155
- camouflage, 257
 - deception, 255–258
 - foragers' CHCs different vs. workers inside nest, 41, 130
 - social insects
 - colony odors, sources, 41
 - colony recognition, 126–130
- Cydia pomonella* (codling moth)
- pest management, 264
- Cynictis penicillata* (yellow mongoose)
- scent marking, 114
- Cynops ensicauda* (newt)
- peptide pheromone, 25
- Cynops pyrrhogaster* (red-bellied newt)
- peptide pheromone, 25
 - population differences in male peptide pheromone, 92
 - species-specific male peptide pheromone, 91
- Cyprinus carpio* (common carp)
- multicomponent sex pheromones, 27
 - pest management, 267
- Cypris larvae (barnacles), 52, 109
- Cysteine-glutathione, 70
- Danaus gilippus* (queen butterfly)
- male transfers pheromone direct to female, 23
- Darcin (mouse Mup20), 35
- binds 2-sec-butyl 4,5 dihydrothiazole (thiazole), 219
 - interaction of pheromone and signature mixture, 219
 - prompts learning male signature mixture and location, 16, 118, 124, 209, 219
- Darwin, Charles
- sexual selection on pheromones, 5, 65
- Dauer larva *Caenorhabditis elegans* (nematode), 25, 210–11
- Dear-enemies (territorial defense), 118, 123, 133
- Decanal, 75
- Decapod shrimps
- mate guarding, 71
- Deception, 251–258
- aggressive chemical mimicry, 251–258
 - blister beetle, 253–255
 - bolas spiders, 251
 - pollination by sexual deception, 252–253
 - imperfect mimicry more attractive, orchid, 252
 - aided by shared biochemistry in all life, 244
 - ants and lycaenid butterflies (see also Mutualism), 250
 - code breaking, 244
 - cost of not responding to deceivers, 251
 - mechanisms (insignificance, chemical camouflage, chemical mimicry), 250
 - of social insects, 255–258
 - arms race, 258
 - camouflage, 257
 - chemical insignificance, 257
 - counterfeiting, 257
- Decyl acetate, 255
- Deer. See *Odocoileus hemionus columbianus*
- Dehydroepiandrosterone, 290
- Dehydro-*exo*-brevicomine, 27, 193, 210
- Dendraster excentricus* (sand dollar)
- feeding currents, benefits of aggregation, 110
 - planktonic larval settlement, 110
- Dendroctonus* (bark beetles)
- Dendroctonus brevicomis*, eavesdropped by predators, 246
 - Dendroctonus frontalis*, 108
 - Dendroctonus micans*, 108
 - larval aggregation pheromones, 108

- Dendroctonus ponderosae*, 107
Dendroctonus pseudotsugae, 108
Dendroctonus rufipennis, 107
Dendroctonus valens, 108
Desaturase enzymes, 94
Desaturase 1 gene, 101
Desmognathus ocoee (Ocoee salamander)
transdermal transfer of pheromone to female bloodstream, 45
Development
caste differences in social insects pheromone production and perception, 207
effects on response to pheromone, 207
learning
maternal behavior, 216
response to semiochemicals, 16
sensory system, periphery, 16
social control of reproduction
social insects and social mammals, parallels, 147–148
switch
dauer pheromone, 25, 210–211
social insects, 45, 211–213
Diaeretiella rapae (parasitoid wasp), arrested by aphid alarm pheromone, 270
Diastereoisomers, 309, 310
Dictyostelium (slime mold)
greenbeards, 40
Differential analysis by 2D-NMR spectroscopy (DANS), 58
2-(sec-Butyl)-dihydrothiazole, 210
1,5-Dimethyl-6,8-dioxabicyclo[3.2.1]octane (frontalin), 11
Dimethyl disulphide, 283
4,8-Dimethyldecanal, 89
2,5-Dimethylpyrazine, 193
Dinoponera quadricaps (ant)
alpha female badge, 7
Diomedea exulans (albatross)
navigation, food location, 226, 242
Dipsosaurus dorsalis (desert iguana)
multimodal signal with ultraviolet cue, 44
scent marks with ultraviolet cue, 123
Disassortative mating, 85
Disruptive selection, 94
DNA methylation, 213, 215
Dodecadienol, 157
Dodecatrienol, 157
(Z)-7-Dodecen-1-yl acetate, 11, 24
Dodecyl acetate, 255
Dogs. *See* *Canis lupus familiaris*
Dolichotis patagonum (mara), males directly urine mark females, 124
Dopamine, 16, 142, 218, 219, 221
Doublesex (*dsx*), 101, 201–204
Drosophila grimshawi (Hawaiian fruit fly)
benefits of female choice unclear, 81
Drosophila melanogaster (fruit fly)
adult flight behavior, 226, 240
changes in CHCs with age, 57
cis-vaccenyl acetate (cVA), anti-aphrodisiac, 22, 88, 179, 202
forager and *malvolio* gene expression, comparison in honeybee, 215
gustatory (as well as olfactory) receptors for pheromones, 175
males, response to CHC pheromones, 15
larval orientation behavior, 226
learning
discriminating receptive partners, 204, 209
macroglomerular complex, 181
male adjusts ejaculate and mating time vs. sperm competition, 89
multimodal courtship signals, 44, 54, 201–204
olfactory receptors
polymorphisms, 191
regional races differ in CHC blends, 92
sex peptide, 45, 88–89, 204
sex-specific neural circuits, 201–204
Drosophila melanogaster subgroup
rapid speciation
cuticular hydrocarbons, 99–103
Drosophila sechellia (fruit fly)
cuticular hydrocarbons in species isolation, 103
olfactory binding proteins and taste responses, 198
Drosophila serrata (fruit fly)
speciation, involves more than sexual selection
alone, 90
Drosophila simulans (fruit fly)
cuticular hydrocarbons in species isolation, 103
repelled by *Morinda citrifolia* (plant), 198
Dufour's gland
host-marking by parasitoids, 111
Dulotic ants, 255
Eavesdropping, 244–249
aggregation pheromones, 246–248
ant alarm pheromones, 246
arms race, 247
egg dumping, 248–249
intra-specific
hormones evolving into pheromones, 18
sex pheromones as aggregation pheromones, 68, 105, 107, 246, 248
of predator odors by prey, 248
sex pheromones, 245–246
“bridge-in-time”, 245
territorial markings, 246
Ecdysteroid hormones, 89, 139, 212, 213
Eciton (army ants), foraging models, 161
Efficacy cost, 32
Egg dumping, 248–249
Eicosanoids. *See* PUFAs
(Z)-11-Eicosen-1-ol, 21, 80
Elater ferrugineus (beetle), monitor for rare prey, 263
Electronic noses, 283
Elephants. *See* *Elephas maximus*; *Loxodonta africana*

- Elephas maximus* (Asian elephant)
 detects pheromones in female urine, 71
 dominant males and musth, 76
 pheromones shared with moths and beetles, 24
- Elminius modestus* (barnacle), long penis, 109
- Enantiomers, 25, 306
- Endocrine-disrupting chemicals, 46
- β -Endorphin
- Ephestia elutella* (tobacco moth)
 females choose larger males, 75
- Epidermal club cells, 166
- Epidermal growth factor receptor (EGFR), 213
- Epigenetic effects, 62, 106, 213
- Epiphyas postvittana* (light brown apple moth)
 pest management, 264
- Epistatic interactions, 98, 103
- (3Z,9Z)-(6S,7R)-Epoxy-heneicosadiene, 251
- Equus caballus* (horse)
 female sex pheromone, *p*-cresol, 261
- Esox lucius* (pike)
 prey learn to associate odor with alarm cue, 167, 262
- 17 β -estradiol, 46
- Estratetraenol, "putative human pheromone", 297
- Estrous cycle synchrony
 doubts, 210
- 4-Ethyl-2-methoxyphenol, 72
- Ethyl oleate, 142, 145, 214
- Ethyl palmitate, 145, 214
- 4-Ethyloctanoic acid, 290
- Eucalyptol, 155
- Euglossine bees
 collect species-specific plant perfume oils as pheromone, 31
- Eulemur* spp. (lemurs)
 greater complexity of glandular secretions in more social species, 42
- Euproctis taiwana* (moth), 245
- Eusociality. *See also* Social insects; Social mammals
 conflict over reproduction in societies, 136–137
 continuum (reproductive skew), 136
 co-operative broodcare, 136
 social insects and social mammals, parallels, 147–148
- Evolution. *See* Pheromones, evolution
- Evolvability, 191
- Exocrine gland-secreting peptide 1 (ESP1), 11, 179, 184, 194, 204–206
- Expectancy violation, mate choice, 78
- Experience. *See* Development; Learning
- Experimental methods. *See* Semiochemical identification
- E-Z* isomers, 310
- Fabre, Jean-Henri, 5
- Falco tinnunculus* (kestrel)
 visual eavesdropping, 245
- (α)-Farnesene, 210, 270
- (*E*)- β -Farnesene, 24, 168, 210, 255, 270
- Fat body, 213
- Felis catus* (domestic cat)
 "feline facial pheromone"(FeliwayTM), 262
 flehmen, 193
 odors invoke fear in rats, 248
- Female choice. *See* Mate choice, Sexual selection
- Fertility signal, 36, 137–146
- Fish
 androgen effects on electro-olfactogram (EOG), 207
 brain
 neural circuits for response to alarm cues, 166
 processing of pheromones, 187
 chemosensory disruption from ocean acidification, changes
 ligand interaction, 46
 internally fertilizing, cryptic female choice, 89
 learn predator odor association with alarm cue, 167, 262
 male pheromones, 66
 males advertise parental care, 77
 orientation behavior
 lateral line, use in rheotaxis, 241
 rheotaxis, 241
 tropotaxis, 242
 pheromones
 evolution, 19
 for management of invasive species, 267
 multicomponent, 19
 sexual selection and imprinting, 74
 sniff, 186
 tracking plumes, 226, 241–242
- Fixatives
 2-phenoxyethanol, 23
 fatty acid esters, 23
- Flavor, 175, 278, 302
- Flehmen, 193
- Flicks, antennal
 behavior of smelling, 186–187
 scaling with increasing size in lifetime, 186
- Fluctuating asymmetry, 81, 300
- Follicle-stimulating hormone, 89, 199
- Foraging (for)* gene, 215
- Formica* ant spp.
 formic acid as defense and alarm pheromone, 21, 170
- Formica cunicularia* (ant), propaganda victim, 255
- Formica exsecta* (ant)
 CHCs in colony recognition, 127
 colony-specific cuticular hydrocarbons, 13, 41
- Formica fusca* (ant)
 diversity of CHCs driven by social parasitic ants, 258
- Formica japonica* (ant)
 colony odor successfully synthesized, 15
- Formica lemni* (ant), host to syrphid fly, 258
- Formica subintegra* (slave-making ant), use of
 propaganda, 255
- Formicoxenus* (shampoo ant), social parasite of *Myrmica*
 ants, 257

- Formyl peptide receptors (FPRs), 78, 195
 co-option from immune system, 188
 Fright reaction. *See* Alarm cues, fish
 Frontalin, 76
Fruitless (fru), 201–204
- GABA, 216–217
Galaxias fasciatus (banded kokopu fish), lateral line for
 rheotaxis, 241
Gallus gallus (red junglefowl)
 cryptic female choice, 89
Gammarus roeselii (crustacean), avoids predator odor, 248
Gargaphia solani (aubergine lace bug)
 intra-specific egg dumping, 248
 subsocial care, alarm pheromone, 165
 Garter snake. *See* *Thamnophis sirtalis parietalis*
 Gas liquid chromatography (GC). *See* Semiochemical
 identification
Gasterosteus aculeatus (stickleback, fish)
 mate choice
 diversity of MHC optimized, 85
 major histocompatibility complex (MHC), 85
Gasterosteus spp. (sticklebacks, fish)
 imprinting, basis of sexual isolation of two species, 16, 74
 GC. *See* Semiochemical identification
 Gene expression, 214
 brain
 induced by pheromone exposure, 208
 differences related to roles in honeybee workers, 214–215
 Genomics
 RNA-seq, 62–63
 sociogenomics, 212
 Geometrical isomers (*cis-trans*, *E-Z*), 310
 Geranial, 255
 Geraniol (3,7-dimethylocta-2,6-dien-1-ol), 165
 Gestalt model
 colony odor, 127, 128
 Giant panda. *See* *Ailuropoda melanoleuca*
 Glands
 anal, 77, 120, 132
 areola
 humans, 295
 castoreum, 120
 cephalic, 80
 Dufour's, 37, 129, 142, 255
 humans
 apocrine, 287
 apoeccrine, 287
 areola, 295
 eccrine, 287
 sebaceous, 287
 sweat glands, 287–288
 hypopharyngeal, 213
 labial, 155
 mandibular, 142, 213
 metatarsal, 167
 post-pharyngeal, 129
 pre-orbital, 121
 pre-putial, 210
 pygidial, 153
 salivary, 145
 social insects
 summarized, 134
 sternal, 129
 subcaudal, 132
 temporal, 76
 tergal, 142, 155
 van der Vecht, 170
 venom, 212
 Glutamic acid, 70
 Glycoprotein
 barnacle, 109
 GnRH pulse generator, 201
Gobius niger (fish)
 chemically inconspicuous sneaker males, 87
 Gonadotropin-releasing hormone (GnRH) neurons, 199–200
 Good genes, 75
 G-protein-coupled receptors (GPCRs), 192
Grapholita molesta (oriental fruit moth)
 flight track, 240
 Greenbeards, 40 *See also* Recognition
 Grueneberg ganglion, 195
Grylodes sigillatus (cricket)
 Coolidge effect, 86
 Guaiacol (2-methoxyphenol), 31
 Guanine, 107
 Guanylate cyclases, 188
 Gustation
 compared with smell (olfaction), 175
 olfactory binding proteins
 specificity of response, 198
 Gustatory receptors (GRs)
 insect GRs
 “pickpocket” ion channels, 203
 ionotropic (unrelated to vertebrate GRs), 197
 Gustatory sensory neurons (GSNs)
 insects, 197
 Gynandromorphs, 201
- Habituation
 aphids, to alarm pheromone, 270
Habronestes bradleyi (spider), attracted to
 fighting ants, 246
Habropoda pallida (solitary bee), duped by blister
 beetle larvae, 253
 Hamilton, selfish herd, 249
 Hamilton's rule, kin selection, 133, 145
 Handicap theory, 32, 33–34, *See also* Signals
 immunocompetence handicap hypothesis, 79
 reliable signals without handicap, 32–37

- HapMap, 281
- Hawaiian drosophilid speciation, 91–92
- Headspace sampling, 57
- Heliconid butterflies
- anti-aphrodisiacs, 88
- Heliconius charithonia* (butterfly)
- males guard female pupae, 71
- Heliconius melpomene* (butterfly)
- male pheromones
 - fixative extends signal life, 23
- Helicoverpa armigera* (cotton bollworm moth)
- seminal fluid molecules, 89
- Helicoverpa zea* (corn ear worm moth)
- sensitive to sympatric species' pheromone components, 55
- Heliothis subflexa* (moth) and *Heliothis virescens* (tobacco budworm moth)
- directional pressures on pheromone blends in sympatric species, 90
 - male pheromones in courtship prevent hybrid matings, 97–98
 - polygenic divergence in signal and male detection in
 - sympatric species, 29, 97–98
- Heliothis virescens* (tobacco budworm moth)
- flight track, 240
 - seminal fluid molecules, 89
- Helogale undulata* (dwarf mongoose)
- individual recognition, 132
 - can discriminate anal gland secretions, not cheek gland, 133
- (3Z,6Z,9Z)-Heneicosatriene, 251
- 9-Hentriacontene, 7
- 7,11-Heptacosadiene, 88, 100, 203
- 2-Heptanone, 193, 194, 199
- Herpestes aurospunctatus* (Indian mongoose)
- individual recognition, 132
- Heterocephalus glaber* (naked mole rat)
- colony odor, parallels with social insects, 132
 - queen suppression of worker reproduction, by physical dominance, 146
 - recruitment of foragers, 155
- Heterodera glycines* (soybean cyst nematode)
- pest management, 266
- (Z,E)-7,11-Hexadecandienyl acetate, 305
- (Z)-9-Hexadecenal, 270
- Hexanoic acid, 290
- Hexenoic acid, 290
- (Z)-3-Hexen-1-ol (*cis*-3-Hexen-1-ol), 295
- Hexyl decanoate, 155
- Hidden preferences (receiver psychology), 21
- High performance liquid chromatography (HPLC), 55
- Homarus americanus* (lobster)
- contest, 72
 - flicks antennae to smell, 186
 - pre-copulatory mate guarding, 71
- Homo sapiens*. *See* Humans
- Homovanillyl alcohol (HVA), 142
- Honest signals. *See* Signals
- Honeybees. *See* *Apis mellifera*
- Hormones. *See also* Primer effects
- cortisol and recognition, human mothers, 279
 - definition, 6
 - influence on central/peripheral responses to
 - pheromones, 207
 - male body condition
 - mate choice by females, 75–76
 - primer effects, 43–44
 - similarities between mammals and insects, 43
 - social insect caste development, 139
 - testosterone androgen effects
 - male hamster brain needs to respond, 208
- Horse. *See* *Equus caballus*
- Host-discrimination, 110
- Host-marking pheromones (HMP)
- evolution of, 111
 - host-discrimination, 110
 - individual recognition, 111
- Human leucocyte antigen (HLA). *See* Humans, mate choice;
- Major histocompatibility complex (MHC);
 - Mammals
- Humans (*Homo sapiens*)
- advertisement of ovulation?, 299–300
 - “copulins”, 299–300
 - male ability to detect, 300
 - age, changes with
 - perception, 283
 - secretions, 284
 - androstadienone (androsta-4,16-dien-3-one)
 - perception, 294–295, 297–298
 - androstenone (5 α -androsta-16-en-3-one)
 - perception, 294–295, 297–298
 - anosmias
 - many molecules induce changes on
 - exposure, 298
 - antiperspirants and deodorants
 - modes of action, 289
 - sales, 282
 - areola glands, 295
 - axilla (armpit) hair and odor-creating
 - bacteria, 288
 - chemical profiles and signature mixtures
 - mate choice, 280
 - mothers and babies, 278–279, 282
 - cleanliness not necessarily a virtue, 282
 - earwax and armpits, 290–291
 - estratetraenol
 - “putative human pheromone”, 297
 - follow chocolate trail, 233
 - genetic polymorphisms, 291
 - perception of smell, 292–295
 - smell production, 290–291
 - mammary pheromone?, 295

- mate choice
 - body symmetry, 300
 - evidence from 250 years' data, Iceland, 84
 - kibbutz or Westermarck effect, 280
 - major histocompatibility complex (MHC), 279–284
 - oral contraceptives, effects?, 281
 - steroid “putative human pheromones”, 296–299
 - T-shirt sniff tests, 281
- microbes
 - contribution to odor, 288–290
 - variation over body and between people, 288–290
- mother–infant olfactory recognition, 217
- mothers and babies
 - areola gland secretion, mammary pheromone?, 295
 - mother cortisol and baby recognition, correlation, 279
 - recognition, 278–279, 282
- odor preferences
 - genetics, olfactory receptor variation, 292–295
 - learning, 277
 - foods, 277
 - neonates different from older children and adults, 277
- odors and memory
 - experimental tests, 280
 - Proust, smells bring back memories, 280
- odors, cultural and social aspects, 277–278
 - Napoleon, 277
 - osmologies, 303
 - perfumes, 277
 - sexual attraction, 277
- olfaction, human
 - anosmias
 - specific, 292–295
 - compared, dogs, 185
 - compared, mice and primates, 275
 - distinguishing isomers, 309
 - genetic polymorphisms, 191, 278, 292–295
 - loss, color vision hypothesis, 291
 - olfactory bulb, size, 275
 - unique individual worlds, 191, 278, 292
- primer effects
 - menstrual synchrony?, 300–301
- recognition, 278–279
 - children, sibs, parents, 279
 - mother–child, 278–279
 - phenotypic matching, 279
 - security blankets (comforters), 279
 - self, partners, 279
 - T-shirt sniff tests, 278
- sexual orientation, 287
- signature mixtures, 278
- steroid “putative pheromones”, 10, 54, 296–299
 - androstadienone, 296–299
 - androst-enol (5 α -androst-16-en-3 α -ol), 296–297
- sweat glands
 - apocrine, apoeccrine, eccrine, 287–288
 - odorless precursors, 290
 - using human odors
 - forensics, 282
 - problems, 282
 - mammary pheromones, 282
 - medical diagnosis, 283
 - smell loss, neurodegenerative diseases, 283
 - smells produced by disease, 283
- dogs as detectors, 283
- electronic noses, 283
 - mood changers and “aromatherapy”, 283
 - vomer-nasal organ (VNO), vestigial, 291
- Hutterites, MHC and partner choice, 281
- Hyaena brunnea* (brown hyena)
 - pasting marks, 123, 132
- Hyaena hyaena* (striped hyena), social
 - greeting, 123
- Hydrobates pelagicus* (storm petrel, bird)
 - avoid kin as mates, 84
- Hydroides* (marine worm)
 - larval settlement, bet-hedging, 110
- Hydroquinone (1,4-dihydroxybenzene), 158
- 3-Hydroxy-2-butanone, 72
- (*E*)-10-Hydroxy-2-decenoic acid, 135, 213
- 6-Hydroxy-6-methyl-3-heptanone, 193, 210
- 3-Hydroxy-3-methyl-hexanoic acid, 290
- Hyperosmias, 293
- Hyposmias, 293
- Iberolacerta cyreni* (lizard)
 - male pheromones depend on body condition, 36, 75
 - pre-existing sensory bias, food lipids, 21
- Ichneumon eumerus* (parasitoid wasp), uses propaganda, 255
- Iguana iguana* (lizard)
 - male femoral gland stimulated by testosterone, 75
- Immunocompetence handicap hypothesis, 79
- Imprinting, 38–40
 - can lead to sexual isolation of two species, 16, 74
 - definition, 215
 - similarities between mammals and social insects, 129
 - species-specific mate recognition, 16
- Index signals, 35
- Indices (unfakeable signals), 35–36
- Individual recognition, 38, 72, 118, 132, 133
 - challenges for, demonstrating, 132–133
- Indole, 88
- Infochemicals, 6
- Infodisruption by pollution, 45
- Information, 18, *See also* Pheromones; Signals
- Innate behavior, 16
- Inosine, 70
- Insect vectors, 263, 270
- Insulin signaling, 139, 211, 212

- Integrated pest management (IPM), 263, 264
 push-pull (stimulo-deterrent diversionary) strategies, 270
- Interaction between modalities, 44
- Interspecific interaction, semiochemical mediated. *See* Deception; Eavesdropping; Mutualisms
- Ips pini* (pine engraver bark beetle), 246
- Ipsdienol, 246
- Iridomyrmex purpureus* (ant), alarm pheromone attracts spider, 246
- Isomers, 305–310
 biological importance, 49, 246, 304
 chirality, 306
 human smell distinguishes, 309
 naming conventions, 309
 (–/+), (d/l), (D/L), 305, 307
 absolute configuration, (R), (S), 309
 optical, 307
 racemic mixture (racemate), 307
 stereoisomers (spatial isomers), 306
 diastereoisomers (*not* mirror images), 309
 enantiomers (mirror images), 306
 structural (constitutional), 305
 chain, 305
 functional group, 305
 geometrical isomers (*cis-trans*, *E-Z*), 310
 positional, 306
- Isopentyl acetate, 43, 208
- Isovaleric acid (3-methylbutanoic acid), 294
- Ixodes neitzi* (tick)
 responds to scent marks of host antelope, 246
- Ixodes scapularis* (tick), clusters on host deer trails, 246
- Jacobson's organ. *See* Vomeronasal organ (VNO)
- (R)-Japonilure, 25
- (S)-Japonilure, 25
- Juvenile hormone (JH), 34, 89, 139, 212, 213, 214, 215
- Kafue lechwe* (Kobus lechwe, antelope)
 lekking, 80
- Kairomones, 136, 166, 244–249
- Kaloterms flavicollis* (termite)
 sex differences, 138
- Kangaroos
 mate guarding by male, 71
- Keiferia lycopersicella* (tomato pin worm)
 pest management, 266
- 3-Keto petromyzonol sulphate, 22
- 15-Keto-prostaglandin-F_{2a}, 207
- Kibbutz or Westernmarck effect
 human mate choice, 279
- Kin selection, 36, 37, 133
- Kinesis, 224, 225
- Kisspeptin, 200–201
- Kobus kob* (Uganda kob, antelope)
 lekking, 80
- Labeled line, 179, 183
- Lacerta monticola*. *See* *Iberolacerta cyreni*
- Lacinipolia renigera* (moth)
 victim of bolas spider, 251
- Ladybugs. *See* Coccinellidae
- Lanierone, 246
- Lasioderma serricorne* (cigarette beetle)
 aggregation pheromone, 109
- Lasioglossum zephyrum* (sweatbee)
 guard bees discriminate relatedness, 128
- Lasius alienus* (ant)
 panic response to its alarm pheromone, 171
- Lasius fuliginosus* (ant), 255
- Lasius neoniger* (ant)
 co-operative foraging, 153
- Lasius niger* (ant)
 marking behavior on colony home range, 160
 modeling foraging, 160
 queen marking of eggs, 146
 queen pheromone, 3-methylhentriacontane (3-MeC₃₁), 141
 queen pheromones, 213
- Lateral protocerebrum, 183
- Learning. *See also* Development
 alarm cues
 gives flexibility of response, 167
 hatchery trout learn predator odors, 262
 discriminating receptive partners, 204, 209
 family and kin recognition
 mammals, 130–133
 mammals
 maternal ewes bond quicker on subsequent births, 209, 217
 needed to distinguish estrous from diestrous female odors, 209
 mitral cells in main olfactory bulb, 216–217
 parasitoid insects
 generalist learns host pheromones, 245
 sensitive period, 131, 216
 signature mixtures, 38–41, 215–218
- Lee-Boot effect, 210
- Leks, 79–81
- Lemur catta* (ring-tailed lemur), “stink fights”, 113, 114
- Lepomis cyanellus* (green sunfish), alloparental care, 248
- Leptogenys peuqueti* (ant)
 multicomponent trail pheromone, 28, 157
- Leptothorax acervorum* (ant)
 tandem running recruitment, 153
- Leptothorax gredleri* (ant)
 colony-specific CHCs may help inbreeding avoidance, 42
- Lestrimelitta* (stingless bee), uses propaganda pheromones for cleptobiosis, 255
- Lévy walks, 226
- Limonene, 31
- Linalool, 71
- Linalool oxide (furanoid), 71

- Linepithema humile* (Argentine ant)
 bioassay design affects aggression result, 53
 learning to distinguish branched CHCs easier, 41
 trail pheromone, disrupting, 270
- Linyphia litogiosa* (spider), conflict between the sexes, 69
- Lipaphis* (*Hyadaphis*) *erysimi* (aphid)
 alarm pheromone, 168
- Liriomyza sativae* (leafminer fly)
 pest management, 266
- Litoria splendida* (magnificent tree frog)
 male pheromone, peptide, 66
 low cost, 34
- L-kynurenine, 23
- Locust. *See* *Schistocerca gregaria*
- Loligo pealeii* (squid)
 aggressive contests, prompted by egg mass protein
 pheromone, 72
- Lordosis, 206, 261
- L-Ovothiol-A, 70
- Loxodonta africana* (African elephant)
 dominant males and musth, 76
 mate guarding, 71
- Lure and kill, pheromone control, 266
- Luteinizing hormone, 89, 199
- Lutzomyia longipalpis* (sandfly)
 lekking, 79
- Macrotermes* (termite)
 building queen chamber, 163
 nest building, 161
- Maculinea alcon* (Alcon blue butterfly)
 arms race of deception, 258
- Maculinea rebeli* (lycaenid butterfly), 255
- Major histocompatibility complex (MHC)
 familial imprinting, 218
 odor preferences for difference in mates, 85
 human leukocyte antigen (HLA), 280
 humans
 mate choice
 evidence from 250 years' data, Iceland, 84
 interaction with individual's microbial flora, 82
 kin recognition, 84
 mate choice, 81–86, 279–284
 for greater diversity of MHC alleles, 85
 MHC diversity, 85
 mutual distinguishing MHC of odors in rats, mice, and
 humans, 280
 not always found in wild populations, 85
 self-reference, 85
 preferences
 women taking oral contraceptives, 281
 selection for polymorphism to avoid inbreeding?, 42
 selection pressures maintaining diversity of MHC alleles, 84
 disassortative mating, 281
 greater resistance to disease or parasites, 85
 higher implantation rates if dissimilar, 85, 281, 284
 reduced chance of inbreeding if dissimilar, 85
 sources of the odors, 82
- Major urinary proteins (MUPs)
 add longevity to volatile signal, 23
 cue to avoid inbreeding, 218
 darcin (Mup20) (mouse), 124, 219
 high metabolic cost of use in scent marking, 35
 mate choice, inbreeding avoidance, 86
 mice
 dominant males, 118
- Male effect (goats and sheep), 201, 221
- Malvolio* gene, 215
- Mammals. *See also* Humans; Primates; Social mammals
 family and kin recognition, 130–133
 clan or group, 131–132
 individual recognition, 132–133
 kin, 131
 mother–offspring, 130–131
- learning
 needed to distinguish estrous from diestrous female odors,
 209
 pheromones, 12
 primer effects, 209–210
 scent marks to census, 263–264
- Man. *See* Humans
- Mandrillus sphinx* (mandrill)
 no estrous synchrony found, 210
- Manduca sexta* (tobacco horn moth)
 octopamine influences peripheral response to
 pheromone, 207
 sexual dimorphism, antennae, receptors, brain, behavior, 201
- Mastophora hutchinsoni* (bolus spider)
 aggressive chemical mimicry of moth pheromones, 251
- Mastotermes darwiniensis* (termite)
 labial gland pheromones, 158
- Mate choice
 benefits, 73
 body condition, 35–36
 effect via testosterone, 75
 by males, 76
 fluctuating asymmetry, 81, 300
 for health/avoid infection, 77–79
 genetic compatibility, 81–86, 279–284
 imprinting, 74
 leks, 79–81
 major histocompatibility complex (MHC), 81–86, 279–284
 mechanisms
 “runaway sexual selection” (“Fisherian sexy sons”), 74
 antagonistic coevolution (chase-away sexual selection), 74
 compatible genes, 74
 direct benefits, 74
 importance of learning/imprinting, 74
 indicators, 74
 interaction with environmental factors, 74

- Mate choice (cont.)
 operational sex ratio, 73
 overview, 73–75
 paternal investment, 76–77
 quality signals, 75–76
 to avoid inbreeding
 familial imprinting, 218
 Mate guarding. *See* Sperm competition and mate guarding
 Maternal behavior and recognition, 130–131, 278–279
 Medial amygdala
 integration of individual identity cues from
 VNO and MOE, 216
 Medial pre-optic hypothalamus (MPOA)
 sensitization by experience, 209
 testosterone androgen effects
 male hamster brain needs to respond, 208
Meles meles (European badger)
 clan recognition, 132
 latrine distribution, variation by habitat, 123
 scent marking, potential population management, 269
 squat marking (allomarking), 40, 132
Meloe franciscanus (blister beetle), triungulin larvae, 253
 Menstrual synchrony, 300–301
p-Menth-1-en-8-ol, 302
Mesocricetus auratus (hamster)
 contribution of different glands to individual odor,
 generalizations, 133
 female advertisement, vaginal secretion trails, 66
 learning own identity from litter mates, 38
 recognition of individuals of other hamster species, 40
Messor barbarus (harvester ant)
 matching trail patterns to food distribution, 159
 Metabolomics, 57–58, 301
 Methoprene, 34
 Methyl linolenate, 214
 3-Methylbutanoic acid (isovaleric acid), 294
 2-Methyl-1-butanol, 139, 212
 2-Methylbut-2-enal, 11, 25, 55
 Methyl 4-methylpyrrole-2-carboxylate, 154
 Methyl eugenol, 75
 (Z)-16-Methyl-9-heptadecenyl isobutyrate, 245
 4-Methyl-3-heptanol, 246
 4-Methyl-3-heptanone, 229, 246
 (R)-(-)-5-Methyl-3-heptanone, 70
 (S)-(+)-5-Methyl-3-heptanone, 70
 6-Methyl-5-hepten-2-one, 246, 255
 Methyl ketones, 76, 84
 Methyl oleate, 145
 Methyl palmitate, 145
 Methyl salicylate, 88
 3-Methylhentriacontane (3-MeC₃₁), 141
 2-Methylthiazolidine, 72
 (Methylthio)methanethiol, 11, 187, 199
 MHC. *See* Major histocompatibility complex
 Mice. *See* *Mus musculus domesticus*
 Microarrays, 62
 Microbes
 contribution to odor
 effect of MHC variation, 82
 human odors, 288–290
 mammals, individual and clan odor, 40, 132
 changes over time, 132
 fish, antimicrobial secretions to protect eggs, 77
 quorum sensing, 48
Microdon mutabilis (syrphid fly), matches its ant host, 258
Micromys minutus (harvest mouse)
 captive breeding, using scent marks, 262
Microtus ochrogaster (prairie vole)
 partner recognition, 133, 218
 recognition of family males prevents pick up of male
 pheromones by female juveniles, 147, 219
Microtus pennsylvanicus (meadow vole)
 adjusts ejaculate if sperm competition likely, 89
 male scent glands graded response to testosterone, 75
 males on better diets, more attractive, 36, 75
 Mimicry. *See* Deception
 Minimal-cost signals, 36
 Mitral/tufted (M/T) cells, 183, 187
 involvement in memory, 216–217
 Modulation of response to pheromones and multimodal inputs, 45
Monomorium minimum (ant), mass recruitment to exclude
 competitors, 157
Monomorium pharaonis (Pharaoh's ant)
 multiple trail pheromones, 160
 role specialization, trail laying, 160
 trail polarity, 233
Morinda citrifolia (plant)
 host to *Drosophila sechellia*, 198
 Morpholino oligonucleotides, 60
 Mosaic signal, 6, 10
 Mother–infant recognition, 130–131
 Moths
 male pheromones
 host plant molecules, evolved via sensory exploitation, 99
 multiple independent evolution, 99
 male scramble competition, 69
 pheromones
 behavioral antagonists, 95
 brain wiring flipped in species using opposite ratios of same
 molecules, 97
 multicomponent, male response wider than range females
 produce, 29
 multicomponent, redundancy, 54
 overview, 94–96
 preventing cross-attraction between sympatric species
 time of day, host plants, blends, 95
 saltational shifts, 29
 signal change in speciation, 28–30
 synergy as natural consequence of combinatorial
 processing, 28

- whole blend acts at a distance, 27, 95
- Mouse. *See Mus musculus domesticus*
- Multicomponent pheromones, 19
 - alarm
 - social insects, 170
 - challenges to identification, 54–55
 - definition, 25
 - male response wider than range females produce, 29
 - mechanisms for evolutionary change of blends, 31
 - synergy, 210
 - inevitable outcome of combinatorial processing, 28
 - whole blend acts at a distance, 27, 95
- Multimodal signals. *See* Signals, multimodal
- Mungos mungo (banded mongoose)
 - nasty neighbors, 123
 - over-marking and mate choice, no evidence, 124
- Mup20. *See* Darcin
- Mus macedonicus (mouse)
 - little MUP diversity vs. high density living house mouse, 42
- Mus musculus domesticus (house mouse)
 - contests for ownership of territories, 72
 - co-operative breeders, 146
 - darcin, 219
 - estrus induction and puberty acceleration in females (Whitten & Vandenberg effects), 209–210
 - exocrine gland-secreting peptide 1 (ESP1), 204–206
 - familial imprinting, odor preferences for difference in mates, 85
 - individual recognition, 118, 132
 - major histocompatibility complex (MHC)
 - selection for polymorphism to avoid inbreeding?, 42
 - major urinary proteins (MUPs)
 - add longevity to volatile signal, 23
 - darcin, prompts learning male signature mixture and location, 16, 118
 - haplotype, cue to avoid inbreeding, 218
 - high metabolic cost of use in scent marking, 35
 - mate choice, inbreeding avoidance, 86
 - male pheromones, 72
 - mate choice
 - MHC diversity, 85
 - MHC related odors, 82
 - multicomponent male mouse pheromone, 27
 - Peg3, paternally imprinted gene, odor learning, 209
 - pup odors increase mother response to pup ultrasonic distress calls, 44
 - pups learn individual odor cue of mother as cue for suckling, 66
 - scent marking, 118
 - as reliable signal of territory ownership, 35
 - darcin, 118
 - marking rates, dominants and subordinates, 118
 - mate choice, 118
 - punishment of subordinates if challenge scent marks of dominant, 37
 - reduce cleaning for lab animal welfare, 262
 - response varies by competitive ability, 118
 - subordinates do not secrete pheromones, do not scent mark, 87
 - territories
 - over-marking
 - freshness, mate choice, 124
 - scent matching hypothesis, 117–118
- Mus musculus subspecies (mouse)
 - divergence in odors, 90
- Mushroom body, 183, 207, 216
- Mustela erminea (stoat)
 - anal gland secretions repel herbivore prey, 269
- Mustela putorius furo (ferret)
 - sex differences in response to pheromones, 206
- Mustela vison (American mink)
 - scent baited traps, 263
- Mustelus canis (dogfish), plume tracking, 226
- Musth. *See* Elephas maximus; Loxodonta africana
 - male elephants, 76
- Mutualism, 249–250
 - ants and lycaenid butterflies (*see also* Deception), 250
 - aphids tended by ants, 250
 - egg dumping as, 248
 - sea anemones and anemonefish, 249–250
- Myrmecia gulosa (bulldog ants)
 - cheating workers immobilized, 36
- Myrmica rubra (ant)
 - host to lycaenid butterfly, 258
- Myrmica sabuleti (ant)
 - matching foraging effort to food value, 160
- Myrmica schencki (ant), 255
- Myzus persicae (aphid)
 - response to alarm pheromone, 270
- Naked mole rats. *See* Heterocephalus glaber (naked mole rat)
- Nasogenital grooming, 219
- Nasty neighbors, 123
- Nasutitermes (termite)
 - soldiers, alarm pheromones, 171
- Nasutitermes takasagoensis (termite)
 - queen-specific volatiles, 139
- Natural selection, definitions, 65
- Nauphoeta cinerea (cockroach)
 - body condition as reliable signal, 35
 - body condition reflected in pheromones, 75
 - male contest and female choice, 72
 - silent satellite males, 87
- Near-infrared spectroscopy (NIRS), 57
- Nematodes. *See also* Caenorhabditis elegans
 - multicomponent pheromones
 - ascarosides, 27
 - parasitic, potential control, 266

- Nematodes (cont.)
 pheromones for pest and disease management, 266
 sex pheromones, 266
Neocapritermes taracua (termite), suicidal defense, 171
 Neocembrene, 157, 304
Neofem2, 139
 Nepotism, 128, 131
 Neral, 255
Nereis succinea (polychaete worm)
 pheromones co-ordinate external fertilization, 70, 110
 Next generation sequencing (NGS), 62–63
Nicrophorus defodiens (burying beetle)
 conflict between the sexes, 69
Nicrophorus vespilloides (burying beetle)
 Coolidge effect, 86
Niveoscincus microlepidotus (lizard), partner recognition, 133
 Noise, 95, 178
 Noradrenaline. *See* Norepinephrine
 Norepinephrine
 release in olfactory bulbs
 learning, 216–217
Notropis umbratili (redfin shiner fish) inter-specific egg dumper, 248
 Nuclear magnetic resonance (NMR) spectrometry, 58
Nycticebus pygmaeus (pygmy loris)
 captive breeding, 262
 over-marking, female choice, 124, 262
 Ocean acidification
 dangers to whole ecosystem by interference with chemosensory systems, 46
 (*E*)- β -Ocimene, 23, 145, 214
 Octanal, 75
 Octopamine, 89, 207, 208
 Octyl decanoate, 155
Odocoileus hemionus columbianus (black-tailed deer)
 alert signals, 165
 Odor space, 174, 193
 Odorant binding proteins (OBPs), 198
 specificity of response
 gustation, 198
 Odorants
 can be any size, 173
Oecophylla longinoda (African weaver ant)
 alarm pheromone response, 229
 recruitment systems, 150
 territories
 ‘owner advantage’, 116
Oecophylla smaragdina (weaver ant)
 territories, nasty neighbors, 123
 Oleic acid, 36, 75
 Olfaction. *See also* Gustation, humans
 accessory olfactory bulb (AOB)
 combinatorial processing, 179
 amygdala
 integration of main olfactory and VNO outputs, 199–201
 behavior of smelling (sniffs and flicks), 186
 combinatorial coding, 173, 178–181
 compare lateral inputs, 233, 240
 eavesdroppers as sensitive as legitimate target, 244
 genetic polymorphisms, 191, 292–295
 glomeruli, 176
 links to higher levels of brain, 186
 no simple chemotropic map, 178
 similarity in vertebrates and insects, 176–178
 insect macroglomerular complex (MGC)
 moth male brain, 179
 processing of many pheromones is without, 181–183
 size reflects OSN numbers, 181
 integration of olfactory and hydrodynamic cue inputs, 44
 integration of olfactory and visual inputs, 44
 integration of vertebrate olfactory systems, 198–201
 labeled line, 179, 183
 main olfactory system (MOS)
 functional overlap with vomeronasal olfactory system (VNO), 198–201
 mapping brain activity
 fluorescing dyes, 183
 odorant-binding proteins (OBPs), 197
 odorants
 can be any size, 173
 olfactory bulb
 modulation
 by lateral interactions, 183
 top-down input, 185
 olfactory cortex
 compare lateral inputs, 233
 olfactory receptors. *See* Olfactory receptors
 olfactory sensory neurons (OSNs). *See* Olfactory sensory neurons
 one neuron – one receptor, 177
 possible separation between general olfaction and pheromone processing, 187
 projections to higher brain, 183–187
 response to predator odor, 187
 temporal coding in brains, 187
 fast oscillations, synchronization, 187
 vomeronasal olfactory system (VNO)
 functional overlap with main olfactory system (MOS), 198–201
 Olfactory cortex, 183
 Olfactory organs
 functional design, 185
 sensitivity thresholds compared, dogs, humans, 185
 sensitivity thresholds compared, mice and primates, 275
 Olfactory receptor co-receptor (ORCO) (OR83b), 197
 Olfactory receptors (ORs). *See also* Gustatory receptors (GRs)
 binding sites, 179
 investigated by site-directed mutagenesis, 179
 evolution
 “birth-and-death” process, 189

Cambridge University Press

978-0-521-11290-1 - Pheromones and Animal Behavior: Chemical Signals and Signatures: Second Edition

Tristram D. Wyatt

Index

[More information](#)

- inhibitors (antagonists), 179
- insects
 - co-expressed with olfactory receptor co-receptor (ORCO), 197
 - unrelated to vertebrate receptors, 196
 - use ionotropic signaling pathways, 196
- mammals
 - evolvability facilitated by glomerular development mechanisms, 190
 - odorant binding sites, 190
 - odorant characteristics, 179
 - pH changes ligand interaction, danger from ocean acidification, 46
 - tuning and specificity, 176, 178–181, 190, 191, 292
- Olfactory recess, nasal flow design for sensitivity, 186
- Olfactory sensory neurons (OSNs)
 - odorant effects on development, honeybee, 16
 - renewal (mammal), 191
- Oncorhynchus masou* (masu salmon)
 - female sex pheromone, L-kynurenine, 23
- Oncorhynchus mykiss* (rainbow trout)
 - learn predator odor, 262
- Oncorhynchus tshawytscha* (salmon)
 - MHC mate choice prevented by male harassment, 85
- Ondata zibethicus* (muskrats) monitoring, 264
- Ontogeny. *See* Development
- Ophrys exaltata* (orchid)
 - dupes solitary bee
 - imperfect mimicry more attractive, 252
- Ophrys sphegodes* (orchid) dupes solitary bee, 252
- Optimal outbreeding, 84, 252
- Optogenetics, 60
- OR11H7P, 294
- OR1D2, 298
- OR2J3, 295
- OR7D4, 294
- OR83b (olfactory receptor co-receptor, ORCO), 197
- OR912–93, 193
- Orchids
 - pollination by sexual deception
 - aggressive chemical mimicry, 252–253
- ORCO, 197
- Orconectes rusticus* (crayfish)
 - distributed chemosensory system on claws, 175
- Oreochromis mossambicus* (tilapia, fish)
 - urine release by dominant males during contests on leks, 72, 79
- Oreotragus oreotragus* (klipspringer antelope)
 - scent marks eavesdropped by ticks, 246
 - territories, 121
 - economics of scent marking, 121
- Oriental fruit moth. *See* *Grapholita molesta*
- Orientation behavior
 - active space, 227
 - diffusion, 227–229
 - arrestment, 224
- chemical plumes, 223
 - response to odor filaments, 238
- chemical specificity, 223
- combining information from different senses, 230, 241
 - crustaceans, 238
- directly guided (taxis), 224, 225
- idiothetic, 224
- indirectly guided (kinesis), 224, 225
- insects
 - tracking plumes, 240–241
- mechanisms, 223–226
 - kinesis, 224, 225
 - klinokinesis, 224, 226
 - orthokinesis, 224
 - optomotor anemotaxis, moths, 240
 - taxis, 224, 225
 - chemotaxis, 230, 239
 - klinotaxis, 224, 225, 226, 230, 239
 - rheotaxis, 241
 - teleotaxis, 224
 - tropotaxis, 224, 225, 230, 231, 232, 239, 242
- odor “landscape”, 223
- odor concentration gradient, 225
- odor stimulus
 - plumes, 235–242
 - short range diffusion, 227–229
 - trails, 229–235
- pheromones allow low-density species to find mates, 223
- ranging, 226–227
 - crosswind, 226
 - Lévy walks, 226
- scale, 223
- self-steered, 224
 - counter-turning
 - moths, 240
- teleology, 223
- trails, detecting direction of
 - ants, 233
 - copepods, 230
 - dogs, 233
 - snails, 233
 - snakes, 233
 - stingless bees, 233
- upwind orientation by male moths, 179
 - stop if wrong species, 179
- Oryctolagus cuniculus* (rabbit)
 - chinning, marking behavior, 114
 - fixatives extend signal life, 23
 - mammary pheromone, 11
 - prompts learning other odors, 16, 209
- Osmetrichia, 66
- Osmia rufa* (red mason bee)
 - chooses mates from own population, 84

- Osmoderma eremite* (scarab beetle), rare species monitored via predator, 263
- Ostariophysi, 166
- Ostrinia furnacalis* (Asian corn borer moth)
reconstructing speciation events, 29, 97
- Ostrinia nubilalis* (European corn borer moth)
E & Z strains also isolated by male pheromones, 99
older male pheromones attractive, 76
pheromone signal genetics, 29, 30, 96–97
- Ourebia ourebi* (oribi antelope)
territories, group defense, 121
- Over-marking, 120
mate choice, 118, 123–124, 262
- Oviposition pheromones, 106
Schistocerca gregaria (desert locust), 106
- Oviposition-marking pheromone (OMP). *See* Host-marking pheromone (HMP)
- Ovis aries* (sheep)
male (ram) effect, 221
use in animal husbandry, 261
maternal ewes bond quicker on subsequent births, 209, 217
mother–infant recognition, 130–131, 218
no disassortative mating by MHC, feral sheep, 85
- Ovulin, 89
- (4Z, 11Z)-Oxacyclotrideca-4,11-dien-2-one, 28
- 9-Oxo-2-decenoic acid (9-ODA), 215
- Oxytocin, 78, 216–217, 218
- Pachycondyla laevigata* (ant)
mass recruitment, 150
- Pachycondyla obscuricornis* (ant)
individual hunter, 150
- Panulirus interruptus* (spiny lobster)
aggregation for defense, 105
- Paraponera clavata* (ant), eavesdropped by parasitoid fly, 246
- Parcoblatta lata* (cockroach)
monitoring as proxy for rare bird, 263
multicomponent sex pheromone, 28
- Parkinson's disease
sense of smell, loss, 283
- Péclet number (*Pe*), definition, 228
- Peg3*, paternally imprinted gene, odor learning (mice), 209
- Pella funesta* (staphylinid beetle), uses propaganda, 255
- Pentacosane, 155
- Pentacosene, 155, 254
- People. *See* Humans
- Peptide pheromones, 30–31, 45, 72, 138, 213
Cynops spp. (newts)
species specificity, 25
darcin (mice), 124
exocrine gland-secreting peptide 1 (ESP1), 204–206
Rhithropanopeus harrisii (mud crab), pumping pheromone (egg release), 110
- Perfumes, 277, 281, 282
- Periplaneta* spp. (cockroach)
multicomponent pheromones, 28
- Pesticides, resistance to, 266
- Petromyzon marinus* (sea lamprey)
larval (migratory) pheromone, 24, 267
pest management, 267
pheromone evolution, 21
sex pheromone, 267
3-keto petromyzonol sulphate, 22
tracking plumes, 241
- Petromyzonol sulfate, 24
- pgFAR* fatty acyl reductase gene, 96
- pH changes ligand interaction, danger from ocean acidification, 46
- Pharmacophagy, 31
- Pheidole dentata* (ant)
response to fire ant pheromone, 172
- Pheidole oxyops* (ant)
trail life, short, 158
- Pheidole pallidulato* (ant), recruits for heavy prey, 159
- Phenol, 31
- Phenolics, 246
- Phenotypic matching, 130, 279
- 2-Phenoxyethanol, 23
- Phenylacetoneitrile, 271
- Phenylethanol, 139
- Pheromonal parsimony, 24–25, 43, 157
- Pheromone biosynthesis-activating neuropeptide (PBAN), 89
- Pheromone-binding proteins (PBPs), 198
LUSH (OBP76a), 198
- Pheromone-degrading enzymes in the sensillum lymph, 201
- Pheromones. *See also* Aggregation; Alarm; Applications; Bioassays; Costs; Glands; *Apis mellifera*; Host-marking; Semiochemical identification; Primer effects; Recruitment; Releaser effects; Scent marking; Signals applications. *See* Applications of semiochemicals
aquatic, soluble polar, 167
brain wiring flipped in species using opposite ratios of same molecules, 97
caste differences
production, 32
combinatorial processing in brain, 178–183
control of reproduction, 133–148
convergent, 24
costs of signaling. *See* Signals, costs
definition, 6
differences from signature mixtures, 9, 14
direct transfer, 23
into bloodstream, 30
evolution, 18–23
enabled by olfactory receptor proteins and combinatorial brain circuits, 18

- from pre-existing chemical cues
 - alarm pheromones, 165
 - genomics approaches, social insects, 63
 - how demonstrate, 21–23
 - males responding to cues of previous mates, 22
 - match signal characteristics to requirements, function, habitat, 23
 - receiver pre-adaptations (sensory bias model), 21
 - ritualization, 18, 19, 21, 113
 - salamander pheromones, 30–31
 - sender-precursor model, from cues, 18–21
- evolutionary change in multicomponent pheromones, 31
- excreted in urine, 19
- glands
 - social insects, summarized, 135
- gustatory response
 - invertebrates, 15
- history, 5, 49
- humans. *See* Humans
- identification. *See* Semiochemical identification
- induce gene expression in brain, 43, 62, 172
- interaction with signature mixtures, 218–221
- multicomponent
 - alarm
 - social insects, 170
 - challenges to identification, 54–55
 - social insects
 - trail, 157
 - solitary bees, 254
 - synergy, inevitable outcome of combinatorial processing, 28
 - whole blend acts at a distance, 27, 95
- neural processing (ON/OFF/AND) for YES/NO, 28
- operational definition, 9–10, 23
- primer effects. *See* Primer effects
- production, 31–32
 - by microbes, 31
 - collect from plants, 31
 - synthesized by sender, 31
- prompting learning other odors, 16
- prompting learning, flexibility in behavior, 208–209
- recruitment. *See* Recruitment pheromones
- response, 43
 - conditional, 206
 - context dependent
 - aphids, 168
 - aphids tended by ants, 249
- saltational shift, 29, 97
- same molecules shared across species, 24
- signals vs cues, 18
- sources. *See* Pheromones, production
- species share molecules if no selection against, 24
- specificity
 - exploring chemical space for related molecules, 28
 - multicomponent, 28
 - unique molecule, 25–27
- synergy, inevitable outcome of combinatorial processing, 28
- trail. *See* Recruitment pheromones
- Philanthus triangulum* (beewolf wasp)
 - males lek, 80
 - pre-existing female sensory bias for prey pheromones, 21
- Phodopus campbelli* (Djungarian hamster)
 - recognition of individuals of other hamster species, 40
- Phthorimaea operculella* (potato tuber moth)
 - pheromone blend ratio varies with rearing temperature, 92
- Picoides borealis* (red-cockaded woodpecker)
 - monitoring cockroach prey by pheromone, 263
- Pieris brassicae* (large cabbage white butterfly)
 - anti-aphrodisiac, 88
 - eavesdropped, 245
- Pieris napi* (green-veined white butterfly)
 - anti-aphrodisiac, 88
- Pieris rapae* (small cabbage white butterfly)
 - anti-aphrodisiac, 88
- Pigs. *See* *Sus scrofa*
- Piperidine alkaloids, 212
- Plankton, 230
 - larval settlement and metamorphosis, 110
- Plants. *See also* Pheromones, production
 - “calling for help”, 244
 - feedstock precursors for synthetic semiochemicals, 273
 - orchids, pollination by sexual deception
 - aggressive chemical mimicry, 252–253
 - self-defense by producing herbivore alarm pheromone, 255
 - transgenics produce aphid alarm pheromone, 270
- Platynereis dumerilii* (polychaete worm)
 - pheromones co-ordinate external fertilization, 70, 110
- Platysoma cylindrica* (predatory beetle), eavesdropping, 246
- Pleiotropy, 94
- Plethodon shermani* (red-legged salamander)
 - direct transfer of peptide pheromone to female’s nostrils, 23, 45
- Plethodontid salamanders
 - pheromone evolution, 30–31
- Plumes
 - eddy chemotaxis, 241
 - turbulence, visualization, 228, 235–238
- Podisus maculiventris* (spined soldier bug), eavesdropped by parasitoid, 245
- Poecilia reticulata* (guppy, fish)
 - cryptic female choice, 89
- Pogonomyrmex badius* (harvester ant) alarm pheromone signal, 229
- Pogonomyrmex barbatus* (harvester ant)
 - CHCs of foragers recruit others, 160
- Pogonomyrmex* spp. (harvester ants), trail species
 - specificity, 157

- Polistes* (paper wasps)
 colony recognition, 129
 evaded by cuckoo species, 257
Polistes dominulus (social wasp), 257
Polistes fuscatus (paper wasp)
 kin recognition mechanisms, 129
 visual recognition to distinguish individual colony members, 40
Polistes metricus (paper wasp)
 lacks specialized face learning abilities, 40
Polistes semenowi (social wasp, parasite), 257
 Pollution, 45–46
 anthropogenic carbon dioxide and ocean acidification
 dangers to whole ecosystem by interference with chemosensory systems, 46
 endocrine-disrupting effects on olfaction, 45–46
 pheromones in aquaculture, risk, 262
Polyergus rufescens (slave-making ant), use of propaganda, 255
Popilla japonica (Japanese beetle), 25
 Pre-copulatory mate guarding. *See* Sperm competition and mate guarding
Premnas spp. (anemonefish)
 symbiotic with sea anemones, 249
 Primates
 chemical profile differences in related species, 91
Homo sapiens. *See* Humans
 over-marking
 female choice, 262
 reproductive suppression, 146
 rhesus monkeys
 copulins, 299
 scent marking, 124
 manipulate mate choice, 262
 Primer effects, 43–44, 300, *See also* Releaser effects
 can be fast acting, 44
 dauer pheromone (*Caenorhabditis elegans* (nematode)), 25, 210–211
 intergenerational
 alarm pheromones in aphids, 169
 desert locusts, 106
 mammals, 209–210
 by either VNO or main olfactory system, 210
 sexually dimorphic hypothalamic neurons releasing kisspeptins, 200–201
 menstrual synchrony, 300–301
 modulation of hormone signaling and gene expression, 209
 social insects
 caste development, 211–213
 transcriptomics, to investigate, 62
 use in pest management, 271
 Projection neurons, 183
 Propaganda (use of alarm pheromones to manipulate other species), 255
See also Deception
 3-Propyl-1,2-dithiolane, 269
Prorhinotermes simplex (termite)
 sex differences, 138
 Prostaglandins, 45, 70
Prostephanus truncatus (larger grain borer beetle), 108
 aggregation pheromone
 monitoring, 263
 males only signal until females arrive, 109
Proteles cristata (aardwolf), 123
 neighbor recognition, 133
 Protocerebrum, 183
 Proust, smells bring back memories, 280
Pseudacanthotermes spiniger (termite)
 trail and sex pheromones, same molecule at different concentrations, 24
Pseudacteon tricuspidis (phorid fly), eavesdrops host alarm pheromone, 246
 Pseudogates (termite nymphs), 138
 Puberty acceleration. *See* Hormones
 Puberty delay
 rodents
 questions about ecological relevance, 147
 Public chemical information, 165
 PUFAs (polyunsaturated fatty acids)
 barnacle egg hatching pheromone, 110
Puntius (fish) (androgen effects on electro-olfactogram (EOG)), 207
 Purines, 107
 Pyranones, 212
 Pyrrolizidine alkaloids, 31
 Quantitative trait locus (QTL), 29, 98, 103
 Queen butterflies. *See* *Danaus gilippus*
 Queen egg-marking pheromone, 36
 Queen mandibular pheromone (QMP). *See* *Apis mellifera*
 Queen pheromones. *See also* Ants; *Apis mellifera*; Pheromones; Termites
 as honest signal, 36, 214
 developmental switches
 mechanisms, 211–213
 social insects, 137–146
 Quorum sensing, 48
 Rabbits. *See* *Oryctolagus cuniculus*
 Racemic mixture (racemate), 307
 Ram effect. *See* Male effect
Rattus norvegicus (Norway rat)
 behavior manipulated by protozoan parasite *Toxoplasma gondii*, 248
 follows trails from good food sources, 155
 sniffing in stereo, compare in olfactory cortex, 233
 Receiver psychology (hidden preferences), 21
 Recognition, 215–218
 dominants, reproductive suppression, 146
 imprinting, 38–40
 learning, 38–41

- mammals, 130–133
 - clan or group, 131–132
 - individual recognition, 132–133
 - kin, 131
 - mother–offspring, 130–131
 - prairie vole partners, mechanisms, 218
 - some glands more variable than others, 133
- mechanisms, 37
 - bioassays to study, 52
 - greenbeards, 40
 - individual recognition, challenges in demonstrating, 132–133
 - phenotypic matching, 38, 130, 131, 279
 - self-matching (armpit effect), 38
 - template, 13
- mothers and babies, 278–279
- social insects
 - colony and kin recognition, 126–130
 - learning needed before discrimination, 126–130
- social species
 - have more complex chemical profiles?, 41–42
- template, 13
 - brain imaging, social insects, 216
- Recruitment pheromones. *See also* Alarm pheromones; Self-organizing systems
 - alarm (social insects), 169–172
 - Apis cerana japonica* (Japanese honeybee), 170–171
 - coordinate defense, 170–171, 255
 - marking enemies for further attack, 169, 170, 171
 - termites, 171
- ants
 - alarm, 170
 - competition strategy, 157
 - group recruitment, 153
 - mass communication, 153
 - modeling foraging, 160–161, 162
 - multiple pheromones, flexibility, 159
 - tandem running, 153
 - trail components with different roles, 50
 - trail specificity, 157
- bumblebees
 - foraging recruitment pheromone, 155
- colony specificity
 - stingless bees, 155
- expand colony diet, 153
- food distribution, recruitment type, 150–153
- Heterocephalus glaber* (naked mole rat)
 - recruitment of foragers, 155
- honeybee
 - with waggle dance, 155
- longevity related to food supply, 158
- Malacosoma americanum* (tent caterpillar), 152
- matching foraging effort to food value, 159–160
- Rattus norvegicus* (Norway rat) follow trails from good food
 - sources, 155
- social insects to co-ordinate attack, 169
- species specificity
 - ants, 157
 - stingless bees, 155
 - termites, 155–157
- stingless bees (Meliponini)
 - odor beacons, 154–155
- termites
 - foraging patterns, 158
- Trigona* spp. (stingless bees)
 - odor beacons, 156
- Redundancy (signal), 54
- Reinforcement (selection against hybridization), 90
- Releaser effects, 43–44. *See also* Primer effects
- Reliable signals. *See* Signals, honest
 - without handicap
 - individual or colony identity, 36
- Replication
 - common lack of, 53
 - good example of value, 210
 - vs. quasireplication, 53
- Reproductive skew, 136, 137
- Reproductive suppression
 - primates, 146
- Reticulitermes santonensis* (termite)
 - sex differences, 138
- Reticulitermes flavipes* (termite)
 - host to beetles, 257
 - soldier terpenes change workers' gene expression, 212
- Reticulitermes speratus* (termite)
 - queen pheromones and caste development, 139, 212
- Reynolds number (*Re*)
 - copepod pheromone trails, 230
 - definition, 228
 - effects on sense organ design, 184
- Rhagoletis pomonella* (apple maggot fly)
 - learning own scent marks, 209
- Rhinella marina* (cane toad)
 - alarm cue, 269
 - management of, 269
- Rhithropanopeus harrisi* (mud crab)
 - pumping pheromone (egg release), 110
- Ritualization, 18, 21
- RNA interference (RNAi), 60, 213
- RNA-seq, 62–63, 91
- Ropalidia marginata* (paper wasp)
 - colony recognition mechanisms, 129
 - non-kin may join colony, 129
- Rotifers
 - do not use diffusable pheromones, 227, 230
 - sex pheromones, 60
- Royal jelly (honeybee), 213

- Royalactin (honeybee), 213
Runaway sexual selection, 74
- Saccolpteryx bilineata* (greater sac-winged bat)
 semiochemicals, microbial contribution, 41
Saguinus fuscicollis (saddle-back tamarin)
 chemical profile, multiple messages in odors, 126
Salaria pavo (peacock blenny, fish)
 male parental care, 77
 honest signal, 35
Saltational shifts, 29, 97
Satellite males. *See* Alternative mating strategies
Saturnia pyri (great peacock moth)
 Farbre's early account, 5
Scale, 229
 communication strategies, 227, 230
 diffusible pheromones in water, 229
 sensory organs, functional design, 185
Scaptotrigona postica (stingless bee), odor beacon
 recruitment, 157
Scent marking, 118
 advertises dominance and/or reproductive status, 114, 115, 124
 captive breeding, exploiting scent marks, 262
 conspicuous signals, 123
 ecological factors and marking patterns, 115, 121, 123
 males directly mark females, 124
 marking rates, dominants and subordinates, 118
 non-territorial mammals, 124
 pest management
 predator odors, as deterrents, 269
 response varies by competitive ability, 118, 115, 118
 scent-fence hypothesis, 115
 scent-matching hypothesis, 115–118
 reducing cost of territorial defense, 116
 self-marking and presentation, 115
 use to manipulate behavior, 262
 using to census, 263–264
 using vertebrate scent marks to manipulate behavior, 269
Schedorhinotermes lamanianus (termite)
 foraging, 158, 159
Schistocerca gregaria (desert locust)
 adult cohesion pheromone, phenylacetonitrile, 271
 aggregation pheromones, 106
 gregarizing factor on eggs, 106
 intergenerational effects, 106
 oviposition pheromones, 106
 possible control of gregarization, 271
 solitary gregarious phases, 106
Schistosoma mansoni (nematode)
 sex pheromone, 266
Schreckstoff, 166. *See* Alarm cues, fish
Scramble competition, 69–71
 moth males, 69
 selection on male moths, 70
Sehirus cinctus (subsocial bug)
 solicitation pheromone, begging, 36
Self-organizing systems, 160–163
 ant foraging, 160–161
 stochastic effects, 161
 stigmergy, 162
 termite nests, 161–163
Self-referent ("armpit") matching, 38
Semibalanus balanoides (barnacle)
 egg hatching pheromone
 eicosanoids (PUFAs), 110
Seminal fluid molecules
 allohormone pheromones, 88–89
 behavioral manipulation, 45
 prostaglandins, 45
Semiochemical identification
 bioassays. *See* Bioassays
 collection and analysis, 55–59
 activity-led fractionation, 56–57
 gas chromatography–mass spectroscopy (GC-MS), 57
 high performance liquid chromatography (HPLC), 55
 liquid-chromatography–MS (LC-MS), 57
 nuclear magnetic resonance spectrometry, 57
 collection and analysis without fractionation, 57–58
 headspace sampling, 57
 in situ analysis, 57
 metabolomics, 57–58
 near-infrared spectroscopy (NIRS), 57
 nuclear magnetic resonance (NMR) spectrometry, 58
 solid-phase microextraction (SPME), 57
 stir-bar sorptive extraction (SBSE), 57
 UV laser desorption/ionization orthogonal time-of-flight MS (UV-LDI-TOF MS), 57
 loss-of-function mutants as tool, 50, 58
 molecular biology and genetics as tools, 59–63
 cautions, misleading effects, 60
 gene manipulations, 60–62
 genomics, 62–63
 knockouts, transgenics, 60
 morpholino oligonucleotides, 60
 optogenetics, 60
 RNA interference (RNAi), 60
 RNA-seq, 63
 visualizing neural circuits, 60–62
 no single ideal approach, 58
 pheromones (operational definition), 9–10
Semiochemicals. *See also* Bioassays
 applications. *See* Applications of semiochemicals
 cues
 definition, 18
 definitions, 1, 6
 allelochemicals, 1
 allomones, 1
 chemical profile, 2
 kairomones, 1

- pheromones, 5
- pheromones (operational definition), 9–10
- signature mixtures, 10
- synomones, 1
- innate, 16
- signature mixtures and pheromones compared, 9, 14
- Sensitive periods, 216
 - behavioral development, 16
 - imprinting, 16
 - learning colony odors (social insects), 129
- Sensory bias
 - food molecules, 21
 - host-plant odors, 21, 99
 - pre-existing sensitivities exploited by signalers, 21
 - prey pheromones, 21
- Sensory organs
 - functional design, 185
- Sensory trap
 - contribution to pheromone evolution, 21
- Septal organ of Maseru, 195
- Serotonin (5-hydroxytryptamine) (5HT), 208
- Settlement inducing protein complex (SIPC), 109
- Sex-specific neural circuits
 - insects, 201–204
 - moths, 201
 - mammals, 204–206
- Sexual conflict, 31
 - anti-aphrodisiac pheromones, 87–88
 - costs of “sensory trap”, 21
 - driving sex peptide variety, 21, 89
 - interference in partner’s signaling to prevent competitors, 69
 - seminal fluid molecules, 88–89
- Sexual dimorphism
 - hypothalamus, kisspeptin neurons, 200–201
 - moths, male antennae and brain, 201
- Sexual selection
 - conflict between the sexes, 69, 87–88, 89
 - contests, 72–73
 - co-ordination, external fertilization, 69
 - definitions, 65–66
 - forms of, 65
 - mate choice, 73–87
 - mechanisms
 - indicators, 74
 - overview, 73–75
 - post-copulatory, 87–90
 - anti-aphrodisiacs, 87–88
 - cryptic choice of sperm by female, 89–90
 - based on MHC, 85
 - seminal fluids, 88–89
 - sperm competition, 89
 - pre-copulatory mate guarding, 71–72
 - scramble competition, 69–71
 - species-specific secretory glands, 66
 - variety of pheromone glands in males, 31
 - which sex should advertise, 66–69
- Sharks
 - lateral line for rheotaxis, 241
 - plume tracking, 226
- Sheep. *See Ovis aries*
- She-males. *See* Alternative mating strategies
- Signals. *See also* Pheromones
 - communication, 18
 - costs
 - acoustic signals (high metabolic cost), 34
 - efficacy cost, 32, 34
 - energy as proxy for fitness costs, 34
 - pheromones
 - most, low costs, 34
 - some, high efficacy costs (time, energy, risk), 34–35
 - risk investigating marks, beavers, 114
 - strategic cost (handicap cost), 32
 - time marking, antelope, 114
 - cues vs. signals, 18
 - definition, 18
 - duration, 23
 - extending signal life, 23
 - evolution
 - co-evolution of sender-receiver, 30–31
 - match signal characteristics to requirements, function, habitat, 23
 - receiver psychology (hidden preferences), 21
 - handicap theory, 32, 33–34
 - honest (reliable) signals without handicap, 32–37
 - body condition and testosterone/androgen effects on males, 75–76
 - indices (unfakeable signals), 35–36
 - other costs including predation, 37
 - shared interest including kin selection, 36
 - signs of paternal investment, poisons and parental care, 76–77
 - social cost (punishment), 36–37
 - social insect queen pheromones, 137–146
 - how receivers follow changes in signal, 93–94
 - minimal cost signals, 32–37
 - multimodal signals, 44–45, 75, 167, 171
 - challenge for bioassays, 54
 - pheromone and tactile signals, 150
 - scent marks with ultraviolet cue, 123
 - sound and odors, 155, 157
 - queen pheromones
 - control or co-operative signal, 137–146
 - redundancy, 44, 54, 167
 - challenge for pheromone identification, 54
 - deer alert signal, 44
 - ritualization, 21
- Signature mixtures, 10–15, 278, 280
 - as “receiver-side” phenomenon, 2, 15
 - definition, 6, 10

- Signature mixtures (cont.)
 differences from pheromones, 9, 14, 37
 identification. *See* Bioassays
 interaction with pheromones, 218–221
 learning, 37, 38–41, 215–218
 olfactory system, 37
 social insect variation in trail marking between colonies,
 stingless bees, 155
 social mammals
 individual recognition and reproductive suppression,
 146–147, 219
- Simuliidae (blackfly)
 oviposition pheromones, 106
- Single nucleotide polymorphisms (SNPs), 191, 294
 olfactory receptors, 292
 secretions (*ABCC11*), 290–291
- Sitophilus* (stored product beetle), 108
- Smell. *See* Olfaction
- Smell-O-Vision, 303
- Snakes
 anti-aphrodisiac pheromones, 88
 Coolidge effect, 86
 show defensive behavior to predatory snake odors, 248
 trail following
 forked tongues, role, 232
 tropotaxis, 232
- Sniffing
 in stereo, 233
 initiates oscillations in brain, 186, 187
- SNPs. *See* Single nucleotide polymorphisms
- Social insects. *See also* Ants; Eusociality; *Apis mellifera*;
 Recruitment pheromones; Self-organising systems;
 Termites
 alarm pheromones, 169–172
 evolved from defensive molecules, 20–21
 caste development, 211–213
 caste differences
 pheromone secretions, 32, 135
 response to pheromones, 135
 colony and kin recognition, 126–130
 colony level selection, 214
 colony odors, 36, 134
 selection by parasites for greater diversity in hosts, 42
 selection by social parasites for greater diversity in hosts, 258
 sources, 41
 communication largely minimal-cost, kin selection, 36
 co-opting genes from solitary ancestors, 215
 differences between queens and workers
 gene expression, 214
 fertility signal, 36, 137–146
 genetic CHC differences between patriline, 130
 guards
 defense of colonies, 126, 170
 importance of colony defense, 170
 individual recognition by founding ant queens, 38
 key roles for pheromones, enabling sociality, 134
 main pheromone glands, summarized, 135
 multicomponent pheromones, 28
 multiple uses of the same pheromone (pheromone parsimony),
 135
 newly eclosed, need to pick up colony odor, 129, 130
 parallels with social mammals, 147–148
 queen egg-marking pheromone, 36
 queen pheromones
 control or co-operative signal, 137–146
 genomics to explore evolution of, 63
 reproductive conflict, 136–137
 suicidal defense, 169, 170, 171
 territories, 116
 worker policing, 36, 142–146
 workers on different tasks
 differences in brain gene expression, 214
- Social mammals. *See also* Eusociality
 clan recognition, 132
 ecological factors, 147
 parallels with social insects, 147–148
 reproductive conflict, 136–137
 reproductive control
 not by pheromones, 146–147, 219
 suppression of helper reproduction, not by pheromone,
 146–147, 219
- Sociogenomics, 212
- Solanum berthaultii* (wild potato), produces aphid alarm
 pheromone, 255
- Solenopsis invicta* (fire ant)
 alarm pheromones eavesdropped by phorid fly, 246
 enemy specification, 172
 greenbeard gene *Gp-9*, 40
 mass communication, 153
 queen mutual inhibition, 212
 queen pheromones, 212
 queen pheromones, as control, 271
 recruitment, 159
 trail pheromone, 231, 270
 alerting and orienting components, 50
- Solid-phase microextraction (SPME), 57
- Speciation, 24
 allopatric, 90
 assortative mating (like with like), 90, 96
 asymmetric tracking, 94
 disruptive selection, 94
 ecological races, 92
 infertile hybrids, 91
 mate choice, 31
 pheromones allow rare morphs to find each other, 94
 pre-mating isolation, role for chemosensory barriers, 91
 process
 signal change, how receivers track changes, 93–94
 reproductive character displacement, 90
 selection against hybridization (reinforcement), 90

- sexual selection
 - role in founding phase of new island populations, 92
- signal change, few genes involved, 278
- signal changes involving many genes, polygenic, 29, 97–98
- signal variation, within and between populations, that selection can act on, 92–93
- sympatric, 90
 - orchids, different pollinators, 253
- Speciation genes, 91, 101
- Sperm competition, 89
- Spermophilus beldingi* (Belding's ground squirrel)
 - kin recognition cues, 131
- Spiders
 - conflict between the sexes, 69
- Spodoptera littoralis* (moth)
 - bat clicks increase male response to pheromone, 44
- Sry gene, sexual differentiation in mammal embryo, 206
- Stabilizing selection, 92, 93
- Standing behavior. *See* Lordosis
- Staphylococci*, 288
- Stereoisomers (spatial isomers), 25, 306
- Stigmergy, 162
- Stingless bees (Meliponini)
 - robber species and resistance, 255
 - trail beacons to food sources, 154–155
- Stir-bar sorptive extraction (SBSE), 57
- Stomatopod
 - antennal flicking rate scales with increasing size in lifetime, 186
- Stored product insects
 - monitoring, 263
- Stored-product beetles, 108
- Strategic cost, 32
- Subesophageal ganglion (SEG). *See* Subesophageal ganglion (SOG)
- Subesophageal ganglion (SOG) (insects)
 - gustatory sensory neurons (GSNs), feed to, 197
- Sulfanyalkanols (thio-alcohols), 290
- Sulfated amino-sterol, 72, 79
- Sulfated steroids, 195
- Supella longipalpa* (brown-banded cockroach)
 - unique molecule as sex pheromone, 25
- Supellapyrone, 25
- Superparasitism
 - host marking, to avoid, 110
- Sus scrofa* (pig)
 - advancing puberty, 261
- BoarmateTM
 - lordosis readiness test for AI, 261
 - reduces aggression, young pigs, 263
 - reduces postpartum interval, 261
- Sweatbees
 - colony recognition, 128–129
- Synergy, 58, 142
 - definition, 28
 - natural outcome of combinatorial processing, 28
 - testing, 28
- Talpa europaea* (mole), territories, 115
- Taste
 - compared with smell (olfaction), 175
 - smell (olfaction) vs. taste (gustation), 15, 173
- Taxis, 224, 225
- Telenomus euproctidis* (parasitoid wasp), phoresy, 245
- Teleogryllus oceanicus* (cricket)
 - male adjusts ejaculate vs. sperm competition, 89
 - uses CHC profiles to avoid genetically similar mates, 81
- Teleology, 223
- Temnochila chlorodia* (predatory beetle), 246
- Template (for recognition), 15, 127
 - brain imaging, social insects, 216
 - definition, 13
- Termites
 - alarm pheromones, 169–172
 - caste-change pheromones, 212
 - colony recognition, 130
 - differences from Hymenoptera, 138
 - differences in gene expression, queens vs. kings and workers, 139
 - nest building as self-organizing system, 161–163
 - pheromonal parsimony, 24, 157
 - pheromones and foraging patterns, 158
 - primer effects, 138–139, 212
 - queen pheromones, 212
 - sex differences in signals
 - polar proteinaceous secretions, 138
 - soldiers, control of numbers, 139, 212
 - suicidal defense, 171
 - trail following, 232
 - trail pheromones, 155–157
 - used as sex pheromones, at high concentration, 157
- Territories, 113, 116. *See also* Scent marking
 - “owner advantage”, 116
 - dear-enemies, 118, 123
 - economics of scent marking, 121–123
 - hinterland and perimeter marking, 121, 123
 - variation by habitat, 121, 123
 - group defense, 113, 114, 120–121
 - nasty neighbors, 123
 - scent marking, 118
 - border maintenance hypothesis, 120–121
 - composite signals with ultraviolet cue, 123
 - lamp-post effect, 44, 123
 - scent fence hypothesis, 115
 - scent matching hypothesis, 115–120
 - social insects, 116

- Testosterone. *See* Hormones
- Tetanolita mynesalis* (moth)
victim of bolas spider, 251
- (*E*)-11-Tetradecenyl acetate, 29, 96
- (*Z*)-9-Tetradecenyl acetate, 251
- (*Z*)-11-Tetradecenyl acetate, 29, 95, 96
- (*Z,E*)-9,12-Tetradecenyl acetate, 251
- Tetradecyl acetate, 255
- Tetragonisca angustula* (stingless bee)
alarm pheromones to activate colony defense, 255
resists raids by robber bees, 255
- Tetramorium caespitum* (ant), foraging, 161
- Thamnophis sirtalis parietalis* (red-sided garter snake)
anti-aphrodisiac, 88
chooses mates from own population, 84
males choose females in better condition, pheromone based, 76
population differences in chemical profiles, 92
semen effects on female behavior, 45
she-males, 87
trails
detecting direction, 233
following, 232
- Thanasimus dubius*, (predatory beetle),
eavesdropping, 246
- Thiazole. *See* Butyl dihydrothiazole
- Ticks
aggregate on scent marks of mammal hosts, 246
assembly pheromone
ecophysiological benefits, 107
guanine, assembly pheromone, 107
pheromones, to control, 266
- Tiger moths. *See under* *Utethesia ornatrix*
- Tomicobia tibialis* (parasitoid wasp), eavesdrops bark beetle host, 247
- Tools to study pheromones. *See* Bioassays; Semiochemical identification
- Toxoplasma gondii* (protozoan parasite)
changes rat's fear of cat odor to attraction, 248
- Trace amine-associated receptors (TAARs), 192, 193
- Trade-offs, of energy allocation, 35
- Trail pheromones. *See* Recruitment pheromones
- Transcriptome, 62
- Transforming growth factor- β (TGF- β) peptides, 211
- Tribolium castaneum* (flour beetle)
aggregation pheromone, 109
cryptic female choice, 89
- Trichogramma brassicae* (parasitoid wasp),
phoresy, 245
- Trichogramma evanescens* (generalist parasitoid), learns host pheromones, 245
- Trichogramma pretiosum* (parasitoid wasp), oviposition marking, 110
- Trichoplusia ni* (cabbage looper moth)
both sexes call, 69
- Trichoplusia ni* (cabbage moth)
multicomponent pheromone
redundancy, 54
pheromone blend shift, males follow, 30, 96
- Trichopsenius frosti* (beetle), matches its termite host, 257
- Tricosane, 155
- Tricosene, 254
- Z*-(9)-Tricosene, 155
- (*E*)-4-Tridecenyl acetate, 266
- Trigeminal system, 195
- Trigona* spp. (stingless bees), 154–155, 255
odor beacon recruitment
Trigona corvina, 155
Trigona hyalinata, 157
Trigona recursa, 154, 156
Trigona spinipes, 154
victim to propaganda pheromones
Trigona subterranea, 255
- Trimethyl-thiazoline, 187
- Tritrophic systems, 244
- Trogoderma glabrum* (khapra beetle)
aggregation pheromone, 109
- Trophallaxis, 129, 138
- TRP2C, 199
- Ultraviolet (UV) cues
Dipsosaurus dorsalis (desert iguana), multimodal signal with UV, 44
- Ultraviolet laser desorption/ionization orthogonal time-of-flight mass spectrometry (UV-LDI-TOF MS), 57
- Unintended bias in experiments, avoiding, 53
- Uric acid, 70
- Uridine diphosphate, 71
- Utethesia ornatrix* (tiger moth)
cryptic female choice, 90
female choice, 76
honest signal, 35, 76
hydroxydanaidal (HD) from diet, 31, 76
male pheromone evolved by sensory exploitation, 99
- cis*-Vaccenyl acetate (cVA) 62, 253
- Vandenbergh effect, 210
- Vanillic acid, 266
- Varroa destructor* (parasitic mite)
response to honeybee larval odors, use as kairomones, 136
- Vasopressin, 218
- Vertebrate pests
Castor canadensis (beaver), 269
Cyprinus carpio (common carp), 267
Mustela vison (American mink), 263
Petromyzon marinus (sea lamprey), 267
Rhinella marina (cane toad), 269
scent marks to census, 263–264

<i>Verticillium lecanii</i> (fungal pathogen), 270	Wikipedia, how scientists can contribute, xiv
<i>Vespa mandarinia japonica</i> (giant hornet), 170–171	Worker policing, 36, 142–146
<i>Vespula germanica</i> (wasp)	
eavesdropping prey leks, 37, 245	<i>Xiphophorus</i> spp. (swordtails, fish), leads to sexual isolation of
<i>Vespula vulgaris</i> (wasp)	two species, 75
marks with sting venom, alarm, 171	<i>Xylotrechus pyrrhoderus</i> (beetle)
<i>Vipera berus</i> (adder, snake), Coolidge effect, 86	stereochemistry of pheromone, 309
Vitellogenin, 214	
Vomeronasal olfactory system	<i>Yponomeuta</i> spp. (small ermine moths)
memory and pregnancy block (Bruce effect), 219–221	sympatric, avoiding cross-attraction
molecular architecture and differences from main	calling time, host plants, pheromone blend, 95
olfactory system (MOE), mapping to glomeruli,	
receptor proteins, 193–195	Zigzag, tracking trail or plumes, 236
Vomeronasal organ (VNO), 193–195	albatross, 242
humans, vestigial, 291	dogs, 233
Vomeronasal receptors VRs, 193–195	humans, 233
	moths, 240
Westermarck or kibbutz effect	snakes, 233
human mate choice, 279	<i>Zootermopsis nevadensis</i> (termite)
Whales and dolphins, 12	fertility signaling, 138
Whitten effect, 146, 210, 220	laying trail, 154