

Index

- accretion rate measurements 45–47, 74–75, 165–166
- Acropora palmata* 31
- Agropyron pungens* 102, 170
- Agrostis stolonifera* 69
- Althaea officinalis* 91
- Aphelia vibernana* 91
- Aphrodes bicinctus* 91
- Armeria maritima* 74, 170
- Artemesia maritima* 90
- Aster tripolium* 74, 86, 90, 91, 169
- Atriplex hastata* 74
- Awre Formation 132, 136
- barrage schemes 109
- Beta maritima* 74
- birds 81, 92
 - Blackwater Estuary 3, 43, 45–46, 57–60
 - Blakeney Point 158, 165, 167, 173
 - Bristol Channel 122
 - Caldicot Level 128
 - Calluna vulgaris* 25
 - Chenopodiaceae 25
 - Cladium mariscus* 25
 - Coleoptera 91
 - coastal defences 115–120
 - competition 68–70
 - Crouch Estuary 3, 43
 - Dee Estuary 3, 90, 104, 106–107
 - Dengie Peninsula 3, 14, 15, 44, 55–57, 98, 109
 - dieback (of *Spartina*) 74
 - Diptera 91
 - Downholland Moss 23–25
 - drainage system development 53–60, 128, 133, 160–161
 - ducks 92
 - Elymus pycnanthus* 74
 - Enteromorpha* 169
 - Ericaceae 25
 - erosion of marshes 11–12, 115
 - Fenland 23, 32
 - Festuca rubra* 52, 55, 69, 86, 98, 102
 - Foulness 3
 - Frankenia laevis* 89
 - Furzey Island 73–75
 - geese 92
 - geoid changes 6, 34
 - grazing 90, 92, 97–98, 100, 102
 - Gwent Levels 81, 98
 - Halimione portulacoides* 73, 75, 89, 100, 169
 - Hamford Water 3, 43
 - Hudson Bay 65
 - Humber Estuary 3, 50
 - hydrodynamics 162–165
 - Inula crithmoides* 90
 - invertebrate fauna 90
 - Iris pseudacorus* 90
 - Juncus gerardii* 74, 86, 170
 - Juncus maritimus* 90, 170
 - Langstone Harbour 3, 89
 - lapwing 92
 - Limonium binervosum* 90
 - Limonium vulgare* 74, 100, 169
 - limus 24, 25, 29
 - loch-head marshes 3, 85–86
 - management of saltmarshes 100–103
 - mangroves 1
 - marsh cliff retreat 4, 49–51
 - Medway Estuary 3, 109
 - Mersey Estuary 3
 - minerogenic sediments 4, 123, 138–140
 - molochite plugs 74
 - mud mounds 11, 13, 51
 - National Nature Reserves 96–97
 - National Rivers Authority 98, 120
 - National Vegetation Classification 86–87, 94
 - niche 63–68
 - niche overlap hypothesis 66
 - Northwick Formation 132, 136
 - Nymphaea* 25
 - Oldbury Surface 128, 130
 - organogenic sediments 4, 123, 138–142
 - oystercatcher 92
 - Pagham Harbour 3
 - Pedicularis* 24
 - Phragmites communis* 24, 25, 86, 89, 92, 170
 - Pilling Marsh 22
 - Plantago maritima* 74

- Poole Harbour 74–76, 89
Portsmouth Harbour 3
productivity 92
Puccinellia maritima 67, 68–9, 73, 86, 92, 98, 102, 170
- RAMSAR Convention 96
reclamation 128–132, 156
redshank 92
restinga 26
Ribble Estuary 3, 82–83, 90
Rio de Janeiro State 16
roddons 23
Ruel Estuary 90
Rumney Formation 127, 135
- Salicornia* spp. 73, 86, 89, 169
salt pans 160
Scot Head Island 158, 164, 167
Severn Estuary 123–137
sea banks 81, 98, 121
sea-level rise 6–11, 19–35, 60–61, 125–6, 155–156
sediment lamination 137
sediment sources 4, 125, 152–3
shingle barriers 3, 158–159
site protection 93–97, 115
Sites of Special Scientific Interest 83, 93–5, 115, 148
Solway Firth 3, 97
Somerset Levels 132
Southwest Lancashire 23–25
Southampton Water 84
- Spartina alterniflora* 88
Spartina anglica 66–67, 74–75, 86, 89, 99, 103, 169
Spartina maritima 88
Spartina townsendii 88
storm surges 22, 34, 98, 171–172
stratigraphic roll-over 127
succession 70–76
Suaeda fruticosa 89
Suaeda maritima 86, 89, 169
- Tay Estuary 3
Tees Estuary 11, 84, 105
Thames Estuary 3
The Wash 81, 84, 89, 108, 151
tidal range 5, 125, 151
tide gauge records 8, 20–21
Tollesbury marshes 59, 61
Triglochin maritima 74, 170
turf cutting 97
turfa 27, 29
Typha angustifolia 25
- Vaucheria* 169
vegetation communities 86–90
- waves 6, 43–47, 63
Wentlooge Formation 127–128, 129, 135, 141
wildfowling 80
wind/wave climate influences 6, 43–47, 136
- zonation 63–64, 169
Zostera 105, 135, 169